

***Crenosoma vulpis* in dog: first case report in Italy and use of the FLOTAC technique for copromicroscopic diagnosis**

L. Rinaldi · G. Calabria · S. Carbone · A. Carrella · G. Cringoli

Published online: 10 January 2008
© Springer-Verlag 2008

**Erratum to: Parasitol Res (2007) 101:1681-1684
DOI 10.007/s00436-006-0713-6**

The original version of this article unfortunately contained a mistake. The presentation of Table 1 was incorrect. The corrected table is given below.

Table 1 LPG values (mean, standard error, and percentiles) of *Crenosoma vulpis* detected by the five copromicroscopic techniques

Copromicroscopic techniques	Multiplication factor	LPG (no.=10 replicates)				
		Mean*	SE	Percentiles		
				25 th	50 th	75 th
FLOTAC	2	91.3 ^a	5.1	82.0	88.0	104.0
McMaster	10	36.7 ^b	6.1	27.5	30.0	52.5
Baermann	0.1	0.7 ^c	0.05	0.6	0.7	0.8
Flotation	1.6	11.1 ^c	0.5	10.0	10.8	12.8
Wisconsin	1.6	36.7 ^b	6.7	31.6	36.7	43.3

*Significant differences for different letters ($P<0.05$).

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00436-007-0713-6>.

L. Rinaldi · S. Carbone · G. Cringoli
Dipartimento di Patologia e Sanità Animale,
Università degli Studi di Napoli "Federico II"-CREMOPAR
Regione Campania,
Naples, Italy

G. Cringoli (✉)
Dipartimento di Patologia e Sanità Animale,
Facoltà di Medicina Veterinaria,
Università degli Studi di Napoli "Federico II",
Via della Veterinaria,
I.80137 Naples, Italy
e-mail: cringoli@unina.it

Present address:
G. Calabria · A. Carrella
Marigliano,
Naples, Italy