


Article

Fractional Reverse Copson's Inequalities via Conformable Calculus on Time Scales

Mohammed Zakarya ^{1,2,*}, Mohamed Altanji ¹, Ghada AlNemer ³ , Hoda A. Abd El-Hamid ⁴, Clemente Cesarano ⁵ and Haytham M. Rezk ⁶

¹ Department of Mathematics, College of Science, King Khalid University, P.O. Box 9004, Abha 61413, Saudi Arabia; maltenji@kku.edu.sa

² Department of Mathematics, Faculty of Science, Al-Azhar University, Assiut 71524, Egypt

³ Department of Mathematical Science, College of Science, Princess Nourah bint Abdulrahman University, P.O. Box 105862, Riyadh 11656, Saudi Arabia; gnnemer@pnu.edu.sa

⁴ Department of Mathematics and Computer Science, Faculty of Science, Beni-Suef University, Beni-Suef 62511, Egypt; Hoda.Ali@science.bsu.edu.eg

⁵ Section of Mathematics, Università Telematica Internazionale Uninettuno, 00186 Rome, Italy; c.cesarano@uninettunouniversity.net

⁶ Department of Mathematics, Faculty of Science, Al-Azhar University, Nasr City 11884, Egypt; haythamrezk@azhar.edu.eg

* Correspondence: mzibrahim@kku.edu.sa

Abstract: This paper provides novel generalizations by considering the generalized conformable fractional integrals for reverse Copson's type inequalities on time scales. The main results will be proved using a general algebraic inequality, chain rule, Hölder's inequality, and integration by parts on fractional time scales. Our investigations unify and extend some continuous inequalities and their corresponding discrete analogues. In addition, when $\alpha = 1$, we obtain some well-known time scale inequalities due to Hardy, Copson, Bennett, and Leindler inequalities.

Keywords: Copson's inequality; Hölder's inequality; time scales; conformable fractional calculus

MSC: 26A15; 26D10; 39A13; 34A40; 34N05



Citation: Zakarya, M.; Altanji, M.; AlNemer, G.; Abd El-Hamid, H.A.; Cesarano, C.; M. Rezk, H. Fractional Reverse Copson's Inequalities via Conformable Calculus on Time Scales. *Symmetry* **2021**, *13*, 542. <https://doi.org/10.3390/sym13040542>

Academic Editors: Ioan Raşa and Wei-Shih Du

Received: 2 March 2021

Accepted: 23 March 2021

Published: 25 March 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The Hardy discrete inequality is known as (see [1]):

$$\sum_{l=1}^{\infty} \left(\frac{1}{l} \sum_{j=1}^l w(j) \right)^h \leq \left(\frac{h}{h-1} \right)^h \sum_{l=1}^{\infty} w^h(l), \quad h > 1, \quad (1)$$

where $w(l) > 0$ for all $l \geq 1$.

In [2], Hardy exemplified the continuous version of (1) by utilizing the calculus of variations, which has the form:

$$\int_0^{\infty} \left(\frac{1}{y} \int_0^y g(s) ds \right)^h dy \leq \left(\frac{h}{h-1} \right)^h \int_0^{\infty} g^h(y) dy, \quad h > 1, \quad (2)$$

where $g \geq 0$, which is integrable over $(0, y)$, g^h is a convergent and integrable function over $(0, \infty)$ and $(h/(h-1))^h$ is a sharp constant in (1) and (2).

In [3,4], Copson outreached the inequalities of Hardy (1) and (2). Particularly, he exemplified that if $h > 1$, $w(j) \geq 0$, $k(j) \geq 0$, $\forall j \geq 1$, $\vartheta(l) = \sum_{j=1}^l k(j)$ and $m > 1$, then:

$$\sum_{l=1}^{\infty} \frac{k(l)}{\vartheta^m(l)} \left(\sum_{j=1}^l w(j)k(j) \right)^h \leq \left(\frac{h}{m-1} \right)^h \sum_{l=1}^{\infty} k(l) \vartheta^{h-m}(l) w^h(l), \quad (3)$$

and if $h > 1$ and $0 \leq m < 1$, then

$$\sum_{l=1}^{\infty} \frac{k(l)}{\vartheta^m(l)} \left(\sum_{j=1}^{\infty} w(j)k(j) \right)^h \leq \left(\frac{h}{1-m} \right)^h \sum_{l=1}^{\infty} k(l) \vartheta^{h-m}(l) w^h(l). \quad (4)$$

The continuous transcriptions of (3) and (4) were exemplified by Copson in [4]. Particularly, he exemplified that if $h \geq 1, m > 1$ and $\vartheta(s) = \int_0^s k(\zeta) d\zeta$, then:

$$\int_0^{\infty} \frac{k(s)}{\vartheta^m(s)} \Theta^h(s) ds \leq \left(\frac{h}{m-1} \right)^h \int_0^{\infty} \frac{k(s)}{\vartheta^{m-h}(s)} g^h(s) ds, \quad (5)$$

where $\Theta(s) = \int_0^{\infty} k(\zeta) g(\zeta) d\zeta$ and if $h > 1, 0 \leq m < 1$, then

$$\int_0^{\infty} \frac{k(s)}{\vartheta^m(s)} \bar{\Theta}^h(s) ds \leq \left(\frac{h}{1-m} \right)^h \int_0^{\infty} \frac{k(s)}{\vartheta^{m-h}(s)} g^h(s) ds, \quad (6)$$

where $\bar{\Theta}(s) = \int_s^{\infty} k(\zeta) g(\zeta) d\zeta$.

Leindler in [5] and Bennett in [6] obtains some generalizations of (3) and (4) by using new weighted function. Specially, Leindler exemplified that if $\vartheta^*(l) = \sum_{j=1}^{\infty} k(j) < \infty, h > 1$ and $0 \leq m < 1$, then:

$$\sum_{l=1}^{\infty} \frac{k(l)}{(\vartheta^*(l))^m} \left(\sum_{j=1}^{\infty} w(j)k(j) \right)^h \leq \left(\frac{h}{1-m} \right)^h \sum_{l=1}^{\infty} k(l) (\vartheta^*(l))^{h-m} w^h(l). \quad (7)$$

Bennett explicated that if $1 < m \leq h$, then:

$$\sum_{l=1}^{\infty} \frac{k(l)}{(\vartheta^*(l))^m} \left(\sum_{j=1}^{\infty} w(j)k(j) \right)^h \leq \left(\frac{h}{m-1} \right)^h \sum_{l=1}^{\infty} k(l) (\vartheta^*(l))^{h-m} w^h(l). \quad (8)$$

Bennett in [7,8] established a converses of the inequalities (3) and (4). Particularly, he exemplified that if $m \leq 0 < h < 1$, then:

$$\sum_{l=1}^{\infty} \frac{k(l)}{\vartheta^m(l)} \left(\sum_{j=1}^{\infty} w(j)k(j) \right)^h \geq \left(\frac{h}{1-m} \right)^h \sum_{l=1}^{\infty} k(l) \left(\sum_{j=1}^l k(j) \right)^{h-m} w^h(l), \quad (9)$$

and if $m > 1 > h > 0, \vartheta(l) \rightarrow \infty$, then

$$\sum_{l=1}^{\infty} \frac{k(l)}{\vartheta^m(l)} \left(\sum_{j=1}^l w(j)k(j) \right)^h \geq \left(\frac{hL}{m-1} \right)^h \sum_{l=1}^{\infty} k(l) \vartheta^{h-m}(l) w^h(l), \quad (10)$$

where

$$L := \inf_{l \in \mathbb{N}} \frac{k(l)}{k(l+1)}.$$

In the last decades, the study of the dynamic equations and inequalities on time scales became a main field in applied and pure mathematics. We refer to the papers [9–15]. In fact, Refs. [16–24] mentions forms of the above inequalities on a time-scale and their extensions.

For example, in [25], Saker et al. exemplified the time scale version of a converse of the inequalities (7) and (8), respectively, as follows:

Assume that \mathbb{T} be a time scale with $w \in (0, \infty)_{\mathbb{T}}$. If $m \leq 0 < h < 1, \vartheta(\zeta) = \int_{\zeta}^{\infty} k(s) \Delta s$ and $\Omega(\zeta) = \int_w^{\zeta} k(s) \eta(s) \Delta s$, then

$$\int_w^{\infty} \frac{k(\zeta)}{\vartheta^m(\zeta)} (\Omega^{\sigma}(\zeta))^h \Delta \zeta \geq \left(\frac{h}{1-m} \right)^h \int_w^{\infty} k(\zeta) \eta^h(\zeta) \vartheta^{h-m}(\zeta) \Delta \zeta. \quad (11)$$

If $0 < h < 1 < m$, $\vartheta(\zeta) = \int_{\zeta}^{\infty} k(s)\Delta s$ and $\bar{\Omega}(\zeta) = \int_{\zeta}^{\infty} k(s)\eta(s)\Delta s$, then

$$\int_w^{\infty} \frac{k(\zeta)}{\vartheta^m(\zeta)} (\bar{\Omega}(\zeta))^h \Delta\zeta \geq \left(\frac{hM^m}{m-1}\right)^h \int_w^{\infty} k(\zeta)\eta^h(\zeta)\vartheta^{h-m}(\zeta)\Delta\zeta, \tag{12}$$

where

$$M := \inf_{\zeta \in \mathbb{T}} \frac{\vartheta^\sigma(\zeta)}{\vartheta(\zeta)} > 0.$$

In the same paper [25], Saker et al. proved the time scale transcript of the Bennet-Leindler inequalities (9) and (10), respectively, as follows: Assume that \mathbb{T} is a time scale with $w \in (0, \infty)_{\mathbb{T}}$. If $m \leq 0 < h < 1$, $\Gamma(\zeta) = \int_w^{\zeta} k(s)\Delta s$ and $\bar{\Omega}(\zeta) = \int_{\zeta}^{\infty} k(s)\eta(s)\Delta s$, then:

$$\int_w^{\infty} \frac{k(\zeta)}{(\Gamma^\sigma(\zeta))^m} (\bar{\Omega}(\zeta))^h \Delta\zeta \geq \left(\frac{h}{1-m}\right)^h \int_w^{\infty} k(\zeta)\eta^h(\zeta) (\Gamma^\sigma(\zeta))^{h-m} \Delta\zeta. \tag{13}$$

If $0 < h \leq m$, $\Gamma(\zeta) = \int_w^{\zeta} k(s)\Delta s$, such that

$$L := \inf_{\zeta \in \mathbb{T}} \frac{\Gamma(\zeta)}{\Gamma^\sigma(\zeta)} > 0,$$

and $\bar{\Phi}(\zeta) = \int_w^{\zeta} k(s)\eta(s)\Delta s$, then

$$\int_w^{\infty} \frac{k(\zeta)}{(\Gamma^\sigma(\zeta))^m} (\bar{\Phi}^\sigma(\zeta))^h \Delta\zeta \geq \left(\frac{hL^{1-m}}{1-m}\right)^h \int_w^{\infty} k(\zeta)\eta^h(\zeta) (\Gamma^\sigma(\zeta))^{h-m} \Delta\zeta. \tag{14}$$

In recent years, a lot of work has been published on fractional inequalities and the subject has become an active field of research with several authors interested in proving the inequalities of fractional type by using the Riemann-Liouville and Caputo derivative (see [26–28]).

On the other hand, the authors in [29,30] introduced a new fractional calculus called the conformable calculus and gave a new definition of the derivative with the base properties of the calculus based on the new definition of derivative and integrals.

The main question that arises now is: Is it possible to prove new fractional inequalities on timescales and give a unified approach of such studies? This in fact needs a new fractional calculus on timescales. Very recently Torres and others, in [31,32], combined a time scale calculus and conformable calculus and obtained the new fractional calculus on timescales. Thus, it is natural to look on new fractional inequalities on timescales and give an affirmative answer to the above question.

In particular, in this paper, we will prove the fractional forms of the classical Hardy, Copson type and its reversed and Leindler inequalities with employing conformable calculus on time scales. The article is structured as follows: Section 2 is an introduction of the basics of fractional calculus on timescales and Section 3 contains the main results.

2. Basic Concepts

In this part, we introduce the essentials of conformable fractional integral and derivative of order $\alpha \in [0, 1]$ on time scales that will be used in this article (see [33–35]). A time scale \mathbb{T} is an arbitrary nonempty closed subset of the real numbers \mathbb{R} . We define the operator $\sigma : \mathbb{T} \rightarrow \mathbb{T}$, as $\sigma(\theta) := \inf \{s \in \mathbb{T} : s > \theta\}$. In addition, we define the function $\mu : \mathbb{T} \rightarrow [0, \infty)$ by $\mu(\theta) := \sigma(\theta) - \theta$. Finally, for any $\theta \in \mathbb{T}$, we refer to the notation $\zeta^\sigma(\theta)$ by $\zeta(\sigma(\theta))$, i.e., $\zeta^\sigma = \zeta \circ \sigma$.

In the following, we define conformable α -fractional derivative and α -fractional integral on \mathbb{T} .

Definition 1 (Definition 1, [31]). Suppose that $\zeta : \mathbb{T} \rightarrow \mathbb{R}$ and $\alpha \in (0, 1]$. Then for $\theta > 0$, we define $D_\alpha^\Delta(\zeta)(\theta)$ to be the number with the property that, for any $\varepsilon > 0$, there is a neighborhood V of θ s.t. $\forall \theta \in V$, we have:

$$\left| [\zeta^\sigma(\theta) - \zeta(s)]\sigma^{1-\alpha(\theta)} - D_\alpha^\Delta(\zeta)(\theta)[\sigma(\theta) - s] \right| \leq \varepsilon|\sigma(\theta) - s|.$$

The conformable α -fractional derivative on \mathbb{T} at 0 as:

$$D_\alpha^\Delta(\zeta(0)) = \lim_{\theta \rightarrow 0} D_\alpha^\Delta(\zeta(\theta)).$$

Theorem 1 (Theorem 51, [31]). Assume $\alpha \in (0, 1]$ and $v, \zeta : \mathbb{T} \rightarrow \mathbb{R}$ be conformable α -fractional derivative on \mathbb{T} , then

(i) The $v + \zeta : \mathbb{T} \rightarrow \mathbb{R}$ is conformable α -fractional derivative and

$$D_\alpha^\Delta(v + \zeta) = D_\alpha^\Delta(v) + D_\alpha^\Delta(\zeta).$$

(ii) For $\lambda \in \mathbb{R}$, then $\lambda v : \mathbb{T} \rightarrow \mathbb{R}$ α -fractional differentiable and

$$D_\alpha^\Delta(\lambda v) = \lambda D_\alpha^\Delta(v).$$

(iii) If v and ζ are α -fractional differentiable, then $v\zeta : \mathbb{T} \rightarrow \mathbb{R}$ is a α -fractional differentiable and

$$D_\alpha^\Delta(v\zeta) = D_\alpha^\Delta(v)\zeta + (v \circ \sigma)D_\alpha^\Delta(\zeta) = D_\alpha^\Delta(v)(\zeta \circ \sigma) + vD_\alpha^\Delta(\zeta). \tag{15}$$

(iv) If v is α -fractional differentiable, then $1/v$ is α -fractional differentiable with:

$$D_\alpha^\Delta\left(\frac{1}{v}\right) = -\frac{D_\alpha^\Delta(v)}{v(v \circ \sigma)}.$$

(v) If v and ζ are α -fractional differentiable, then v/ζ is α -fractional differentiable with:

$$D_\alpha^\Delta\left(\frac{v}{\zeta}\right) = \frac{\zeta D_\alpha^\Delta(v) - vD_\alpha^\Delta(\zeta)}{\zeta(\zeta \circ \sigma)}, \tag{16}$$

valid at all points $\theta \in \mathbb{T}^k$ for which $\zeta(\theta)(\zeta(\sigma(\theta))) \neq 0$.

Lemma 1 (Chain rule [32]). Suppose that $\zeta : \mathbb{T} \rightarrow \mathbb{R}$ is continuous and α -fractional differentiable at $\theta \in \mathbb{T}$, for $\alpha \in (0, 1]$ and $v : \mathbb{R} \rightarrow \mathbb{R}$ is continuously differentiable. Then $(v \circ \zeta) : \mathbb{T} \rightarrow \mathbb{R}$ is α -fractional differentiable and

$$D_\alpha^\Delta(v \circ \zeta)(\theta) = v'(\zeta(d))D_\alpha^\Delta(\zeta(\theta)), \text{ where } d \in [\theta, \sigma(\theta)]. \tag{17}$$

Definition 2 (Definition 26, [31]). For $0 < \alpha \leq 1$, then the α -fractional integral of ζ , is defined as

$$I_\alpha^\Delta(\zeta(s)) = \int \zeta(s)\Delta_\alpha s = \int \zeta(s)s^{\alpha-1}\Delta s.$$

Theorem 2 (Theorem 31, [31]). Suppose that $l, m, n \in \mathbb{T}$ and $\lambda \in \mathbb{R}$. If $v, \zeta : \mathbb{T} \rightarrow \mathbb{R}$, then

- (vi) $\int_l^m [v(s) + \zeta(s)] \Delta_\alpha s = \int_l^m v(s) \Delta_\alpha s + \int_l^m \zeta(s) \Delta_\alpha s.$
- (vii) $\int_l^m \lambda v(s) \Delta_\alpha s = \lambda \int_l^m v(s) \Delta_\alpha s.$
- (viii) $\int_l^m v(s) \Delta_\alpha s = - \int_l^m v(s) \Delta_\alpha s.$

- (ix) $\int_l^m v(s)\Delta_\alpha s = \int_l^n v(s)\Delta_\alpha s + \int_n^m v(s)\Delta_\alpha s.$
- (x) $\int_l^l v(s)\Delta_\alpha s = 0$

Lemma 2 (Integration by parts formula [31]). Suppose that $l, m \in \mathbb{T}$ where $m > l$. If v, ζ are conformable α -fractional differentiable and $\alpha \in (0, 1]$, then:

$$\int_l^m v(s) D_\alpha^\Delta(\zeta(s))\Delta_\alpha s = [v(s)\zeta(s)]_l^m - \int_l^m \zeta^\sigma(s) D_\alpha^\Delta(v(s))\Delta_\alpha s. \tag{18}$$

Lemma 3 (Hölder’s inequality [32]). Let $l, m \in \mathbb{T}$ where $m > l$. If $\alpha \in (0, 1]$ and $F, G : \mathbb{T} \rightarrow \mathbb{R}$, then

$$\int_l^m |F(s)G(s)| \Delta_\alpha s \leq \left(\int_l^m |F(s)|^\lambda \Delta_\alpha s \right)^{\frac{1}{\lambda}} \left(\int_l^m |G(s)|^\mu \Delta_\alpha s \right)^{\frac{1}{\mu}}, \tag{19}$$

where $\lambda > 1$ and $1/\lambda + 1/\mu = 1$.

Through our paper, we will consider the integrals are given exist (are finite i.e., convergent).

3. Results

Here, we will exemplify our main results in this article by utilizing Hölder’s inequality, chain rule, and integration by parts for fractional on time scale.

Theorem 3. Suppose that \mathbb{T} is a time scale with $w \in (0, \infty)_{\mathbb{T}}, k \leq 0 < h < 1$ and $\alpha \in (0, 1]$. Define

$$\vartheta(y) = \int_y^\infty x(s)\Delta_\alpha s \text{ and } \Omega(y) = \int_w^y x(s)\eta(s)\Delta_\alpha s.$$

Then

$$\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\Omega^\sigma(y))^h \Delta_\alpha y \geq \left(\frac{h}{\alpha - k} \right)^h \int_w^\infty x(y)\eta^h(y)\vartheta^{h-k+\alpha-1}(y)\Delta_\alpha y. \tag{20}$$

Proof. By utilizing the formula of integration by parts (18) on

$$\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\Omega^\sigma(y))^h \Delta_\alpha y,$$

with $\zeta^\sigma(y) = (\Omega^\sigma(y))^h$ and $D_\alpha^\Delta(v(y)) = \frac{x(y)}{\vartheta^{k-\alpha+1}(y)}$, we have

$$\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\Omega^\sigma(y))^h \Delta_\alpha y = v(y)\Omega^h(y) \Big|_w^\infty + \int_w^\infty (-v(y))D_\alpha^\Delta(\Omega^h(y))\Delta_\alpha y, \tag{21}$$

where

$$v(y) = - \int_y^\infty \frac{x(s)}{\vartheta^{k-\alpha+1}(s)} \Delta_\alpha s.$$

Using $\Omega(w) = 0$ and $v(\infty) = 0$ in (21), we see that

$$\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\Omega^\sigma(y))^h \Delta_\alpha y = - \int_w^\infty v(y)D_\alpha^\Delta(\Omega^h(y))\Delta_\alpha y. \tag{22}$$

By utilizing chain rule, we get:

$$D_\alpha^\Delta(\Omega^h(y)) = h\Omega^{h-1}(d)D_\alpha^\Delta(\Omega(y)) = \frac{hD_\alpha^\Delta(\Omega(y))}{(\Omega(d))^{1-h}} \geq \frac{hD_\alpha^\Delta(\Omega(y))}{(\Omega^\sigma(y))^{1-h}}.$$

Since $D_\alpha^\Delta(\Omega(y)) = x(y)\eta(y)$, we have

$$D_\alpha^\Delta(\Omega^h(y)) \geq \frac{hx(y)\eta(y)}{(\Omega^\sigma(y))^{1-h}}. \tag{23}$$

Next note $D_\alpha^\Delta\vartheta(y) = -x(y) \leq 0$. By the chain rule, we have (note $k \leq 0$)

$$\begin{aligned} D_\alpha^\Delta(\vartheta^{\alpha-k}(y)) &= (\alpha - k)\vartheta^{\alpha-k-1}(d)D_\alpha^\Delta\vartheta(y) \\ &= \frac{\alpha-k}{\vartheta^{k+1-\alpha}(d)}D_\alpha^\Delta\vartheta(y) \geq \frac{\alpha-k}{\vartheta^{k+1-\alpha}(y)}D_\alpha^\Delta\vartheta(y) \\ &= \frac{\alpha-k}{\vartheta^{k+1-\alpha}(y)}(-x(y)) \geq \frac{-(\alpha-k)x(y)}{\vartheta^{k+1-\alpha}(y)}. \end{aligned}$$

This leads to

$$\vartheta^{-k+\alpha-1}(y)x(y) \geq \frac{-1}{\alpha - k}D_\alpha^\Delta(\vartheta^{\alpha-k}(y)),$$

and then, we have

$$-v(y) = \int_y^\infty \frac{x(s)}{\vartheta^{k-\alpha+1}(y)}\Delta_\alpha s \geq \frac{-1}{\alpha - k} \int_y^\infty D_\alpha^\Delta(\vartheta^{\alpha-k}(s))\Delta_\alpha s = \frac{1}{(\alpha - k)\vartheta^{k-\alpha}(y)} \tag{24}$$

Substituting (23), (24) into (22) yields:

$$\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)}(\Omega^\sigma(y))^h \Delta_\alpha y \geq \left(\frac{h}{\alpha - k}\right) \int_w^\infty \frac{\eta(y)x(y)}{\vartheta^{k-\alpha}(y)(\Omega^\sigma(y))^{1-h}} \Delta_\alpha y. \tag{25}$$

Raises (25) to the factor h , we have:

$$\left(\int_y^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)}(\Omega^\sigma(y))^h \Delta_\alpha y\right)^h \geq \left(\frac{h}{\alpha - k}\right)^h \left(\int_w^\infty \left(\frac{\eta^h(y)x^h(y)}{\vartheta^{h(k-\alpha)}(y)(\Omega^\sigma(y))^{h(1-h)}}\right)^{\frac{1}{h}} \Delta_\alpha y\right)^h. \tag{26}$$

By applying Hölder’s inequality (19) on the term

$$\left(\int_w^\infty \left(\frac{\eta^h(y)x^h(y)}{\vartheta^{h(k-\alpha)}(y)(\Omega^\sigma(y))^{h(1-h)}}\right)^{\frac{1}{h}} \Delta_\alpha y\right)^h,$$

with indices $\lambda = 1/h > 1, \mu = 1/(1 - h)$ (note that $\frac{1}{\lambda} + \frac{1}{\mu} = 1$, where $\lambda > 1$) and

$$F(y) = \frac{\eta^h(y)x^h(y)}{\vartheta^{h(k-\alpha)}(y)(\Omega^\sigma(y))^{h(1-h)}} \text{ and } G(y) = \left(\frac{x(y)}{\vartheta^{k-\alpha+1}(y)}\right)^{1-h} (\Omega^\sigma(y))^{h(1-h)},$$

we see that

$$\begin{aligned} \left(\int_w^\infty F^{\frac{1}{h}}(y) \Delta_\alpha y \right)^h &= \left(\int_w^\infty \left(\frac{\eta^h(y)x^h(y)}{\vartheta^{h(k-\alpha)}(y)(\Omega^\sigma(y))^{h(1-h)}} \right)^{\frac{1}{h}} \Delta_\alpha y \right)^h \\ &\geq \frac{\int_w^\infty F(y)G(y)\Delta_\alpha y}{\left(\int_w^\infty G^{\frac{1}{1-h}}(y)\Delta_\alpha y \right)^{1-h}} \\ &= \left(\int_w^\infty \left(\frac{\eta^h(y)(x(y)\vartheta^{-k+\alpha-1}(y))^{1-h}x^h(y)\Omega^\sigma(y)^{h(1-h)}}{\vartheta^{h(k-\alpha)}(y)(\Omega^\sigma(y))^{h(1-h)}} \right) \Delta_\alpha y \right) \\ &\times \left(\int_w^\infty \frac{x(y)(\Omega^\sigma(y))^h}{\vartheta^{k-\alpha+1}(y)} \Delta_\alpha y \right)^{h-1} \\ &= \left(\int_w^\infty \left(\frac{\eta^h(y)x(y)}{\vartheta^{h(k-\alpha)}(y)(\vartheta^{k-\alpha+1}(y))^{h-1}} \right) \Delta_\alpha y \right) \\ &\times \left(\int_w^\infty \frac{x(y)(\Omega^\sigma(y))^h}{\vartheta^{k-\alpha+1}(y)} \Delta_\alpha y \right)^{h-1}. \end{aligned}$$

This means that

$$\int_w^\infty \left(\frac{\eta^h(y)x^h(y)}{\vartheta^{h(k-\alpha)}(y)(\Omega^\sigma(y))^{h(1-h)}} \right)^{\frac{1}{h}} \Delta_\alpha y \geq \frac{\int_w^\infty \eta^h(y)x(y)\vartheta^{h-k+\alpha-1}(y)\Delta_\alpha y}{\left(\int_w^\infty \frac{x(y)(\Omega^\sigma(y))^h}{\vartheta^{k-\alpha+1}(y)} \Delta_\alpha y \right)^{1-h}}, \tag{27}$$

by substitution (27) into (26), we get

$$\left(\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\Omega^\sigma(y))^h \Delta_\alpha y \right)^h \geq \left(\frac{h}{\alpha-k} \right)^h \frac{\int_w^\infty \eta^h(y)x(y)\vartheta^{h-k+\alpha-1}(y)\Delta_\alpha y}{\left(\int_w^\infty \frac{x(y)(\Omega^\sigma(y))^h}{\vartheta^{k-\alpha+1}(y)} \Delta_\alpha y \right)^{1-h}}.$$

This means that

$$\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\Omega^\sigma(y))^h \Delta_\alpha y \geq \left(\frac{h}{\alpha-k} \right)^h \int_w^\infty x(y)\eta^h(y)\vartheta^{h-k+\alpha-1}(y)\Delta_\alpha y,$$

which the wanted inequality (20). \square

Corollary 1. *If we put $\alpha = 1$ in Theorem 6, then we get*

$$\int_w^\infty \frac{x(y)}{\vartheta^k(y)} (\Omega^\sigma(y))^h \Delta y \geq \left(\frac{h}{1-k} \right)^h \int_w^\infty x(y)\eta^h(y)\vartheta^{h-k}(y)\Delta(y), \tag{28}$$

where $w \in (0, \infty)_{\mathbb{T}}, k \leq 0 < h < 1$,

$$\vartheta(y) = \int_v^\infty x(s)\Delta s \text{ and } \Omega(y) = \int_w^y x(s)\eta(s)\Delta s,$$

which is (11) in the Introduction.

Remark 1. *If we take $\mathbb{T} = \mathbb{R}$ in Theorem 6, then:*

$$\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\Omega(y))^h y^{\alpha-1} dy \geq \left(\frac{h}{\alpha-k} \right)^h \int_w^\infty x(y)\eta^h(y)\vartheta^{h-k+\alpha-1}(y) y^{\alpha-1} dy, \tag{29}$$

where $w \in (0, \infty), k \leq 0 < h < 1, \vartheta(y) = \int_y^\infty x(s)s^{\alpha-1} ds$ and $\Omega(y) = \int_w^y x(s)\eta(s)s^{\alpha-1} ds$.

Remark 2. *Clearly, for $\alpha = 1$ and $w = 1$, Remark 1 coincides with Remark 1 in [25].*

Remark 3. As a result, if $\mathbb{T} = \mathbb{Z}$ in (20), and $k \leq 0 < h < 1$, then:

$$\sum_{y=w}^{\infty} \frac{x(w)}{\vartheta^{k-\alpha+1}(w)} (\Omega^\sigma(y))^h y^{\alpha-1} (y+1)^{\alpha-1} \geq \left(\frac{h}{\alpha-k}\right)^h \sum_{y=w}^{\infty} x(w) \vartheta^{h-k+\alpha-1}(y) \eta^h(w) (y+1)^{\alpha-1}, \tag{30}$$

where, $\Omega^\sigma(y) = \Omega(y+1) = \sum_{s=w}^y x(s) \eta(s) (s+1)^{\alpha-1}$ and $\vartheta(y) = \sum_{s=y}^{\infty} x(s) (s+1)^{\alpha-1}$.
 If $\alpha = 1$, then (30) becomes

$$\sum_{y=w}^{\infty} \frac{x(w)}{\vartheta^k(w)} (\Omega^\sigma(y))^h \geq \left(\frac{h}{1-k}\right)^h \sum_{y=w}^{\infty} x(w) \vartheta^{h-k}(y) \eta^h(w), \tag{31}$$

where, $\Omega^\sigma(y) = \Omega(y+1) = \sum_{s=w}^y x(s) \eta(s)$ and $\vartheta(y) = \sum_{s=y}^{\infty} x(s)$. which is Remark 2 in [25].

Theorem 4. Suppose that \mathbb{T} be a time scale with $w \in (0, \infty)_{\mathbb{T}}$, $0 < h < 1 < k$ and $\alpha \in (0, 1]$. Assume that $\vartheta(y)$ is defined as in Theorem 6 such that:

$$M := \inf_{y \in \mathbb{T}} \frac{\vartheta^\sigma(y)}{\vartheta(y)} > 0, \tag{32}$$

and define $\bar{\Omega}(y) = \int_y^\infty x(s) \eta(s) \Delta_\alpha s$. Then

$$\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\bar{\Omega}(y))^h \Delta_\alpha y \geq \left(\frac{hM^{k-\alpha+1}}{k-\alpha}\right)^h \int_w^\infty x(y) \eta^h(y) \vartheta^{h-k+\alpha-1}(y) \Delta_\alpha y. \tag{33}$$

Proof. Utilizing the formula of integration by parts (18) on

$$\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\bar{\Omega}(y))^h \Delta_\alpha y,$$

with $(y) = (\bar{\Omega}(y))^h$ and $D_\alpha^\Delta \zeta(y) = \frac{x(y)}{\vartheta^{k-\alpha+1}(y)}$, we have

$$\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\bar{\Omega}(y))^h \Delta_\alpha y = \zeta(y) (\bar{\Omega}(y))^h \Big|_w^\infty + \int_w^\infty \zeta^\sigma(y) \left(-D_\alpha^\Delta (\bar{\Omega}(y))^h\right) \Delta_\alpha y,$$

where $\zeta(y) = \int_w^y \frac{x(s)}{\vartheta^{k-\alpha+1}(s)} \Delta_\alpha s$. This with $\bar{\Omega}(\infty) = 0$ and $\zeta(w) = 0$ implies that

$$\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\bar{\Omega}(y))^h \Delta_\alpha y = \int_w^\infty \zeta^\sigma(y) \left(-D_\alpha^\Delta (\bar{\Omega}(y))^h\right) \Delta_\alpha y. \tag{34}$$

But utilizing chain rule, we obtain:

$$-D_\alpha^\Delta (\bar{\Omega}(y))^h = -h \bar{\Omega}^{h-1}(d) D_\alpha^\Delta \bar{\Omega}(y) = \frac{hx(y)\eta(y)}{(\bar{\Omega}(d))^{1-h}} \geq \frac{hx(y)\eta(y)}{(\bar{\Omega}(y))^{1-h}}. \tag{35}$$

Since $D_\alpha^\Delta \bar{\Omega}(y) = -x(y)\eta(y) \leq 0$ and $d \geq y$, we find that $\bar{\Omega}(y) \geq \bar{\Omega}(d)$. By substituting (35) into (34) and using that $D_\alpha^\Delta \zeta(y) \geq 0$, we get

$$\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\bar{\Omega}(y))^h \Delta_\alpha y \geq h \int_w^\infty \zeta(y) \frac{x(y)\eta(y)}{(\bar{\Omega}(y))^{1-h}} \Delta_\alpha y. \tag{36}$$

Next note $D_\alpha^\Delta \vartheta(y) = -x(y) \leq 0$. By the chain rule, we have (note $k \leq 0$)

$$\begin{aligned} D_\alpha^\Delta \left(\vartheta^{\alpha-k}(y) \right) &= (\alpha - k) \vartheta^{\alpha-k-1}(d) D_\alpha^\Delta \vartheta(y) \\ &= \frac{\alpha-k}{\vartheta^{k+1-\alpha}(d)} D_\alpha^\Delta \vartheta(y) \leq \frac{\alpha-k}{\vartheta^{k+1-\alpha}(y)} D_\alpha^\Delta \vartheta(y) \\ &\leq \frac{k-\alpha}{\vartheta^{k+1-\alpha}(d)} x(y) \leq \frac{k-\alpha}{(\vartheta^\sigma(y))^{k+1-\alpha}} x(y) \\ &= (k - \alpha) \frac{x(y)}{(\vartheta^\sigma(y))^{k+1-\alpha}} \frac{\vartheta^{k+1-\alpha}(y)}{\vartheta^{k+1-\alpha}(y)} \\ &\leq \frac{(k-\alpha)x(y)}{M^{k+1-\alpha}(\vartheta(y))^{k+1-\alpha}}. \end{aligned}$$

This implies that

$$\zeta(y) = \int_w^y \frac{x(x)}{\vartheta^{k+1-\alpha}(s)} \Delta_\alpha s \geq \left(\frac{M^{k+1-\alpha}}{k - \alpha} \right) \int_w^y D_\alpha^\Delta \left(\vartheta^{\alpha-k}(s) \right) \Delta_\alpha s = \left(\frac{M^{k+1-\alpha}}{k - \alpha} \right) \vartheta^{\alpha-k}(y). \tag{37}$$

By substituting (37) into (36) yields

$$\int_w^\infty \frac{x(y)}{\vartheta^{k+1-\alpha}(y)} (\overline{\Omega}(y))^h \Delta_\alpha s \geq \left(\frac{hM^{k+1-\alpha}}{k - \alpha} \right) \int_w^\infty \frac{x(y)\eta(y)}{\vartheta^{\alpha-k}(y)\overline{\Omega}^{(1-h)}(y)} \Delta_\alpha y. \tag{38}$$

Raising (38) to the factor h , we get:

$$\left(\int_w^\infty \frac{x(y)}{\vartheta^{k+1-\alpha}(y)} (\overline{\Omega}(y))^h \Delta_\alpha y \right)^h \geq \left(\frac{hM^{k+1-\alpha}}{k - \alpha} \right)^h \left(\int_w^\infty \left(\frac{x^h(y)\eta^h(y)}{\vartheta^{h(k-\alpha)}(y)\overline{\Omega}^{h(1-h)}(y)} \right)^{\frac{1}{h}} \Delta_\alpha y \right)^h. \tag{39}$$

The rest of the proof is identical to the proof of Theorem 6 and hence is deleted. \square

Corollary 2. *If we put $\alpha = 1$ in Theorem 8, then:*

$$\int_w^\infty \frac{x(y)}{\vartheta^k(y)} (\overline{\Omega}(y))^h \Delta y \geq \left(\frac{hM^k}{k-1} \right)^h \int_w^\infty x(y)\eta^h(y)\vartheta^{h-k}(y)\Delta(y), \tag{40}$$

where $w \in (0, \infty)_{\mathbb{T}}$, $0 < h < 1 < k$, $\overline{\Omega}(y) = \int_y^\infty x(s)\eta(s)\Delta s$ and $\vartheta(y) = \int_y^\infty x(s)\Delta s$ such that

$$M := \inf_{y \in \mathbb{T}} \frac{\vartheta^\sigma(y)}{\vartheta(y)} > 0,$$

which is (12) in the Introduction

Remark 4. *If we take $\mathbb{T} = \mathbb{R}$ ($i, e, \sigma(y) = y$) in Theorem 8, then*

$$\int_w^\infty \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\overline{\Omega}(y))^h y^{\alpha-1} dy \geq \left(\frac{h}{k - \alpha} \right)^h \int_w^\infty x(y)\eta^h(y)\vartheta^{h-k+\alpha-1}(y)y^{\alpha-1} dy, \tag{41}$$

where $0 < h < 1 < k$, $\alpha \in (0, 1]$, $\vartheta(y) = \int_y^\infty x(s)s^{\alpha-1} ds$, $\overline{\Omega}(y) = \int_y^\infty x(s)\eta(s)s^{\alpha-1} ds$ and

$$M := \inf_{y \in \mathbb{T}} \frac{\vartheta^\sigma(y)}{\vartheta(y)} = \inf_{y \in \mathbb{R}} \frac{\vartheta(y)}{\vartheta(y)} = 1.$$

If $\alpha = 1$ and $w = 1$, then (41) becomes:

$$\int_1^\infty \frac{x(y)}{\vartheta^k(y)} (\overline{\Omega}(y))^h dy \geq \left(\frac{h}{k-1} \right)^h \int_1^\infty x(y)\eta^h(y)\vartheta^{h-k}(y)dy, \tag{42}$$

which is Remark 3 in [25].

Remark 5. As a special case of (33), when $\mathbb{T} = \mathbb{Z}$ (i.e. $\sigma(y) = y + 1$) and $0 < h < 1 < k$, we get:

$$\sum_{y=w}^{\infty} \frac{x(y)}{\vartheta^{k-\alpha+1}(y)} (\overline{\Omega}(y))^h \geq \left(\frac{hM^k}{k-\alpha}\right)^h \sum_{y=w}^{\infty} x(y)\eta^h(y)\vartheta^{h-k+\alpha-1}(y), \tag{43}$$

where $\vartheta(y) = \sum_{s=y}^{\infty} x(s) (s + 1)^{\alpha-1}$, and $\overline{\Omega}(y) = \sum_{s=w}^{\infty} x(s)\eta(s) (s + 1)^{\alpha-1}$.

$$M := \inf_{y \in \mathbb{T}} \frac{\vartheta^\sigma(y)}{\vartheta(y)} = \inf_{y \in \mathbb{Z}} \frac{\vartheta(l+1)}{\vartheta(l)} > 0,$$

which is Remark 4 in [25], when $\alpha = 1$.

Theorem 5. Suppose that \mathbb{T} is a time scale with $w \in (0, \infty)_{\mathbb{T}}$, $k \leq 0 < h < 1$ and $\alpha \in (0, 1]$. Assume that $\Gamma(y) = \int_w^y x(s)\Delta_\alpha s$ and $\overline{\Omega}(y) = \int_w^y x(s)\eta(s)\Delta_\alpha s$. Then

$$\int_w^\infty \frac{x(y)}{(\Gamma^\sigma(y))^{k-\alpha+1}} (\overline{\Omega}(y))^h \Delta_\alpha y \geq \left(\frac{h}{\alpha-k}\right)^h \int_w^\infty x(y)\eta^h(y)(\Gamma^\sigma(y))^{h-k+\alpha-1} \Delta_\alpha y. \tag{44}$$

Proof. Utilizing the formula of integration by parts (18) on

$$\int_w^\infty \frac{x(y)}{(\Gamma^\sigma(y))^{k-\alpha+1}} (\overline{\Omega}(y))^h \Delta_\alpha y,$$

with $v(y) = (\overline{\Omega}(y))^h$ and $D_\alpha^\Delta \zeta(y) = \frac{x(y)}{(\Gamma^\sigma(y))^{k-\alpha+1}}$, we get

$$\int_w^\infty \frac{x(y)}{(\Gamma^\sigma(y))^{k-\alpha+1}} (\overline{\Omega}(y))^h \Delta_\alpha y = \zeta(y)(\overline{\Omega}(y))^h \Big|_w^\infty + \int_w^\infty \zeta^\sigma(y) (-D_\alpha^\Delta (\overline{\Omega}(y))^h) \Delta_\alpha y,$$

where $\zeta(y) = \int_w^y \frac{x(s)}{(\Gamma^\sigma(s))^{k-\alpha+1}} \Delta_\alpha s$. This with $\overline{\Omega}(\infty) = 0$ and $\Gamma(w) = 0$ imply that

$$\int_w^\infty \frac{x(y)}{(\Gamma^\sigma(y))^{k-\alpha+1}} (\overline{\Omega}(y))^h \Delta_\alpha y = \int_w^\infty \zeta^\sigma(y) (-D_\alpha^\Delta (\overline{\Omega}(y))^h) \Delta_\alpha y. \tag{45}$$

By utilizing chain rule, we get:

$$-D_\alpha^\Delta (\overline{\Omega}(y))^h = -h\overline{\Omega}^{h-1}(d)D_\alpha^\Delta \overline{\Omega}(y) = \frac{hx(y)\eta(y)}{\overline{\Omega}^{1-h}(d)} \geq \frac{hx(y)\eta(y)}{\overline{\Omega}^{1-h}(y)}. \tag{46}$$

Since $D_\alpha^\Delta \overline{\Omega}(y) = -x(y)\eta(y) \leq 0$. By substituting (46) into (45) and using that $D_\alpha^\Delta \zeta(y) \geq 0$, we have:

$$\int_w^\infty \frac{x(y)}{(\Gamma^\sigma(y))^{k-\alpha+1}} (\overline{\Omega}(y))^h \Delta_\alpha y \geq h \int_w^\infty \zeta^\sigma(y) \frac{x(y)\eta(y)}{\overline{\Omega}^{1-h}(y)} \Delta_\alpha y. \tag{47}$$

Next note $D_\alpha^\Delta \Gamma(y) = x(y) \geq 0$. By the chain rule, we have (note $k \leq 0$)

$$\begin{aligned} D_\alpha^\Delta (\Gamma^{\alpha-k}(y)) &= (\alpha-k)\Gamma^{\alpha-k-1}(d)D_\alpha^\Delta \Gamma(y) \\ &= \frac{\alpha-k}{\Gamma^{k+1-\alpha}(d)} D_\alpha^\Delta \Gamma(y) \leq \frac{\alpha-k}{\Gamma^{k+1-\alpha}(y)} D_\alpha^\Delta \Gamma(y) \\ &\leq \frac{\alpha-k}{\Gamma^{k+1-\alpha}(y)} x(y) \leq \frac{\alpha-k}{(\Gamma^\sigma(y))^{k+1-\alpha}} x(y). \end{aligned}$$

This implies that

$$\zeta^\sigma(y) = \int_w^{\sigma(y)} \frac{x(s)}{(\Gamma^\sigma(s))^{k-\alpha+1}} \Delta_\alpha s \geq \frac{1}{\alpha-k} \int_w^{\sigma(y)} D_\alpha^\Delta (\Gamma(s))^{\alpha-k} \Delta_\alpha s = \frac{1}{\alpha-k} (\Gamma^\sigma(y))^{\alpha-k}. \tag{48}$$

By substituting (48) into (47) yields

$$\int_w^\infty \frac{x(y)}{(\Gamma^\sigma(y))^{k-\alpha+1}} (\overline{\Omega}(y))^h \Delta_\alpha y \geq \frac{h}{\alpha-k} \int_w^\infty (\Gamma^\sigma(y))^{\alpha-k} \frac{x(y)\eta(y)}{\overline{\Omega}^{1-h}(y)} \Delta_\alpha y. \tag{49}$$

Raises (49) to the factor h , we get:

$$\left(\int_w^\infty \frac{x(y)}{(\Gamma^\sigma(y))^{k-\alpha+1}} (\overline{\Omega}(y))^h \Delta_\alpha y \right)^h \geq \left(\frac{h}{\alpha-k} \right)^h \left(\int_w^\infty \left((\Gamma^\sigma(y))^{h(\alpha-k)} \frac{x^h(y)\eta^h(y)}{\overline{\Omega}^{h(1-h)}(y)} \right)^{\frac{1}{h}} \Delta_\alpha y \right)^h. \tag{50}$$

The rest of the proof is identical to the proof of Theorem 6 and hence is deleted. \square

Corollary 3. *If we put $\alpha = 1$ in Theorem 10, then:*

$$\int_w^\infty \frac{x(y)}{(\Gamma^\sigma(y))^k} (\overline{\Omega}(y))^h \Delta y \geq \left(\frac{h}{1-k} \right)^h \int_w^\infty x(y)\eta^h(y) (\Gamma^\sigma(y))^{h-k} \Delta y, \tag{51}$$

where $w \in (0, \infty)_{\mathbb{T}}, k \leq 0 < h < 1$,

$$\Gamma(y) = \int_w^y x(s) \Delta s \text{ and } \overline{\Omega}(y) = \int_y^\infty x(s)\eta(s) \Delta s,$$

which is (13) in the Introduction.

Remark 6. *In Theorem 10, if we take $\mathbb{T} = \mathbb{R}$ (i.e. $\sigma(y) = y$), then:*

$$\int_w^\infty \frac{x(y)}{(\Gamma(y))^{k-\alpha+1}} (\overline{\Omega}(y))^h y^{\alpha-1} dy \geq \left(\frac{h}{\alpha-k} \right)^h \int_w^\infty x(y)\eta^h(y) (\Gamma(y))^{h-k+\alpha-1} y^{\alpha-1} dy, \tag{52}$$

where $w \in (0, \infty), k \leq 0 < h < 1, \alpha \in (0, 1]$

$$\Gamma(y) = \int_w^y x(s) s^{\alpha-1} ds \text{ and } \overline{\Omega}(y) = \int_y^\infty x(s)\eta(s) s^{\alpha-1} ds.$$

If $\alpha = 1$ and $w = 1$, then (52) becomes

$$\int_w^\infty \frac{x(y)}{(\Gamma(y))^k} (\overline{\Omega}(y))^h dy \geq \left(\frac{h}{1-k} \right)^h \int_w^\infty x(y)\eta^h(y) (\Gamma(y))^{h-k} dy, \tag{53}$$

where

$$\Gamma(y) = \int_1^y x(s) ds \text{ and } \overline{\Omega}(y) = \int_y^\infty x(s)\eta(s) ds,$$

which is Remark 5 in [25]

Remark 7. *As a special case of (44) when $\mathbb{T} = \mathbb{Z}$ (i.e. $\sigma(y) = y + 1$), we get:*

$$\sum_{y=w}^\infty \frac{x(y)}{(\Gamma^\sigma(y))^{k-\alpha+1}} (\overline{\Omega}(y))^h (y+1)^{\alpha-1} \geq \left(\frac{h}{\alpha-k} \right)^h \sum_{y=w}^\infty x(y)\eta^h(y) (\Gamma^\sigma(y))^{h-k+\alpha-1} (y+1)^{\alpha-1}, \tag{54}$$

where $\Gamma^\sigma(y) = \Gamma(y+1) = \sum_{s=w}^y x(s) (s+1)^{\alpha-1}, \overline{\Omega}(y) = \sum_{s=y}^\infty x(s)\eta(s) (s+1)^{\alpha-1}, k \leq 0 < h < 1$.

For $\alpha = 1$ in (54), then we get the inequality in Remark 6 in [21].

Theorem 6. Suppose that \mathbb{T} is a time scale with $w \in (0, \infty)_{\mathbb{T}}, 0 < h \leq 1 < k$ and $\alpha \in (0, 1]$. Assume that $\Gamma(y) = \int_w^y x(s) \Delta_{\alpha} s$ such that

$$L := \inf_{y \in \mathbb{T}} \frac{\Gamma(y)}{\Gamma^{\sigma}(y)} > 0, \tag{55}$$

and define $\bar{\Phi}(y) = \int_w^y x(s) \eta(s) \Delta_{\alpha} s$

$$\int_w^{\infty} \frac{x(y)}{(\Gamma^{\sigma}(y))^{k-\alpha+1}} (\bar{\Phi}(\sigma(y)))^h \Delta_{\alpha} y \geq \left(\frac{hL^{\alpha-k}}{k-\alpha} \right)^h \int_w^{\infty} x(y) \eta^h(y) (\Gamma^{\sigma}(y))^{h-k+\alpha-1} \Delta_{\alpha} y. \tag{56}$$

Proof. Utilizing the formula of integration by parts (18) on

$$\int_w^{\infty} \frac{x(y)}{(\Gamma^{\sigma}(y))^{k-\alpha+1}} (\bar{\Phi}(\sigma(y)))^h \Delta_{\alpha} y,$$

with $\zeta^{\sigma}(y) = (\bar{\Phi}(\sigma(y)))^h$ and $D_{\alpha}^{\Delta} v(y) = \frac{x(y)}{(\Gamma^{\sigma}(y))^{k-\alpha+1}}$, we have

$$\int_w^{\infty} \frac{x(y)}{(\Gamma^{\sigma}(y))^{k-\alpha+1}} (\bar{\Phi}(\sigma(y)))^h \Delta_{\alpha} y = v(y) (\bar{\Phi}(y))^h \Big|_w^{\infty} + \int_w^{\infty} (-v(y)) (D_{\alpha}^{\Delta} (\bar{\Phi}(y))^h) \Delta_{\alpha} y,$$

where $v(y) = \int_w^y \frac{x(s)}{(\Gamma^{\sigma}(s))^{k-\alpha+1}} \Delta_{\alpha} s$. This with $\bar{\Phi}(\infty) = 0$ and $\Gamma(w) = 0$ imply that

$$\int_w^{\infty} \frac{x(y)}{(\Gamma^{\sigma}(y))^{k-\alpha+1}} (\bar{\Phi}(\sigma(y)))^h \Delta_{\alpha} y = \int_w^{\infty} (-v(y)) (D_{\alpha}^{\Delta} (\bar{\Phi}(y))^h) \Delta_{\alpha} y. \tag{57}$$

By utilizing chain rule, we obtain:

$$D_{\alpha}^{\Delta} (\bar{\Phi}(y))^h = h (\bar{\Phi}(y))^{h-1} (d) D_{\alpha}^{\Delta} \bar{\Phi}(y) = \frac{hx(y)\eta(y)}{\bar{\Phi}^{1-h}(d)} \geq \frac{hx(y)\eta(y)}{(\bar{\Phi}(\sigma(y)))^{1-h}}. \tag{58}$$

Since $D_{\alpha}^{\Delta} \bar{\Phi}(y) = x(y)\eta(y) \geq 0$. By substituting (58) into (57) and using that $D_{\alpha}^{\Delta} \bar{\Phi}(y) \geq 0$, we have:

$$\int_w^{\infty} \frac{x(y)}{(\Gamma^{\sigma}(y))^{k-\alpha+1}} (\bar{\Phi}(\sigma(y)))^h \Delta_{\alpha} y \geq h \int_w^{\infty} (-v(y)) \frac{x(y)\eta(y)}{(\bar{\Phi}(\sigma(y)))^{1-h}} \Delta_{\alpha} y. \tag{59}$$

Next note $D_{\alpha}^{\Delta} \Gamma(y) = x(y) \geq 0$. By the chain rule, we have (note $k \leq 0$)

$$\begin{aligned} D_{\alpha}^{\Delta} (\Gamma^{\alpha-k}(y)) &= (\alpha - k) \Gamma^{\alpha-k-1} (d) D_{\alpha}^{\Delta} \Gamma(y) \\ &= \frac{\alpha-k}{\Gamma^{k+1-\alpha}(d)} D_{\alpha}^{\Delta} \Gamma(y) \leq \frac{\alpha-k}{\Gamma^{k+1-\alpha}(y)} D_{\alpha}^{\Delta} \Gamma(y) \\ &\leq \frac{\alpha-k}{\Gamma^{k+1-\alpha}(y)} x(y) \leq \frac{\alpha-k}{(\Gamma^{\sigma}(y))^{k+1-\alpha}} x(y), \end{aligned}$$

And

$$\begin{aligned} -v(y) &= - \int_w^y \frac{x(s)}{(\Gamma^{\sigma}(s))^{k-\alpha+1}} \Delta_{\alpha} s \\ &\geq \frac{1}{k-\alpha} \int_w^y D_{\alpha}^{\Delta} (\Gamma^{\alpha-k}(s)) \Delta_{\alpha} s \\ &= \left(\frac{1}{k-\alpha} \right) - \left(\frac{\Gamma(y)}{\Gamma^{\sigma}(y)} \right)^{\alpha-k} (\Gamma^{\sigma}(y))^{\alpha-k} \\ &\geq \left(\frac{L^{\alpha-k}}{k-\alpha} \right) (\Gamma^{\sigma}(y))^{\alpha-k}, \end{aligned} \tag{60}$$

by substituting (60) into (59), we get

$$\int_w^\infty \frac{x(y)}{(\Gamma^\sigma(y))^{k-\alpha+1}} (\overline{\Phi}(\sigma(y)))^h \Delta_\alpha y \geq \left(\frac{hL^{\alpha-k}}{k-\alpha}\right) \int_w^\infty (\Gamma^\sigma(y))^{\alpha-k} \frac{x(y)\eta(y)}{(\overline{\Phi}(\sigma(y)))^{1-h}} \Delta_\alpha y. \tag{61}$$

Raises (61) to the factor h , we get

$$\left(\int_w^\infty \frac{x(y)}{(\Gamma^\sigma(y))^{k-\alpha+1}} (\overline{\Phi}(\sigma(y)))^h \Delta_\alpha y\right)^h \geq \left(\frac{hL^{\alpha-k}}{k-\alpha}\right)^h \left(\int_w^\infty \left(\Gamma^\sigma(y)\right)^{h(\alpha-k)} \frac{x^h(y)\eta^h(y)}{(\overline{\Phi}^\sigma(y))^{h(1-h)}} \Delta_\alpha y\right)^{\frac{1}{h}}.$$

The rest of the proof is identical to the proof Theorem 6 and hence is deleted. \square

Corollary 4. *If we put $\alpha = 1$ in Theorem 12, then:*

$$\int_w^\infty \frac{x(y)}{(\Gamma^\sigma(y))^k} (\overline{\Phi}(\sigma(y)))^h \Delta y \geq \left(\frac{hL^{\alpha-k}}{k-\alpha}\right)^h \int_w^\infty x(y)\eta^h(y)(\Gamma^\sigma(y))^{h-k} \Delta y, \tag{62}$$

where $w \in (0, \infty)_{\mathbb{T}}, 0 < h \leq 1 < k, \overline{\Phi}(y) = \int_w^y x(s)\eta(s)\Delta s$ and $\Gamma(y) = \int_w^y x(s)\Delta_\alpha s$ such that

$$L := \inf_{y \in \mathbb{T}} \frac{\Gamma(y)}{\Gamma^\sigma(y)} > 0,$$

which is (14) in the Introduction

Remark 8. *If we take $\mathbb{T} = \mathbb{R}$ (i.e., $\sigma(y) = y$) in Theorem 12, then:*

$$\int_w^\infty \frac{x(y)}{(\Gamma(y))^{k-\alpha+1}} (\overline{\Phi}(y))^h y^{\alpha-1} dy \geq \left(\frac{h}{k-\alpha}\right)^h \int_w^\infty x(y)\eta^h(y)(\Gamma(y))^{h-k+\alpha-1} y^{\alpha-1} dy, \tag{63}$$

where $w \in (0, \infty)_{\mathbb{T}}, 0 < h \leq 1 < k, \overline{\Phi}(y) = \int_w^y x(s)\eta(s)s^{\alpha-1} ds$ and $\Gamma(y) = \int_w^y x(s)s^{\alpha-1} ds$ such that

$$L := \inf_{y \in \mathbb{T}} \frac{\Gamma(y)}{\Gamma^\sigma(y)} = \inf_{y \in \mathbb{R}} \frac{\Gamma(y)}{\Gamma(y)} = 1.$$

If $\alpha = 1$ and $w = 1$, then (63) becomes

$$\int_1^\infty \frac{x(y)}{(\Gamma(y))^k} (\overline{\Phi}(y))^h dy \geq \left(\frac{h}{k-1}\right)^h \int_1^\infty x(y)\eta^h(y)(\Gamma(y))^{h-k} dy, \tag{64}$$

where

$$\Gamma(y) = \int_1^y x(s)ds \quad \text{and} \quad \overline{\Phi}(y) = \int_1^y x(s)\eta(s)ds,$$

which is Remark 7 in [25]

Remark 9. *As a special case of (56), when $\mathbb{T} = \mathbb{Z}$ (i.e. $\sigma(y) = y + 1$) and $0 < h \leq 1 < k$, we get:*

$$\sum_{w=y}^\infty \frac{x(y)}{(\Gamma^\sigma(y))^{k-\alpha+1}} (\overline{\Phi}(\sigma(y)))^h (y+1)^{\alpha-1} \geq \left(\frac{hL^{\alpha-k}}{k-\alpha}\right)^h \sum_{w=y}^\infty x(y)\eta^h(y)(\Gamma^\sigma(y))^{h-k+\alpha-1} (y+1)^{\alpha-1}, \tag{65}$$

where, $\Gamma^\sigma(y) = \Gamma(y+1) = \sum_{s=w}^y x(s) (s+1)^{\alpha-1}, \overline{\Phi}(\sigma(y)) = \overline{\Phi}(y+1) = \sum_{s=w}^y x(s)\eta(s)(s+1)^{\alpha-1}.$

$$L := \inf_{y \in \mathbb{T}} \frac{\Gamma(y)}{\Gamma^\sigma(y)} = \inf_{y \in \mathbb{Z}} \frac{\Gamma(y)}{\Gamma(y+1)},$$

which is Remark 8 in [25], when $\alpha = 1$.

Applications

The applications of quantum calculus play an important role in mathematics and the field of natural sciences, such as physics and chemistry. It has many applications in orthogonal polynomials, number theory, quantum theory, etc. In this section, some example for Reverse Copson’s Inequalities in fractional quantum calculus are selected to fulfil the applicability of the obtained results.

Now, we give an example using the time scale $\mathbb{T} = q^{\mathbb{N}_0}, q > 1$ which is a time scale with interesting applications in quantum calculus.

Example 1. (Quantum calculus case 1.): Let $\mathbb{T} = q^{\mathbb{N}_0} = \{t : t = q^n, n \in \mathbb{N}_0, q > 1\}$. Then for all $t \in q^{\mathbb{N}_0}$, we have

$$\sigma(t) = qt, \mu(t) = (q - 1)t \text{ and } \int_a^b f(t)\Delta_\alpha t = \sum_{n=\log_q a}^{\log_q b-1} f(q^n)\mu(q^n)(q^n)^{\alpha-1}, \forall a, b \in q^{\mathbb{N}_0} \tag{66}$$

Now, with the help of Theorem 3 and the above identities in (66), we can deduce

$$\sum_{n=\log_q w}^\infty \frac{x(q^n)}{\vartheta^{k-\alpha+1}(q^n)} (\Omega^\sigma(q^n))^h \mu(q^n)(q^n)^{\alpha-1} \geq \left(\frac{h}{\alpha - k}\right)^h \sum_{n=\log_q w}^\infty x(q^n)\eta^h(q^n)(\vartheta(q^n))^{h-k+\alpha-1} \mu(q^n)(q^n)^{\alpha-1}.$$

where,

$$\Omega^\sigma(q^n) = \Omega(\sigma(q^n)) = \int_w^{\sigma(q^n)} x(s)\eta(s)\Delta_\alpha s = \sum_{m=\log_q w}^{\log_q \sigma(q^n)-1} x(q^m)\eta(q^m)\mu(q^m)(q^m)^{\alpha-1},$$

and

$$\vartheta(q^n) = \int_{q^n}^\infty x(s)\Delta_\alpha s = \sum_{m=\log_q q^n}^\infty x(q^m)\mu(q^m)(q^m)^{\alpha-1}$$

For an application of Theorem 4, we give the following example.

Example 2. (Quantum calculus case 2.): Let $\mathbb{T} = q^{\mathbb{N}_0}, q > 1$, then the relation (66) is satisfy. Hence, we have:

$$\sigma(t) = qt, \mu(t) = (q - 1)t \text{ and } \int_a^b f(t)\Delta_\alpha t = \sum_{n=\log_q a}^{\log_q b-1} f(q^n)\mu(q^n)(q^n)^{\alpha-1}, \forall a, b \in q^{\mathbb{N}_0}$$

Now, with the help of Theorem 4 and the above identities in (66), we can deduce:

$$\sum_{n=\log_q w}^\infty \frac{x(q^n)}{\vartheta^{k-\alpha+1}(q^n)} (\bar{\Omega}(q^n))^h \mu(q^n)(q^n)^{\alpha-1} \geq \left(\frac{hM^{k-\alpha+1}}{k - \alpha}\right)^h \sum_{n=\log_q w}^\infty x(q^n)\eta^h(q^n)(\vartheta(q^n))^{h-k+\alpha-1} \mu(q^n)(q^n)^{\alpha-1}.$$

where, $\bar{\Omega}(q^n) = \int_{q^n}^\infty x(s)\eta(s)\Delta_\alpha s = \sum_{m=\log_q q^n}^\infty x(q^m)\eta(q^m)\mu(q^m)(q^m)^{\alpha-1}$. $M := \inf_{q^n \in \mathbb{T}} \frac{\vartheta^\sigma(q^n)}{\vartheta(q^n)} > 0$, $\vartheta(q^n)$ is defined in the above example.

Note that. By using theorems 10 and 12, we can apply the technique used in the above examples to obtain different applications. In addition, the above result is important not only for arbitrary time scales, but also for quantum calculus.

4. Conclusions and Future Work

The new fractional calculus on timescales is presented with applications in new fractional inequalities on timescales like Hardy, Bennett, Copson, and Leindler types. In-

equalities are considered in rather general forms and contain several special integral and discrete inequalities. The technique is based on the applications of well-known inequalities and new tools from fractional calculus. In future research, we will continue to generalize more dynamic inequalities by using Specht's ratio, Kantorovich's ratio, functional generalization, and n-tuple fractional diamond- α integral. It will be interesting to find the inequalities in α, β -symmetric quantum and stochastic calculus.

Author Contributions: M.Z. and H.M.R. contributed by preparing the introduction, preliminaries, and formulating (theorem 6, theorem 8) its proof and corollaries (cor.7, cor. 9 and cor.10). M.A. contributed by preparing the introduction, preliminaries, and formulating (theorem 10) its proof and corollaries (cor.11). G.A. contributed by preparing the introduction, preliminaries, and formulating theorem 12, its proof and corollaries (cor.7). C.C. and H.A.A.E.-H. contributed by preparing the introduction, preliminaries, and formulating theorem 6, its proof and corollaries (cor. 13). All authors contributed equally to the writing of the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding: The authors extend their appreciation to the Deanship of Scientific Research at King Khalid University for funding this work through the General Research Project, grant number GRP/56/42.

Data Availability Statement: No data were used to support this study.

Acknowledgments: The authors extend their appreciation to the Deanship of Scientific Research at King Khalid University for funding this work through the General Research Project, grant number GRP/56/42.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Abd El-Hamid, H.A.; Rezk, H.M.; Ahmed, A.M.; AlNemer, G.; Zakarya, M.; El Saify, H.A. Dynamic inequalities in quotients with general kernels and measures. *J. Funct. Spaces* **2020**, *2020*, 5417084. [[CrossRef](#)]
2. Ahmed, A.M.; AlNemer, G.; Zakarya, M.; Rezk, H.M. Some dynamic inequalities of Hilbert's Type. *J. Funct. Spaces* **2020**, *2020*, 4976050. [[CrossRef](#)]
3. Ahmed, A.M.; Saker, S.H.; Kenawy, M.R.; Rezk, H.M. Lower bounds on a generalization of Gesaro operator on time scale. *J. DCDIS Ser. A* **2020**. accepted.
4. AlNemer, G.; Zakarya, M.; Abd El-Hamid, H.A.; Agrwal, P.; Rezk, H.M. Some dynamic Hilbert's-type, inequalities on time scales. *Symmetry* **2020**, *12*, 1410. [[CrossRef](#)]
5. AlNemer, G.; Zakarya, M.; El-Hamid, H.A.; Kenawy, M.; Rezk, H. Dynamic Hardy-type inequalities with non-conjugate parameters. *Alex. Eng. J.* **2020**, *59*, 4523–4532. [[CrossRef](#)]
6. Benkhattou, N.; Hassani, S.; Torres, D.F. A conformable fractional calculus on arbitrary time scales. *J. King Saud Univ. Sci.* **2016**, *28*, 93–98. [[CrossRef](#)]
7. Bennett, G. Some elementary inequalities. *Q. J. Math.* **1987**, *38*, 401–425. [[CrossRef](#)]
8. Bemmatt, G. Some elementary inequalities II. *Q. J. Math.* **1988**, *39*, 385–400. [[CrossRef](#)]
9. Bohner, M.; Peterson, A. *Dynamic Equations on Time Scales: An Introduction with Applications*; Birkhause: Boston, MA, USA, 2001.
10. Bohner, M.; Peterson, A. *Advances in Dynamic Equation on Time Scales*; Birkhause: Boston, MA, USA, 2003.
11. Copson, E.T. Note on series of positive terms. *J. Lond. Math. Soc.* **1928**, *3*, 49–51. [[CrossRef](#)]
12. Copson, E.T. Some integral inequalities. *Prof. R. Soc. Edinburg Sect. A Math.* **1976**, *75*, 157–164. [[CrossRef](#)]
13. Hardy, G.H. Notes on a theorem of Hilbert. *Math. Z.* **1920**, *6*, 314–317. [[CrossRef](#)]
14. Hardy, G.H. Notes on some points in the integral calculus, LX. An inequality between integrals. *Mass. Math.* **1925**, *54*, 150–156.
15. Leindler, L. Generalization of inequalities of Hardy and Littlewood. *Acta Sci. Math.* **1970**, *31*, 285–297.
16. Leindler, L. Some inequalities pertaining to Bennetts results. *Acta Sci. Math.* **1993**, *58*, 261–269.
17. Nwaeze, E.R.; Torres, D.F.M. Chain rules and inequalities for the BHT fractional calculus on arbitrary timescales. *Arab. J. Math.* **2017**, *6*, 13–20. [[CrossRef](#)]
18. O'Regan, D.; Rezk, H.M.; Saker, S.H. Some dynamic inequalities involving Hilbert and Hardy-Hilbert operators with kernels. *Results Math.* **2018**, *73*, 146. [[CrossRef](#)]
19. Rezk, H.M.; El-Hamid, H.A.A.; Ahmed, A.M.; AlNemer, G.; Zakarya, M. Inequalities of Hardy type via superquadratic functions with general kernels and measures for several variables on time scales. *J. Funct. Spaces* **2020**, *2020*, 6427378. [[CrossRef](#)]
20. Rezk, H.M.; AlNemer, G.; El-Hamid, H.A.A.; Abdel-Aty, A.-H.; Nisar, K.S.; Zakarya, M. Hilbert-type inequalities for time scale nabla calculus. *Adv. Differ. Equ.* **2020**, *2020*, 3079. [[CrossRef](#)]
21. Saker, S.H.; O'Regan, D.; Agarwal, R.P. Converses of Copson's inequalities on time scales. *J. Math. Inequal. Appl.* **2015**, *18*, 241–254. [[CrossRef](#)]

22. Saker, S.H.; Ahmed, A.M.; Rezk, H.M.; O'Regan, D.; Agarwal, R.P. New Hilbert's dynamic inequalities on time scale. *J. Math. Inequal. Appl.* **2017**, *20*, 1017–1039.
23. Saker, S.H.; El-Deeb, A.A.; Rezk, H.M.; Agarwal, R.P. On Hilbert's inequality on time scales. *Appl. Anal. Discret. Math.* **2017**, *11*, 399–423. [[CrossRef](#)]
24. Saker, S.H.; Rezk, H.M.; O'Regan, D.; Agarwal, R.P. A variety of inverse Hilbert type inequality on time scales. *J. DCDIS Ser. A* **2017**, *24*, 347–373.
25. Saker, S.H.; Rabie, S.; AlNemer, G.; Zakarya, M. On structure of discrete Muchenhaupt and discrete Gehring classes. *J. Inequal. Appl.* **2020**, *233*, 1–18. [[CrossRef](#)]
26. Saker, S.H.; Sayed, A.; AlNemer, G.; Zakarya, M. Half-linear dynamic equations and investigating weighted Hardy and Copson inequalities. *Adv. Differ. Equ.* **2020**, *519*, 1–19. [[CrossRef](#)]
27. Saker, S.H.; Rezk, H.M.; Krnić, M. More accurate dynamic Hardy-type inequalities obtained via superquadraticity. *Rev. Real Acad. Cienc. Exactas Fís. Nat. Ser. A Mat.* **2019**, *113*, 2691–2713. [[CrossRef](#)]
28. Saker, S.H.; Rezk, H.M.; Abohela, I.; Baleanu, D. Refinement multidimensional dynamic inequalities with general kernels and measures. *J. Inequal. Appl.* **2019**, *2019*, 1–16. [[CrossRef](#)]
29. Saker, S.; Kenawy, M.; AlNemer, G.; Zakarya, M. Some fractional dynamic inequalities of Hardy's type via conformable calculus. *Mathematics* **2020**, *8*, 434. [[CrossRef](#)]
30. Zakarya, M.; Abdd El-Hamid, H.A.; AlNemer, G.; Rezk, H.M. More on Hölder's inequality and it's reverse via the diamond-alpha integral. *Symmetry* **2020**, *12*, 1716. [[CrossRef](#)]
31. Bogdan, K.; Dyda, B. The best constant in a fractional Hardy inequality. *Math. Nachr.* **2011**, *284*, 629–638. [[CrossRef](#)]
32. Jleli, M.; Samet, B. Lyapunov-type inequalities for a fractional differential equation with mixed boundary conditions. *Math. Inequal. Appl.* **2015**, *18*, 443–451. [[CrossRef](#)]
33. Yildiz, C.; Ozdemir, M.E.; Onelan, H.K. Fractional integral inequalities for different functions. *New Trends Math. Sci.* **2015**, *3*, 110–117.
34. Abdeljawad, T. On conformable fractional calculus. *J. Comput. Appl. Math.* **2015**, *279*, 57–66. [[CrossRef](#)]
35. Khalil, R.; Al Horani, M.; Yousef, A.; Sababheh, M. A new definition of fractional derivative. *J. Comput. Appl. Math.* **2014**, *264*, 65–70. [[CrossRef](#)]