

Erratum to: Effects of Global Postural Reeducation on gait kinematics in parkinsonian patients: a pilot randomized three-dimensional motion analysis study

Valeria Agosti^{1,2} · Carmine Vitale^{1,2} · Dario Avella^{1,2} · Rosaria Rucco^{2,5} ·
Gabriella Santangelo³ · Pierpaolo Sorrentino⁴ · Pasquale Varriale^{1,2} ·
Giuseppe Sorrentino^{1,2}

Published online: 4 February 2016
© Springer-Verlag Italia 2016

Erratum to: Neurol Sci
DOI 10.1007/s10072-015-2433-5

Unfortunately, the Fig. 3 was published incorrectly in the original publication. The correct figure is given below:

The online version of the original article can be found under
doi:[10.1007/s10072-015-2433-5](https://doi.org/10.1007/s10072-015-2433-5).

✉ Giuseppe Sorrentino
giuseppe.sorrentino@uniparthenope.it

Carmine Vitale
cavit69@hotmail.com

¹ Department of Motor Sciences and Wellness, University of Naples Parthenope, via Medina 40, 80133 Naples, Italy

² Institute Hermitage-Capodimonte, Naples, Italy

³ Department of Psychology, Second University of Naples, Caserta, Italy

⁴ Department of Neurosciences, Reproductive and Odontostomatological Sciences, University of Naples Federico II, Naples, Italy

⁵ Department of Sciences and Technologies, University of Naples Parthenope, Naples, Italy




Fig. 3 The effect of 4 weeks GPR program on the RoM of the thigh (T), knee (K), and ankle (A) joints on the sagittal plane in PD vs. control group at different times. The RoMs for each joint were normalized for the 100 % of the gait cycle. Δ s value represents the

difference between two consecutive peaks in the gait cycle. Details in the text. *Statistical significance as compared to baseline evaluation (pre-test) $p < 0.005$