

Alpha-enhancement in the MW: results from the SDSS spectroscopic stellar database

M. Franchini¹, C. Morossi¹, P. Di Marcantonio¹,
M. L. Malagnini^{1,2} and M. Chávez^{3,4}

¹INAF - Osservatorio Astronomico di Trieste, Via G.B. Tiepolo, 11, I-34131 Trieste, Italy
email: franchini@oats.inaf.it morossi@oats.inaf.it dimarcan@oats.inaf.it

²Dip. di Astronomia, Univ. degli Studi di Trieste, Via G.B. Tiepolo, 11, I-34131 Trieste, Italy
email: malagnini@oats.inaf.it

³Instituto Nacional de Astrofisica, Optica y Electronica, A.P. 51 y 216, 72000 Puebla, Mexico

⁴IAM-Universidad de Guadalajara, Av. Vallarta 2602, C.P. 44130 Guadalajara, Mexico

Abstract. We analyzed a sample of about 2500 stars extracted from the Sloan Digital Sky Survey with T_{eff} in the range 4750–6500 K and $\log g$ greater than 1.5. Atmospheric parameter estimates are obtained by comparing observed and synthetic spectral indices. The dependence of the α -enhancement phenomenon on stellar metallicity and on Galactic position is investigated.

Keywords. Galaxy: stellar content, stars: abundances, stars: late-type

1. The observational data-set

The observational data-set was extracted from SDSS-DR5 (Adelman-McCarthy *et al.* 2007) which contains 154,925 spectra classified as STARS. Since we are interested in finding G and early K stars belonging to the Galactic disks and halo, we selected from the total set of spectra classified as STARS those with $0.7 < g - r < 1.3$ mag and $(g + r)/2 < 21$ mag. Colors were dereddened by using the extinction map by Schlegel *et al.* (1998). A set of constraints based on SNR, accuracy of radial velocity determination, consistency of the calculated SDSS spectrophotometric color indices $(g - r)$ and $(r - i)$ with the photometric values was applied to select stars with specific quality characteristics.

2. Results and discussion

We present preliminary results for a sample of 2347 stars whose atmospheric parameter values T_{eff} , $\log g$, and $[\text{Fe}/\text{H}]$, are derived by comparing observed indices with synthetic ones. The latter are derived from an *ad hoc* library of high resolution synthetic spectra (Malagnini *et al.* 2005) computed either with Solar-Scaled Abundances (SSA grid) or α -enhanced ones (NSSA grid). The models and the synthetic spectra are at fixed $[\text{Fe}/\text{H}]$ in the range $+0.5 \div -2.5$ with $[\alpha/\text{Fe}] = +0.0$ or $+0.4$. In order to derive the atmospheric parameters we use the following set of Lick/SDSS indices which are only marginally dependent on α -element abundances: G4300, Fe4383, Ca4455, H β , Fe5335, Fe5406, Fe5709, Fe5783 and NaD (definitions in Worthey *et al.* 1994). Then, we estimate the $[\alpha/\text{Fe}]$ values by comparing the observed α -dependent Lick/SDSS indices Ca4227, Mgb, Mg₂ with the corresponding theoretical values computed by using the derived atmospheric parameters and $[\alpha/\text{Fe}] = 0$ or $[\alpha/\text{Fe}] = 0.4$. A linear interpolation was applied to derive the value of $[\alpha/\text{Fe}]$ for each star. In Franchini *et al.* (2004) we introduced four Lick/IDS index-index diagrams, (NaD vs Ca4227, NaD vs Mg₂, NaD vs Mgb, and NaD vs CaMg), which allow us to identify different loci representative of SSA and NSSA stars regardless of T_{eff} , $\log g$,




Figure 1. Mg_2 vs NaD Lick/SDSS index-index diagrams. Upper panels: theoretical predictions; lower panels: Lick/SDSS indices for stars in different ranges of $[Fe/H]$ (see text).

and $[Fe/H]$. Equivalent diagrams for NaD vs Mg_2 built with Lick/SDSS indices are shown in Figure 1. The upper panels show the positions of synthetic points computed for $\log g$ from 3.5 to 5.0, T_{eff} in the range 4750–6500 K, and $[Fe/H] = 0.0, -0.5, \text{ and } -1.0$. Indices computed with $[\alpha/Fe]=0.4$ are represented by red points while those with $[\alpha/Fe]=0.0$ are indicated by green points. The lower panels show observational indices for 3 sub-samples of our SDSS stars at different $[Fe/H]$ values. Red points correspond to stars with derived $[\alpha/Fe]>+0.3$ while green points represent stars with $[\alpha/Fe]<0.1$. The consistency of the lower panels with the predictions of the upper panels states the statistical reliability of our determinations of atmospheric parameters and $[\alpha/Fe]$. In each panel the inset plot shows the distribution of $[\alpha/Fe]$ of the corresponding stars: as can be seen, a higher percentage of stars with $[\alpha/Fe]>+0.3$ is found, as expected, at lower $[Fe/H]$ value. A similar analysis using sub-samples of our SDSS stars grouped according to different distances from the Galactic plane, $|Z_{gal}|$, shows an increase in the percentage of NSSA stars ($[\alpha/Fe]>+0.3$) with increasing $|Z_{gal}|$.

Acknowledgements

This work received partial financial support from the Mexican CONACyT via grant 36547-E, from MIUR COFIN-2003028039 and COFIN-2004025729 and from PRIN-INAF 2005 (P.I. M. Bellazzini). Funding for the creation and distribution of the SDSS Archive has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the Nat. Aeronautics and Space Administration, the Nat. Science Found., the U.S. Dep. of Energy, the Japanese Monbukagakusho, and the Max Planck Society.

References

- Adelman-McCarthy, J. K. *et al.* 2007, *ApJS* submitted
Franchini, M., Morossi, C., Di Marcantonio, *et al.* 2004, *ApJ* 601, 485
Malagnini, M. L., Franchini, M., Morossi, *et al.* 2005, *ESA SP-576* 595
Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ* 500, 525
Worthey, G., Faber, S. M., Gonzalez, J. J., & Burstein, D. 1994, *ApJ* 94, 687



F.l.t.r.: Margaret Meixner, Bengt Gustafsson, Eline Tolstoy and Annette Ferguson.