

CD16^{low/negative} Tumor-Infiltrating Lymphocyte: Lymphoid or Myeloid in Origin?

To the Editors: Although tumor-infiltrating lymphocytes (TIL) are predominantly composed of T lymphocytes, other immune cells also occur within the tumor and may be important in tumor control. We therefore read with interest the study of Schleypen et al. (1) who identified a proportion of TIL in renal cell carcinomas as natural killer (NK) cells. In some tumors, the proportion of such NK-TIL exceeded 20% (high NK-TIL), whereas others had fewer than 20% (low NK-TIL). High NK-TIL were mostly CD16^{positive}. In contrast, low NK-TIL were CD16^{low or negative}. These NK cell populations were not cytotoxic to the HLA class I antigen-negative cell line K562; they showed a diversity of CD16 expression and variable ability to become cytotoxic after stimulation. Interestingly, low NK-TIL were cytotoxic^{negative} and perforin^{dim/negative}. Our own *in vitro* studies on human NK cell development shed further light on the nature of these NK TIL cells.

We stimulated granulocyte-colony stimulating factor mobilized CD34^{positive} cells from healthy donors with stem cell factor and interleukin-2. After 3 weeks, the cultures generated a heterogeneous CD56^{positive}CD16^{positive/negative} cell population, which phenotypically and functionally resembled the low NK-TIL described in Schleypen et al. article. These cultured cells included two populations: a CD56^{bright}CD16^{negative} and a CD56^{low}CD16^{low/negative} population. They had few or no cytotoxic granules and showed negligible cytotoxicity against K562 and P815 cell lines. Interestingly, the CD56^{low} population expressed the myeloid marker CD33 (2, 3). To further investigate the myeloid origin of CD56^{low}CD33^{positive} cells, these cells were electronically sorted (Fig. 1A) and cultured in the presence of stem cell factor, interleukin-2, interleukin-4, and granulocyte macrophage colony-stimulating factor. After a 10-day culture, these cells underwent a profound morphologic change, rounding up, losing adherence, and acquiring numerous digitations, closely resembling immature dendritic cells (Fig. 1B). Because of technical limitations, we could not analyze their phenotype, but we showed that both lymphoid (CD33^{negative}; ref. 4) and myeloid (CD33^{positive}) populations (2, 3) had a strong antiproliferative effect on the myeloid cell lines K562 and P815.

Given the phenotypic and functional similarities between our *in vitro* cultured cells and low NK-TIL, we propose (a) that low NK-TIL may represent the myeloid CD56^{low} population we found in culture and (b) that NK-TIL cells can control tumor proliferation through a cytostatic effect, rather than by exerting direct cytotoxicity against the tumor. To determine whether the NK-TIL cells described by Schleypen et al. share identity with the myeloid NK cells we found in culture, further phenotypic and functional analysis (including tests of cytostatic function) on TIL cells would be worthwhile not only in renal cell carcinomas but also in other solid malignancies because infiltrating CD56^{positive} cells have been shown in infiltrating a variety of malignancies to be associated with improved survival rate (5–7).




Fig. 1. CD56⁺ myeloid cells acquired dendritic cell characteristics on interleukin-4 and GM-CSF stimulation.

Giuseppe Sconocchia

Consiglio Nazionale delle Ricerche Institute for Organ Transplantation and Immunocytology, Rome, Italy

Domenico Del Principe

Department of Pediatrics, School of Medicine, University of Tor Vergata, Rome, Italy

A. John Barrett

Allogeneic Stem Cell Transplantation, Hematology Branch, National Heart, Lung, and Blood Institute, NIH, Bethesda, Maryland

References

- Schleypen JS, Baur N, Kammerer R, et al. Cytotoxic markers and frequency predict functional capacity of natural killer cells infiltrating renal cell carcinoma. *Clin Cancer Res* 2006;12:718–25.
- Sconocchia G, Fujiwara H, Rezvani K, et al. G-CSF-mobilized CD34⁺ cells cultured in interleukin-2 and stem cell factor generate a phenotypically novel monocyte. *J Leukoc Biol* 2004;76:1214–9.
- Sconocchia G, Keyvanfar K, El OF, et al. Phenotype and function of a CD56⁺ peripheral blood monocyte. *Leukemia* 2005;19:69–76.
- Sconocchia G, Provenzano M, Rezvani K, et al. CD34⁺ cells cultured in stem cell factor and interleukin-2 generate CD56⁺ cells with antiproliferative effects on tumor cell lines. *J Transl Med* 2005;3:15.
- Coca S, Perez-Piqueras J, Martinez D, et al. The prognostic significance of intratumoral natural killer cells in patients with colorectal carcinoma. *Cancer* 1997;79:2320–8.
- Villegas FR, Coca S, Villarrubia VG, et al. Prognostic significance of tumor infiltrating natural killer cells subset CD57 in patients with squamous cell lung cancer. *Lung Cancer* 2002;35:23–8.
- Ishigami S, Natsugoe S, Tokuda K, et al. Prognostic value of intratumoral natural killer cells in gastric carcinoma. *Cancer* 2000;88:577–83.

Clinical Cancer Research

CD16^{low}/negative Tumor-Infiltrating Lymphocyte: Lymphoid or Myeloid in Origin?

Giuseppe Sconocchia, Domenico Del Principe and A. John Barrett

Clin Cancer Res 2007;13:1620.

Updated version Access the most recent version of this article at:
<http://clincancerres.aacrjournals.org/content/13/5/1620>

Cited articles This article cites 7 articles, 1 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/13/5/1620.full#ref-list-1>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/13/5/1620>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.