

1-1-2013

## The alien spreading of *Chama pacifica* Broderip, 1835 (Mollusca: Bivalvia: Chamidae) in the Mediterranean Sea

FABIO CROCETTA

PAOLO RUSSO

Follow this and additional works at: <https://dctubitak.researchcommons.org/zoology>



Part of the [Zoology Commons](#)

---

### Recommended Citation

CROCETTA, FABIO and RUSSO, PAOLO (2013) "The alien spreading of *Chama pacifica* Broderip, 1835 (Mollusca: Bivalvia: Chamidae) in the Mediterranean Sea," *Turkish Journal of Zoology*: Vol. 37: No. 1, Article 12. <https://doi.org/10.3906/zoo-1110-28>

Available at: <https://dctubitak.researchcommons.org/zoology/vol37/iss1/12>

This Article is brought to you for free and open access by TÜBİTAK Academic Journals. It has been accepted for inclusion in Turkish Journal of Zoology by an authorized editor of TÜBİTAK Academic Journals.

## The alien spreading of *Chama pacifica* Broderip, 1835 (Mollusca: Bivalvia: Chamidae) in the Mediterranean Sea

Fabio CROCETTA<sup>1\*</sup>, Paolo RUSSO<sup>2</sup>

<sup>1</sup>Stazione Zoologica Anton Dohrn, Villa Comunale, I-80121 Naples, Italy

<sup>2</sup>Società Italiana di Malacologia, c/o Museo di Scienze Planetarie, via Galcianese, 20H, I-59100 Prato, Italy

Received: 31.10.2011 • Accepted: 04.07.2012 • Published Online: 24.12.2012 • Printed: 21.01.2013

**Abstract:** A detailed Mediterranean synonymy (= list of published records) and distributional map is offered for the Pacific jewel-box *Chama pacifica* Broderip, 1835, an alien mollusk invasive in the deep eastern Mediterranean Sea. A more detailed examination of the single specimen published from the Thermaikos Gulf led us to consider it as a juvenile of the native congeneric *C. gryphoides* Linnaeus, 1758. Concomitantly, the presence of *C. pacifica* along the Greek shores is here first recorded on the basis of 4 adult live specimens sampled at Fokià Bay (Kápathos Island). Dates of first record in the Mediterranean for each country in which the species was recorded are reviewed with respect to the recent bibliography. On the basis of the specimens directly examined or reported in the literature, the first record date for Turkey and Lebanon is 1999, and for Greece, 2011. Finally, a discussion regarding the Mediterranean spreading of *C. pacifica* is offered.

**Key words:** *Chama pacifica*, Chamidae, alien Mollusca, Mediterranean Sea, Greece

### 1. Introduction

Biological pollution is a common phenomenon worldwide (Molnar et al., 2008) and although marine invasions have been widely documented all over the world (Molnar et al., 2008), they are very conspicuous in the Mediterranean Sea, as pointed out in several recent contributions (e.g., Zenetos et al., 2010). The large Pacific jewel-box *Chama pacifica* Broderip, 1835 is a very peculiar species usually considered as one of the most variable cosmopolitan chamids (Huber, 2010) recently included in the *CIESM Atlas of Exotic Species in the Mediterranean* (Zenetos et al., 2004). Its shell shows a strongly irregular outline from suboval to subcircular, including reports of narrower or elongated forms. It is a quite large bivalve species (shells growing up to about 100 mm in length or width), solid and inequivalve, with the lower valve (left valve, LV) usually bigger and deeper than the upper one (right valve, RV), which is usually flat. The ligament is external and the umbones are spirally coiled prosogyrate. The shell sculpture is typically different on each valve, with short-to medium-sized spines that are often more pronounced on the left side (seen from the external) of each valve. Internally with a hinge plate thick with characteristic pachyodont teeth and finely crenulated margins (sometimes not fully visible). Two large, subequal, ovate, and dorsoventrally elongate adductor muscles scars. Pallial line entire and with no sinus.

External color highly variable, from white to pink-red, spines often white. Internal color usually half white and half rose red (half rose red on the right side in the RV and on the left side in the LV, seen from the internal).

In the present paper, the pattern of geographical and temporal spread of *Chama pacifica* Broderip, 1835 in the Mediterranean is reexamined following assembly of additional records based on published and as of yet unpublished data, offering a detailed Mediterranean synonymy (= list of published records) and distributional map, reviewing first record dates and reporting the first record in Greece on the basis of sampled specimens and reevaluation of the sole previous record from the area.

### 2. Materials and methods

*Chama pacifica* Broderip, 1835 is considered an alien in accordance with the definition proposed by the International Union for Conservation of Nature (IUCN) (<http://www.iucn.org/themes/pbia/wl/docs/biodiversity/cop6/invasives.doc>).

Literature was surveyed for Mediterranean records and first record dates of *C. pacifica*, particularly that concerning faunistics, taxonomy, and biogeography. Gathered data were used to build an updated Mediterranean distribution and synonymy.

\* Correspondence: [fabio.crocetta@szn.it](mailto:fabio.crocetta@szn.it)

In this study, specimens (spm, spms) and shells (RV, LV) belonging to *C. pacifica* were collected by snorkeling and scuba diving, mainly within the frame of the French–Lebanese program of scientific cooperation CEDRE (material from Lebanon, legit Ghazi Bitar, Jean-Georges Harmelin, and Helmut Zibrowius; material from Greece, legit Paolo Russo; material from Turkey, legit Giovanni Buzzurro and Panayotis Ovalis; material from Cyprus, legit unknown). Specimens from Greece and Turkey are currently stored in the private collection of Paolo Russo (Venice, Italy), the specimen from Cyprus is in the private collection of Fabio Crocetta (Naples, Italy), and specimens from Lebanon are currently stored at the Charles Darwin Department of Biology and Biotechnology, La Sapienza University (Rome, Italy) under the care of Marco Oliverio. High-resolution color photos of the controversial specimen from Greece (see below: Mediterranean published distribution and relevant notes) were also examined. Samples recorded were measured with vernier calipers. Sizes reported are in millimeters and given as shell height (SH, from the umbone area to the opposing valve margin)  $\times$  shell length (SL, perpendicular from the height line).

### 3. Results

#### 3.1. *Chama pacifica* Broderip, 1835

##### 3.1.1. Published records

*Chama broderipii* Reeve - Tillier and Bavay, 1905: 177; *Chama broderipi* [sic!] Reeve - Pallary, 1912: 166, tab. XVII (III), fig. 8; *Chama broderipi* Rve - Moazzo, 1939: 71; *Chama reflexa* Reeve - Moazzo, 1939: 71; *Chama reflxa* [sic!] Reeve 1846 - Çeviker, 1999: 26; *Chama reflexa* Reeve 1846 - Çeviker, 1999: 27; *Chama reflexa* - Çeviker, 2001: 40, fig. 3; *Chama pacifica* Broderip, 1834 [sic!] - Mienis et al., 1993: 13–18; Bogi and Galil, 1997: 43–44, figs: 8–9; Çeviker, 2001: 45–46, fig. 21; Zibrowius and Bitar, 2003: 70; Zenetos et al., 2004: 262–263, figure; Mienis, 2004: 119; Türkmen et al., 2005: 107–110; Buzzurro and Cecalupo, 2006: 29; Delongueville and Scaillet, 2006: 30–31, fig. 2; Delongueville and Scaillet, 2007: 58, 62, 64, 66, 68, fig. 40; Zenetos et al., 2009: 2, 6; Delongueville and Scaillet, 2010: 3–5, fig. 6; Bakır et al., 2012: 180; *Chama pacifica* Broderip [sic!], 1834 - Çeviker, 2001: 41; *Chama pacifica* (Broderip, 1834) [sic!] - Bitar et al., 2005: 313; Delongueville and Scaillet, 2006: 29; *Chama pacifica* - Bitar et al., 2007: 437; Bitar, 2010: 452; *Chama pacifica* Broderip, 1835 - Zurel et al., 2011: 1–9.

Not *Chama pacifica* sensu Manousis et al. (2010).

Unpublished material examined

Cyprus:

Larnaca Bay: June 2009 - 5 m on stones - 1 spm: 39.2  $\times$  31.5.

Greece:

Fokià Bay (Kàrpathos Island): August 2011 - 1.5 m on

stones - 4 spms: 52  $\times$  31.9 (Figure 1a); 49.5  $\times$  50.6 (Figure 1b); 45.8  $\times$  32.3; 44.6  $\times$  31.9.

Lebanon:

Ramkine Island: May 2000 - 3 m - 2 RV: 40  $\times$  38.5; 43.5  $\times$  33.3. May 2000 - 13 m, sediment - 6 RV: 31.6  $\times$  29.6; 31  $\times$  32.3; 29  $\times$  26; 28.5  $\times$  21; 15  $\times$  11; 14.5  $\times$  12; 2 LV: 18.9  $\times$  13; 17.5  $\times$  13.1. June 2000 - 13–14 m, sediment - 1 RV: 5.8  $\times$  7.2. July 2003 - 7 m, cave - 1 RV: 49.3  $\times$  43.2; 1 LV: 26  $\times$  21. July 2003 - 15 m, cave - 2 spms: 19.2  $\times$  11; 18  $\times$  11.2.

Anfeh: October 1999 - 10 m - 1 spm: 29  $\times$  32.5.

Ras El Chakaa: October 1999 - 3–5 m, cave - 1 RV: 5  $\times$  4.5.

Chak El Hatab: July 2003 - 14 m, cave - 1 RV: 4  $\times$  3.9.

Selaata: October 1999 - 5–6 m, concretions - 1 spm: 9  $\times$  8. October 1999 - 5–6 m, cave - 2 spms: 9.5  $\times$  6.1; 9  $\times$  6.5. October 1999 - 6–7 m - 1 spm: 15.8  $\times$  11. October 1999 - 9 m, cave - 6 RV: 9  $\times$  7.2; 6  $\times$  6.9; 5.5  $\times$  4.9; 4.2  $\times$  4.2; 3.5  $\times$  3.2. May 2001 - 30 m - 2 RV: 12.5  $\times$  14.5; 9.5  $\times$  7.5. September 2002 - 21 m, cave - 3 spms: 58  $\times$  55; 27.5  $\times$  19.5; 18.5  $\times$  16; 1 RV: 48.5  $\times$  41.5; 1 LV: 56  $\times$  49; 49.2  $\times$  46. July 2003 - 20 m, cave - 7 spms: 61.5  $\times$  52; 51.5  $\times$  47; 50.5  $\times$  43; 19.5  $\times$  13.5; 13.5  $\times$  1.1; 13.5  $\times$  8.1; 10.5  $\times$  5; 4 RV: 42  $\times$  36.2; 34.5  $\times$  34; 32.4  $\times$  27.2; 5  $\times$  4.5; 2 LV: 29  $\times$  23.2; 21.7  $\times$  20.5.

Kfar Abida: April 2000 - 7–8 m, cave - 1 spm: 51  $\times$  5.

El Barbara: June 2000 - 26–28 m - 2 spms: 17  $\times$  13.5; 11.2  $\times$  11.2; 4 RV: 6.9  $\times$  6; 6.2  $\times$  6.1; 5.5  $\times$  5; 4.5  $\times$  4.2; 4 LV: 9  $\times$  5.5; 7  $\times$  5.1; 5  $\times$  4.2; 4.2  $\times$  3.

Jbail: October 1999 - 16 m, cave sediment - 1 RV: 5.9  $\times$  5.1.

Beirut: June 2000 - 1 RV: 7.6  $\times$  6.5. September 2002 - 3–10 m - 3 spms: 35.5  $\times$  32.5; 17  $\times$  12; 14  $\times$  10. September 2002 - 10 m - 3 spms: 15.9  $\times$  11.1; 7  $\times$  5.5; 5.1  $\times$  4.9. September 2002 - around 15 m - 1 spm: 34.2  $\times$  23. September 2002 - around 20 m - 1 spm: 19.5  $\times$  15; July 2003 - 3–11 m - 1 spm: 9.3  $\times$  7.

El Zahrani: June 2000 - 14 m - 1 RV: 5.1  $\times$  5.2. June 2000 - 24 m - 3 spms: 18.1  $\times$  13; 7  $\times$  7; 4.2  $\times$  4.

El Kassmieh: October 1999 - 44 m, sediment - 14 RV: 18.9  $\times$  19; 10.5  $\times$  8.5; 8.2  $\times$  8.1; 8  $\times$  7.7; 8  $\times$  7; 7.5  $\times$  7.5; 6.1  $\times$  5.5; 6  $\times$  5.5; 6  $\times$  5.1; 6  $\times$  5; 6  $\times$  6.5; 4.5  $\times$  4.5; 4.1  $\times$  5; 2.2  $\times$  2.2; 1 LV: 9.5  $\times$  11.

Turkey:

İskenderun: June 2005 - 1.5–2 m of depth on stones - 1 spm: 54.8  $\times$  36.4. August 2009 - 5 m on stones - 3 spms: 50.3  $\times$  32.6; 41.8  $\times$  23.4; 22.7  $\times$  14.2.

##### 3.1.2. Mediterranean published distribution, first record dates, and relevant notes

So far, *Chama pacifica* Broderip, 1835 records are limited to the eastern Mediterranean Sea (see Sciberras and Schembri, 2007; Zenetos et al., 2010a; Antit et al., 2011; Katsanevakis et al., 2011; Crocetta, 2012). A recent record of one *C. pacifica* juvenile specimen from the Thermaikos Gulf (Manousis et al., 2010) has been a matter of debate,


leading to divergent opinions. Zenetos et al. (2011) considered it as a misidentification (of another alien species - *Chama aspersa* Reeve, 1846), while Zurel et al. (2011) and Tzomos et al. (2012) accepted it as *C. pacifica*. Tzomos et al. (2012) stated the first record date from Greece in 2005 on the basis of the above-cited paper. Here, a more detailed examination of high-resolution color photos of this specimen allowed us to clarify the issue. Even if intraspecific variation within *Chama* species is high, the external sculpture with very weak and regular spines (usually more pronounced and less regular in *C. pacifica*), the internal yellow color pattern (usually purplish in *C. pacifica*), and the shape of the adductor muscle scars of the right valve (usually narrower and more elongated in *C. pacifica*) clearly point to a juvenile of the native *Chama gryphoides* Linnaeus, 1758. Published distribution, first record dates, and relevant notes are summarized in the Table and Figure 1c.

#### 4. Discussion

Widespread and presumably native from the Indo-West Pacific, *Chama pacifica* Broderip, 1835 invaded the Mediterranean Sea at least at the beginning of the last century (Tillier and Bavay, 1905), becoming first established and then invasive among the fouling communities from the intertidal to the infralittoral zone of the deep eastern Mediterranean Sea (Zenetos et al., 2010a). *C. pacifica*, in fact, constitutes a leading taxa in its habitat, both in biomass and numbers (Fishelson, 2000; Zurel et al., 2011), mainly found on hard substrates and able to thrive in harbor environments. It outnumbered the native low water congener *Chama gryphoides* Linnaeus, 1758 (Mienis, 2003; Galil, 2007). The surface of its shell provides strongholds for a diverse community of algae and invertebrates (Fishelson, 2000).

Considering the conspicuous records of *C. pacifica* in the Suez Canal (Moazzo, 1939), from where it presumably invaded the Mediterranean via natural dispersal, and its published distribution, where records in Cyprus paradoxically precede those in Lebanon and Syria (see Table and Figure 1c), 2 hypotheses of Mediterranean spreading may be suggested. The first is that this alien bivalve has followed the main pathway of Lessepsian immigrants spreading within the Mediterranean by prevailing currents; in this view, *C. pacifica* was likely present in Lebanon before 1999, and in Syria before 2003, the dates of its first published records, but it went unrecorded due to lack of specific samplings. The second is that its records from Cyprus and Turkey could be man-mediated (e.g., via shipping). The Gulf of İskenderun, in fact, is not only one of Turkey's largest ports on the Mediterranean Sea but it also serves as an important naval training base (Zenetos et al., 2010b). A molecular approach may solve the question. Regarding the presence of *C. pacifica* along the Greek coasts, although this is just speculation, *C. pacifica* may have colonized Greek coasts via natural dispersal from neighboring Turkish shores. This seems to be confirmed by the fact that K rpathos Island is quite far from any commercial route.

In addition, our record raises to 49 the number of alien Mollusca species accepted as reported from Greece (Eleftheriou et al., 2011; Zenetos et al., 2011). Consequently, the presence of a third alien *Chama* species in the area (*C. pacifica* together with *C. aspersa* Reeve, 1846 and *C. asperella* Lamarck, 1819; see Ovalis and Zenetos, 2007) locally outnumbers native *Chama* spp. (*C. gryphoides* Linnaeus, 1758 and *C. circinata* Monterosato, 1878). Finally, besides the recent arrival of *C. pacifica* on K rpathos Island (previous research held by one of us [P.R.] in 2007 did not reveal its presence) and



**Figure 1.** *Chama pacifica* Broderip, 1835. a) and b) Specimens from Foki  Bay (K rpathos Island, Greece): 52.0 × 31.9 and 49.5 × 50.6. c) Mediterranean spreading and first record dates.

**Table.** *Chama pacifica* Broderip, 1835 Mediterranean published distribution, first record dates, and relevant notes.

| Country | First record date | Main bibliographic references from the area  | Relevant notes  |
|---------|-------------------|--|---|
| Egypt   | 1905              | Tillier and Bavay, 1905; Pallary, 1912.  | First recorded from Egypt without an exact finding date.  |
| Israel  | 1988              | Mienis et al., 1993; Bogi and Galil, 1997; Delongueville and Scaillet, 2007; Galil, 2007.  |   |
| Lebanon | 1999              | Zibrowius and Bitar, 2003; Bitar et al., 2007; present paper.  | First recorded from Lebanon without an exact finding date (research held “since 1999”: Zibrowius and Bitar, 2003). The material here analyzed allows us to determine its first record as in 1999.   |
| Syria   | 2003              | Bitar et al., 2005.  |   |
| Cyprus  | 1998              | Zenetos et al., 2009; Delongueville and Scaillet, 2010.  |   |
| Turkey  | 1999              | Çeviker, 1999; Çeviker, 2001; Buzzurro and Cecalupo, 2006; Zenetos et al., 2010b; Albayrak, 2011; Çinar et al., 2011; present paper. | A recent review of alien species from Turkey stated the first record of <i>C. pacifica</i> in 2000 (Çinar et al., 2011) on the basis of Çeviker (2001). This finding date has been accepted by Tzomos et al. (2012). Bibliographic research allows us to state the first record at least to 1999, when the species was first figured from Turkey (without an exact finding date) by Çeviker (1999). |
| Greece  | 2011              | Present paper.   | See “Mediterranean published distribution, first record dates, and relevant notes.”   |

taking into account what is reported for the deep eastern Mediterranean shores, it is possible to envisage that this alien bivalve will soon become established along the Greek shores. Future observations are needed to confirm our hypotheses.

### Acknowledgments

Material from Lebanon was collected during a field trip funded by the French–Lebanese cooperation program

CEDRE: Ghazi Bitar, Marco Oliverio, and Helmut Zibrowius allowed us to study it. Sofia Galinou-Mitsoudi sent the high-resolution color photos of the controversial specimen from Greece. Serhat Albayrak, Christiane Delongueville, Henk K. Mienis, Italo Nofroni, Bilal Öztürk, and Roland Scaillet sent requested bibliographies. Dimitris Poursanidis provided the template of the Mediterranean map. Paolo Sordino critically reviewed the manuscript. We are grateful to all of them.

### References

- Albayrak, S. 2011. Alien marine bivalve species reported from Turkish seas. *Cah. Biol. Mar.* 52(1): 107–118.
- Antit, M., Gofas, S., Salas, C. and Azzouna, A. 2011. One hundred years after *Pinctada*: an update on alien Mollusca in Tunisia. *Medit. Mar. Sci.* 12(1): 53–73.
- Bakır, B.B., Öztürk, B., Doğan, A. and Önen, M. 2012. Mollusc fauna of Iskenderun Bay with a checklist of the region. *Turk. J. Fish. Aquat. Sci.* 12: 171–184.
- Bitar, G. 2010. Impact des changements climatiques et des espèces exotiques sur la biodiversité et les habitats marins au Liban. *Rapp. Comm. int. Mer Médit.* 39: 452.
- Bitar, G., Dupuy De La Grandrive, R., Foulquie, M. and Verlaque, M. 2005. *Neomeris annulata* Dickie (Dasycladales, Chlorophyceae): a potential new invader from the Red Sea to the Mediterranean Sea. *Cryptogamie, Algol.* 26(3): 309–317.
- Bitar, G., Ocaña, O. and Ramos-Esplá, A.A. 2007. Contribution of the Red Sea alien species to structuring some benthic biocenosis in the Lebanon coasts (Eastern Mediterranean). *Rapp. Comm. int. Mer Médit.* 38: 437.
- Bogi, C. and Galil, S.B. 1997. Ritrovamenti lungo le coste israeliane. *La Conchiglia* 29(284): 42–45.
- Buzzurro, G. and Cecalupo, A. 2006. I molluschi lessepsiani di Taşucu (Turchia sud-orientale): descrizione di *Parviturbo dibellai* n. sp. (Gastropoda: Trochoidea: Skeneidae). *Boll. Malac.* 42(1–4): 27–32.
- Çeviker, D. 1999. Lessepsians from Indian Ocean and the Red Sea. *Sualtı Dünyası* 45: 22–27.
- Çeviker, D. 2001. Recent immigrant bivalves in the Northeastern Mediterranean off Iskenderun. *La Conchiglia* 298: 39–46.



- Çinar, M.E., Bilecenoglu, M., Öztürk, B., Katağan, T., Yokeş, M.B., Aysel, V., Dağlı, E., Açıık, S., Özcan, T. and Erdoğan, H. 2011. An updated review of alien species on the coasts of Turkey. *Medit. Mar. Sci.* 12(2): 257–315.
- Crocetta, F. 2012. Marine alien Mollusca in Italy: a critical review and state of the knowledge. *J. Mar. Biol. Assoc. U.K.* 92: 1357–1365.
- Delongueville, C. and Scaillet, R. 2006. Mollusques associés à *Spondylus spinosus* Schreibers, 1793 dans le golfe d'Iskenderun (Turquie). *Novapex* 7(2–3): 29–33.
- Delongueville, C. and Scaillet, R. 2007. Les espèces invasives de mollusques en Méditerranée. *Novapex* 8(2): 47–70.
- Delongueville, C. and Scaillet, R. 2010. Echantillonnage de mollusques invasifs et première signalisation de *Chama aspersa* Reeve, 1846 à Chypre Nord. *Novapex* 11(1): 3–7.
- Eleftheriou, A., Anagnostopoulou-Visilia, E., Anastasopoulou, E., Ateş, S.A., Bachari, N. El I., Cavas, L., Cevik, C., Çulha, M., Cevik, F., Delos, A.L., Derici, O.B., Erguden, D., Fragopoulou, N., Giangrande, A., Göksan, T., Gravili, C., Gurlek, M., Hattour, A., Kapiris, K., Kouraklis, P., Lamouti, S., Prato, E., Papa, L., Papantoniou, G., Parlapiano, I., Poursanidis, D., Turan, C. and Yaglioglu, D. 2011. New Mediterranean biodiversity records (December 2011). *Medit. Mar. Sci.* 12(2): 491–508.
- Fishelson, L. 2000. Marine animal assemblages along the littoral of the Israeli Mediterranean seashore: The Red-Mediterranean Seas communities of species. *Ital. J. Zool.* 67(3): 393–415.
- Galil, B.S. 2007. Seeing red: alien species along the Mediterranean coast of Israel. *Aquat. Invasions* 2(4): 281–312.
- Huber, M. 2010. Compendium of Bivalves. A Full-Color Guide to 3,300 of the World's Marine Bivalves. A Status on Bivalvia after 250 Years of Research. ConchBooks, Hackenheim.
- Katsanevakis, S., Zenetos, A., Vesna Macic, V., Beqiraj, S., Poursanidis, D. and Kashta, L. 2011. Invading the Adriatic: spatial patterns of marine alien species across the Ionian-Adriatic boundary. *Aquatic Biology* 13: 107–118.
- Manousis, T., Mpardakis, G., Paraskevopoulos, C. and Galinou-Mitsoudi, S. 2010. The Bivalvia Mollusca of Thessaloniki & Thermaikos Gulf (North Aegean Sea, Greece) with emphasis on new species for Hellenic waters. *J. Biol. Res. Thessalon.* 14: 161–179.
- Mienis, H.K. 2003. Native marine molluscs replaced by Lessepsian migrants. *Tentacle* 11: 15–16.
- Mienis, H.K. 2004. New data concerning the presence of Lessepsian and other Indo-Pacific migrants among the molluscs in the Mediterranean Sea with emphasis on the situation in Israel. *Turk. J. Aquat. Life* 2(2): 117–131.
- Mienis, H.K., Galili, E. and Rapoport, J. 1993. On the presence of the Indo-Pacific bivalve *Chama pacifica* in the Eastern Mediterranean (Mollusca: Bivalvia: Chamidae). *Gloria Maris* 32(2): 13–18.
- Moazzo, P.G. 1939. Mollusques testacés marins du Canal de Suez. *Mém. Inst. Egypte* 38: 1–283.
- Molnar, J.L., Gamboa, R.L., Revenga, C. and Spalding, M.D. 2008. Assessing the global threat of invasive species to marine biodiversity. *Front. Ecol. Environ.* 6: 485–492.
- Ovalis, P. and Zenetos, A. 2007. The establishment of two more alien mollusca (*Chama aspersa* and *Chama asperella*) in the eastern Mediterranean. *Medit. Mar. Sci.* 8(2): 97–100.
- Pallary, P. 1912. Catalogue des mollusques du littoral Méditerranéen de l'Égypte. *Mém. Inst. Egypte* 7: 69–207.
- Sciberras, M. and Schembri, P.J. 2007. A critical review of records of alien marine species from the Maltese Islands and surrounding waters (Central Mediterranean). *Medit. Mar. Sci.* 8(1): 41–66.
- Tillier, L. and Bavay, A. 1905. Les mollusques testacés du Canal de Suez. *B. Soc. Zool. Fr.* 30: 170–181.
- Türkmen, A., Türkmen, M. and Tepe, Y. 2005. Biomonitoring of heavy metals from Iskenderun Bay using two bivalve species *Chama pacifica* Broderip, 1834 and *Ostrea stentina* Payraudeau, 1826. *Turk. J. Fish. Aquat. Sci.* 5: 107–111.
- Tzomos, T., Kitsos, M.S., Koutsoubas, D. and Koukouras, A. 2012. Evolution of the entrance rate and of the spatio-temporal distribution of Lessepsian Mollusca in the Mediterranean Sea. *J. Biol. Res. Thessalon.* 17: 81–96.
- Zenetos, A., Gofas, S., Russo, G. and Templado, J. 2004. CIESM Atlas of Exotic Species in the Mediterranean. Vol. 3. CIESM Publishers, Monaco.
- Zenetos, A., Gofas, S., Verlaque, M., Çinar, M.E., Garcia Raso, J.E., Bianchi, C.N., Morri, C., Azzurro, E., Bilecenoglu, M., Frogli, C., Siokou, I., Violanti, D., Sfriso, A., San Martin, G., Giangrande, A., Katağan, T., Ballesteros, E., Ramos-Esplà, A., Mastrototaro, F., Ocaña, O., Zingone, A., Gambi, M.C. and Strefaris, N. 2010a. Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Medit. Mar. Sci.* 11(2): 381–493.
- Zenetos, A., Katsanevakis, S., Poursanidis, D., Crocetta, F., Damalas, D., Apostolopoulos, G., Gravili, C., Vardala-Theodorou, E. and Malaquias, M. 2011. Marine alien species in Greek Seas: additions and amendments by 2010. *Medit. Mar. Sci.* 12(1): 95–120.
- Zenetos, A., Konstantinou, F. and Konstantinou, G. 2009. Towards homogenization of the Levantine alien biota: additions to the alien molluscan fauna along the Cypriot coast. *Mar. Biod. Records* 2: e156.
- Zenetos, A., Ovalis, P. and Çeviker, D. 2010b. On some Indo-Pacific boring endolithic Bivalvia species introduced into the Mediterranean Sea with their host – spread of *Sphenia rueppelli* A. Adams, 1850. *Medit. Mar. Sci.* 11(1): 201–207.
- Zibrowius, H. and Bitar, G. 2003. Invertébrés marins exotiques sur la côte du Liban. *Leb. Sci. J.* 4(1): 67–74.
- Zurel, D., Gophna, U. and Benayahu, Y. 2011. Parity and disparity between two *Chama* oysters: the reproductive biology of the Indo-Pacific *C. pacifica* Broderip, invasive to the Mediterranean Sea; and *C. savignyi* Lamy, indigenous to the Red Sea. *Mar. Ecol.* 33: 261–271.