

# Varusna deformacija kolen nogometašev najstnikov

## Varus knee deformity in adolescent soccer players

**Avtor / Author**

**Ustanova / Institute**

**Zmago Krajnc<sup>1,2</sup> Matjaž Vogrin<sup>1,2</sup>**

<sup>1</sup>Univerzitetni klinični center Maribor, Klinika za kirurgijo, Oddelek za ortopedijo, Maribor, Slovenija; <sup>2</sup>Univerza v Mariboru, Medicinska fakulteta, Katedra za ortopedijo, Maribor, Slovenija;

<sup>1</sup>University Medical Centre Maribor, Clinic for Surgery, Department of Orthopedics, Maribor, Slovenia; <sup>2</sup>University of Maribor, Faculty of Medicine, Department of Orthopedics, Maribor, Slovenia

**Ključne besede:**

Nogomet, rastna cona, varusna deformacija kolena, šport, najstnik

**Key words:**

soccer, growth plate, varus knee, bowlegs, sports-related, adolescent

**Članek prispel / Received**

9. 7. 2018

**Članek sprejet / Accepted**

4. 10. 2018

**Naslov za dopisovanje /**

**Correspondence**

assist. dr. Zmago Krajnc, dr. med.,  
Univerzitetni klinični center Maribor,  
Oddelek za ortopedijo, Ljubljanska 5,  
2000 Maribor, Slovenija  
Telefon: +386 41374263  
E-pošta: zmago.krajnc@guest.arnes.si

**Izvleček**

Osi spodnje okončine se dinamično razvijajo in oblikujejo v prvih dveh desetletjih življenja. V zadnjih letih opažamo znatno povečano intenzivnost vadbe med treningi in tekmovanji (čas, frekvenca, moč) in posledično povečano število akutnih in kroničnih poškodb med mladimi športniki. Med športe z večjo incidenco poškodb uvrščamo tudi nogomet. Pregled literature zadnjih 15 let, ki preučuje deformacijo kolenskih sklepov mladih nogometašev, nakazuje pogostejši pojav varusne deformacije spodnjih okončin med nogometaši v primerjavi s splošno populacijo in športniki drugih športnih panog, najverjetneje zaradi specifičnih obremenitev nogometne igre. Varusna deformacija vpliva na spremenjeno biomehaniko kolenskega sklepa s povečano obremenitvijo medialne sklepne špranje in posledično zgodnejšim pojavom obrabnih sprememb kolenskega sklepa. Varusna deformacija kolenskih sklepov dodatno

**Abstract**

The lower limb axis develops dynamically throughout the first and second decades of life. In recent years, with respect to youth athletes, we have been facing a progressive increase in the intensity of exercise during training and competition (involving increased time, frequency, and power), and, as a result, there has been an increase in the number of acute and chronic overuse injuries. There are some sports with a higher incidence of injuries, one of which is soccer. A review of literature for the past 15 years with a focus on varus knee deformities in adolescent soccer players showed that varus axial deformities of the lower limbs occur more frequently in soccer players than in the general population and in athletes from other sport, probably due to the specific loading pattern in soccer training. Varus deformities alter knee biomechanics, which results in medial joint space overloading, and consequently an early onset of degenerative joint

vpliva na pogostejši pojav sindroma patelo-femoralne bolečine in poškodb meniskusov med športniki. Etiologija varusne deformacije spodnjih udov med nogometaši je posledica kombinacije različnih dejavnikov: od naravne selekcije nogometašev z genetsko predispozicijo za varusno deformacijo spodnjih okončin do mehanske preobremenitve medialnega dela rastne cone proksimalne golenice, ki je posledica igranja nogometa. Klinične študije in raziskave na živalih nakazujejo, da lahko intenzivna športna vadba negativno vpliva na normalen razvoj rastne cone kolenskih sklepov in posledično na pojav nepravilnosti v rasti.

disease. In addition, bowlegs predisposes athletes to patello-femoral pain syndrome and meniscal lesions. The etiology of genu varum in soccer is thought to be multifactorial, ranging from the natural selection of players to mechanical overload of the proximal medial tibial growth plate. A growing number of clinical reports and animal studies have indicated that intensive sport training may precipitate pathologic changes of the knee joint growth plate, and even produce a growth disturbance.

## INTRODUCTION

More and more people (both children and adults) worldwide are participating in sport on a daily basis (1,2). Sports provide fun and satisfaction; however, sports also pose an increased risk of musculoskeletal injuries (3,4).

Physical activities and involvement in sports significantly influence the wellbeing, health, and personal development of children (5). Over recent years, these beneficial effects resulting from sports activities have been accompanied by a progressive increase in the intensity of training, competitiveness at an increasingly early age, and a premature focus on single-sport activities. As a result of increased overload, a number of sports-related adverse consequences manifest as injuries. Such injuries are attributed to incorrect preparation of athletes for intensive sport activities, as well as overload in training and competition (1,6). Several authors have indicated that 0.5-35 injuries occur per 1000 hours of exposure to sports activities (7,8). The incidence of injuries varies and depends on the age, gender, and type of sport. Hockey, American football, and soccer have the highest incidence of injuries in boys, while soccer, basketball, and gymnastics account for the highest incidence in girls (7,8).


Soccer is the most popular sport worldwide with respect to registered players and fans. According to the Fédération Internationale de Football Association (FIFA), a total of 270 million people are registered soccer players, of which 200,000 are professional

soccer players (9-11). More than 50% of registered soccer players are < 18 years of age (12). Soccer has specific rules and regulations, and a unique method of play (10). Soccer combines walking, jogging, and sprinting. Soccer is characterized by frequent contact among players, high speed, quick and sudden changes in direction, and tackling maneuvers (10-12). Unlike other sports, soccer is mainly characterized by kicking the ball (12,13). During a soccer game, players and their musculoskeletal system are frequently exposed to major loading. Therefore, soccer has the greatest risk of injuries among all sports (3). Many authors agree that the majority of injuries that occur in adult and adolescent soccer players are limited to the lower extremities, with knee and ankle injuries accounting for most of the injuries (9,12). According to the literature, the incidence of injuries in football players increases with age and is significantly higher during games and lower during training (9,12).

### Varus knee deformities in soccer players

During the first decade of life, the axes of the lower limb undergo dynamic changes; however, changes in the shape of the bone and the lower limb axis can occur due to physiologic bone remodeling and under the influence of external and internal factors throughout the entire life. It is well known that lower limb varus deformities (so called bowlegs) are more frequent in soccer players compared to their peers. Chantraine

(14,15) reported that > 80% knees of retired soccer players have lower limb varus deformities, which is significantly higher than the 33% of the normal male population with constitutional varus knees (> 3° of varus). An increased incidence of varus knees among adolescent soccer players has also been reported when compared to peers who did not play soccer (13,16,17) [Figure 1].



**Figure 1.** Bow legs of 18-year old professional soccer player, varus deformation of both knees

A review of the literature for the past 15 years involving varus knee deformities in adolescent soccer players revealed only a few relevant studies, and showed that varus axial deformities of the lower limbs occur more frequently in soccer players than in the general population and other athletes, probably due to a specific loading pattern in soccer training (13,16–19). Thijs et al. (17) compared varus development in male adolescents 13–15 years of age who did or did not participate in sports, and established that varus occurred in both groups. This is in agreement with the physiologic development of the lower limb axis described by Cahuzac et al. (20), who observed physiologic varus in boys at the end of the growth spurt period. Thijs et al. (17) also reported that varus was more common

in the group of boys 13- to 15- years-old who were engaged in sports. Yaniv et al. (13), Witvrouw et al. (16), and Asadi et al. (18) compared groups of adolescent soccer players and other athletes and es-

tablished that soccer players between 13 and 16 years of age have a significantly higher degree of knee varus deformities compared to non-soccer players who were engaged in other sports involving increased impact on knee joints, such as tennis, volleyball, basketball, and handball (4,13,16). Athletes engaged in these sports are exposed to similar loads characterized by slow and fast running, quick and sudden changes in the direction of movement, and jumping. The only differences between soccer players and athletes in other sports is tackling and kicking the ball, which might have a significant influence on the frequent lower limb varus deformities in soccer players (13,16,17). In a growth plate study using diffusion-weighted imaging (DWI), Krajnc et al. (19) concluded that the growth plates in adolescent soccer players (the medial proximal tibia in particular) are greatly affected by playing soccer, which might result in more frequent knee varus deformities in soccer players.

Due to the specific nature of soccer, the knee joints of soccer players are exposed to impact forces and torsional loading (13,15). In a growing and developing skeleton, these forces are further transferred onto growth plates, which are excessively and asymmetricaly loaded. This feature may lead to disturbed growth plate functioning and deformation of growing bones and joints.

#### Reasons for varus knee deformities in adolescent soccer players

It has been established that the increased intensity of sport activities during the growth spurt period has an effect on the incidence of increased varus in a specific group of adolescent boys, especially among soccer players, who are exposed to specific training on a daily basis (13,16–18). Repeated loading in soccer players during the growth spurt phase is believed to affect lower limb varus angulation. A possible explanation for varus deformation in soccer players might be provided by the Hueter-Volkman Law, which states that increased pressure parallel to the axis of the epiphysis inhibits growth, while decreased pressure promotes growth and changes in compressive forces causing asymmetric growth of bone, or by Frost's theory of chondral remodeling which suggests that physiologic loading stimulates growth, whereas loads outside

this range (lower or higher) inhibit growth (21–23). When analyzing specific activities that might give rise to varus in soccer players, many authors agree that it is the kicking of the ball that is unique to soccer in comparison to other sports (12,13,16). Witvrouw et al. (13,24) showed knee adduction moment during ball kicking, which is the primary determinant of medial-lateral load distribution on the knee joint during gait, is enhanced. The adduction moment is related to the magnitude of the compressive load on the medial knee compartment, which is additionally increased during intensive training (17,24). The varus and valgus loading to individual knee joint parts during sudden cutting maneuvers are considerably larger than those measured during normal running activities; however, the applied loads are believed to be medial-lateral symmetric, and thus do not affect the development of varus deformities (25). Sudden changes in direction are typical for other sport activities as well and are not unique to soccer; however, kicking the ball is a unique characteristic specific to soccer compared with other sports. Authors have also suggested a theory of natural selection for soccer players with a genetic predisposition to varus, which plays an important role in infant balance and stability during the first steps and possibly when playing football (16).

According to the Hueter-Volkman Law of remodeling, this enhanced adduction moment, resulting from repeated loads to knee joints that mostly occur during kicking actions, might have a negative impact on growth plates in the medial knee compartment. Eventually, this effect might have a negative impact on the growth and development of knees in adolescent soccer players (17).

#### **Growth plate disturbance: A reason for varus deformation?**

The growth plates around the knee joints (distal femur and proximal tibia) contribute 55%–70% of the growth of the lower limb. Even minimal growth disturbances in this area, therefore, can have a significant impact on the occurrence of possible deformations (26,27). Studies conducted to date indicate a correlation between increased sport intensity and pathologic changes in growth plates that lead to growth

disturbance in affected bones and limbs (1,2,5,7,28). The susceptibility of the growth plate and the resulting increased risk of injuries is believed to be highest during the growth spurt in adolescents (1,27,29). This finding is consistent with an animal study that showed reduced growth plate resistance during the rapid growth phase (30). A thorough understanding of the pathophysiologic processes at the cellular level in growth plate overloads requires a histologic evaluation of tissue. Radiologic imaging remains the basic tool used to study the changes occurring in growth plates in humans. The results may be compared with the findings from animal studies that allow the comparison of radiologic changes in growth plates under the impact of different loads with the histologic samples of the same tissue. When evaluating growth plate injuries in humans, diagnostics are limited to conventional radiography, computer tomography (CT), and magnetic resonance imaging (MRI), but typical radiologic changes become radiologically-visible only after an interval of symptomatic pain not observed in asymptomatic athletes (1,27,31,32). It is known that increased compression loads cause changes in the growth plate width, which can be identified on an MRI scan and later, on X-ray images; however, only after a long-lasting increased load or short period of significant overload of growth plates (acute injuries), which has been confirmed with animal studies. Increased loads within physiologic limits are supposed to cause a transitional width of growth plates. If these limits are exceeded and irreparable damage has already occurred, the growth plate narrows and gradually closes (1,2,27,32–34).


Krajnc et al. (19) have used a more recent radiologic tool (Diffusion Weighted Imaging - DWI) to introduce a new method of evaluating subtle changes in growth plates resulting from repeated loads that have not yet caused clinical symptoms and/or detectable changes in growth plates that are visible using standard radiologic methods. During the study, adolescent soccer players (14 years-of-age) underwent two months of intensive football training; however, a difference in growth plate width compared to baseline measurements was not demonstrated, but only changes at the cellular level obtained by calculating the apparent diffusion coefficient (ADC), especially



in the medial proximal tibial growth plate, which is supposed to reflect greater tissue cellularity in this area (Figure 2). The results of this research are consistent with the histologic findings in animal growth plates exposed to different external loads. The authors of histologic animal studies established that increased compression load on growth plates causes the expansion of hypertrophic layers in growth plates due to an increased number of chondrocytes (increased cellular-

ity) and simultaneous gradual disorganization of the layers and columns (33,35–37).

It can be concluded that the more frequent varus deformation of knee joints among athletes, in particular soccer players, is the result of changes occurring in the growth plate as a response to repeated rotation and compression forces affecting the growth plates of knee joints.



**Figure 2.** A sample of DWI analysis of growth plate in an adolescent soccer player before (A, A1) and two months after (B, B1) sport activity. There were no significant differences in growth plate surface area measured in MR images (A, B). Calculated ADC values after 2 months were decreased and may reflect an increased cellularity of the growth plate (A1, B1) (reproduced with permission) (19).

## CONCLUSION

Higher incidence of varus deformations in the lower extremities of soccer players is related to several causes; specifically, several biological and mechanical factors, affecting the overloaded medial part of the growth plate in knee joints, are the most likely cause. Varus deformation changes the mechanical load placed on the knee joint. The center of mechanical axes is medialized and the medial compartment load is consequently increased several times compared to the lateral compartment (38–40). Genu varum predisposes athletes to degenerative, overuse, and traumatic lesions of the knee joints, patellofemoral pain syndrome, and meniscal lesions (13,17,41). The prob-

ability of developing a degenerative knee joint in a varus knee deformation is two-fold compared to neutral and valgus knees. The probability and progression of knee cartilage degeneration increases with the extent of varus deformation, and thus negatively affects the quality of life in athletes, even after the sporting career ends (41).

Despite the efforts of numerous researchers, the exact mechanism underlying an increased incidence of varus in soccer players remains unknown. Further research is required to gain more insight into the causes of varus deformation of the knee in order to enable millions of young soccer players to play soccer safely without negative consequences for their health and life.

## REFERENCES

1. Caine D, DiFiori J, Maffulli N. Physeal injuries in children's and youth sports: reasons for concern? *Br J Sports Med* [Internet]. 2006 Sep;40(9):749–60. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16807307>.
2. Brenner JS, Kelly AW. Overuse and Overtraining Injuries in Teenage Athletes. *Adolesc Med State Art Rev* [Internet]. 2015 Apr;26(1):79–99. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26514033>.
3. Buckwalter JA. Sports, joint injury, and posttraumatic osteoarthritis. *J Orthop Sport Phys Ther*. 2003;33: 578-88.
4. Buckwalter JA MJ. Sports and osteoarthritis. *Curr Opin Rheumatol*.. 2004;16: 634-39.
5. Shanmugam C, Maffulli N. Sports injuries in children. *Br Med Bull* [Internet]. 2008;86:33–57. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18285352>.
6. Soomro N, Sanders R, Hackett D, Hubka T, Ebrahimi S, Freeston J et al. The Efficacy of Injury Prevention Programs in Adolescent Team Sports: A Meta-analysis. *Am J Sports Med* [Internet]. 2015 Dec 16; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26673035>.
7. Frisch A, Croisier JL, Urhausen A, Seil R, Theisen D. Injuries, risk factors and prevention initiatives in youth sport. *Br Med Bull*. 2009;92(1):95–121.
8. Caine D, Maffulli N, Caine C. Epidemiology of Injury in Child and Adolescent Sports: Injury Rates, Risk Factors, and Prevention. *Clin Sports Med*. 2008;27(1):19–50.
9. Mohib M, Moser N, Kim R, Thillai M, Gringmuth R. A four year prospective study of injuries in elite Ontario youth provincial and national soccer players during training and matchplay. *J Can Chiropr Assoc* [Internet]. 2014;58(4):369–76. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4262805&tool=pmcentrez&rendertype=abstract>.
10. Manning MR, Levy RS. Soccer. *Phys Med Rehabil Clin N Am* [Internet]. 2006 Aug;17(3):677–95, vii. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16952758>
11. Wong P, Hong Y. Soccer injury in the lower extremities. *Br J Sports Med* [Internet]. 2005 Aug;39(8):473–82. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16046325>.

12. Rössler R, Junge A, Chomiak J, Dvorak J, Faude O. Soccer Injuries in Players Aged 7 to 12 Years: A Descriptive Epidemiological Study Over 2 Seasons. *Am J Sport Med.* 2015;
13. Witvrouw E, Danneels L, Thijs Y, Cambier D, Bellemans J. Does soccer participation lead to genu varum? *Knee Surgery, Sport Traumatol Arthrosc.* 2009;17(4):422–7.
14. Bellemans J, Colyn W, Vandenneucker H, Victor J. The Chitranjan ranawat award. Is neutral mechanical alignment normal for all patients? The concept of constitutional varus. *Clin Orthop Relat Res.* 2012;470(1):45–53.
15. Chantraine A. Knee joint in soccer players: osteoarthritis and axis deviation. *Med Sci Sports Exerc* [Internet]. 1985 Aug;17(4):434–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/3839890>.
16. Yaniv M, Becker T, Goldwirt M, Khamis S, Steinberg DM, Weintraub S. Prevalence of Bowlegs Among Child and Adolescent Soccer Players. *Clin J Sport Med* [Internet]. 2006 Sep;16(5):392–6. Available from: <http://content.wkhealth.com/linkback/openurl?sid=WKPTLP:landingpage&an=00042752-200609000-00003>.
17. Thijs Y, Bellemans J, Rombaut L, Witvrouw E. Is high-impact sports participation associated with bowlegs in adolescent boys? *Med Sci Sports Exerc.* 2012;44(6):993–8.
18. Asadi K, Mirbolook A, Heidarzadeh A, Kivi MM, Meybodi MKE, Rad MR. Association of soccer and genu varum in adolescents. *Trauma Mon.* 2015;20(2):47–51.
19. Krajnc Z, Ruprecht M, Drobnič M. Quantitative Evaluation of Growth Plates around the Knees of Adolescent Soccer Players by Diffusion-Weighted Magnetic Resonance Imaging. *Biomed Res Int* [Internet]. 2015;2015:482017. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26693482>
20. Cahuzac J, Vardon D S de gauzy J, Sales de Gauzy J. Development of the clinical tibiofemoral angle in normal adolescents. *Surgery.* 1995;77(5):729–32.
21. Frost HM. A chondral modeling theory. *Calcif Tissue Int* [Internet]. 1979 Nov 6;28(3):181–200. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/92358>.
22. C. H. Anatomical study of the limb joints of newborns and adults. *Virchows Arch.* 1862;25:572–99.
23. R. V. Impairments of the musculoskeletal system. In: *Handbook for common and special surgery.* Stuttgart: Ferdinand Enkle; 1869. p. 845–920.
24. Sharma L, Hurwitz DE, Thonar EJ, Sum JA, Lenz ME, Dunlop DD et al. Knee adduction moment, serum hyaluronan level, and disease severity in medial tibiofemoral osteoarthritis. *Arthritis Rheum* [Internet]. 1998 Jul;41(7):1233–40. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/9663481>.
25. Besier TF, Lloyd DG, Cochrane JL, Ackland TR. External loading of the knee joint during running and cutting maneuvers. *Med Sci Sports Exerc* [Internet]. 2001 Jul;33(7):1168–75. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11445764>.
26. Ecklund K, Jaramillo D. Patterns of premature physal arrest: MR imaging of ill children. *Am J Roentgenol.* 2002;178(4):967–72.
27. Jawetz ST, Shah PH, Potter HG. Imaging of physal injury: overuse. *Sports Health* [Internet]. 2015;7(2):142–53. Available from: <http://sph.sagepub.com/content/early/2014/11/17/1941738114559380.abstract>.
28. Rosendahl K, Strouse PJ. Sports injury of the pediatric musculoskeletal system. *Radiol Med* [Internet]. 2016 Feb 2; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26838592>.
29. Sasaki T, Ishibashi Y, Okamura Y, Toh S, Sasaki T. MRI evaluation of growth plate closure rate and pattern in the normal knee joint. *J Knee Surg* [Internet]. 2002;15(2):72–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12013076>.
30. Bright RW, Burstein AH, Elmore SM. Epiphyseal plate cartilage. A biomechanical and histological analysis of failure modes. *J Bone Joint Surg Am* [Internet]. 1974 Jun;56(4):688–703. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/4835816>.
31. Laor T, Wall EJ, Vu LP. Physal widening in the knee due to stress injury in child athletes. *Am J Roentgenol.* 2006;186(5):1260–4.
32. Forriol F, Shapiro F. Bone development: interaction of molecular components and biophysical forces. *Clin Orthop Relat Res* [Internet]. 2005 Mar;432:14–33. Available from: <http://www>.

- ncbi.nlm.nih.gov/pubmed/15738800.
33. Stokes IA, Mente PL, Iatridis JC, Farnum CE, Aronsson DD. Enlargement of growth plate chondrocytes modulated by sustained mechanical loading. *J Bone Joint Surg Am* [Internet]. 2002 Oct;84-A(10):1842–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12377917>.
  34. Ogden JA. *Skeletal injury in the child*. New York: Springer Verlag; 2000.
  35. Arriola F, Forriol F, Cañadell J. Histomorphometric study of growth plate subjected to different mechanical conditions (compression, tension and neutralization): an experimental study in lambs. Mechanical growth plate behavior. *J Pediatr Orthop B* [Internet]. 2001 Oct;10(4):334–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11727379>.
  36. TRUETA J, TRIAS A. The vascular contribution to osteogenesis. IV. The effect of pressure upon the epiphysial cartilage of the rabbit. *J Bone Joint Surg Br* [Internet]. 1961 Nov;43-B:800–13. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/14039826>.
  37. Jaramillo D, Laor T, Zaleske DJ. Indirect trauma to the growth plate: results of MR imaging after epiphyseal and metaphyseal injury in rabbits. *Radiology* [Internet]. 1993 Apr;187(1):171–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8451408>.
  38. Espandar R, Mortazavi SM BT. *Angular Deformities of the Lower Limb in Children*. 2010;
  39. Cook SD, Lavernia CJ, Burke SW, Skinner HB, Haddad RJ. A biomechanical analysis of the etiology of tibia vara. *J Pediatr Orthop* [Internet]. 1983 Sep;3(4):449–54. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/6630488>.
  40. Sabharwal S. Blount Disease. *J Bone Jt Surg* [Internet]. 2009 Jul 1;91(7):1758-76. Available from: <http://jbjs.org/cgi/doi/10.2106/JBJS.H.01348>.
  41. Brouwer GM, van Tol AW, Bergink AP, Belo JN, Bernsen RMD, Reijman M, et al. Association between valgus and varus alignment and the development and progression of radiographic osteoarthritis of the knee. *Arthritis Rheum* [Internet]. 2007 Apr;56(4):1204–11. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17393449>.