

# Signs of Overload After an Intensified Training

## Authors

G. Bresciani<sup>1</sup>, M. J. Cuevas<sup>1</sup>, O. Molinero<sup>1</sup>, M. Almar<sup>1</sup>, F. Suay<sup>2</sup>, A. Salvador<sup>2</sup>, J. A. de Paz<sup>1</sup>, S. Marquez<sup>1</sup>, J. González-Gallego<sup>1</sup>

## Affiliations

<sup>1</sup>Institute of Biomedicine (IBIOMED), University of León, Spain

<sup>2</sup>Department of Psychobiology, University of Valencia, Spain

## Key words

- intensive aerobic training
- biomarkers of fatigue
- stress-recovery
- mood states
- overtraining

## Abstract

▼ This study investigated effects of a 9-week intensified aerobic training and 3-weeks of recovery on signs of overload in 9 healthy active young males. Blood and saliva samples were collected and psychological questionnaires were administered during baseline (T1), intermediate load (T2), maximal load (T3), and recovery (T4) periods. Maximal oxygen uptake increased and blood lactate concentration decreased in T3, while running time in a 3000m track field test was significantly shorter. No significant changes were found in hematocrit, haemoglobin concentration, white blood cell count, lactate dehydrogenase, transaminases, interleukin-6, tumour necrosis factor- $\alpha$ , myeloperoxidase and markers of oxida-

tive stress in plasma, or salivary cortisol and testosterone. Increases in different negative affect scales and in the total mood disturbance score of the Profile of Mood States were observed during T3. Scores in the stress scales of the Recovery-Stress Questionnaire for Athletes and in the State Anxiety Scale of the State-Trait Anxiety Inventory also showed significant increases during T3. The lack of effects in biomarkers together with the changes observed in psychological assessment indicates that an intensified training can produce psychological disturbances prone to early overreaching development. Additionally, it seems that psychological parameters are sensitive markers to detect stress produced by load increases.

## Introduction

▼ Intensive aerobic training can produce beneficial effects on aerobic performance, running economy, prevention of bone and muscle loss, immune system, or the plasmatic concentration of growth factors [3,33]. However, training volumes below what can be considered optimal do not result in the desired adaptation, whereas training volumes above the optimum may lead to an undesired condition usually referred to as *overtraining syndrome* [24]. In competitive sports hard training is essential to improve performance. So, one should be able to not just train hard, but also to recover after training [18]. Unfortunately, there is relatively little research on valid and practical tools to monitor the recovery and stress states.

There are also evidences which suggest that exercise may induce alterations in different organic systems. Thus, it is well known that long-duration, high-intensity exercise is associated with immunosuppression, a higher susceptibility to infection [34], microinjuries and a local inflammatory reaction in the musculature causing

changes in serum markers of inflammation [5,26]. Overload training can also induce oxidative stress, which is also related to inflammation, leading to an impaired antioxidant defence and a lack of anticipated adaptations to training, as well as a distortion of the redox balance [40]. It is also well known that exercise alters hormonal responses and concentrations. Actually, the free testosterone to cortisol (fTCR) ratio has been suggested as a potential endocrine biomarker to monitor the training status of an athlete [41]. Increased exercise stress is not only manifested in physiological and biochemical changes, but it is often presented in conjunction with psychological alterations. The symptoms of overtraining include depressed mood, general apathy, decreased self-esteem, emotional instability, disturbed sleep and others [11]. It has been suggested that examination of the athlete's performance and mood state at different periods during a competitive season is necessary for a better understanding of performance outcomes [9]. A symptom of a disturbed stress-regeneration caused by exercise increased demands is a

accepted after revision  
January 05, 2011

## Bibliography

DOI <http://dx.doi.org/10.1055/s-0031-1271764>  
Published online:  
March 4, 2011  
Int J Sports Med 2011; 32:  
338–343 © Georg Thieme  
Verlag KG Stuttgart · New York  
ISSN 0172-4622

## Correspondence

**Dr. Javier González-Gallego**  
University of León  
Institute of Biomedicine  
Campus Vegazana s/n  
24071 León  
Spain  
Tel.: +34/987/291 258  
Fax: +34/987/291 267  
jgonga@unileon.es

worsened mood state, as measured by the Profile of Mood States (POMS) [31]. In recent years it has also been indicated that the avoidance of overtraining and the achievement of optimal performance can be only realized when athletes are able to recover and optimally balance training stress and subsequent recovery [21]. The complex effects of stress and recovery may be measured through the Recovery-Stress Questionnaire for Athletes (RESTQ-Sport), which assesses the frequency of experienced stressors and regeneration related activities [32]. The RESTQ-Sport has been shown to be a valuable tool for monitoring the reactions to changes in training load [12,22], the prevention of illness, and the optimization of training and recovery programs [6,7].

To understand which mechanisms are behind exercise performance is essential in order to be able to give sport coaches scientific feedbacks so as to facilitate a better background for training outcomes and to determine the point at which training becomes maladaptive [6]. Therefore, in this study a longitudinal design was used to determine if a 9-week intensified aerobic training can induce signs of overload. Functional, biochemical, haematological, hormonal and psychological measurements were performed in order to obtain a wide-ranging effect of such type of training.

## Materials and Methods



### Participants

9 healthy active young males participated in the study ( $22.3 \pm 1.4$  yr), which was approved by the Ethics Committee of the University of León and carried out according to the Declaration of Helsinki and Ethical Standards in Sport and Exercise Science Research [17]. Written informed consent was then obtained from participants, who were not training regularly before the beginning of the study.

### Experimental design

The training protocol consisted of an intensive aerobic training based on outside running followed by a 3-week recovery period. The training frequency was 3 times a week, with an initial duration of the sessions of 40, 30 and 40 min respectively, increasing weekly training volume by 5 min for each session. Training intensity was controlled by training impulse (TRIMPS) [2] and participants were fully encouraged to not slow down the intensity. A global positioning system device (GPS) was also used for training monitoring (Garmin Edge 305, New Zealand). Heart rate (HR) was monitored with a short-range telemetry system monitor (Polar Team System, Polar Electro Oy, Kempele, Finland).

Moreover, participants had to perform several physical and psychological tests at 4 different time periods during the training schedule: baseline (T1), intermediate load (T2), maximal load (T3), and recovery (T4). During periods T1 and T4 subjects did not train. During periods T3 and T4 measurements were performed 24 h after the last training session to avoid major sample bias.

### Testing protocols

The participants filled in the psychological questionnaires and a blood sample was obtained at the laboratory. Venous blood and saliva samples were collected while participants were resting. Blood samples were centrifuged in order to obtain the plasma. Plasma aliquots and saliva samples were stored at  $-80^{\circ}\text{C}$  for

further analysis. Next, anthropometric data were obtained according to the American College of Sports Medicine. After the blood sampling and the anthropometric assessment, a maximal oxygen uptake test was performed in order to assess training adaptation. Participants walked on a motorized treadmill (Cosmos, Pulsar, Germany) for 3 min at  $6\text{ km}\cdot\text{h}^{-1}$ . Speed was then increased by  $1\text{ km}\cdot\text{h}^{-1}$  every minute until exhaustion. Achievement of  $\text{VO}_2\text{max}$  was considered as the attainment of at least one of the following criteria: (a) participant's volitional exhaustion, (b) a plateau in  $\text{VO}_2$ , or (c) a  $\text{HR} \pm 10\text{ beats}\cdot\text{min}^{-1}$  of age-predicted maximal HR. Expired gases were collected and analyzed using a breath-by-breath automated gas analysis system (CPX plus, Medical Graphics, St. Paul, MN, USA). Heart rate was monitored and a capillary blood sample was obtained from the earlobe of each participant 3 min after the end of the test, and blood lactate concentration measurements were performed using a miniphotometer (Miniphotometer Plus LP20, Dr. Lange, Germany). Perceived exertion was verified every time the speed was increased using the Borg scale of 20 items [4].

Finally, a period of 48 h after the maximal oxygen uptake test was respected before participants performed the track field test. The test was performed individually and consisted of running 3000 m as fast as possible on a track of Olympic measures. HR data and wind speed was controlled using an anemometer (Anemo Cup Anemometer, BSRIA Instrument Solutions, UK). A chronometer was used to control lap and total times. Participants were verbally encouraged to run as fast as possible on each of the 4 track trial tests during the training schedule.

### Haematological, Biochemical and Hormonal Analysis

Hematocrit (Hct), haemoglobin concentration (Hb), and total and differential counts of white blood cells (neutrophils, lymphocytes, monocytes, basophils and eosinophils) were determined using an automated haematology analyzer (Coulter Juniors JS, Coulter Electronics, Delkenheim, Germany).

Creatinine, lactate dehydrogenase (LDH), alanine aminotransferase (ALT), aspartate aminotransferase (AST), bilirubin direct, albumin, and uric acid were determined in plasma by a Cobas Integra 400 analyzer (Hoffmann-La Roche, Basel, Switzerland).

Myeloperoxidase (MPO) protein levels and cytokines IL-6 and  $\text{TNF-}\alpha$  were measured in plasma using an enzyme immune assay (OxisResearchTM, BioCheck, Inc MPO-EIA, USA and Human IL-6 and  $\text{TNF-}\alpha/\text{TNFSF1A}$  immunoassays, Quantikine® HS, R&D Systems, USA, respectively). Malondialdehyde (MDA) was assayed by Western blot analysis using an appropriate antibody [19]. Protein carbonyls content was measured using a commercial kit (Oxyblot™ protein oxidation kit, Millipore, USA).

Salivary cortisol concentrations were analyzed using a competitive solid phase with the commercial kit Coat-A-Count C adapted to salivary levels (DPC, Siemens Medical Solutions Diagnostics). Salivary testosterone concentrations were analyzed by a competitive chemiluminescence immunoassay (Diagnostics Biochem Canada Inc.).

### Psychological assessment

The Profile of Mood State (POMS) [28] is a 65-item questionnaire which provides a method of assessing transient, fluctuating mood states. It includes 5 negative affect scales: fatigue, depression, tension, anger, confusion and a positive affect scale: vigour scale. Subjects are given a score for each of the 6 mood states. The total mood disturbance score (TMD) is calculated by summation of the negative scales and subtraction of the positive

scale. The Recovery-Stress Questionnaire for Athletes (RESTQ-Sport) [23] is a questionnaire consisting of 76 items which indicate how often the respondent participated in various activities during the past 3 days and nights. The measure includes 12 scales which assess various stressing agents of a general nature and general recovery activities and 7 additional sports-specific scales, with 4 questions per scale and a warm-up question. The scores of the stress-related scales were summed up and divided by the number of scales to obtain a total stress score. The same procedure was used for the recovery-oriented scales, resulting in a total recovery score. Scores were also obtained for general stress, sport-specific stress, general recovery and sport-specific recovery. The Spanish versions of the POMS and the RESTQ-Sport have been previously demonstrated to be valid and reliable instruments [13, 14]. Participants also filled in the Spanish version [38] of the 20 items State-Trait Anxiety Inventory State-Anxiety scale (STAI-S), which measures the temporary condition of state anxiety [37].

### Statistical analysis

ANOVA with repeated measures was used to determine the differences between Hb, Hct, creatinine, MDA, protein carbonyls, MPO, TMD and training load. In the rest of the data (i.e., POMS and RESTQ-Sport) MANOVA main effects were analyzed. Sphericity was checked using the Mauchly sphericity test. Post-hoc multiple comparisons were used to identify the location of the pair-wise significant differences between training phases corrected using the Bonferroni adjustment. An alpha level of 0.05 was chosen to represent statistical significance. The Statistical Package for Social Sciences (SPSS, Ins, Chicago, I), Version 17, was used for all analyses. Training impulses were calculated using a spreadsheet (Microsoft Office Excel 2007, Microsoft Corporation, USA).

### Results

Table 1 shows the anthropometric assessment and participants' characteristics. No significant changes were found in the different time periods of the training protocol for weight or body fat. Maximal oxygen uptake increased significantly during the period T3 when compared to T1 ( $P=0.019$ ). Maximal HR and blood lactate decreased significantly in T3 compared to the T1 ( $P=0.039$ ;  $P=0.016$ , respectively). Concerning track trial test, the time performance in the test showed a decrease during the T3 period compared to T1 ( $P=0.033$ ).

Fig. 1 illustrates the work load registered in each of the 9 weeks. Training load (TRIMPS) increased progressively during the 9-week intensified training protocol.

Table 2 presents the haematological parameters, which showed no differences during the different periods of the training program.

Table 3 shows that biochemical data did not significantly differ among the various training periods.

Table 4 presents the hormonal data obtained during the study. Testosterone concentration tended to decrease and that of cortisol to increase, resulting in progressive non significant lower values for the free testosterone to cortisol ratio (fTCR).

Table 5 shows POMS scales, STAI-S and RESTQ-Sport scales measured during the entire training protocol. Several scores for the different scales tended to increase in T3 when compared to T1, reaching significance in the tension ( $P=0.024$ ), fatigue

( $P=0.016$ ) and confusion ( $P=0.021$ ) scales, and in the total mood ( $P=0.037$ ) of the POMS, in the STAI-S score ( $P=0.005$ ), and in the general stress ( $P=0.008$ ), sport specific stress ( $P=0.011$ ), and total stress ( $P=0.008$ ) scales of the RESTQ-Sport.

### Discussion

In the current study  $VO_{2max}$  increased significantly during the period of maximal load when compared to the baseline values. Gormley et al. [15] also found an increase in  $VO_{2max}$  after a training period of similar intensity with match-aged participants, concluding that high intensity training is more effective than a moderate-intensity program in order to improve the  $VO_{2max}$  of young participants. Confirming previous data, detraining caused a decrease in the  $VO_{2max}$  measurements [29, 30].


Mujika et al. found higher values for lactate concentration during an intensified training period [30]. However, the decrease in blood lactate levels observed in our study suggests a proper training adaptation to the load imposed during the training, assuring that participants were more able to cope with the acid load produced during the test protocol. Changes in the time to perform the track field test appear to confirm the physiological adaptation to training. Therefore the data presented here support the beneficial effects of the training program on the functional measurements performed. In fact, Swain and Franklin [39] stated that vigorous-intensity exercise confers greater cardio-protective health benefits than moderate-intensity exercise. Moreover, clinical trials have found that higher-intensity exercise results in greater reductions in resting blood pressure than lower intensity exercise [1, 20].

Haematological parameters did not show differences over the training period. Previous studies which focused on intensive training also failed to show changes in the immune cells

**Table 1** Anthropometric and training data (mean  $\pm$  SD) obtained during the entire training protocol.

	T1	T2	T3	T4
weight (kg)	78.9 $\pm$ 7.4	77.2 $\pm$ 7.8	74.8 $\pm$ 6.8	78.0 $\pm$ 8.3
body fat (%)	11.9 $\pm$ 5.2	9.9 $\pm$ 5.3	7.9 $\pm$ 5.1	11.8 $\pm$ 5.0
$VO_{2max}$ (ml/kg/min <sup>-1</sup> )	45.2 $\pm$ 2.3	49.3 $\pm$ 7.2	63.6 $\pm$ 6.6*	44.9 $\pm$ 2.5
HRmax (bpm)	191 $\pm$ 5	189 $\pm$ 11	183 $\pm$ 7*	188 $\pm$ 7
blood lactate (mmol/l)	11.5 $\pm$ 1.4	11.0 $\pm$ 2.6	8.3 $\pm$ 2.1*	11.1 $\pm$ 1.5
track trial test (s)	785 $\pm$ 44	787 $\pm$ 59	751 $\pm$ 49*	793 $\pm$ 44

T1 to T4: test sessions. \* Significantly different from T1 ( $P<0.05$ )



**Fig. 1** Training load increment along 9 weeks of training (mean  $\pm$  SD). TRIMPS: training impulses; T1: baseline; T2: intermediate load; T3: maximal load; T4: recovery.

	T1	T2	T3	T4
erythrocytes (cells × 10 <sup>12</sup> l <sup>-1</sup> )	4.9 ± 0.3	4.8 ± 0.2	4.9 ± 0.3	5.1 ± 0.2
Hb (g dl <sup>-1</sup> )	15.2 ± 0.8	14.9 ± 0.4	15.3 ± 0.6	15.7 ± 0.7
Hct (%)	43.7 ± 2.3	43.1 ± 1.7	43.6 ± 1.6	45.8 ± 2.0
total leukocytes (cells × 10 <sup>9</sup> l <sup>-1</sup> )	6.0 ± 1.5	6.4 ± 2.5	5.8 ± 1.5	6.2 ± 1.9
neutrophils (%)	53.5 ± 9.8	54.3 ± 10.8	55.0 ± 10.6	49.7 ± 10.0
neutrophils (cells × 10 <sup>9</sup> l <sup>-1</sup> )	3.2 ± 1.0	3.6 ± 2.3	3.2 ± 1.2	3.2 ± 1.4
lymphocytes (%)	35.5 ± 9.2	33.9 ± 9.9	34.2 ± 9.5	37.1 ± 6.9
lymphocytes (cells × 10 <sup>9</sup> l <sup>-1</sup> )	2.1 ± 0.7	2.0 ± 0.4	1.9 ± 0.7	2.2 ± 0.7
monocytes (%)	8.1 ± 1.4	8.6 ± 1.2	7.8 ± 1.2	9.6 ± 3.0
monocytes (cells × 10 <sup>9</sup> l <sup>-1</sup> )	0.5 ± 0.1	0.5 ± 0.2	0.4 ± 0.0	0.5 ± 0.2
basophils (%)	0.3 ± 0.1	0.5 ± 0.1	0.4 ± 0.1	0.4 ± 0.2
basophils (cells × 10 <sup>9</sup> l <sup>-1</sup> )	0.0 ± 0.1	0.0 ± 0.1	0.0 ± 0.0	0.0 ± 0.0
eosinophils (%)	2.4 ± 1.0	2.6 ± 1.1	2.3 ± 1.0	2.9 ± 2.7
eosinophils (cells × 10 <sup>9</sup> l <sup>-1</sup> )	0.1 ± 0.0	0.1 ± 0.0	0.1 ± 0.0	0.2 ± 0.3

Hb: haemoglobin; Hct: hematocrit, T1 to T4: test sessions

**Table 2** Haematological parameters (mean ± SD) analyzed during the entire training protocol.

**Table 3** Biochemical data (mean ± SD) obtained during the entire training protocol.

	T1	T2	T3	T4
LDH (U/l)	243 ± 81	275 ± 51	214 ± 56	200 ± 76
ALT (U/l)	22.9 ± 8.9	20.4 ± 10.6	18.5 ± 6.4	19.0 ± 6.0
AST (U/l)	33.7 ± 21.2	31.9 ± 32.3	25.9 ± 5.7	24.4 ± 6.6
albumin (g/dl)	4.3 ± 0.2	4.0 ± 0.3	4.3 ± 0.2	4.3 ± 0.1
direct bilirubin (mg/dl)	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.1	0.1 ± 0.1
creatinine (mg/dl)	0.9 ± 0.1	0.8 ± 0.1	0.8 ± 0.1	0.9 ± 0.1
uric acid (μmol/l)	356 ± 68	314 ± 34	334 ± 51	362 ± 73
IL-6 (pg/mL)	0.9 ± 0.8	1.2 ± 3.7	0.4 ± 0.6	0.9 ± 1.2
TNF-alpha (pg/mL)	1.1 ± 0.3	0.9 ± 0.4	1.0 ± 0.2	1.0 ± 0.4
MPO (ng/mL)	4.0 ± 2.4	4.5 ± 3.3	4.7 ± 3.5	4.5 ± 1.9
-SH (nmol/mg prot)	45.6 ± 7.8	44.4 ± 7.1	46.5 ± 5.3	45.6 ± 7.9
protein carbonyls (%)	100 ± 4	99 ± 5	98 ± 3	100 ± 2
MDA (%)	100 ± 6	101 ± 4	97 ± 5	100 ± 2

LDH: lactate dehydrogenase, ALT: alanine transaminase, AST: aspartate transaminase, IL-6: interleukine 6, TNF-alpha: tumour necrosis factor alpha, MPO: myeloperoxidase, -SH: sulfhydryl groups, MDA: malondialdehyde. T1 to T4: test sessions

**Table 4** Hormonal data (mean ± SD) obtained during the entire training protocol.

	T1	T2	T3	T4
testosterone (pmol/l)	45.9 ± 17.5	43.8 ± 10.8	40.0 ± 9.7	39.2 ± 14.8
cortisol (nmol/l)	3.4 ± 1.5	3.5 ± 1.6	3.8 ± 1.3	4.8 ± 2.8
FTCR	18.3 ± 13.9	15.2 ± 7.7	12.0 ± 4.2	11.3 ± 7.0

FTCR: Free Testosterone to Cortisol ratio. T1 to T4: test sessions

measurements [8, 16, 25, 36]. Along the same lines, no changes were observed for the biochemical parameters measured in this study. The cytokine hypothesis of overtraining proposes that trauma to the musculoskeletal system leading to a local inflammatory response is the initiating event in the development of overtraining [36]. Finally, inadequate recovery and a continuation of the athletes' training regimen result in the release of inflammatory mediators and the subsequent release of pro-inflammatory cytokines. Confirming previous research, plasma IL-6 and TNF-α concentrations remained unchanged [16]. So, although the underlying causative mechanism/s of overreaching and overtraining is still unclear, it does not appear that elevations in circulating cytokines are primarily responsible for the fatigue associated with long-duration, high-intensity exercise. Impaired antioxidant capacity and increased oxidative stress

**Table 5** POMS scales, STAI-S, and RESTQ-Sport scales (mean ± SD) measured during the entire training protocol.

	T1	T2	T3	T4
<b>POMS</b>				
tension	6.0 ± 3.1	2.9 ± 3.9	10.7 ± 5.1*	3.6 ± 3.2
depression	8.0 ± 9.4	5.2 ± 4.4	10.6 ± 3.8	5.1 ± 4.3
anger	6.0 ± 8.1	4.9 ± 3.9	10.8 ± 5.5	6.8 ± 4.9
vigour	18.3 ± 6.9	18.7 ± 6.5	13.7 ± 6.8	21.5 ± 2.9
fatigue	6.2 ± 4.8	7.6 ± 3.7	16.3 ± 6.5*	4.0 ± 2.4
confusion	2.9 ± 3.7	1.7 ± 2.2	6.5 ± 2.4*	1.1 ± 3.3
total mood disturbance	111 ± 27	104 ± 13	141 ± 16*	99 ± 15
<b>STAI-S</b>				
STAI-S	14.0 ± 7.6	15.1 ± 6.3	23.0 ± 4.3*	15.5 ± 5.9
<b>RESTQ-Sport</b>				
total stress	1.2 ± 0.7	1.5 ± 0.7	2.2 ± 0.7*	1.1 ± 0.7
total recovery	3.4 ± 0.8	3.5 ± 0.7	2.9 ± 0.6	3.9 ± 0.7
general stress	1.4 ± 0.6	1.4 ± 0.5	2.3 ± 0.5*	1.2 ± 0.6
general recovery	3.8 ± 1.0	3.9 ± 0.8	3.2 ± 0.7	3.6 ± 0.9
sport-specific stress	1.0 ± 0.9	1.5 ± 1.0	2.2 ± 1.1*	1.2 ± 0.9
sport-specific recovery	2.8 ± 0.8	3.0 ± 0.7	2.6 ± 0.8	3.2 ± 1.2

T1 to T4: test sessions. \* Significantly different from T1 ( $P < 0.05$ )

could be also associated with fatigue [35], and might be related to the overtraining syndrome [10]. Previous studies found that overtraining status was associated with an increase in resting oxidative stress, reflected in higher protein carbonyl concentrations in overtrained athletes compared to trained controls. Furthermore, responses to acute exercise-induced stress were diminished in overtrained athletes [40]. However, the intense training program here proposed had no effects on MDA, protein carbonyls, and SH groups. Moreover, the lack of significant changes in serum enzymes observed in the present research seems to indicate that the organic systems involved in the training adaptation/regeneration cope well with the training demands of an intensified training period in the young healthy participants.

A similar trend was observed for the hormonal responses. Although cortisol tended to increase and testosterone tended to decrease, no statistically significant differences were found over the whole training season. Several studies have previously investigated the hormonal response in overtraining, but widely conflicting results have been reported and no useful quantitative criteria have been defined [42]. In fact, it is difficult to compare these studies, since there are many variables, such as the defini-

tion of training status, the diurnal variation, and the glycogen status, to consider when interpreting hormonal data. Our results, in agreement with others, suggest that hormonal markers should not be used as an early marker of overreaching in a practical training environment [8].

If functional parameters showed improvements and there were no significant modifications in biomarkers, changes in different scales of the STAI, POMS and RESTQ-Sport indicated the existence of psychological alterations and suggested a risk of overtraining. In 1999, McKenzie stated that overtrained athletes may be firstly identified by a change in psychological markers, followed by medical condition and then performance outcome [27]. Coutts et al. [8] also found changes in the RESTQ-Sport, with a significant increase in total stress and a reduction in total recovery, after an intensive training in comparison to a normal training. These authors proposed that the overtraining syndrome may be related to other factors rather than large training loads, and suggested that psychological measures, together with performance, are useful indicators of overreaching. Nevertheless, although Coutts et al. [8] registered lower performance outcomes, in the present study the performance improved at the end of the training. This discrepancy is possibly due to the fact that in the mentioned study there were 3 different types of exercises during training: swimming, cycling and running, which involves more exercise-related stress. Apart from that, Coutts et al. designed a training protocol aiming to purposely overreach the participants, while our study was focused on investigating the impact of an intensified training protocol [8]. In fact, the unfavourable pattern of stress and recovery was mainly apparent in both the general and the sport-specific stress scales. This provides an indication that the focus should be on helping athletes with subjective stress. In any case, although psychometric instruments provide useful and rapid information, they do not provide the final diagnosis that someone is overtrained, which requires under-recovery over a longer period of time and other indicators such as a chronic performance plateau that cannot be positively influenced by short amounts of rest [21].

A limitation of our research is the absence of a control group which enabled us to 'filter' out threats to experimental validity like learning and habituation effects on tests. Moreover, the number of subjects participating in the study was small. However, in spite of the possibility of type II errors development, sample size was enough to confirm that training was becoming maladaptive and to demonstrate that psychological instruments were sensitive enough to detect signs of overload. From the data obtained it became obvious that a short-period intensified training program can induce several performance improvements in a few weeks time, however, the participant's perceived effort should be taken into account in order to avoid an undesirable stress-related situation. In summary, we can conclude that psychological questionnaires may be a useful means to detect early overreaching even though when it is disguised by positive performance outcomes, and that longitudinal monitoring of changes in stress and recovery may be useful for detection of overtraining in its early stages.

### Acknowledgements

▼  
This work was supported by the Acción Estratégica Sobre el Deporte, Spain (grants n° 2006-56141-C03-01 to J. G., n° 2006-56141-C03-02 to S.M., and n° 2006-56141-C03-03 to F.S.).

### References

- 1 Asikainen TM, Miilunpalo S, Kukkonen-Harjula K, Nenonen A, Pasanen M, Rinne M, Uusi-Rasi K, Oja P, Vuori I. Walking trials in postmenopausal women: effect of low doses of exercise and exercise fractionation on coronary risk factors. *Scand J Med Sci Sports* 2003; 13: 284–292
- 2 Banister EW, Calvert TW, Savage MV, Bach TM. A systems model of training for athletic performance. *Aust J Sports Med* 1975; 7: 57–61
- 3 Blair SN, Morris JN. Healthy hearts and the universal benefits of being physically active: physical activity and health. *Ann Epidemiol* 2009; 19: 253–256
- 4 Borg G. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc* 1982; 14: 377–381
- 5 Bresciani G, Cuevas MJ, Garatachea N, Molinero O, Almar M, de Paz JA, Márquez S, González-Gallego J. Monitoring biological and psychological measures during an entire season in male handball players. *Eur J Sport Sci* 2010; 6: 377–384
- 6 Brink MS, Visscher C, Arends S, Zwerver J, Post W, Lemmink K. Monitoring stress and recovery: New insights for the prevention of injuries and illnesses in elite youth soccer players. *Br J Sports Med* 2010; 44: 809–815
- 7 Brink MS, Visscher C, Coutts A, Lemmink K. Changes in perceived stress and recovery in overreached young elite soccer players. *Scand J Med Sci Sports* doi: 10.1111/j.1600-0838.2010.01237.x
- 8 Coutts AJ, Wallace LK, Slatery K. Monitoring changes in performance, physiology, biochemistry, and psychology during overreaching and recovery in triathletes. *Int J Sports Med* 2007; 28: 125–134
- 9 Filaire E, Bernain X, Sagnol M, Lac G. Preliminary results on mood state, salivary testosterone: cortisol ratio and team performance in a professional soccer team. *Eur J Appl Physiol* 2001; 86: 179–184
- 10 Finaud J, Lac G, Filaire E. Oxidative stress: Relationship with exercise and training. *Sports Med* 2006; 36: 327–358
- 11 González-Boto R, Molinero O, Márquez S. Psychological models of overtraining: a perspective and a new model. *Ansiedad Estres* 2008; 14: 221–237
- 12 González-Boto R, Salguero A, Tuero C, González-Gallego J, Márquez S. Monitoring the effects of training load changes on stress and recovery in swimmers. *J Physiol Biochem* 2008; 64: 19–26
- 13 González-Boto R, Salguero A, Tuero C, Márquez S, Kellmann M. Spanish adaptation and analysis by structural equation modelling of an instrument for monitoring overtraining: The Recovery-Stress Questionnaire (RESTQ-Sport). *Soc Behav Personality* 2008; 36: 636–650
- 14 González-Boto R, Salguero A, Tuero C, Márquez S. Validez concurrente de la versión española del Cuestionario de Recuperación-Estrés para Deportistas (RESTQ-Sport) [Concurrent validity of the Spanish version of the Recovery-Stress Questionnaire for Athletes (RESTQ-Sport)]. *Rev Psicol Dep* 2009; 18: 53–72
- 15 Gormley SE, Swain DP, High R, Spina RJ, Dowling EA, Kotipalli US, Gandrakota R. Effect of intensity of aerobic training on VO2max. *Med Sci Sports Exerc* 2008; 40: 1336–1343
- 16 Halson SL, Lancaster GI, Jeukendrup AE, Gleeson M. Immunological responses to overreaching in cyclists. *Med Sci Sports Exerc* 2003; 35: 854–861
- 17 Harriss DJ, Atkinson G. International Journal of Sports Medicine – Ethical Standards in Sport and Exercise Science Research. *Int J Sports Med* 2009; 30: 701–702
- 18 Hynynen E, Uusitalo A, Kontinen N, Rusko H. Heart rate variability during night sleep and after awakening in overtrained athletes. *Med Sci Sports Exerc* 2006; 38: 313–317
- 19 Jiménez-Jiménez R, Cuevas MJ, Almar M, Lima E, García-López D, De Paz JA, González-Gallego J. Eccentric training impairs NF-kappaB activation and over-expression of inflammation-related genes induced by acute eccentric exercise in the elderly. *Mech Ageing Dev* 2008; 129: 313–321
- 20 Kang HS, Gutin B, Barbeau P, Owens S, Lemmon CR, Allison J, Litaker MS, Le NA. Physical training improves insulin resistance syndrome markers in obese adolescents. *Med Sci Sports Exerc* 2002; 34: 1920–1927
- 21 Kellmann M. Preventing overtraining in athletes in high-intensity sports and stress-recovery monitoring. *Scand J Med Sci Sports* 2010; 20: 95–102
- 22 Kellmann M, Günther KD. Changes in stress and recovery in elite rowers during preparation for the Olympic Games. *Med Sci Sports Exerc* 2000; 32: 676–683
- 23 Kellmann M, Kallus KW. Recovery-Stress Questionnaire for Athletes: User Manual. Champaign, IL: Human Kinetics; 2001
- 24 Kenttä G, Hassmén P. Overtraining and recovery. A conceptual model. *Sports Med* 1998; 26: 1–16
- 25 Mackinnon LT. Immunity in athletes. *Int J Sports Med* 1997; 1: S62–S68

- 26 Main LC, Dawson B, Heel K, Grove JR, Landers GJ, Goodman C. Relationship between inflammatory cytokines and self-report measures of training overload. *Res Sports Med* 2010; 18: 127–139
- 27 McKenzie DC. Markers of excessive exercise. *Can J Appl Physiol* 1999; 24: 66–73
- 28 McNair DM, Lorr M, Droppleman LF. Manual for Profile of Mood States Questionnaire. San Diego California: Educational and Industrial Testing Service; 1971
- 29 Mujika I, Padilla S. Cardiorespiratory and metabolic characteristics of detraining in humans. *Med Sci Sports Exerc* 2001; 33: 413–421
- 30 Mujika I, Goya A, Ruiz E, Grijalba A, Santisteban J, Padilla S. Physiological and performance responses to a 6-day taper in middle-distance runners: influence of training frequency. *Int J Sports Med* 2002; 23: 367–373
- 31 Nederhof E, Lemmink K, Zwerver J, Mulder T. The effect of high load training on psychomotor speed. *Int J Sports Med* 2007; 28: 595–601
- 32 Nederhof E, Zwerver J, Brink M, Meeusen R, Lemmink K. Different diagnostic tools in nonfunctional overreaching. *Int J Sports Med* 2008; 29: 590–597
- 33 O'Donovan G, Blazevich AJ, Boreham C, Cooper AR, Crank H, Ekelund U, Fox KR, Gately P, Giles-Corti B, Gill JM, Hamer M, McDermott I, Murphy M, Mutrie N, Reilly JJ, Saxton JM, Stamatakis E. The ABC of Physical Activity for Health: a consensus statement from the British Association of Sport and Exercise Sciences. *J Sports Sci* 2010; 28: 573–591
- 34 Radak Z, Chung HY, Goto S. Systemic adaptation to oxidative challenge induced by regular exercise. *Free Radic Biol Med* 2008; 44: 153–159
- 35 Smirnova IV, Pall ML. Elevated levels of protein carbonyls in sera of chronic fatigue syndrome patients. *Mol Cell Biochem* 2003; 248: 93–95
- 36 Smith LL. Tissue trauma: the underlying cause of overtraining syndrome? *J Strength Cond Res* 2004; 18: 185–193
- 37 Spielberger CD, Gorsuch RL, Lushene RE. STAI manual for the State-Trait Anxiety Inventory (Self-Evaluation questionnaire). Palo Alto, CA: Consulting Psychology Press; 1970
- 38 Spiebelger CD, Gorsuch RL, Lushene RE. Cuestionario de ansiedad estado-rasgo [State-trait anxiety questionnaire]. TEA Ediciones. Madrid; 1982
- 39 Swain DP, Franklin BA. Comparison of cardioprotective benefits of vigorous versus moderate intensity aerobic exercise. *Am J Cardiol* 2006; 97: 141–147
- 40 Tanskanen M, Atalay M, Uusitalo A. Altered oxidative stress in overtrained athletes. *J Sports Sci* 2010; 28: 309–317
- 41 Urhausen A, Gabriel H, Kindermann W. Blood hormones as markers of training stress and overtraining. *Sports Med* 1995; 20: 251–276
- 42 Urhausen A, Kindermann W. Diagnosis of overtraining. What tools do we have? *Sports Med* 2002; 32: 95–102