

Kansas Agricultural Experiment Station Research Reports

Volume 10
Issue 2 *Southeast Research and Extension*
Center Agricultural Research

Article 8

2024

Role of Soil Management in Control of Soil-Borne Diseases

Gretchen F. Sassenrath
Kansas State University, gsassenrath@ksu.edu

Christopher R. Little
Kansas State University, crlittle@ksu.edu

Xiaomao Lin
Kansas State University, xlin@ksu.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Agronomy and Crop Sciences Commons](#), and the [Plant Pathology Commons](#)

Recommended Citation

Sassenrath, Gretchen F.; Little, Christopher R.; and Lin, Xiaomao (2024) "Role of Soil Management in Control of Soil-Borne Diseases," *Kansas Agricultural Experiment Station Research Reports*: Vol. 10: Iss. 2. <https://doi.org/10.4148/2378-5977.8577>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2024 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Role of Soil Management in Control of Soil-Borne Diseases

G.F. Sassenrath, C. Little,¹ and X. Lin²

Summary

Soil-borne diseases are a significant cause of crop yield reduction. Alternative soil-management methods can enhance the soil's natural disease-controlling organisms. This study explores the effect of alternative production methods on a primary soybean disease, charcoal rot, caused by the fungus *Macrophomina phaseolina*. Treatments that could potentially enhance or reduce the disease pressure were implemented, and soil tests were conducted for nutrients, soil properties, and disease presence. Manure increased the nutrient levels in the soil, as expected, but did not influence disease control. Solarization increased the temperature within the plots and increased the number of colony-forming units (CFUs) of *M. phaseolina*.

Introduction

Suppressive soils have been defined as soils that can inhibit the growth of naturally occurring soil-borne diseases. These soils can suppress or control disease-causing organisms, including fungi (e.g., *Fusarium virguliforme* (cause of sudden death syndrome, SDS), *Macrophomina phaseolina* (cause of charcoal rot), *Phytophthora* root rot (*Phytophthora sojae*), and nematodes (e.g., soybean cyst nematode, *Heterodera glycines*)). How the native soil microbial communities reduce disease is not known. Knowledge of factors that contribute to and support these beneficial microbial communities is also unknown.

One example of this natural improvement in the soil microbial community reducing disease was demonstrated in previous research (Sassenrath et al., 2017, 2019), which demonstrated that a high-glucosinolate mustard (*Brassica juncea*) reduced fungal populations that caused charcoal rot in both soil and soybean plants. Here, those results are expanded on by exploring the interaction between soil health and disease pressure. Management practices that increase, decrease, or maintain disease pressure were tested in field studies to determine the influence of those practices on soil health, fungal pathogen presence, and soybean growth and yield.

Impact of Soil Health on Soybean Disease

Crop plants that are disease hosts increase the number of disease-causing organisms in the soil. The increase in colony forming units (CFU) of *M. phaseolina* in the soil after

¹ Department of Plant Pathology, College of Agriculture, Kansas State University.

² Department of Agronomy, College of Agriculture, Kansas State University.

soybean production has been shown previously (Sassenrath et al., 2019). Other factors that reduce soil-borne diseases include high-glucosinolate mustard as a cover crop (Sassenrath et al., 2017, 2019) and increasing the soil temperature (e.g., “solarization”). Use of animal manures greatly increases the diversity of the soil microbial community, and beneficial microorganisms in particular. In addition to improving the nutrient balance of the soil, manure may contribute to reduced disease pressure (Graham et al., 2014) and soybean cyst nematode populations (Bao et al., 2013).

Diseases are primary factors that reduce the yield and quality of soybeans in Kansas and throughout the world. Soil-borne diseases are prevalent in eastern Kansas crop fields. Certain plants have been shown to produce chemicals that act as biofumigants that control or reduce harmful soil fungi. Animal manures have also been used to alter the soil microbiome to improve control of disease organisms. The working hypothesis of this research is that improving the overall soil health by supporting healthy soil microbial communities can reduce disease pressure, i.e., creating suppressive soils by altering management practices will reduce disease pressure. This research explores the relationship between soil health and disease pressure. The research tests the ability of cover crops, animal manure, and solarization to control or reduce charcoal rot in soybean production through improved soil microbial communities.

Experimental Procedures

Replicated plots were established at the Southeast Research and Extension Center in Parsons in the spring, 2023. Plots included: fallow, mustard cover crop, soybean, corn stubble, cow manure, and plastic sheets. Temperature and moisture sensors (HOBO, Onset, Inc., Bourne, MA) were installed at 2-in. depth in the soil, and temperature and moisture were recorded continuously. Plastic sheets provide a “solarization” treatment, increasing soil temperature and potentially reducing soil microbes. Plastic sheets were placed on plots and held in place with concrete blocks. Corn stubble was spread to about a 2-in. layer; corn stubble provides more carbon for soil microbes, increasing their abundance, but is also a host for *M. phaseolina*, potentially increasing the disease prevalence. Animal manure provides an additional food source for the microbes and adds additional microbes to the soil; manure has been shown to reduce some pathogens in soil. Cow manure was spread to about a 2-in. layer on the plots. The use of a mustard cover crop has been shown to reduce the number of CFUs of *M. phaseolina*, while soybeans are a host and increase the CFUs of *M. phaseolina*. The high-glucosinolate mustard, Might Mustard Pacific Gold (Johnny’s Selected Seeds, Winslow, ME) was broadcast in plots in early April. The fallow treatment was left unplanted and served as a control. Five cultivars of soybeans, ranging from maturity group (MG) 3.2 to 5.2, were also planted to test for variation in charcoal rot sensitivity.

Soil samples to a depth of 6 in. were collected in spring prior to implemented treatments, mid-season, and after harvest. Soils were analyzed for nutrients and microbial activity, and for the number of CFUs of *M. phaseolina*. Soil microbial activity was measured with the CO₂-burst test (Solvita, Woods End, ME).

Results and Discussion

Nutrients changed in response to treatments (Table 1). As anticipated, the cow manure treatment had the highest levels of organic matter (OM), potassium (K), and phos-

phorus (P). Surprisingly, the solarization treatment (plastic film) had the highest total nitrogen (N) and microbiologically active carbon.

Interesting differences were observed in the environment within the soil in the different treatments (Figure 1). The soil with animal manure retained the most moisture (Figure 1), most likely due to the coverage of the soil, reducing evaporation from the soil surface. Similarly, corn stalks and plastic reduced evaporative losses and retained soil moisture. Mustard seed and soybeans removed soil moisture, resulting in soil moisture levels that were lower than those measured in the fallow treatment. The solarization treatment (plastic) increased soil temperature an average of 15°F above that of bare soil (fallow) (Figure 2). Cow manure and corn stubble kept the soil an average 8°F cooler. Plants (soybeans and mustard seed cover crop) also kept the soil cooler, but not as cool as the corn stubble and manure treatments.

The soil microbial activity, as measured by the Solvita CO₂-burst test, was very high in the corn stubble and manure, as would be expected from the high carbon content of those materials (Figure 3). The soil treated with cow manure also had much higher microbial activity, but the activity under the corn stubble was less than that planted to soybeans. Microbial activity with the mustard seed cover crop was also low. The fallow and solarization treatments had the lowest microbial activity at mid-season.

Conclusions

The soil microbiome controls many of the functions of the soil. It is possible, through alternative management practices, to alter the soil microbiome to support helpful organisms, such as arbuscular mycorrhizae, while controlling disease-causing organisms. Preliminary evidence from this study showed a minor change in microbial composition and activity with treatments. However, the unusually hot and dry weather may have compromised the results, as soybean yields were greatly reduced.

Acknowledgements

This research is supported by funding from the Kansas Soybean Commission and the U.S. Department of Agriculture National Institute of Food and Agriculture, Hatch project 1018005.

References

- Bao, y., Chen, S., Vetsch, J., Randall, G. 2013. Soybean yield and *Heterodera glycines* responses to liquid swine manure in nematode suppressive soil and conducive soil. *J. Nematology*. 45(1):21-29.
- Graham, E., Grandy, S., Thelen, M. 2014. Manure effects on soil organisms and soil quality. Michigan State Extension. <https://www.canr.msu.edu>
- Sassenrath, G.F., Lingenfelter, J., Lin, X. 2023. Corn and soybean production – 2022 summary. *Kansas Agricultural Experiment Station Research Reports*: Vol. 9: Iss. 2.
- Sassenrath, G.F., Little, C., Roozeboom, K., Lin, X., Jardine, D. 2019. Controlling soil-borne disease in soybean with a mustard cover crop. *Kansas Agricultural Experiment Station Research Reports*: Vol. 5: Iss. 2. <https://doi.org/10.4148/2378-5977.7740>
- Sassenrath, G.F., Little, C.R., Hsiao, C.-J., Shoup, D.E., Lin, X. 2017. Cover crop system to control charcoal rot in soybeans. *Kansas Agricultural Experiment Station Research Reports*: Vol. 3: Iss. 2. <https://doi.org/10.4148/2378-5977.1383>

Table 1. Changes in soil nutrients with treatment

	Organic matter	Potassium	Mehlich P	Total N
	%	----- ppm -----		
Mid-season				
Corn stubble	1.9	74.5	23.0	13.4
Cow manure	2.4	150.5	47.3	16.1
Fallow	2.0	66.8	18.5	13.2
Mustard seed	2.1	78.5	23.0	15.6
Plastic film	2.0	62.3	19.5	21.2
Soybeans	2.0	73.8	20.3	12.9
At harvest				
Corn stubble	2.0	80.3	27.3	19.6
Cow manure	2.2	160.3	53.0	27.8
Fallow	2.0	60.8	25.8	20.1
Mustard seed	2.0	74.5	29.0	21.8
Plastic film	2.0	60.3	29.0	33.3
Soybeans	2.0	71.8	25.5	19.2




Figure 1. Difference in soil moisture at 2-in. soil depth between fallow and different treatments.




Figure 2. Difference in soil temperature (°F) at 2-in. soil depth between fallow and treatments.




Figure 3. CO₂ burst test for different treatments.