



**Michigan
Technological
University**

Michigan Technological University
Digital Commons @ Michigan Tech

Michigan Tech Publications, Part 2

1-19-2024

Non-invasive methods for monitoring weasels: emerging technologies and priorities for future research

David S. Jachowski
Clemson University

Scott M. Bergeson
Purdue University Fort Wayne

Stacy Cotey
Michigan Technological University, srcotey@mtu.edu

Elizabeth Croose
The Vincent Wildlife Trust

Tim R. Hofmeester
Sveriges lantbruksuniversitet

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.mtu.edu/michigantech-p2>



Part of the [Forest Sciences Commons](#)

Recommended Citation

Jachowski, D., Bergeson, S., Cotey, S., Croose, E., Hofmeester, T., MacPherson, J., & et al. (2024). Non-invasive methods for monitoring weasels: emerging technologies and priorities for future research. *Mammal Review*. <http://doi.org/10.1111/mam.12344>
Retrieved from: <https://digitalcommons.mtu.edu/michigantech-p2/434>

Follow this and additional works at: <https://digitalcommons.mtu.edu/michigantech-p2>



Part of the [Forest Sciences Commons](#)

Authors

David S. Jachowski, Scott M. Bergeson, Stacy Cotey, Elizabeth Croose, Tim R. Hofmeister, Jenny MacPherson, and et al.

REVIEW

Non-invasive methods for monitoring weasels: emerging technologies and priorities for future research

David S. JACHOWSKI*†  Department of Forestry and Environmental Conservation, Clemson University, Clemson, SC, USA. Email: djachow@clemson.edu

Scott M. BERGESON† Department of Biological Sciences, Purdue University Fort Wayne, 2101 E. Coliseum Blvd, Fort Wayne, IN 46818, USA. Email: bergesos@pfw.edu

Stacy R. COTEY† College of Forest Resources and Environmental Science, Michigan Technological University, 1400 Townsend Drive, Houghton, MI 49931, USA. Email: srcotey@mtu.edu

Elizabeth CROOSE† Vincent Wildlife Trust, 3–4 Bronsil Courtyard, Eastnor, Ledbury, Herefordshire HR8 1EP, UK and British Small Animal Veterinary Association, Woodrow House, 1 Telford Way, Waterwells Business Park, Gloucester GL2 2AB, UK. Email: lizziecroose1@gmail.com

Tim R. HOFMEESTER† Department of Wildlife, Fish, and Environmental Studies, Swedish University of Agricultural Sciences, Skogsmarksgränd 7, SE-907 36 Umeå, Sweden and Stichting Kleine Marters/ Small Mustelid Foundation, Goudreinette 10, 6922 AE, Duiven, The Netherlands. Email: tim.hofmeester@slu.se

Jenny MACPHERSON† Vincent Wildlife Trust, 3–4 Bronsil Courtyard, Eastnor, Ledbury, Herefordshire HR8 1EP, UK. Email: jenny.macpherson@vwt.org.uk

Patrick WRIGHT† Vincent Wildlife Trust, 3–4 Bronsil Courtyard, Eastnor, Ledbury, Herefordshire HR8 1EP, UK. Email: patrick.wright@vwt.org.uk

Camilo A. CALDERÓN-ACEVEDO Department of Environmental and Forest Biology, College of Environmental Science and Forestry, State University of New York, 1 Forestry Drive, Syracuse, NY 13210, USA. Email: cacalder@esf.edu

Stephen P. CARTER Vincent Wildlife Trust, 3–4 Bronsil Courtyard, Eastnor, Ledbury, Herefordshire HR8 1EP, UK. Email: steve.carter@vwt.org.uk

Andrin C. DÜRST Division of Conservation Biology, Institute of Ecology and Evolution, University of Bern, Baltzerstrasse 6, 3012 Bern, Switzerland. Email: andrin.duerst@unibe.ch

Gregory B. EGLOFF Division of Conservation Biology, Institute of Ecology and Evolution, University of Bern, Baltzerstrasse 6, 3012 Bern, Switzerland. Email: gregory.egloff@students.unibe.ch

M. Kevin HAMED Department of Fish and Wildlife Conservation, Virginia Tech, 101 Cheatham Hall, Blacksburg, VA 24061, USA. Email: khamed@vt.edu

Paul HAPEMAN Department of Biology, Central Connecticut State University, 343 Copernicus Hall, New Britain, CT 06050, USA. Email: hapeman@ccsu.edu

Stephen N. HARRIS Department of Forestry and Environmental Conservation, Clemson University, Clemson, SC, USA. Email: sh2@g.clemson.edu

Kendyl HASSLER Florida Fish and Wildlife Conservation Commission, 1105 SW Williston Rd, Gainesville, FL 32601, USA. Email: kendyl.hassler@gmail.com

Jean-Yves HUMBERT Division of Conservation Biology, Institute of Ecology and Evolution, University of Bern, Baltzerstrasse 6, 3012 Bern, Switzerland. Email: jean-yves.humbert@unibe.ch

Denise KARP Artenspürhunde Schweiz, Fährweg 45, CH-4600 Olten, Switzerland. Email: karpdenise@gmail.com

Roland KAYS North Carolina Museum of Natural Sciences, 11 West Jones St, Raleigh, NC 27601, USA and Department of Forestry and Environmental Resources, North Carolina State University, Raleigh, NC 27606, USA. Email: rwkays@ncsu.edu

Jelena MAUSBACH Artenspürhunde Schweiz, Fährweg 45, CH-4600 Olten, Switzerland. Email: jelena.mausbach@gmx.eu

Dana J. MORIN *Department of Wildlife, Fisheries and Aquaculture, Mississippi State University Forest and Wildlife Research Center, Box 9690, Mississippi State, MS 39762, USA. Email: dm3090@msstate.edu*

Jeroen MOS *Stichting Kleine Marters/Small Mustelid Foundation, Goudreinette 10, 6922 AE, Duiven, The Netherlands. Email: info@mos-ecologie.nl*

Salvador Salvador ALLUÉ *Department of Environmental Science, University of Girona, Maria Aurèlia Capmany 69, 17003 Girona, Spain. Email: ssallue@gmail.com*

Lisa SMITH *Florida Fish and Wildlife Conservation Commission, 1105 SW Williston Rd, Gainesville, FL 32601, USA. Email: lisa.smith@myfwc.com*

Joshua P. TWINING *New York Cooperative Fish and Wildlife Research Unit, Department of Natural Resources and the Environment, Cornell University, Fernow Hall, Ithaca, NY 14853, USA. Email: jpt93@cornell.edu*

Chad R. WILLIAMSON *Department of Biology, Minot State University, 500 University Ave. W, Minot, ND 58707, USA. Email: chad.williamson@minotstateu.edu*

Karol ZUB *Mammal Research Institute Polish Academy of Sciences, Stoczek 1, 17-230 Białowieża, Poland. Email: karolzub@ibs.bialowieza.pl*

Keywords

camera trap, *Mustela*, *Neogale*, non-invasive monitoring

*Correspondence

†These authors shared equally in leading the manuscript.

Received: 28 August 2023

Accepted: 20 December 2023

Editor: DR

doi: 10.1111/mam.12344

ABSTRACT

1. Weasels (genus *Mustela* and *Neogale*) are of management concern as declining native species in some regions and invasive species in others. Regardless of the need to conserve or remove weasels, there is increasingly a need to use non-invasive monitoring methods to assess population trends.
2. We conducted a literature review and held the first ever International Weasel Monitoring Symposium to synthesise information on historical and current non-invasive monitoring techniques for weasels. We also explored current limitations, opportunities, and areas of development to guide future research and long-term monitoring.
3. Our literature search revealed that in the past 20 years, camera traps were the most commonly used non-invasive monitoring method (62% of studies), followed by track plates or scent stations designed to collect footprints (23%) and walking transects for tracks in snow or soil (8.7%).
4. Experts agreed that the most promising non-invasive monitoring techniques available include use of citizen scientist reporting, detection dogs, detecting tracks, non-invasive genetic surveys, and enclosed or unenclosed camera trap systems. Because each technique has benefits and limitations, using a multi-method approach is likely required.
5. There is a need for strong commitment to dedicated monitoring that is replicated over space and time such that trend data can be ascertained to better inform future management action. The diversity of non-invasive monitoring methods now available makes such monitoring possible with relatively minor commitments of funding and effort.

INTRODUCTION

Weasels (members of the genus *Mustela* and *Neogale*, here primarily focusing on the smaller species long-tailed weasel *Neogale frenata*, stoat *Mustela erminea*, and least weasel *Mustela nivalis*) are small mustelids distributed

across much of the northern hemisphere. Weasels play an important role in ecosystem function (King & Powell 2007), although data from North America (Jachowski et al. 2021) and Europe (Wright et al. 2022) suggest weasels are in decline through portions of their historical range. Where weasels have been introduced,

they can have detrimental impacts through predation and displacement of native species (Pech & Maitland 2016, Rodrigues *et al.* 2017). Whether to conserve or remove weasels, there is a need for standardised, long-term monitoring to determine factors important to their presence, distribution, and use of habitat across multiple spatial scales over time (Jachowski *et al.* 2021).

Broad-scale surveys and long-term monitoring for weasels have largely been limited to regions where there are intensive, sustained efforts to eradicate weasels (Jones *et al.* 2004), or to the northern extent of the range through annual track surveys or lemming nest occupancy rates (Feige *et al.* 2012, Schmidt *et al.* 2012). Trapping harvest data were historically used to monitor weasel (and other furbearer) population trends (Tapper & Reynolds 1996, Aebischer *et al.* 2011), but harvest data are not always proportional to abundance (McDonald & Harris 2002, Maunder *et al.* 2006, Fukasawa *et al.* 2020). In North America, decline of fur markets and regulation of harvest for conservation, further limit availability of these data (Jachowski *et al.* 2021).

A primary challenge in monitoring weasels is that they are difficult to detect. Traditionally, invasive methods such as live-trapping were used to study weasels (King 1980, Zub *et al.* 2008), but high cost and effort required for such methods makes them impractical at spatial or temporal scales meaningful to conservation. Non-invasive methods for small carnivore monitoring are typically less labour-intensive and can be more easily replicated across sites and repeated over time. Historically limited to surveys based on animal tracks (Quick 1944), camera traps have become a common non-invasive method for monitoring small carnivores globally over the past several decades and setups have recently been developed specifically for weasels (Mos & Hofmeester 2020). Other techniques useful for detecting weasels non-invasively include the use of detection dogs (Steury 2012), citizen science (Linzey & Hamed 2016), and most recently, non-invasive molecular techniques (Zielinski *et al.* 2020, Broadhurst *et al.* 2021). With this rapid emergence of multiple potential non-invasive monitoring methods, there is a need for information on which methods are best suited to future research and monitoring.

Here, we summarise recent trends in non-invasive weasel monitoring techniques and highlight priorities for future development. In 2023, we conducted a literature review and held the first International Weasel Monitoring Symposium to synthesise information on historical and current non-invasive monitoring techniques for weasels. We also explored current limitations, opportunities, and areas of development for each technique to guide future research in broad-scale surveys and long-term monitoring.

MATERIALS AND METHODS

We conducted a Web of Science search on 17 April 2023 for published literature on non-invasive monitoring methods for weasels in the wild over a 20-years period (2003–2023) using the following query: TS=(‘Mustela’ OR ‘Neogale’) and TS=(‘monitoring’) AND DT=(Article OR Book OR Book Chapter OR Review) AND PY=(2003–2023). We reviewed titles and abstracts of the 673 papers generated and determined that 69 papers were relevant (i.e. documented weasels in a field setting using a non-invasive method) for subsequent scoring. From these studies we extracted the technique used, year of study, country, focal species, bait type (if any), and population metric calculated from each technique.

On 24 February 2023, we hosted a virtual International Weasel Monitoring Symposium with experts from Europe and North America presenting talks in sessions structured around widely used non-invasive weasel monitoring methods: citizen science, detection dogs, tracks, non-invasive genetic surveys, unenclosed camera traps, and enclosed camera traps. Sessions concluded with structured discussions on history of development, advantages and strengths, current limitations, and areas of future development for each technique. Subsequently, subgroups of experts drafted sections of this manuscript following the symposium topics.

RESULTS

Breadth of non-invasive weasel monitoring techniques

In the past 20 years, camera traps (or video recorders) were the most commonly used non-invasive monitoring method (62% of studies, Fig. 1). Other common methods were track plates or scent stations designed to collect footprints (23% of studies) and walking transects for tracks in snow or soil (8.7%). Less-commonly used methods included transect sampling for scat/faeces, interviews with local residents or use of historical records, hair snares, roadkill surveys, and checking lemming nests for weasel occupancy. Nine of the 69 studies involved multiple non-invasive methods (Fig. 1).

Nearly one third (33%) of studies used some type of bait. Of those studies, a common practice for monitoring weasels (either intentionally or incidentally in monitoring nest fate) was placing cameras at active or artificial bird nests (23% of bait studies), followed by use of rabbit (12%), fish (12%), chicken (7.7%) or some unnamed type of meat (12%). Various other baits used less commonly were eggs, peanuts or peanut butter, mayonnaise, and lure-infused poison blocks. While




Fig. 1. Literature search results for non-invasive weasel monitoring methods used globally over the past 20 years highlighting the frequency with which each type of method was used (top panel) and which focal species were detected (bottom panel). If a specific species was not mentioned in a study, it was tallied as species or 'sp'. The portion of each bar in orange represents publications produced in the past 10 years (since 2013).

numerous lures were commonly used across camera and track-based survey techniques, no individual study in our literature search explicitly compared multiple lure or bait types on detection probability (although see Buyaskas et al. 2020).

Most studies were in North America (22%), Europe (43%) or New Zealand (17%; Fig. 2), and most (38%) reported on *Mustela erminea* (Fig. 1), although several studies targeted multiple species simultaneously. There

were very few ($n < 3$) studies on Asian weasel species *Mustela altaica*, *Mustela itatsi* and *Mustela strigidorsa*, and no studies on weasel species endemic to Africa or South America (although see Cepeda-Duque et al. 2023, published after our literature search). Most studies used methods to estimate species presence or to assess distribution (45%), followed by estimating relative abundance (44%), and species occupancy and/or detection probability (11%).




Fig. 2. Global map with countries highlighted where dedicated studies have taken place on non-invasive weasel monitoring methods in the past 20 years based on our literature review.

Current non-invasive monitoring methods

CITIZEN SCIENCE

HISTORY OF DEVELOPMENT

Citizen science is a collaborative approach to scientific research that involves active participation from the public. This has emerged as a powerful tool, bridging the gap between scientific research and society, with numbers of projects having increased substantially over the last three decades (Davis et al. 2023). Citizen science and participatory volunteer networks have increasingly been utilised to assist with small carnivore research and monitoring (Silvertown 2009). Examples include interviews and questionnaire surveys of local residents to gather data on distribution of weasel species (Lau et al. 2010, Bolduc et al. 2023, Hayder et al. 2023), including an ongoing survey of the Irish stoat (*Mustela erminea hibernica*) based on reported sightings of live or dead animals throughout Ireland (<https://biodiversityireland.ie/surveys/irish-stoat-survey/>).

Data gleaned from social media posts can yield information on species' distribution (Wright et al. 2023), and social media group pages are emerging as a way for members of the public to share information about small carnivores. One example is the Small Carnivore Conservation Project Thailand Facebook Group (<https://www.facebook.com/groups/128334450533090/>), which has generated many valuable records of weasel species from northern Thailand.

ADVANTAGES AND STRENGTHS

A key benefit of citizen science is its ability to increase the capacity of a project. With collective efforts of

numerous volunteers, scientists can gather vast amounts of data over large geographical areas and extended periods, which would be impossible for traditional research teams alone (Conrad & Hilchey 2011). Sightings can be used to refine estimates of population size in combination with other monitoring techniques such as trapping data via integrated removal models (Zub et al. 2022). Wide scale public engagement can potentially enhance the scope of data collection and strengthen the connection between scientific research and society.

CURRENT LIMITATIONS

Challenges of citizen science include ensuring data quality and integrating systematically and non-systematically collected data. For weasels in particular, standardised protocols for data reporting and expert ID are important given the high probability of false positives and species misidentification (Fig. 3). Requiring pictures to be submitted along with observations is critical, as >90% of reported weasel sightings in Florida, USA that included photos have been incorrectly identified (L. Smith, unpublished data).

Citizen science projects face a trade-off between opportunistic sampling, which may generate more samples with few restrictions, and involving citizens in a structured sampling design that may discourage their participation (Shirk et al. 2012). To address habitat and spatial sampling gaps, Lasky et al. (2021) adopted a hybrid sampling design, monitoring progress towards sampling goals during the study and supplementing volunteer efforts with additional field work by project staff to fill gaps. In addition, emerging statistical models, such as integrated species distribution models, enable




Fig. 3. Outreach material produced by Alabama Nongame Wildlife Program (USA) to gain sightings information for long-tailed weasels, which are infrequently reported in their state (illustration credit: Shari Amsel).

integration of presence-only records from citizen science with detection/non-detection data from structured surveys (e.g. Koshkina et al. 2017).

FUTURE DEVELOPMENT

Reliable, user-friendly software is one of the most important aspects of scaling up citizen science. Examples include a mobile phone application for citizen science projects that provides a user interface for administrators without the need for programming skills (Ellul et al. 2013) and SciStarter, which can help project coordinators manage their volunteers (Hoffman et al. 2017).

While some platforms are built for specific projects, others have evolved into more integrated platforms that serve generic projects (Liu et al. 2021). iNaturalist is an online portal and smartphone app that anyone can use to upload photos of mammals for identification, verification and reporting. This currently has 3034 research grade observations for *Neogale frenata*, 1876 for *Mustela nivalis*, and 1679 for *Mustela erminea* worldwide, although these observations likely require further verification given some species cannot be told apart without full body photos in areas of overlap (Kays et al. 2022). Regardless, new analytical approaches such as integrated species distribution

models offer ways to utilise citizen science data to improve our understand of species distribution and environmental relationships (e.g. Gilbert et al. 2021) that could be expanded to weasels.

DETECTION DOGS

HISTORY OF DEVELOPMENT

The earliest use of detection dogs for conservation is thought to be the 1890s in New Zealand for kiwis *Apteryx australis*, *Apteryx owenii* and kakapos *Strigops habroptilus* (Hill & Hill 1987). Since 1930, detection dogs have been used for a minimum of 408 animal species and 42 plants (and fungi and bacteria; Grimm-Seyfarth et al. 2021). In at least 102 cases (72 scientific), dogs were used for detecting mustelids (Grimm-Seyfarth et al. 2021), including long-tailed weasels (Stuery 2012), fishers *Pekania pennanti* (Long et al. 2007, Thompson et al. 2012, Zielinski et al. 2013) and black-footed ferrets *Mustela nigripes* (Reindl-Thompson et al. 2006).

ADVANTAGES AND STRENGTHS

Detection dogs (Fig. 4) are useful for surveying for weasel presence and resulting data have primarily been used to



Fig. 4. Left, Cowboy the detection dog and Kendyl Hassler searching for long-tailed weasel in Florida, USA (photo credit: Lisa Smith). Right, Django the mustelid detection dog (*Lutra lutra*, *Mustela erminea*, *Mustela nivalis*) in Switzerland (photo credit: Denise Karp, Artenspürhunde Schweiz).

understand small carnivore habitat associations (Smith et al. 2006, Zielinski et al. 2013). Scats detected by dogs can be analysed to determine sex, individual identification, physiological measures (such as faecal cortisol and progesterone metabolites), some parasites, and diet (Wasser et al. 2004). Results from detection dog surveys are used to target subsequent trapping efforts, both for conservation and eradication. Dogs can cover a large area systematically in a single visit (compared to point locations of track plates or camera traps), are often able to find scats in areas missed by human observers and can follow scent trails and guide towards spots with increased activity (Egloff et al. 2022, Schenker et al. 2023).

CURRENT LIMITATIONS

While detection dogs are highly effective at locating their targets, there are limitations to their use. As with many other target species (Bennett et al. 2019), there are no standardised guidelines or protocols for training detection dogs or conducting weasel scat surveys. Different dogs and dog-handlers may have different detection rates. While these factors can be included as covariates in subsequent modelling, the diverse attributes of each dog and handler make comparability of survey results among sites and over time difficult. Sourcing of scats for training detection dogs can be difficult as scats used for training are ideally from multiple wild individuals (DeMatteo et al. 2019). Dogs should ideally be trained in a wild setting on untouched scat samples after they have been trained on 'put out' samples, requiring knowledge of areas where weasels are present and the phenotype of weasel scats, which can vary seasonally and regionally. Dogs require at least 6 months of specialised training (Schenker et al. 2023). Unlike passive monitoring approaches like camera traps, dogs in a field setting can only be used for limited hours per day. They require maintenance and health care, and continued motivation if detection rates are low. Climate and terrain may impact the length of time a dog is able to work, as well as its ability to locate scats (Leigh & Dominick 2015). Scat

can degrade quickly in warm and wet conditions and can be removed quickly by dung beetles or other animals (Livingston et al. 2005, Norris & Michalski 2010). Detection distance for weasel scat is low, especially when it has dried out, so dogs need to cover an area extensively. Reed et al. (2011) found that dogs in controlled trials detected >75% of carnivore scats within 10 m of transects, with detection rates decreasing with distance. For weasel scats, maximum detection distance is estimated to be around 3 m, and more often 50 cm with dried scats (D. Karp, personal communication). Collectively, given these issues and the cost involved, it is important to evaluate the return on investment detection dogs provide in monitoring for weasels (Long et al. 2007).

FUTURE DEVELOPMENT

Development of protocols is needed for training, running trials, and search methodology used when working with dogs. There is a need to assess the influence of site-specific conditions (habitat, terrain, weather) on scat detection probability, and account for this variation in modelling of data. Strategic integration of detection dogs with other techniques is another area for further development. For example, where the small carnivorous marsupial *Antechinus arktos* was not previously detected using live-trapping and passive camera-trapping efforts, Thomas et al. (2020) used camera traps at locations where detection dogs had alerted on target odour and confirmed the species in 100% of detections. Finally, more natural history information on weasel scent-marking behaviour is needed to guide targeted searching for scats during detection dog surveys.

TRACKS

HISTORY OF DEVELOPMENT

Footprint track methods were among the first non-invasive techniques for studying weasels (Quick 1944) and were

adapted for surveying stoats in the 1970s (King & Edgar 1977). Methods include observing tracks made by animals in snow and soil (Sundell et al. 2013) or actively setting track plates or footprint tunnels where animals walk through a substance such as ink, graphite, or carbon paper before stepping onto a clean surface to leave prints (Tempero et al. 2007, Červinka et al. 2014).

ADVANTAGES AND STRENGTHS

Track monitoring is mostly used for surveying weasel distribution, occupancy, and habitat use (King & Edgar 1977). Track techniques are generally cheap and easy to implement. Despite the recent growth of camera-based monitoring, a study in northeast North America found that weasels were detected more frequently using track plates than with standard open baited camera traps (Gompper et al. 2006). Thus, track surveys still provide a valuable method for small mammal monitoring, particularly when funding is limited.

CURRENT LIMITATIONS

Track quality can be influenced by weather or field conditions. It is important to quantify factors that could impact detection such as prey density and time of year (Graham 2002), or snow cover and depth if conducting snow tracking (Forsey & Baggs 2001). Tracking tunnels protect tracks from weather, but require regular checks, especially where small rodents are abundant. Footprints of *Mustela nivalis* and *Mustela erminea* overlap in size, so where they co-occur it is not always possible to identify tracks to species.

FUTURE DEVELOPMENT

Track-based surveying could be advanced by automated track identification software. In addition to faster processing of tracks within track plates for dedicated studies, programs such as WildTrack (<https://www.wildtrack.org/our-work/fit-technology>) offer potential crowdsourced track-based locational data for weasels. Individual identification based on weasel tracks has been limited to studies that practice toe clipping (King & Edgar 1977), although evidence of successful individual identification of fishers using high-quality track plate images (Herzog et al. 2007) suggests it could be possible for weasels.

NON-INVASIVE GENETIC SAMPLING

HISTORY OF DEVELOPMENT

Molecular techniques have provided information on multiple topics of weasel ecology, such as population structure (McDevitt et al. 2013), ancestral relationships (Masuda & Yoshida 1994,

Kurose et al. 2005a), and levels of inbreeding (Huang et al. 2007). These objectives typically require numerous samples of high-quality DNA from blood or tissue cells of live or dead animals. Although non-invasive genetic sampling of faeces, dead animals or hair samples is well established for many species (Taberlet & Luikart 1999), very few studies on weasels have been published (see Kurose et al. 2005b, Zielinski et al. 2020, Schenker et al. 2023). Hair snaring with subsequent DNA analysis has been used with mixed success for stoats and weasels. In New Zealand, Wales and Ireland, 20 cm lengths of 45 mm diameter plastic pipe with a sampling strip covered in adhesive across the entrance at each end have had limited success for collecting hair samples from stoats (Clayton et al. 2011, McAney 2011) and weasels (MacPherson, unpublished data). However, García and Mateos (2009) found that hair snaring was more successful at detecting least weasels than track censuses or scat sampling.

Recently, environmental DNA (eDNA) surveys have gained popularity (Ruppert et al. 2019) and become a viable tool for detection of rare species (Leempoel et al. 2020, Sales et al. 2020). To identify the presence of a species, samples of soil, water or air are screened for mitochondrial DNA shed by animals (Thomsen & Willerslev 2015). Surveys using these techniques have successfully detected several mustelid species, including the critically endangered European mink *Mustela lutreola* (Croose et al. 2023), least weasel, stoat (Broadhurst et al. 2021), European pine marten *Martes martes* and Eurasian otter *Lutra lutra* (Sales et al. 2020). Yet their use for detection of weasels remains limited, likely due to the species' relative rarity, with few metabarcoding studies reporting successful detection (e.g. Bolton 2021), and others failing to detect weasels by eDNA despite detection by camera traps in the same study (e.g. Leempoel et al. 2020, Sales et al. 2020).

ADVANTAGES AND STRENGTHS

Non-invasive genetic sampling of faeces or hair can be used to identify individuals and thus estimate population size and density using mark-recapture or spatial capture-recapture models (Royle et al. 2013). This has yet to be tested on weasels (but see Fuller et al. 2016 for an example with American mink *Neogale vison*). A benefit of eDNA sampling from soil has been improved detection of stoats, which tend to be undetected by surveying devices (e.g. Mostela, footprint tunnels; Bolton 2021).

CURRENT LIMITATIONS

Along with their relatively high costs, eDNA analyses can result in false negatives with soil (Leempoel et al. 2020) and water samples (Sales et al. 2020). As weasels occur in relatively low densities the probability of randomly collecting their DNA is lower than for other species. However,

targeted sampling and baited devices can improve detection rates of low-density and cryptic species using eDNA (Ichu 2022). The success of eDNA also depends on sampling design and animal behaviour (Leempoel *et al.* 2020). Probability of detection can be reduced during the PCR process if poor quality or small quantity of eDNA results in a very low number of reads (Bolton 2021). Use of targeted primers instead of metabarcoding can improve detection probabilities, especially of rare species, for which small amounts of DNA can be swamped out early in the PCR process.

FUTURE DEVELOPMENT

Improvements are required in eDNA techniques specifically for the detection of weasels or small carnivores. Species-specific assays would greatly increase sensitivity and reduce costs of qPCR techniques over metabarcoding (Harper *et al.* 2018). Development of optimal sampling protocols using baited stations or predated nests would be useful as these can likely increase the probability of detection. Linking eDNA sampling with detection dogs might be considered when dogs alert on target species activity but no scat can be found. It also could be linked with track tunnels, enclosed camera traps or hair traps, the latter potentially used to identify individuals.

UNENCLOSED CAMERA TRAPS

HISTORY OF DEVELOPMENT

Most research using camera traps has targeted mammals in the order Carnivora (Burton *et al.* 2015), including some early studies targeting mustelids (e.g. González & Lara 2007, Rosellini *et al.* 2008). However, small mustelids such as weasels and stoats often were not detected in camera trap surveys (Zielinski & Kucera 1995, Kelly & Holub 2008) or were removed from analyses due to small sample sizes (Johnson *et al.* 2009). Ineffectiveness of early camera trap methods at detecting small mustelids, even with bait, is likely due to their small body size and fast movement (Kelly & Holub 2008) and low trigger sensitivities of early camera traps (Wearn & Glover-Kapfer 2019). Recent increases in sensitivity of camera traps have resulted in an increasing number of unenclosed camera trap studies reporting detection of weasels (Moser *et al.* 2017, Ghose *et al.* 2018) and even targeting weasels specifically (Evans & Mortelliti 2022).

ADVANTAGES AND STRENGTHS

Camera traps are likely more cost- and time-efficient than other methods because they passively collect data and require little time and few supplies. Camera traps can be

deployed for prolonged periods, making them effective for broad spatial- or temporal-scale studies (Hsing *et al.* 2022). The cost of using camera traps can be 2–5 times lower than that of live traps (De Bondi *et al.* 2010, White *et al.* 2023). Costs were similar between camera trap surveys and wildlife detection dogs (Glen *et al.* 2016), eDNA surveys (Lyet *et al.* 2021), and non-invasive genetic sampling (Twining *et al.* 2022). However, much less technical expertise and training is required to employ camera traps, and the cost of supplies and equipment is relatively low after a large initial investment to purchase the cameras. Lengthy surveys are likely more cost-effective with camera traps than with methods that require lab equipment/supplies (e.g. eDNA) or specialist practitioners (e.g. detection dogs).

CURRENT LIMITATIONS

Detection rates (0.02–0.44 detections/100 camera days; Ross *et al.* 2013, Kolowski & Forrester 2017) and detection probabilities (0.05–0.24; Croose *et al.* 2022, Evans & Mortelliti 2022) of weasels are relatively low and vary greatly across camera trap studies with differing weasel species and survey methods. The speed, agility and small body size of weasels can enable them to pass through the small zone of detection, or become blocked from view, before a camera is triggered (Evans *et al.* 2019). This is especially evident as distance of the animal from the camera increases. However, with addition of bait, unenclosed camera traps have successfully collected detailed data on weasel occupancy patterns (Evans & Mortelliti 2022).

Sympatric weasel species are often difficult to differentiate using external physical traits (King & Powell 2007), limiting species identification in camera trap surveys. The ratio of tail length to combined head and body length has been confirmed with DNA to distinguish between some weasel species (a ratio >44% distinguishing long-tailed weasels from stoats; Hall 1951, Elsasser & Parker 2008). This ratio can be calculated from morphometric data using camera images if a reference of scale is within view (Evans & Mortelliti 2022). However, there is doubt if this ratio is effective throughout the species' shared range (St-Pierre *et al.* 2006). Additional work is needed to find effective ways of distinguishing sympatric weasel species in camera images. Excepting the *Mustela nivalis vulgaris* morph, which has unique patterns of gular spots and irregular back-belly margins, weasels typically lack markings that can be used to identify individuals (King & Powell 2007). Analyses that require individual recognition (e.g. spatial capture–recapture analyses) are likely to need other techniques (e.g. PIT-tags, hair samples for DNA analyses, collaring), or using structures that funnel animals close to the

camera (e.g. Mostela; Mos & Hofmeester 2023). Statistical models that avoid the need to identify individuals may be an option (e.g. unmarked spatial capture–recapture; Santini et al. 2022).

While camera traps are relatively time-efficient, it takes time to check them regularly to download data, change batteries and clear vegetation from the field of view. Frequency of checks depends on memory card capacity and battery life, which vary with activity recorded by the camera and environmental conditions. Also, image management and analysis can be very time consuming, especially if done manually. Image management and processing software can help (e.g. Niedballa et al. 2016), but the usual need for human skill to identify species limits overall efficiency of the method.

FUTURE DEVELOPMENT

Studies employing unenclosed camera traps would greatly benefit from advances in automated species identification, methods to estimate body condition, and development and standardisation of best practices. Software has been used to identify small mustelid species from images collected by camera traps (Yu et al. 2013) and is now widely available through platforms such as Wildlife Insights (Ahumada et al. 2019). Further development of these methods (i.e. training using species-specific data) is needed to generate more reliable species identification. Calculation of body condition is difficult using camera traps due to the requirement of body mass data (Krebs & Singleton 1993, Schulte-Hostedde et al. 2001). A method to simultaneously collect body size and mass data from individual weasels or an alternative method using visible characteristics that are highly correlated to body condition (e.g. Pérez-Flores et al. 2016) would be valuable.

Developing best practices that enhance detection of weasels on unenclosed camera trap surveys would benefit the design of site- or species-specific as well as landscape-scale camera-trapping protocols. This includes determining which camera models and settings are better, where to place cameras and which lure(s) are most attractive. While best practices will likely vary among species, landscapes and project objectives, such information is critical to designing long-term monitoring strategies using unenclosed and enclosed camera traps (Jachowski et al. 2021).

ENCLOSED CAMERA TRAPS

HISTORY OF DEVELOPMENT

Methods that confine camera traps in enclosed spaces have been developed to survey smaller-bodied vertebrates more effectively. By directing animals closer to the camera trap

they improve detectability and get better photographs. The Mostela system, designed for small mustelids, has a camera trap aimed horizontally at an opening in the side of a tracking tunnel in a box (Fig. 5). Mostela has mainly been used for studying least weasel and stoat in the Netherlands (Westra 2019, Mos & Hofmeester 2020), England (Croose & Carter 2019), and Poland (Hofmeester et al. in press). After publication of its design (Mos & Hofmeester 2020), Mostela has been used in additional studies including least weasel in the USA (Holloway et al. 2022) and the endemic Irish stoat (Croose et al. 2022). Other studies adopted similar camera trap designs, such as foldable plastic boxes for weasels and stoats at high altitudes in Spain (Fig. 5; Salvador et al. 2022) and metal boxes for long-tailed weasel and Colombian weasel *Neogale felipei* in Colombia (Cepeda-Duque et al. 2023). Another boxed camera trap, developed to monitor small mammals under snow (Soininen et al. 2015), has shown success in detecting *Mustela nivalis nivalis* and *Mustela erminea* in arctic systems (Kleiven et al. 2023).

The AHDriFT system (Fig. 6) uses camera traps in boxes, with each camera aiming downward from inside the top of a box. Animals are funnelled to box entrances by drift fence(s) with a camera box at each end. This method, originally described by Martin et al. (2017), was developed from the Hunt trap designed by McCleery et al. (2014). AHDriFT systems are typically used to survey herpetofauna and rodent communities (Martin et al. 2017, Boynton et al. 2021). However, substantial weasel bycatch has been reported (Amber et al. 2021a, White et al. 2023).

ADVANTAGES AND STRENGTHS

Data from enclosed camera traps have been used to estimate occurrence, relative activity, and daily and seasonal activity patterns (Mos & Hofmeester 2020, Amber et al. 2021b, Croose et al. 2022), and predator–prey dynamics (Kleiven et al. 2023). Hofmeester et al. (in press) compared relative abundance estimated from Mostela data using Royle-Nichols models (Royle & Nichols 2003) with the minimum number of weasels known to occur in the area based on live-trapping. They found Mostela to have potential for tracking yearly fluctuations in weasel abundance. An attempt to estimate density using Mostela data for individually identified least weasels in the Netherlands has promising results (Fig. 5; Mos & Hofmeester 2023). Mos and Hofmeester (2020) suggested that absolute densities can be estimated from Mostela data using capture–mark–recapture and spatially explicit capture–recapture methods for species with fur patterns that allow individual identification (e.g. the pattern of spots between dorsal and ventral fur colours of *Mustela nivalis vulgaris*; Fig. 5).

While enclosed camera traps share advantages with unenclosed systems, they can be deployed in a wider range



Fig. 5. Top: Mostela boxes transported by foot in the Catalan Pyrenees, Spain (photo credit: S. Salvador). Bottom: an example of four individual least weasels (*Mustela nivalis*) recorded within 1 week on one camera trap location (photo credit: Small Mustelid Foundation).

of habitats, such as under vegetation, under snow, and among rocks in areas where trees are scarce. *Mustela* seems to work best for the least weasel, likely because it can be placed in microhabitats that are frequented by this species. AHDriFT systems have only been reported to detect long-tailed weasels (Amber et al. 2021a, White et al. 2023), likely because studies were in regions where long-tailed weasels were the dominant or only weasel species.

Enclosed camera trap systems are able to detect both weasels and their small mammal prey. These ‘bycatch’

data can be valuable, especially where limited information is available on the small mammal community or where methods such as live-trapping, are hard to implement (see, e.g. Soininen et al. 2015). Simultaneous sampling of weasels and their potential prey can reveal interactions and provide data to analyse prey-related demographic fluctuations (Hamed MK, Holloway AW, Watts C, Webster A, Tanner K, and Moore T, unpublished data, Kleiven et al. 2023). Because drift fences associated with AHDriFT systems increase the overall



Fig. 6. Top: AHDriFT system deployment showing drift fence with modified overturned trash can at end to direct animals into the trash can. Bottom: Inside the trash can a downward facing camera takes a photo of a weasel in Indiana, USA (photo credit: Scott Bergeson and Carsten White).

area sampled by the device, they might make these systems more effective at recording weasels and their prey.

CURRENT LIMITATIONS

Application of enclosed camera systems to multiple weasel species is not yet properly tested. Stoats avoid *Mostela* in some cases (Croose et al. 2022). When using bait, and in certain locations, enclosed systems can collect numerous photos or videos of non-target species (especially small rodents), making data analysis difficult, particularly without an effective data management system. Using bait also increases the risk of attracting bears, cows or other large animals that can disturb equipment. Enclosed camera traps are relatively cumbersome to transport. It is unclear what the effective sampling area of a single enclosed camera system is (with the possible exception of the AHDriFT system), a measure that is important to estimate as a basis for interpreting data. We also need to test if estimates of occurrence and (relative) abundance based on enclosed camera systems are accurate and sensitive enough to detect trends over time.

FUTURE DEVELOPMENT

Full-scale application of enclosed camera systems as monitoring devices would benefit from lighter and cheaper designs that allow deployment of a larger number of units. Finding materials other than PVC for cameras would reduce potential problems caused by animals (e.g. bears) being attracted by PVC. Addition of a hair snag within or at the entrance of the device would enable collection of samples for analyses based on DNA. Wider testing of cameras with white flash would be helpful (Herrera et al. 2021). White flash might be more disturbing to animals, but it would enable easier identification of species and individuals (e.g. Mos & Hofmeester 2020). Comparison of enclosed systems to other methods is needed to test their suitability for monitoring weasel occurrence and density at larger scales. A comparison of *Mostela* data to live-trapping data of least weasels in Poland shows promising results (Hofmeester et al. *in press*) but needs to be replicated under different circumstances and for different species. Comparisons of AHDriFT data to live-trapping data from midwestern USA are promising (Amber et al. 2021a, White et al. 2023), but more comparisons are required. Because ongoing studies indicate that availability of alternative underground structures and holes might influence the chance that a weasel enters an enclosed camera system, investigation of this relationship would be beneficial. Differences among species and individuals in their tendency to enter enclosed systems needs to be tested. Simultaneous deployment of enclosed and unenclosed camera traps in proximity enables comparison of their detection probabilities (Croose et al. 2022, Cepeda-Duque et al. 2023). Finally, there is a need to train a machine learning algorithm to automate labour-intensive identification of species from images similar to that which exists for unenclosed camera trap photos (Böhner et al. 2023).

TOWARDS IMPROVED WEASEL MONITORING

Whether attempting a first survey for weasels or sustaining a long-term monitoring program, important contributions are needed to advance weasel monitoring. Where long-term weasel monitoring data exist, it makes sense to sustain those efforts using similar methodologies. However, traditional methods should be tested against other emerging non-invasive techniques reviewed here to evaluate the rigour of those results and utility of emerging techniques (Smith & Weston 2017). For example, long-term studies in northern latitudes using lemming nest occupation and snow track surveys as

indices of weasel abundance (Sundell *et al.* 2013) allow for the comparative evaluation of other weasel monitoring approaches (e.g. baited cameras or track tunnels). Where new monitoring is planned, it is important to trial multiple techniques reviewed here simultaneously. Such comparative studies are critical to providing insight into the utility of differing monitoring techniques for a given species and site characteristics (e.g. habitat, range of densities, non-target bait consumption/decomposition rate).

Where weasels are of management concern and little is known, there is an emerging pattern of using an iterative, multi-method approach to gain an understanding of weasel distribution and population ecology. First, citizen science and historical records are often used to identify where weasels persist. Second, baited camera traps (and to a lesser extent detection dogs or molecular approaches) are used to gather information on spatial distribution and factors influencing occupancy of weasels within those focal areas (e.g. Ghose *et al.* 2018, Cepeda-Duque *et al.* 2023). Third, where weasels are known to be resident, researchers use enclosed camera trap setups (Mostela and/or AHDriFT systems) to gain insights into weasel behaviour and factors influencing their relative activity or abundance. Formalising this process within management plans and across political boundaries could help build towards large scale, comparable trend data needed to information management.

CONCLUSIONS

The management concerns surrounding weasels across many portions of their range globally, either as a declining or invasive species, necessitates improved monitoring methods. While our review highlights the diversity of methods used to non-invasively monitor weasels, there are clear trends in certain methods becoming more commonly used than others. Rather than relying on any single method, there is great promise in using a multi-method approach to long-term weasel monitoring that combines citizen science with adaptations of both common and emerging technologies. Regardless of the approach used, there is a need for strong commitment to dedicated weasel monitoring that is replicated over space and time to inform future management action.

ACKNOWLEDGEMENTS

We thank all the attendees of the 1st International Weasel Monitoring Symposium for contributing to discussions that informed the drafting of this manuscript. We thank M. Keating for assistance in creating a figure, and R. Jachowski and two anonymous reviewers for comments that helped improve this manuscript.

DATA AVAILABILITY STATEMENT

Data sharing not applicable as no new data generated.

REFERENCES

- Aebischer NJ, Davey PD, Kingdon NG (2011) *National Gamebag Census: Mammal Trends to 2009*. Game & Wildlife Conservation Trust, Fordingbridge, UK.
- Ahumada JA, Fegraus E, Birch T, Flores N, Kays R, Brien TGO *et al.* (2019) Wildlife insights: a platform to maximize the potential of camera trap and other passive sensor wildlife data for the planet. *Environmental Conservation* 47: 1–6.
- Amber ED, Lipps GJ Jr, Peterman WE (2021a) Evaluation of the AHDriFT camera trap system to survey for small mammals and herpetofauna. *Journal of Fish and Wildlife Management* 12: 197–207.
- Amber ED, Myers JM, Lipps GJ Jr, Peterman WE (2021b) Small mammal daily activity periods derived using AHDriFT camera traps. *Mammal Research* 66: 289–300.
- Bennett EM, Hauser CE, Moore JL (2019) Evaluating conservation dogs in the search for rare species. *Conservation Biology* 34: 314–325.
- Böhner H, Kleiven EF, Ims RA, Soininen EM (2023) A semi-automatic workflow to process images from small mammal camera traps. *Ecological Informatics* 76: 102150.
- Bolduc D, Fauteux D, Gagnon CA, Gauthier G, Bêty J, Legagneux P (2023) Testimonials to reconstruct past abundances of wildlife populations. *Basic and Applied Ecology* 68: 23–34.
- Bolton PL (2021) *Comparing Fine-Scale Temporal Environmental DNA (eDNA) Sampling to Camera Trapping for Detecting Mammalian Carnivores*. PhD thesis, University of Salford, Salford, UK.
- Boynton MK, Toenies M, Cornelius N, Rich LN (2021) Comparing camera traps and visual encounter surveys for monitoring small animals. *California Fish and Wildlife* 107: 99–117.
- Broadhurst HA, Gregory LM, Bleakley EK, Perkins JC, Lavin JV, Bolton P *et al.* (2021) Mapping differences in mammalian distributions and diversity using environmental DNA from rivers. *Science of the Total Environment* 801: 149724.
- Burton AC, Neilson E, Moreira D, Ladle A, Steenweg R, Fisher JT, Bayne E, Boutin S (2015) Wildlife camera trapping: a review and recommendations for linking surveys to ecological processes. *Journal of Applied Ecology* 52: 675–685.
- Buyaskas M, Evans BE, Mortelliti A (2020) Assessing the effectiveness of attractants to increase camera trap detections of North American mammals. *Mammalian Biology* 100: 91–100.
- Cepeda-Duque JC, Arango-Correa E, Andrade-Ponce GP, Mazariegos L, Hofmeester TR, Ramírez-Chaves HE (2023)

- Expanding the frontiers of camera-trapping in Colombia: application of the “Mostela” system to gain knowledge on small non-volant mammals from an Andean cloud forest. *Mammalia* 87: 419–428.
- Červinka J, Drahníková L, Kreisinger J, Šálek M (2014) Effect of habitat characteristics on mesocarnivore occurrence in urban environment in the Central Europe. *Urban Ecosystems* 17: 893–909.
- Clayton RI, Byrom AE, Anderson DP, Edge K-A, Gleeson D, McMurtrie P, Veale A (2011) Density estimates and detection models inform stoat (*Mustela erminea*) eradication on Resolution Island, New Zealand. In: Veitch CR, Clout MN, Towns DR (eds) *Island Invasives: Eradication and Management*, 413–417. IUCN, Gland, Switzerland.
- Conrad CC, Hilchey KG (2011) A review of citizen science and community-based environmental monitoring: issues and opportunities. *Environmental Monitoring and Assessment* 176: 273–291.
- Croose E, Carter SP (2019) A pilot study of a novel method to monitor weasels (*Mustela nivalis*) and stoats (*M. erminea*) in Britain. *Mammal Communications* 5: 6–12.
- Croose E, Hanniffy R, Hughes B, McAney K, MacPherson J, Carter SP (2022) Assessing the detectability of the Irish stoat *Mustela erminea hibernica* using two camera trap-based survey methods. *Mammal Research* 671: 1–8.
- Croose E, Hanniffy R, Harrington A, Pödra M, Gómez A, Bolton PL et al. (2023) Mink on the brink: comparing survey methods for detecting a critically endangered carnivore, the European mink *Mustela lutreola*. *European Journal of Wildlife Research* 69: 34.
- Davis LS, Zhu L, Finkler W (2023) Citizen science: is it good science? *Sustainability* 15: 4577.
- De Bondi N, White JG, Stevens M, Cooke R (2010) A comparison of the effectiveness of camera trapping and live trapping for sampling terrestrial small-mammal communities. *Wildlife Research* 37: 456–465.
- DeMatteo KE, Davenport B, Wilson LE (2019) Back to the basics with conservation detection dogs: fundamentals for success. *Wildlife Biology* 1: 1–9.
- Egloff G, Arlettaz R, Humbert J-Y (2022) *Comparing Detection Dogs and Camera Traps as Sampling Methods to Estimate Mustela Erminea and Mustela Nivalis Occurrence*. BS thesis, University of Bern, Bern, Switzerland.
- Ellul C, Gupta S, Haklay MM, Bryson K (2013) A platform for location based app development for citizen science and community mapping. In: Krisp JM (ed) *Progress in Location-Based Services*, 71–90. Springer, Heidelberg, Germany.
- Elsasser SC, Parker GH (2008) Morphometric criteria for distinguishing species and age-cohorts of ermine (*Mustela erminea*) and long-tailed weasel (*M. frenata*). *Acta Zoologica Academiae Scientiarum Hungaricae* 54: 75–88.
- Evans BE, Mortelliti A (2022) Forest disturbance and occupancy patterns of American ermine (*Mustela richardsonii*) and long-tailed weasel (*Neogale frenata*): results from a large-scale natural experiment in Maine, United States. *Journal of Mammalogy* 103: 1338–1349.
- Evans BE, Mosby CE, Mortelliti A (2019) Assessing arrays of multiple trail cameras to detect north American mammals. *PLoS ONE* 14: e0217543.
- Feige N, Ehrlich D, Popov IY, Broekhuizen S (2012) Monitoring least weasels after a winter peak of lemmings in Taimyr: body condition, diet and habitat use. *Arctic* 65: 273–282.
- Forsey ES, Baggs EM (2001) Winter activity of mammals in riparian zones and adjacent forests prior to and following clear-cutting at copper Lake, Newfoundland, Canada. *Forest Ecology and Management* 145: 163–171.
- Fukasawa K, Osada Y, Iijima H (2020) Is harvest size a valid indirect measure of abundance for evaluating the population size of game animals using harvest-based estimation? *Wildlife Biology* 2020: 1–7.
- Fuller AK, Sutherland CS, Royle JA, Hare MP (2016) Estimating population density and connectivity of American mink using spatial capture–recapture. *Ecological Applications* 26: 1125–1135.
- García P, Mateos I (2009) Evaluation of three indirect methods for surveying the distribution of the least weasel *Mustela nivalis* in a Mediterranean area. *Small Carnivore Conservation* 40: 22–26.
- Ghose M, Sharma D, Murali NS (2018) First record of yellow-bellied weasel (*Mustela kathiah*) from Buxa Tiger Reserve, West Bengal, India. *NeBIO* 9: 212–214.
- Gilbert NA, Pease BS, Anhalt-Depies CM, Clare JD, Stenglein JL, Townsend PA, Van Deelen TR, Zuckerberg B (2021) Integrating harvest and camera trap data in species distribution models. *Biological Conservation* 258: 109147.
- Glen AS, Anderson D, Veltman CJ, Garvey PM, Nichols M (2016) Wildlife detector dogs and camera traps: a comparison of techniques for detecting feral cats. *New Zealand Journal of Zoology* 43: 127–137.
- Gompper ME, Kays RW, Ray JC, Lapoint SD, Bogan DA, Cryan JR (2006) A comparison of noninvasive techniques to survey carnivore communities in northeastern North America. *Wildlife Society Bulletin* 34: 1142–1151.
- González CAL, Lara DRA (2007) Noteworthy record of the Tayra (Carnivora: Mustelidae: *Eira barbara*) in the Sierra Gorda biosphere reserve, Querétaro, México. *Western North American Naturalist* 67: 150–151.
- Graham IM (2002) Estimating weasel *Mustela nivalis* abundance from tunnel tracking indices at fluctuating field vole *Microtus agrestis* density. *Wildlife Biology* 8: 279–287.
- Grimm-Seyfarth A, Harms W, Berger A (2021) Detection dogs in nature conservation: a database on their

- world-wide deployment with a review on breeds used and their performance compared to other methods. *Methods in Ecology and Evolution* 12: 568–579.
- Hall ER (1951) American weasels. *University of Kansas Publications, Museum of Natural History* 4: 1–466.
- Harper LR, Lawson Handley L, Hahn C, Boonham N, Rees HC, Gough KC et al. (2018) Needle in a haystack? A comparison of eDNA metabarcoding and targeted qPCR for detection of the great crested newt (*Triturus cristatus*). *Ecology and Evolution* 8: 6330–6341.
- Hayder F, Madikiza ZJK, Do Linh San E (2023) Updated distribution and current population status of the least weasel (*Mustela nivalis*) in Tunisia: a countrywide interview survey. *African Journal of Wildlife Research* 53: 11–18.
- Herrera DJ, Moore SM, Herrmann V, McShea WJ, Cove MV (2021) A shot in the dark: white and infrared LED flash camera traps yield similar detection probabilities for common urban mammal species. *Hystrix* 32: 72–75.
- Herzog CJ, Kays RW, Ray JC, Gompper ME, Zielinski WJ, Higgins R, Tymeson M (2007) Using patterns in track-plate footprints to identify individual fishers. *Journal of Wildlife Management* 71: 955–963.
- Hill S, Hill J (1987) *Richard Henry of Resolution Island*. John McIndoe Ltd, Dunedin, New Zealand.
- Hoffman C, Cooper CB, Kennedy EB, Farooque M, Cavalier D (2017) Scistarter 2.0: a digital platform to foster and study sustained engagement in citizen science. In: Ceccaroni L, Piera J (eds) *Analyzing the Role of Citizen Science in Modern Research*, 50–61. IGI Global, Hershey, Pennsylvania, USA.
- Hofmeester TR, Mos J, Zub K (in press) Comparing direct (live-trapping) and indirect (camera-trapping) approaches for estimating the abundance of weasels (*Mustela nivalis*). *Mammalian Biology*.
- Holloway A, Owen J, Hamed K (2022) Utilisation of Mostela boxes and citizen science to update the distribution of least weasels (*Mustela nivalis*) in Virginia, USA. *34th European Mustelid Colloquium*.
- Hsing P-Y, Hill RA, Smith GC, Bradley S, Green SE, Kent VT et al. (2022) Large-scale mammal monitoring: the potential of a citizen science camera-trapping project in the United Kingdom. *Ecological Solutions and Evidence* 3: e12180.
- Huang C-C, Lin R-C, Li S-H, Lee L-L (2007) Characterization of polymorphic tetranucleotide microsatellite loci from the Siberian weasel (*Mustela sibirica*). *Molecular Ecology Notes* 7: 483–485.
- Ichu IG (2022) *Evaluating Non-invasive Environmental Methods for Detecting Tropical African Pangolin Species to Inform Conservation Actions*. MS thesis, Mississippi State University, Mississippi State, USA.
- Jachowski D, Kays R, Butler A, Hoylman AM, Gompper ME (2021) Tracking the decline of weasels in North America. *PLoS ONE* 16: e0254387.
- Johnson A, Vongkhamheng C, Saithongdam T (2009) The diversity, status and conservation of small carnivores in a montane tropical forest in northern Laos. *Oryx* 43: 626–633.
- Jones C, Moller H, Hamilton W (2004) A review of potential techniques for identifying individual stoats (*Mustela erminea*) visiting control or monitoring stations. *New Zealand Journal of Zoology* 31: 193–203.
- Kays R, Lasky M, Allen ML, Dowler RC, Hawkins MT, Hope AG et al. (2022) Which mammals can be identified from camera traps and crowdsourced photographs? *Journal of Mammalogy* 103: 767–775.
- Kelly MJ, Holub EL (2008) Camera trapping of carnivores: trap success among camera types and across species, and habitat selection by species, on Salt Pond Mountain, Giles County, Virginia. *Northeastern Naturalist* 15: 249–262.
- King CM (1980) Population biology of the weasel *Mustela nivalis* on British game estates. *Ecography* 3: 160–168.
- King CM, Edgar RL (1977) Techniques for trapping and tracking stoats (*Mustela erminea*); a review, and a new system. *New Zealand Journal of Zoology* 4: 193–212.
- King CM, Powell RA (2007) *The Natural History of Weasels and Stoats: Ecology, Behavior, and Management*. Oxford University Press, New York, New York, USA.
- Kleiven EF, Barraquand F, Gimenez O, Henden J-A, Ims RA, Soininen EM, Yoccoz NG (2023) A dynamic occupancy model for interacting species with two spatial scales. *Journal of Agricultural, Biological and Environmental Statistics* 28: 466–482.
- Kolowski JM, Forrester TD (2017) Camera trap placement and the potential for bias due to trails and other features. *PLoS ONE* 12: e0186679.
- Koshkina V, Wang Y, Gordon A, Dorazio RM, White M, Stone L (2017) Integrated species distribution models: combining presence-background data and site-occupancy data with imperfect detection. *Methods in Ecology and Evolution* 8: 420–430.
- Krebs CJ, Singleton GR (1993) Indexes of condition for small mammals. *Australian Journal of Zoology* 41: 317–323.
- Kurose N, Abramov AV, Masuda R (2005a) Comparative phylogeography between the ermine *Mustela erminea* and the least weasel *M. nivalis* of Palaearctic and Nearctic regions, based on analysis of mitochondrial DNA control region sequences. *Zoological Science* 22: 1069–1078.
- Kurose N, Masuda R, Tatara M (2005b) Fecal DNA analysis for identifying species and sex of sympatric carnivores: a noninvasive method for conservation on the Tsushima Islands, Japan. *Journal of Heredity* 96: 688–697.
- Lasky M, Parsons A, Schuttler S, Mash A, Larson L, Norton B, Pease B, Boone H, Gatens L, Kays R (2021) Candid critters: challenges and solutions in a large-scale citizen science camera trap project. *Citizen Science: Theory and Practice* 6: 1–17.

- Lau MWN, Fellowes JR, Chan BPL (2010) Carnivores (Mammalia: Carnivora) in South China: a status review with notes on the commercial trade. *Mammal Review* 40: 247–292.
- Leempoel K, Hebert T, Hadly EA (2020) A comparison of eDNA to camera trapping for assessment of terrestrial mammal diversity. *Proceedings of the Royal Society B: Biological Sciences* 287: 20192353.
- Leigh KA, Dominick M (2015) An assessment of the effects of habitat structure on the scat finding performance of a wildlife detection dog. *Methods in Ecology and Evolution* 6: 745–752.
- Linzey DW, Hamed MK (2016) Distribution of the least weasel (*Mustela nivalis*) in the southeastern United States. *Southeastern Naturalist* 15: 243–258.
- Liu H-Y, Dörler D, Heigle F, Grossberndt S (2021) Citizen science platforms. In: Vohland K, Land-Zandstra A, Ceccaroni L, Lemmens R, Perelló J, Ponti M, Samson R, Wagenknecht K (eds) *The Science of Citizen Science*, 439–459. Springer, London, UK.
- Livingston TR, Gipson PS, Ballard WB, Sanchez DM, Krausman PR (2005) Scat removal: a source of bias in feces-related studies. *Wildlife Society Bulletin* 33: 172–178.
- Long RA, Donovan TM, MacKay P, Zielinski WJ, Buzas JS (2007) Effectiveness of scat detection dogs for detecting forest carnivores. *Journal of Wildlife Management* 71: 2007–2017.
- Lyet A, Pellissier L, Valentini A, Dejean T, Hehmeyer A, Naidoo R (2021) eDNA sampled from stream networks correlates with camera trap detection rates of terrestrial mammals. *Scientific Reports* 11: 11362.
- Martin SA, Rautsaw RM, Robb F, Bolt MR, Parkinson CL, Seigel RA (2017) Set AHDriFT: applying game cameras to drift fences for surveying herpetofauna and small mammals. *Wildlife Society Bulletin* 41: 804–809.
- Masuda R, Yoshida MC (1994) Nucleotide sequence variation of cytochrome b genes in three species of weasels *Mustela itatsi*, *Mustela sibirica*, and *Mustela nivalis*, detected by improved PCR product-direct sequencing technique. *Journal of the Mammalogical Society of Japan* 19: 33–43.
- Maunder MN, Sibert JR, Fonteneau A, Hampton J, Kleiber P, Harley SJ (2006) Interpreting catch per unit effort data to assess the status of individual stocks and communities. *ICES Journal of Marine Science* 63: 1373–1385.
- McAney K (2011) *A Pilot Study to Test the Use of Hair Tubes to Detect the Irish Stoat along Hedgerows in County Galway*. Vincent Wildlife Trust, Eastnor, UK.
- McCleery RA, Zweig CL, Desa MA, Hunt R, Kitchens WM, Percival HF (2014) A novel method for camera-trapping small mammals. *Wildlife Society Bulletin* 38: 887–891.
- McDevitt AD, Oliver MK, Piertney SB, Szafrńska PA, Konarzewski M, Zub K (2013) Individual variation in dispersal associated with phenotype influences fine-scale genetic structure in weasels. *Conservation Genetics* 14: 499–509.
- McDonald RA, Harris S (2002) Population biology of stoats *Mustela erminea* and weasels *Mustela nivalis* on game estates in Great Britain. *Journal of Applied Ecology* 39: 793–805.
- Mos J, Hofmeester TR (2020) The *Mostela*: an adjusted camera trapping device as a promising non-invasive tool to study and monitor small mustelids. *Mammal Research* 65: 843–853.
- Mos J, Hofmeester TR (2023) Year-round monitoring of common weasels (*Mustela nivalis*) using *Mostela* boxes in The Netherlands. *1st International Weasel Monitoring Symposium*.
- Moser AM, Diggins CA, Ford WM (2017) High-elevation observations of long-tailed weasel and eastern chipmunk in North Carolina. *Southeastern Naturalist* 16: N40–N42.
- Niedballa J, Sollmann R, Courtiol A, Wilting A (2016) camtrapR: an R package for efficient camera trap data management. *Methods in Ecology and Evolution* 7: 1457–1462.
- Norris D, Michalski F (2010) Implications of faecal removal by dung beetles for scat surveys in a fragmented landscape of the Brazilian Amazon. *Oryx* 44: 455–458.
- Pech R, Maitland M (2016) Conservation of native fauna in highly invaded systems: managing mammalian predators in New Zealand. *Restoration Ecology* 24: 816–820.
- Pérez-Flores J, Calmé S, Reyna-Hurtado R (2016) Scoring body condition in wild Baird's tapir (*Tapirus bairdii*) using camera traps and opportunistic photographic material. *Tropical Conservation Science* 9: 1940082916676128.
- Quick HF (1944) Habits and economics of the New York weasel in Michigan. *Journal of Wildlife Management* 8: 71–78.
- Reed SE, Bidlack AL, Hurt A, Getz WM (2011) Detection distance and environmental factors in conservation detection dog surveys. *Journal of Wildlife Management* 75: 243–251.
- Reindl-Thompson SA, Shivik JA, Whitelaw A, Hurt A, Higgins KF (2006) Efficacy of scent dogs in detecting black-footed ferrets at a reintroduction site in South Dakota. *Wildlife Society Bulletin* 34: 1435–1439.
- Rodrigues M, Bos AR, Schembri PJ, de Lima RF, Lymberakis P, Parpal L et al. (2017) Origin and introduction history of the least weasel (*Mustela nivalis*) on Mediterranean and Atlantic islands inferred from genetic data. *Biological Invasions* 19: 399–421.
- Rosellini S, Osorio E, Ruiz-González A, Piñeiro A, Barja I (2008) Monitoring the small-scale distribution of sympatric European pine martens (*Martes martes*) and stone martens (*Martes foina*): a multievidence approach using faecal DNA analysis and camera-traps. *Wildlife Research* 35: 434–440.

- Ross J, Hearn AJ, Macdonald DW (2013) Recent camera-trap records of Malay weasel *Mustela nudipes* in Sabah, Malaysian Borneo. *Small Carnivore Conservation* 49: 20–24.
- Royle JA, Nichols JD (2003) Estimating abundance from repeated presence–absence data or point counts. *Ecology* 84: 777–790.
- Royle JA, Chandler RB, Sollmann R, Gardner B (2013) *Spatial Capture-Recapture*. Academic Press, Waltham, Massachusetts, USA.
- Ruppert KM, Kline RJ, Rahman MS (2019) Past, present, and future perspectives of environmental DNA (eDNA) metabarcoding: a systematic review in methods, monitoring, and applications of global eDNA. *Global Ecology and Conservation* 17: e00547.
- Sales NG, McKenzie MB, Drake J, Harper LR, Browett SS, Coscia I et al. (2020) Fishing for mammals: landscape-level monitoring of terrestrial and semi-aquatic communities using eDNA from riverine systems. *Journal of Applied Ecology* 57: 707–716.
- Salvador S, Vila A, Palazón S (2022) Monitoring stoats (*Mustela erminea*) and other small mammals at high altitude under a scenario of climate change. *34th European Mustelid Colloquium*.
- Santini G, Abolaffio M, Ossi F, Franzetti B, Cagnacci F, Focardi S (2022) Population assessment without individual identification using camera-traps: a comparison of four methods. *Basic and Applied Ecology* 61: 68–81.
- Schenker L, Arletta R, Humbert J-Y (2023) *Assessing the Ecological Value of Small Structures for Mustelid Populations in Agricultural Landscapes*. MS thesis, University of Bern, Bern, Switzerland.
- Schmidt NM, Ims RA, Høye TT, Gilg O, Hansen LH, Hansen J, Lund M, Fuglei E, Forchhammer MC, Sittler B (2012) Response of an arctic predator guild to collapsing lemming cycles. *Proceedings of the Royal Society B: Biological Sciences* 279: 4417–4422.
- Schulte-Hostedde AI, Millar JS, Hickling GJ (2001) Evaluating body condition in small mammals. *Canadian Journal of Zoology* 79: 1021–1029.
- Shirk JL, Ballard HL, Wilderman CC, Phillips T, Wiggins A, Jordan R et al. (2012) Public participation in scientific research: a framework for deliberate design. *Ecology and Society* 17: 29.
- Silvertown J (2009) A new dawn for citizen science. *Trends in Ecology & Evolution* 24: 467–471.
- Smith DHV, Weston KA (2017) Capturing the cryptic: a comparison of detection methods for stoats (*Mustela erminea*) in alpine habitats. *Wildlife Research* 44: 418–426.
- Smith DA, Ralls K, Cypher BL, Clark HO Jr, Kelley PA, Williams DF, Maldonado JE (2006) Relative abundance of endangered San Joaquin kit foxes (*Vulpes macrotis mutica*) based on scat-detection dog surveys. *Southwestern Naturalist* 51: 210–219.
- Soininen EM, Jensvoll I, Killengreen ST, Ims RA (2015) Under the snow: a new camera trap opens the white box of subnivean ecology. *Remote Sensing in Ecology and Conservation* 1: 29–38.
- St-Pierre C, Ouellet J-P, Dufresne F, Chaput-Bardy A, Huber F (2006) Morphological and molecular discrimination of *Mustela erminea* (ermine) and *M. frenata* (long-tailed weasels) in eastern Canada. *Northeastern Naturalist* 13: 143–152.
- Stuery TD (2012) *Sampling to Determine the Distribution and Habitat Use of Long-Tailed Weasels*. Final Report. Alabama Endangered Wildlife Projects.
- Sundell J, O'Hara RB, Helle P, Hellstedt P, Henttonen H, Pietiäinen H (2013) Numerical response of small mustelids to vole abundance: delayed or not? *Oikos* 122: 1112–1120.
- Taberlet P, Luikart G (1999) Non-invasive genetic sampling and individual identification. *Biological Journal of the Linnean Society* 68: 41–55.
- Tapper S, Reynolds J (1996) The wild fur trade: historical and ecological perspectives. In: Taylor VJ, Dunstone N (eds) *The Exploitation of Mammal Populations*, 28–44. Springer, Dordrecht, the Netherlands.
- Tempero GW, McDonald RM, King CM (2007) Distribution and activity of small mammals on pastoral farmland and forest in New Zealand. *Wildlife Biology in Practice*, 3, 43–51.
- Thomas ML, Baker L, Beattie JR, Baker AM (2020) Determining the efficacy of camera traps, live capture traps, and detection dogs for locating cryptic small mammal species. *Ecology and Evolution* 10: 1054–1068.
- Thompson CM, Royle JA, Garner JD (2012) A framework for interference about carnivore density from unstructured spatial sampling of scat using detector dogs. *Journal of Wildlife Management* 76: 863–871.
- Thomsen PF, Willerslev E (2015) Environmental DNA – an emerging tool in conservation for monitoring past and present biodiversity. *Biological Conservation* 183: 4–18.
- Twining JP, McFarlane C, O'Meara D, O'Reilly C, Reyne M, Montgomery WI, Helyar S, Tosh DG, Augustine BC (2022) A comparison of density estimation methods for monitoring marked and unmarked animal populations. *Ecosphere* 13: e4165.
- Wasser SK, Davenport B, Ramage ER, Hunt KE, Parker M, Clarke C, Stenhouse G (2004) Scat detection dogs in wildlife research and management: application to grizzly and black bears in the yellowhead ecosystem, Alberta, Canada. *Canadian Journal of Zoology* 82: 475–492.
- Wearn OR, Glover-Kapfer P (2019) Snap happy: camera traps are an effective sampling tool when compared with alternative methods. *Royal Society Open Science* 6: 181748.
- Westra SA (2019) A nine month small mustelid survey across four research sites in The Netherlands. *Lutra* 62: 89–107.
- White CL, Jenkins LJ, Proctor TL, Clements J, Jordan MA, Bergeson SM (2023) Comparing effectiveness of AHDriFT

- systems and Sherman traps for surveying small mammals in northeastern Indiana. *Journal of Fish and Wildlife Management* 14: 108–120.
- Wright PGR, Croose E, Macpherson JL (2022) A global review of the conservation threats and status of mustelids. *Mammal Review* 52: 410–424.
- Wright PGR, Croose E, Hunter SB, MacPherson J, Çoraman E, Yarotskiy V et al. (2023) Can social media be used to inform the distribution of the marbled polecat, *Vormela peregusna*? *Mammal Research* 68: 295–304.
- Yu X, Wang J, Kays R, Jansen PA, Wang T, Huang T (2013) Automated identification of animal species in camera trap images. *EURASIP Journal on Image and Video Processing* 2013: 52.
- Zielinski WJ, Kucera TE (1995) *American Marten, Fisher, Lynx, and Wolverine: Survey Methods for their Detection*. U.S. Department of Agriculture, General Technical Report PSW-GTR-157.
- Zielinski WJ, Thompson CM, Purcell KL, Garner JD (2013) An assessment of fisher (*Pekania pennanti*) tolerance to forest management intensity on the landscape. *Forest Ecology and Management* 310: 821–826.
- Zielinski WJ, Linnell MA, Schwartz MK, Pilgrim K (2020) Exploiting the winter trophic relationships between weasels (*Mustela* spp.) and their microtine prey as a survey method for weasels in meadow ecosystems. *Northwest Science* 93: 185–192.
- Zub K, Sönnichsen L, Szafrńska PA (2008) Habitat requirements of weasels *Mustela nivalis* constrain their impact on prey populations in complex ecosystems of the temperate zone. *Oecologia* 157: 571–582.
- Zub K, García-Díaz P, Sankey S, Eisler R, Lambin X (2022) Using a modeling approach to inform progress towards stoat eradication from the Orkney Islands. *Frontiers in Conservation Science* 2: 780102.