



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Step-Down Gradient Improves the Chromatographic Separation of Sucrose Monocaprato Regioisomers

Lie, Aleksander; Pedersen, Lars Haastrup

Publication date:
2011

Document Version
Early version, also known as pre-print

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Lie, A., & Pedersen, L. H. (2011). *Step-Down Gradient Improves the Chromatographic Separation of Sucrose Monocaprato Regioisomers*. Poster presented at 6th Danish Conference on Biotechnology and Molecular Biology. Syntehtic biology and cell factories, Vejle, Denmark.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Step-Down Gradient Improves the Chromatographic Separation of Sucrose Monocaprato Regioisomers

Aleksander Lie*, Lars H. Pedersen

Department of Biotechnology, Chemistry and Environmental Engineering,
Aalborg University

*alie@bio.aau.dk



Introduction

Sugar fatty acid esters have a broad range of industrial applications and are used in the food, cosmetic and pharmaceutical industries. These compounds can be synthesized by conventional chemical processes or by enzymatic methods in organic solvents. Their physical and chemical properties depend on fatty acid chain length and both position and degree of esterification. The sugar monoesters are the most important types of these compounds because of higher solubility in water compared to the corresponding oligoesters.

In analysing sugar fatty acid monoester syntheses, it is necessary to achieve separation of the regioisomers. Ritthitham et al. (2009) previously demonstrated that when using reversed-phase high-pressure liquid chromatography (RP-HPLC), a step down in the concentration of acetonitrile in the eluent, below the initial concentration, improved the separation of sucrose fatty acid monoesters significantly. They also provided the elution order for seven regioisomers of sucrose monocaprato. Sucrose, shown in Figure 1, has eight hydroxyl substituents, and can form up to eight monoester regioisomers.




Figure 1: Haworth perspective formula of sucrose with carbon atom numbering.

Aim

The aim of the present study was to improve the chromatographic separation of sucrose monocaprato regioisomers using reversed-phase high-pressure liquid chromatography.

Method

A commercial sample of sucrose monocaprato (>95 %) was analysed on an HPLC-system (HP Series 1100) with evaporative light-scattering detector (Alltech ELSD 800). The stationary phase was a C18 column (Waters Symmetry, 5 μ m, 4.6x250mm) and gradients of acetonitrile (CH₃CN) in water (H₂O) were used as mobile phase.




Figure 2: Chromatograms and elution profiles for elution programmes 1 and 2.

I) Elution Factors: A 42 %, B 3 min., C 30 %. II) Elution Factors: A 42 %, B 3 min., C 33 %. All other elution parameters equal.

In tables: t_R – retention time (min.), R_s – peak resolution, H – peak height.

Results

In the present investigation of sucrose monocaprato regioisomers analysed using RP-HPLC, a systematic study of the separation effects was performed. Three elution factors important for separation were identified:

- Initial concentration of CH₃CN in the eluent mixture
- Duration of the initial isocratic elution
- Step-down concentration of CH₃CN in the eluent mixture

In one experiment elution factor C was varied while all other parameters were kept constant, while the effects of variation of elution factors A and B was investigated separately.

References

Ritthitham S, Wimmer R, Stensballe A, Pedersen LH. (2009) Analysis and purification of O-decanoyl sucrose regio-isomers by reversed phase high pressure liquid chromatography with evaporative light scattering detection. Journal of Chromatography A. 1216(25):4963-4967

Figures 2 and 3 show that elution factor C has a significant effect on the separation of the sucrose monoester regioisomers. In figure 2, a change in elution factor C from 30 % to 33 % between elution programmes 1 and 2 resulted in retention times for individual regioisomers being reduced with up to 21 % (peak 4). In addition, the band broadening of individual regioisomers – expressed as the peak height (or height equivalent to a theoretical plate) – was reduced by a factor of up to about 3.5, or increased by a factor of up to about 7.9, with this change in elution factor C (peaks 4 and 5, respectively). Similar results were observed for the change in elution factor C between elution programmes 3 and 4, shown in figure 3.

The results also indicate how the other elution factors affect retention times. Elution factor A has an influence on the retention times of all the sugar monoester peaks. While the effect of elution factor C appear to be dependent on an interaction between elution factors B and C, as the change in elution factor B from 3.5 to 3 minutes significantly reduced the impact of changes in elution factor C on peaks 1-3 (compare figures 2 and 3).




Figure 3: Chromatograms and elution profiles for elution programmes 3 and 4.

I) Elution Factors: A 41 %, B 3.5 min., C 26 %. II) Elution Factors: A 41 %, B 3.5 min., C 33 %. All other elution parameters equal.

In tables: t_R – retention time (min.), R_s – peak resolution, H – peak height.

* These values do not take into account the possible additional peak from this elution programme.

Table 1: Elution factors and separation results for individual elution programmes

Elution Programme	Elution Factors (A-B-C)	Retention time separation of regioisomers**	Standard deviation of peak resolutions	Peak resolution target fit***
1	42-3-30	0.31	1.13	0.56
2	42-3-33	0.38	1.10	0.57
3*	41-3.5-26	0.40	1.59	0.64
4	41-3.5-33	1.22	0.93	1.28
Ritthitham et al.		0.23	1.28	0.81

* The chromatogram suggests the presence of an additional peak. This has not been taken into account for the calculation of these values.

** $\frac{1}{\sum_{i=1}^{n-1} |\Delta t_{R_i} - \Delta t_{R_{i+1}}|}$, where $\Delta t_{R_i} = t_{R_{i+1}} - t_{R_i}$, Δt_{R} is the average of the retention time differences,

and n is the number of regioisomer peaks

*** $\frac{x}{\sum_{i=1}^n |R_{S_i} - 1.5|}$, where n is the number of peak resolutions, and x is the number of $R_S \geq 1.5$

For all the analyses reported here, the separation of the monoester regioisomers was improved compared to previous reports. The optimal elution programme achieved a 400 % improvement in the separation of the sucrose monocaprato regioisomer peaks – based on retention times. In addition, the optimal elution programme shows a reduction in the standard deviation of the peak resolutions, as well as an improvement in how close the peak resolutions are to the goal of 1.5 (baseline separation). In fact, all peak resolutions are above 1.5 for elution programme 4 for the studied sample.

Conclusions

- A step-down gradient can significantly improve the separation of sucrose monocaprato regioisomers in reversed-phase high-pressure liquid chromatography analysis.
- The improvement achieved in the separation of sucrose monocaprato regioisomers in terms of retention times was in the area of 400 %.