



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Comparing the plug-in gait model output to an individually scaled musculoskeletal model using an extensive normative data set

Kersting, Uwe G.; Mogensen, P.; Lund, Morten Enemark; Nielsen, J. F.

Published in:

Proceedings, 4th Annual Meeting of the Danish Society of Biomechanics, 26 October 2012, Aarhus, Denmark

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Kersting, U. G., Mogensen, P., Lund, M. E., & Nielsen, J. F. (2012). Comparing the plug-in gait model output to an individually scaled musculoskeletal model using an extensive normative data set. In *Proceedings, 4th Annual Meeting of the Danish Society of Biomechanics, 26 October 2012, Aarhus, Denmark* (pp. 7). Aarhus University.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Comparing the plug-in gait model output to an individually scaled musculoskeletal model using an extensive normative data set

U.G. Kersting^{1,2}, P. Mogensen², M.E. Lund³, J.F. Nielsen²

¹Center for Sensory-Motor Interaction, Dept. of Health Science and Technology, Aalborg University; ²Gait Laboratory, Hammel Neurocenter, Hammel; ³Dept. of Mechanical and Manufacturing Engineering, Aalborg University

Introduction: Studies have compared walking gait of children to that of adults or between certain age groups, i.e., elderly versus younger adults (e.g., Bleyenheuft & Detrembleur, 2012). Comprehensive reference data bases are hampered by variable marker sets and inter-tester variability being inherent issues (McGinley et al., 2009).

The purpose of this study was to provide a consistent database of healthy controls from childhood to the elderly to be used in clinical studies at our center. Secondly, results obtained from an established gait model were compared to an individually scaled musculoskeletal model.

Methods: A total of 195 subjects, aged from 6 to 86 years, were recruited. Kinematic and kinetic data were recorded by a Vicon system (612, 8 cameras, Vicon, Oxford, UK) in a gait laboratory. Six trials at preferred walking speed were recorded. The Vicon gait marker protocol was used and processed using the plug-in gait (PIG). Additionally, raw data were exported and processed in the AnyBody Modelling System (AMS, AnyBodyTech, Aalborg, DK) using an optimization and scaling algorithm. Age groups were compared by ANOVA and correlations with age and selected anthropometric parameters were performed. Results between models were compared by paired t-tests.

Results: No age dependency of some selected kinematic parameters such as knee angle at TD or the amount of knee flexion during stance was found. Walking velocity did only vary minimally with age while cadence was increased in children compared to all adult age groups. Across all subjects, a significant correlation of leg length and cadence was observed. Integrated ankle joint power increased with age while hip work peaked at age 25 years. Model comparisons revealed off-sets between discrete kinematic parameters while ranges of motion were comparable. This also applies to net joint kinetic parameters. Individual muscle forces or muscle work were not extracted from the PIG calculations but only from the scaled model.

Discussion and Conclusion: In summary, gait velocity was highly consistent despite the fact that subjects chose their preferred walking speed. In healthy subjects, gait kinematics remains reasonably constant over a large age range. There are, however, shifts in total joint work which were opposite to data reported by Karamanidis & Arampatzis (2005). Musculoskeletal models allow for a more detailed analysis of changes in particular muscle groups which correlated well to previous results. It is therefore expected that the inclusion of individualized musculoskeletal models into standard clinical gait analysis will allow for enhanced understanding of alterations in gait function.

References:

- Bleyenheuft C., Detrembleur C. (2012). Clin Biomech Mar 2. [Epub ahead of print].
McGinley J.L., Baker R., Wolfe R., Morris M.E. (2009). Gait Posture, 29(3):360-9.
Karamanidis K., Arampatzis A. (2005). J Exp Biol, 3907-23.




Figure 1: Maximum knee valgus angle during stance vs. age; polynomial fit, $p > 0.05$.




Figure 2: Comparison of total joint work per gait cycle for two age groups. A: Plug-in gait; B: AnyBody Modeling System.