

Southern Illinois University Carbondale OpenSIUC

Miscellaneous (presentations, translations,
interviews, etc)

Department of Mathematics

9-19-2000

Stochastic Dynamics of Infinite-Dimensional Systems (Stochastic and Non-linear Analysis Seminar, University of Illinois)

Salah-Eldin A. Mohammed

Southern Illinois University Carbondale, salah@sfde.math.siu.edu

Follow this and additional works at: http://opensiuc.lib.siu.edu/math_misc

 Part of the [Mathematics Commons](#)

Stochastic and Non-linear Analysis Seminar; Department of Mathematics; University of Illinois;
Urbana-Champaign, Illinois; September 19, 2000

Recommended Citation

Mohammed, Salah-Eldin A., "Stochastic Dynamics of Infinite-Dimensional Systems (Stochastic and Non-linear Analysis Seminar, University of Illinois)" (2000). *Miscellaneous (presentations, translations, interviews, etc)*. Paper 18.
http://opensiuc.lib.siu.edu/math_misc/18

This Article is brought to you for free and open access by the Department of Mathematics at OpenSIUC. It has been accepted for inclusion in Miscellaneous (presentations, translations, interviews, etc) by an authorized administrator of OpenSIUC. For more information, please contact opensiuc@lib.siu.edu.

**STOCHASTIC DYNAMICS
OF
INFINITE-DIMENSIONAL
SYSTEMS**

Urbana: September 19, 2000

Salah-Eldin A. Mohammed

Southern Illinois University

Carbondale, IL 62901-4408

Web site: <http://sfde.math.siu.edu>

Stable Manifolds

Outline

- Smooth cocycles in Hilbert space. Stationary trajectories.
- Linearization of a cocycle along a stationary trajectory.
- Ergodic theory of cocycles in Hilbert space.
- Hyperbolicity of stationary trajectories. Lyapunov exponents.
- Cocycles generated by stochastic systems with memory. Via random diffeomorphism groups.
- *Local Stable Manifold Theorem* for stochastic differential equations with memory (SFDE's): Existence of smooth stable and unstable manifolds in a neighborhood of a hyperbolic stationary trajectory.
- Proof: Ruelle-Oseledec multiplicative ergodic theory + perfection techniques.

The Cocycle

$(\Omega, \mathcal{F}, P) :=$ complete probability space.

$\theta : \mathbf{R}^+ \times \Omega \rightarrow \Omega$ a P -preserving (ergodic) semi-group on (Ω, \mathcal{F}, P) .


$E :=$ real (separable) Hilbert space, norm $\|\cdot\|$, Borel σ -algebra.

Definition.

$k =$ non-negative integer, $\epsilon \in (0, 1]$. A $C^{k,\epsilon}$ *perfect cocycle* (X, θ) on E is a measurable random field $X : \mathbf{R}^+ \times E \times \Omega \rightarrow E$ such that:

- (i) For each $\omega \in \Omega$, the map $\mathbf{R}^+ \times E \ni (t, x) \mapsto X(t, x, \omega) \in E$ is continuous; for fixed $(t, \omega) \in \mathbf{R}^+ \times \Omega$, the map $E \ni x \mapsto X(t, x, \omega) \in E$ is $C^{k,\epsilon}$ ($D^k X(t, x, \omega)$ is C^ϵ in x).
- (ii) $X(t_1 + t_2, \cdot, \omega) = X(t_2, \cdot, \theta(t_1, \omega)) \circ X(t_1, \cdot, \omega)$ for all $t_1, t_2 \in \mathbf{R}^+$, all $\omega \in \Omega$.
- (iii) $X(0, x, \omega) = x$ for all $x \in E, \omega \in \Omega$.

Cocycle Property



Vertical solid lines represent random fibers: copies of E . (X, θ) is a “vector-bundle morphism”.

Definition

A random variable $Y : \Omega \rightarrow E$ is a *stationary point* for the cocycle (X, θ) if

$$X(t, Y(\omega), \omega) = Y(\theta(t, \omega)) \quad (1)$$

for all $t \in \mathbf{R}$ and every $\omega \in \Omega$. Denote stationary trajectory (1) by $X(t, Y) = Y(\theta(t))$.

Linearization. Hyperbolicity.

Linearize a $C^{k,\epsilon}$ cocycle (X, θ) along a stationary random point Y : Get an $L(E)$ -valued cocycle $(DX(t, Y(\omega), \omega), \theta(t, \omega))$. (Follows from cocycle property of X and chain rule.)

Theorem. (*Oseledec-Ruelle*)

Let $T : \mathbf{R}^+ \times \Omega \rightarrow L(E)$ be strongly measurable, such that (T, θ) is an $L(E)$ -valued cocycle, with each $T(t, \omega)$ compact. Suppose that

$$E \sup_{0 \leq t \leq 1} \log^+ \|T(t, \cdot)\|_{L(E)} < \infty,$$

$$E \sup_{0 \leq t \leq 1} \log^+ \|T(1-t, \theta(t, \cdot))\|_{L(E)} < \infty.$$

Then there is a sure event $\Omega_0 \in \mathcal{F}$ such that $\theta(t, \cdot)(\Omega_0) \subseteq \Omega_0$ for all $t \in \mathbf{R}^+$, and for each $\omega \in \Omega_0$,

$$\lim_{t \rightarrow \infty} [T(t, \omega)^* \circ T(t, \omega)]^{1/(2t)} := \Lambda(\omega)$$

exists in the uniform operator norm. $\Lambda(\omega)$ is self-adjoint with a non-random spectrum

$$e^{\lambda_1} > e^{\lambda_2} > e^{\lambda_3} > \dots$$

where the λ_i 's are distinct. Each $e^{\lambda_i} > 0$ has a fixed finite non-random multiplicity m_i and eigen-space $F_i(\omega)$, with $m_i := \dim F_i(\omega)$. Set $i = \infty$ when $\lambda_i = -\infty$. Define

$$E_1(\omega) := E, \quad E_i(\omega) := \left[\bigoplus_{j=1}^{i-1} F_j(\omega) \right]^\perp, \quad i > 1, \quad E_\infty := \ker \Lambda(\omega).$$

Then

$$E_\infty \subset \cdots \subset \cdots \subset E_{i+1}(\omega) \subset E_i(\omega) \cdots \subset E_2(\omega) \subset E_1(\omega) = E,$$

$$\lim_{t \rightarrow \infty} \frac{1}{t} \log \|T(t, \omega)x\| = \begin{cases} \lambda_i & \text{if } x \in E_i(\omega) \setminus E_{i+1}(\omega), \\ -\infty & \text{if } x \in E_\infty(\omega), \end{cases}$$

and

$$T(t, \omega)(E_i(\omega)) \subseteq E_i(\theta(t, \omega))$$

for all $t \geq 0$, $i \geq 1$.


Proof.

Based on discrete version of Oseledec's multiplicative ergodic theorem and the perfect ergodic theorem. ([Ru.1], I.H.E.S Publications, 1979, pp. 303-304; cf. Furstenberg & Kesten (1960), [Mo.1]). \square

Lyapunov Spectrum:

$\{\lambda_1, \lambda_2, \lambda_3, \dots\} := \text{Lyapunov spectrum of } (T, \theta).$

Spectral Theorem



Definition

A stationary point $Y(\omega)$ of (X, θ) is *hyperbolic* if the linearized cocycle $(DX(t, Y(\omega), \omega), \theta(t, \omega))$ has

a non-vanishing Lyapunov spectrum $\{\dots < \lambda_{i+1} < \lambda_i < \dots < \lambda_2 < \lambda_1\}$, viz. $\lambda_i \neq 0$ for all $i \geq 1$.

Let $i_0 > 1$ be s.t. $\lambda_{i_0} < 0 < \lambda_{i_0-1}$.

Assume $X(t, \cdot, \omega)$ *locally compact* and

$$E \log^+ \sup_{0 \leq t_1, t_2 \leq r} \|D_2 X(t_2, Y(\theta(t_1)), \theta(t_1))\|_{L(E)} < \infty.$$

By Oseledec-Ruelle Theorem, there is a sequence of closed finite-codimensional (Oseledec) spaces

$$\dots E_{i-1}(\omega) \subset E_i(\omega) \subset \dots \subset E_2(\omega) \subset E_1(\omega) = E,$$

$$E_i(\omega) = \{x \in E : \lim_{t \rightarrow \infty} \frac{1}{t} \log \|DX(t, Y(\omega), \omega)(x)\| \leq \lambda_i\},$$

$i \geq 1$ and all $\omega \in \Omega^*$, a sure event in \mathcal{F} satisfying $\theta(t, \cdot)(\Omega^*) = \Omega^*$ for all $t \in \mathbf{R}$.

Let $\{U(\omega), S(\omega) : \omega \in \Omega^*\}$ be the unstable and stable subspaces associated with the linearized cocycle (DX, θ) ([Mo.1], Theorem 4, Corollary 2; [M-S.1], Theorem 5.3). Then get a measurable invariant splitting

$$E = \mathcal{U}(\omega) \oplus \mathcal{S}(\omega), \quad \omega \in \Omega^*,$$


$$DX(t, Y(\omega), \omega)(\mathcal{U}(\omega)) = \mathcal{U}(\theta(t, \omega)), \quad DX(t, Y(\omega), \omega)(\mathcal{S}(\omega)) \subseteq \mathcal{S}(\theta(t, \omega)),$$

for all $t \geq 0$, with *exponential dichotomies*

$$\|DX(t, Y(\omega), \omega)(x)\| \geq \|x\|e^{\delta_1 t} \quad \text{for all } t \geq \tau_1^*, x \in \mathcal{U}(\omega),$$

$$\|DX(t, Y(\omega), \omega)(x)\| \leq \|x\|e^{-\delta_2 t} \quad \text{for all } t \geq \tau_2^*, x \in \mathcal{S}(\omega),$$

with $\tau_i^* = \tau_i^*(x, \omega) > 0, i = 1, 2$, random times and $\delta_i > 0, i = 1, 2$, fixed.



Nonlinear Stochastic Systems with Memory

“Regular” Itô SFDE with finite memory:

$$\left. \begin{aligned} dx(t) &= H(x(t), x_t) dt + \sum_{i=1}^m G_i(x(t)) dW_i(t), \\ (x(0), x_0) &= (v, \eta) \in M_2 := \mathbf{R}^d \times L^2([-r, 0], \mathbf{R}^d) \end{aligned} \right\} \quad (I)$$

Solution segment $x_t(s) := x(t + s)$, $t \geq 0, s \in [-r, 0]$.

m -dimensional Brownian motion $W := (W_1, \dots, W_m)$, $W(0) = 0$.

Ergodic Brownian shift θ on Wiener space (Ω, \mathcal{F}, P) . $\bar{\mathcal{F}} := P$ -completion of \mathcal{F} .

State space M_2 , Hilbert with usual norm $\|\cdot\|$.

Can allow for “smooth memory” in diffusion coefficient.

$H : M_2 \rightarrow \mathbf{R}^d$, $C^{k,\delta}$, globally bounded.

$G : \mathbf{R}^d \rightarrow L(\mathbf{R}^m, \mathbf{R}^d)$, $C_b^{k+1,\delta}$; $G := (G_1, \dots, G_m)$.

$B((v, \eta), \rho)$ open ball, radius ρ , center $(v, \eta) \in M_2$;

$\bar{B}((v, \eta), \rho)$ closed ball.

Then (I) has a stochastic semiflow $X : \mathbf{R}^+ \times M_2 \times \Omega \rightarrow M_2$ with $X(t, (v, \eta), \cdot) = (x(t), x_t)$. X is $C^{k, \epsilon}$ for any $\epsilon \in (0, \delta)$, takes bounded sets into relatively compact sets in M_2 . (X, θ) is a perfect cocycle on M_2 ([M-S.4]).

Example

Consider the affine linear sfde

$$\left. \begin{aligned} dx(t) &= H(x(t), x_t) dt + G dW(t), & t > 0 \\ x(0) &= v \in \mathbf{R}^d, & x_0 = \eta \in L^2([-r, 0], \mathbf{R}^d) \end{aligned} \right\} (I')$$

where $H : M_2 \rightarrow \mathbf{R}^d$ is a continuous linear map, G is a fixed $(d \times p)$ -matrix, and W is p -dimensional Brownian motion. Assume that the linear deterministic $(d \times d)$ -matrix-valued FDE

$$dy(t) = H \circ (y(t), y_t) dt$$

has a semiflow

$$T_t : L(\mathbf{R}^d) \times L^2([-r, 0], L(\mathbf{R}^d)) \rightarrow L(\mathbf{R}^d) \times L^2([-r, 0], L(\mathbf{R}^d)), t \geq 0,$$

which is uniformly asymptotically stable. Set

$$Y := \int_{-\infty}^0 T_{-u}(I, 0)G dW(u) \quad (2)$$

where I is the identity $(d \times d)$ -matrix. Integration by parts and

$$W(t, \theta(t_1, \omega)) = W(t + t_1, \omega) - W(t_1, \omega), \quad t, t_1 \in \mathbf{R}, \quad (3)$$

imply that Y has a measurable version satisfying (1). Y is Gaussian and thus has finite moments of all orders. See ([Mo.1], Theorem 4.2, Corollary 4.2.1, pp. 208-217.) More generally, when H is hyperbolic, one can show that a stationary point of (I') exists ([Mo.1]).

In the general white-noise case an invariant measure on M_2 for the one-point motion gives rise to a stationary point provided we suitably enlarge the underlying probability space. Conversely, let $Y : \Omega \rightarrow M_2$ be a stationary random point independent of the Brownian motion $W(t)$, $t \geq 0$. Let $\rho := P \circ Y^{-1}$ be the distribution of Y . By independence of Y and W , ρ is an invariant measure for the one-point motion

Theorem. ([M-S], 2000) (*The Stable Manifold Theorem*)

Assume smoothness hypotheses on H and G . Let $Y : \Omega \rightarrow M_2$ be a hyperbolic stationary point of the SFDE (I) such that $E(\|Y(\cdot)\|^{\epsilon_0}) < \infty$ for some $\epsilon_0 > 0$

Suppose the linearized cocycle $(DX(t, Y(\omega), \omega), \theta(t, \omega), t \geq 0)$ of (I) has a Lyapunov spectrum $\{\dots < \lambda_{i+1} < \lambda_i < \dots < \lambda_2 < \lambda_1\}$. Define $\lambda_{i_0} := \max\{\lambda_i : \lambda_i < 0\}$ if at least one $\lambda_i < 0$. If all finite λ_i are positive, set $\lambda_{i_0} = -\infty$. (This implies that λ_{i_0-1} is the smallest positive Lyapunov exponent of the linearized semiflow, if at least one $\lambda_i > 0$; in case all λ_i are negative, set $\lambda_{i_0-1} = \infty$.)

Fix $\epsilon_1 \in (0, -\lambda_{i_0})$ and $\epsilon_2 \in (0, \lambda_{i_0-1})$. Then there exist

- (i) a sure event $\Omega^* \in \mathcal{F}$ with $\theta(t, \cdot)(\Omega^*) = \Omega^*$ for all $t \in \mathbf{R}$,
- (ii) $\bar{\mathcal{F}}$ -measurable random variables $\rho_i, \beta_i : \Omega^* \rightarrow (0, 1)$, $\beta_i > \rho_i > 0$, $i = 1, 2$, such that for each $\omega \in \Omega^*$, the following is true:

There are $C^{k, \epsilon}$ ($\epsilon \in (0, \delta)$) submanifolds $\tilde{\mathcal{S}}(\omega), \tilde{\mathcal{U}}(\omega)$ of $\bar{B}(Y(\omega), \rho_1(\omega))$ and $\bar{B}(Y(\omega), \rho_2(\omega))$ (resp.) with the following properties:

(a) $\tilde{\mathcal{S}}(\omega)$ is the set of all $(v, \eta) \in \bar{B}(Y(\omega), \rho_1(\omega))$ such that

$$\|X(nr, (v, \eta), \omega) - Y(\theta(nr, \omega))\| \leq \beta_1(\omega) e^{(\lambda_{i_0} + \epsilon_1)nr}$$

for all integers $n \geq 0$. Furthermore,

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \|X(t, (v, \eta), \omega) - Y(\theta(t, \omega))\| \leq \lambda_{i_0}$$

for all $(v, \eta) \in \tilde{\mathcal{S}}(\omega)$. Each stable subspace $\mathcal{S}(\omega)$ of the linearized semiflow DX is tangent at $Y(\omega)$ to the submanifold $\tilde{\mathcal{S}}(\omega)$, viz. $T_{Y(\omega)}\tilde{\mathcal{S}}(\omega) = \mathcal{S}(\omega)$. In particular, $\text{codim } \tilde{\mathcal{S}}(\omega) = \text{codim } \mathcal{S}(\omega)$, is fixed and finite.

$$(b) \limsup_{t \rightarrow \infty} \frac{1}{t} \log \left[\sup \left\{ \frac{\|X(t, (v_1, \eta_1), \omega) - X(t, (v_2, \eta_2), \omega)\|}{\|(v_1, \eta_1) - (v_2, \eta_2)\|} : (v_1, \eta_1) \neq (v_2, \eta_2), (v_1, \eta_1), (v_2, \eta_2) \in \tilde{\mathcal{S}}(\omega) \right\} \right] \leq \lambda_{i_0}.$$

(c) (Cocycle-invariance of the stable manifolds):

There exists $\tau_1(\omega) \geq 0$ such that

$$X(t, \cdot, \omega)(\tilde{\mathcal{S}}(\omega)) \subseteq \tilde{\mathcal{S}}(\theta(t, \omega))$$

for all $t \geq \tau_1(\omega)$. Also

$$DX(t, Y(\omega), \omega)(\mathcal{S}(\omega)) \subseteq \mathcal{S}(\theta(t, \omega)), \quad t \geq 0.$$

(d) $\tilde{\mathcal{U}}(\omega)$ is the set of all $(v, \eta) \in \bar{B}(Y(\omega), \rho_2(\omega))$ with the property that there is a unique “history” process $y(\cdot, \omega) : \{-nr : n \geq 0\} \rightarrow M_2$ such that $y(0, \omega) = (v, \eta)$ and for each integer $n \geq 1$, one has $X(r, y(-nr, \omega), \theta(-nr, \omega)) = y(-(n-1)r, \omega)$ and

$$\|y(-nr, \omega) - Y(\theta(-nr, \omega))\|_{M_2} \leq \beta_2(\omega) e^{-(\lambda_{i_0} - 1 - \epsilon_2)nr}.$$

Furthermore, for each $(v, \eta) \in \tilde{\mathcal{U}}(\omega)$, there is a unique continuous-time “history” process also denoted by $y(\cdot, \omega) : (-\infty, 0] \rightarrow M_2$ such that $y(0, \omega) = (v, \eta)$, $X(t, y(s, \omega), \theta(s, \omega)) = y(t+s, \omega)$ for all $s \leq 0, 0 \leq t \leq -s$, and

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \|y(-t, \omega) - Y(\theta(-t, \omega))\| \leq -\lambda_{i_0} - 1.$$

Each unstable subspace $\mathcal{U}(\omega)$ of the linearized semi-flow DX is tangent at $Y(\omega)$ to $\tilde{\mathcal{U}}(\omega)$, viz. $T_{Y(\omega)}\tilde{\mathcal{U}}(\omega) = \mathcal{U}(\omega)$. In particular, $\dim \tilde{\mathcal{U}}(\omega)$ is finite and non-random.

(e) Let $y(\cdot, (v_i, \eta_i), \omega), i = 1, 2$, be the history processes associated with $(v_i, \eta_i) = y(0, (v_i, \eta_i), \omega) \in \tilde{\mathcal{U}}(\omega), i = 1, 2$. Then

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \log \left[\sup \left\{ \frac{\|y(-t, (v_1, \eta_1), \omega) - y(-t, (v_2, \eta_2), \omega)\|}{\|(v_1, \eta_1) - (v_2, \eta_2)\|} : \right. \right. \\ \left. \left. (v_1, \eta_1) \neq (v_2, \eta_2), (v_i, \eta_i) \in \tilde{\mathcal{U}}(\omega), i = 1, 2 \right\} \right] \\ \leq -\lambda_{i_0-1}.$$

(f) (Cocycle-invariance of the unstable manifolds):

There exists $\tau_2(\omega) \geq 0$ such that

$$\tilde{\mathcal{U}}(\omega) \subseteq X(t, \cdot, \theta(-t, \omega))(\tilde{\mathcal{U}}(\theta(-t, \omega)))$$

for all $t \geq \tau_2(\omega)$. Also

$$DX(t, \cdot, \theta(-t, \omega))(\mathcal{U}(\theta(-t, \omega))) = \mathcal{U}(\omega), \quad t \geq 0;$$

and the restriction

$$DX(t, \cdot, \theta(-t, \omega))|_{\mathcal{U}(\theta(-t, \omega))} : \mathcal{U}(\theta(-t, \omega)) \rightarrow \mathcal{U}(\omega), \quad t \geq 0,$$

is a linear homeomorphism onto.


(g) *The submanifolds $\tilde{\mathcal{U}}(\omega)$ and $\tilde{\mathcal{S}}(\omega)$ are transversal, viz.*

$$M_2 = T_{Y(\omega)}\tilde{\mathcal{U}}(\omega) \oplus T_{Y(\omega)}\tilde{\mathcal{S}}(\omega).$$

Assume, in addition, that H, G are C_b^∞ . Then the local stable and unstable manifolds $\tilde{\mathcal{S}}(\omega), \tilde{\mathcal{U}}(\omega)$ are C^∞ .

Figure summarizes essential features of Stable Manifold Theorem:

Stable Manifold Theorem



A picture is worth a 1000 words!

Outline of Proof

- By definition, a *stationary* random point $Y(\omega) \in M_2$ is invariant under the semiflow X ; viz $X(t, Y) = Y(\theta(t, \cdot))$ for all times t .
- Linearize the semiflow X along the stationary point $Y(\omega)$ in M_2 . By stationarity of Y and the cocycle property of X , this gives a linear perfect cocycle $(DX(t, Y), \theta(t, \cdot))$ in $L(M_2)$, where $D = \text{spatial}$ (Fréchet) derivatives.
- Ergodicity of θ allows for the notion of *hyperbolicity* of a stationary solution of (I) via Oseledec-Ruelle theorem: Use local compactness of the semiflow for times greater than the delay r ([M-S.4]), and apply multiplicative ergodic theorem to get a discrete non-random Lyapunov spectrum $\{\lambda_i : i \geq 1\}$ for the linearized cocycle. Y is *hyperbolic* if $\lambda_i \neq 0$ for every i .
- Assume that $\|Y\|^{\epsilon_0}$ is integrable (for small ϵ_0). Variational method of construction of the semiflow shows that the linearized cocycle satisfies hypotheses of “perfect versions”

of ergodic theorem and Kingman's subadditive ergodic theorem. These refined versions give invariance of the Oseledec spaces under the continuous-time linearized cocycle. Thus the stable/unstable subspaces will serve as tangent spaces to the local stable/unstable manifolds of the non-linear semiflow X .

- Establish continuous-time integrability estimates on the spatial derivatives of the non-linear cocycle X in a neighborhood of the stationary point Y . Estimates follow from the variational construction of the stochastic semiflow coupled with known global spatial estimates for finite-dimensional stochastic flows.
- Introduce the auxiliary perfect cocycle

$$Z(t, \cdot, \omega) := X(t, (\cdot) + Y(\omega), \omega) - Y(\theta(t, \omega)), \quad t \in \mathbf{R}^+, \omega \in \Omega.$$

Refine arguments in ([Ru.2], Theorems 5.1 and 6.1) to construct local stable/unstable manifolds for the discrete cocycle $(Z(nr, \cdot, \omega), \theta(nr, \omega))$ near 0 and hence (by translation) for $X(nr, \cdot, \omega)$

near $Y(\omega)$ for all ω sampled from a $\theta(t, \cdot)$ -invariant sure event in Ω . This is possible because of the continuous-time integrability estimates, the perfect ergodic theorem and the perfect subadditive ergodic theorem. By interpolating between delay periods of length r and further refining the arguments in [Ru.2], show that the above manifolds also serve as local stable/unstable manifolds for the *continuous-time* semiflow X near Y .

- Final key step: Establish the asymptotic invariance of the local stable manifolds under the stochastic semiflow X . Use arguments underlying the proofs of Theorems 4.1 and 5.1 in [Ru.2] and some difficult estimates using the continuous-time integrability properties, and the perfect subadditive ergodic theorem. Asymptotic invariance of the local unstable manifolds follows by employing the concept of a *stochastic history process* for X coupled with similar arguments to the above. Existence of history process compensates for the lack of invertibility of the semiflow.

REFERENCES

- [A-S] Arnold, L., and Scheutzow, M. K. R., Perfect cocycles through stochastic differential equations, *Probab. Th. Rel. Fields*, 101, (1995), 65-88.
- [C] Carverhill, A., Flows of stochastic dynamical systems: Ergodic theory, *Stochastics*, 14 (1985), 273-317.
- [I-W] Ikeda, N., and Watanabe, S., *Stochastic Differential Equations and Diffusion Processes*, Second Edition, North-Holland-Kodansha (1989).
- [Ku] Kunita, H., *Stochastic Flows and Stochastic Differential Equations*, Cambridge University Press, Cambridge, New York, Melbourne, Sydney (1990).
- [Mo.1] Mohammed, S.-E. A., The Lyapunov spectrum and stable manifolds for stochastic linear delay equations, *Stochastics and Stochastic Reports*, Vol. 29 (1990), 89-131.
- [Mo.2] Mohammed, S.-E. A., Lyapunov exponents and stochastic flows of linear and affine hereditary Systems, *Diffusion Processes and Related Problems in Analysis*, Vol. II, edited

by Mark Pinsky and Volker Wihstutz, Birkhauser (1992), 141-169.

- [M-S.1] Mohammed, S.-E. A., and Scheutzow, M. K. R., Lyapunov exponents of linear stochastic functional differential equations driven by semimartingales, Part I: The multiplicative ergodic theory, *Ann. Inst. Henri Poincaré, Probabilités et Statistiques*, Vol. 32, 1, (1996), 69-105. pp. 43.
- [M-S.2] Mohammed, S.-E. A., and Scheutzow, M. K. R., Spatial estimates for stochastic flows in Euclidean space, *The Annals of Probability*, 26, 1, (1998), 56-77.
- [M-S.3] Mohammed, S.-E. A., and Scheutzow, M. K. R., The stable manifold theorem for stochastic differential equations, *The Annals of Probability*, Vol. 27, No. 2, (1999), 615-652.
- [M-S.4] Mohammed, S.-E. A., and Scheutzow, M. K. R., The stable manifold theorem for nonlinear stochastic hereditary systems. Part I: Existence of the semiflow, Part II: The local stable manifold theorem. (preprints).
- [O] Oseledec, V. I., A multiplicative ergodic theorem. Lyapunov characteristic numbers for

dynamical systems, *Trudy Moskov. Mat. Obšč.* 19 (1968), 179-210. English transl. *Trans. Moscow Math. Soc.* 19 (1968), 197-221.

[Ru.1] Ruelle, D., Ergodic theory of differentiable dynamical systems, *Publ. Math. Inst. Hautes Etud. Sci.* (1979), 275-306.

[Ru.2] Ruelle, D., Characteristic exponents and invariant manifolds in Hilbert space, *Annals of Mathematics* 115 (1982), 243–290.

[Wa] Wanner, T., Linearization of random dynamical systems, in *Dynamics Reported*, vol. 4, edited by U. Kirchgraber and H.O. Walther, Springer (1995), 203-269.