

Southern Illinois University Carbondale
OpenSIUC

Honors Theses

University Honors Program

8-1999

Production of an Improved Vector for Synthesis and Purification of Eukaryotic Transcription Factors (GAL4-VP16, GAL4-AD and GAL4-D)

Pedro Oyarbide Valencia
Southern Illinois University Carbondale

Follow this and additional works at: http://opensiuc.lib.siu.edu/uhp_theses

Recommended Citation

Valencia, Pedro Oyarbide, "Production of an Improved Vector for Synthesis and Purification of Eukaryotic Transcription Factors (GAL4-VP16, GAL4-AD and GAL4-D)" (1999). *Honors Theses*. Paper 129.

This Dissertation/Thesis is brought to you for free and open access by the University Honors Program at OpenSIUC. It has been accepted for inclusion in Honors Theses by an authorized administrator of OpenSIUC. For more information, please contact opensiuc@lib.siu.edu.

**Production of an Improved Vector for Synthesis and Purification
of Eukaryotic Transcription Factors (GAL4-VP16, GAL4-AD and GAL4-D)**

Pedro Oyarbide Valencia

August 4, 1999

Undergraduate Honors Thesis, Department of Microbiology

Laboratory of Medical Biochemistry

Southern Illinois University Carbondale

Carbondale, Illinois 62901 U.S.A.

Table of Contents

I.	Introduction	1
II.	Abstract	4
III.	Materials & Methods	5
	A. BMH Competent Cell Preparation (for Electroporation)	5
	B. Dephosphorilation of the 5' End of the pET21d Vector	6
	C. Preparation of the pET21d Vector for the Later Sub-Cloning of the GAL4 Transcription Factors	7
	D. Transformation of Constructs into DH5alfa Competent Cells	8
	E. Preparation of the GAL4-AD and GAL4-D Inserts	10
	F. GAL4-VP16 Sequencing Reaction	11
	G. GAL4-VP16 Transcription Factor DNA Sequencing Reaction	14
	H. BIGGERprep Plasmid DNA Preparation Procedure	16
	I. GAL4-VP16 Transcription Factor Amplification (Taq Polymerase Protocol)	18
	J. GAL4-VP16 Transcription Factor Amplification (Pfu Turbo Protocol)	19
	K. Ligation of the Dephosphorilated pET21d Vector and the GAL4-AD and GAL4-D Transcription Factors	20
	L. Digestion of the GAL4-AD and GAL4-D Inserts with NcoI and XhoI Restriction Enzymes	21
	M. TB Broth & 2 x YT Media Preparation	22
IV.	Experimental Results	23
V.	Discussion	25
VI.	Appendix A, Primer & GAL4-VP16 Sequences and Genetic Maps	29
VII.	Appendix B, Materials & Methods	40
VIII.	Appendix C, Figures	42
	A. Figure 1: pET21d Digestion by XhoI and NcoI	42
	B. Figure 2: GAL4-AD and GAL4-D Digested PCR Inserts	43
	C. Figure 3: GAL4-AD and GAL4-D Ligation Products	44

D. Figure 4: Taq Polymerase Protocol Amplifications of GAL4-VP16	45
E. Figure 5: Pfu Turbo Protocol Amplifications of GAL4-VP16	46
F. Figure 6: GAL4 Series Amplification Products	47
IX. Acknowledgements	48
X. References	49

Introduction

In recent years, the study of eukaryotic gene expression has evolved due to technological developments that have greatly increased knowledge and understanding of the subject. There are many different types of molecules involved in the process of gene expression. Some of them have become critically important due to their enhanced expression activation properties. These are called transcription factors (TF), and the following study is focused on the sub-cloning and expression of the GAL4 TF series for the later study of their particular activities and the nature of their interactions in the chromatin remodeling process.

In general TF as well as other regulatory proteins, possess the following two abilities: (1) the ability to recognize specific target sequences located in the DNA template that belong to enhancers, promoters and other regulatory elements that affect a particular gene, and; (2) the ability, once bound to DNA, to exercise their function by binding to other components of the transcription apparatus. These TFs can be divided into two different groups according to their aminoacid residue composition. The first group includes general transcription factors, such as TFIID, TFIIB, while the second group is made up of specific or acidic transcription factors such as GAL4 series. The latter possesses a higher percentage of acidic aminoacid residues in its primary structure.

Regarding this particular series of acidic transcriptional activators, a wide array of features have been attributed to them. This thesis mainly focuses on the GAL4-AD, GAL4-D and GAL4-VP16 transcriptional activators. The GAL4 series members are transcriptional activators (TA) of fungal origin, composed of several functional domains that include the following:

a characteristic cysteine rich DNA binding domain common to all members of the family, a dimerization domain, various transactivation domains generally exhibiting a high acidic content and a highly variable central region, that is supposed to be involved in regulation and in effector recognition. (Poch 1997: 229)

One hypothesis gives credence to the existence of eight highly conserved functional motifs. These motifs are located in a large domain of approximately 225-405 residues. One researcher found that "its inhibitory activity may be mediated by hydrophobic interactions linked to the presence of amphipathic alpha-helices" (Poch 1997: 229). Therefore, the function of a transcription factor depends directly on the type of interaction of its different motifs. Differences in functional motifs constitute the differences between transcription factors. There is a molecular basis for the differentiation between TFs in terms of their ability to bind sites within chromatin. GAL4 is known to interact with its cognate site when it is assembled into nucleosome core particles, meaning that its affinity for nucleosomally organized target sites is lower than for the same sites in free DNA (Beato 1997: 3561). It has also been demonstrated that GAL4 derivatives can potentiate replication-independent transcription in vitro from preassembled chromatin templates. For instance, the GAL4-VP16 acidic activator is able to enhance transcription of a template assembled into chromatin in vitro if added only during, not after, DNA replication (Xu 1998: 1201). As a general trait, most eukaryotic gene promoters contain multiple binding sites for one or more transcriptional activators that interact in a synergistic manner. This is the manifestation of the need of many contacts between activators and the general transcription machinery. While normally most of the transcription activators, (TA), need several contacts to elicit the maximum level of

possible transcriptional activity, GAL4 can produce the same activation pattern with only one contact under physiological conditions (Xu 1995: 7677).

These features are most useful to researchers in understanding eukaryotic gene regulation. It is necessary to determine sequences within these TF that will interact with DNA and which are the target sequences that they bind preferentially. For this reason, this study investigates segments of TA. The GAL4-AD, a 34-amino acid region at the carboxy terminus end of the GAL4 TF (Melcher 1995: 2839) has been proved to specifically and directly bind to the TATA-binding protein (TBP). Other studies have shown that for GAL4 TF the palindromic CGG triplets at the ends of the 17-bp recognition site in DNA are essential for tight binding, whereas the identity of the remaining internal 11-bp are of much less importance (Liang 1996: 3773).

The study of both primary sequences and more complex structures in GAL4 derivatives has led to suppositions of several other hypotheses regarding their stability and the driving force for their interactions with DNA. For example, the binuclear metal ion structure present in these TF, is shown to have a net negative charge of -2. This fact has allowed for hypothesizing that this metal ion is the recipient of several hydrogen bonds, notably from the main chain amide protons of the ligating cysteine residues, indicating the charge is stabilized in this manner (Gadhavi 1997: 145).

Abstract

The main goal in conducting this series of experiments is to accomplish the construction of an expression vector that contains the coding regions for the GAL4 series transcription factors. This expression vector should possess special characteristics including a histidine tag that helps to easily locate the vector. The other main characteristic is a T7 promoter that allows for a high induction level upon addition of IPTG. The original pET21d expression vector was used in these experiments because it possesses these two intrinsic features. A sub-cloning process of the coding regions of the GAL4 series transcription factors (GAL4-AD, GAL4-D and GAL4-VP16) into the original pET21d expression vector resulted in the pET21d-GAL4 expression vector. The coding regions for the GAL4-AD and GAL4-D transcription factors were extracted from a pGEX-cs vector by Polymerase Chain Reaction (PCR) amplification with specially designed primers. The GAL4-VP16 transcription factor was extracted from the pJL2 plasmid. In the case of the GAL4-VP16 transcription factor, the sequence was not initially available and experimental sequencing of the transcription factor was necessary to acquire enough information about the sequence to design the primers for the PCR amplification¹.

¹ See Appendix A for sequences and genetic maps for the three different transcription factors, primers and expression vectors.

Materials and Methods²

A detailed description of the experimental procedures, the conditions in which the experiments were performed and the pertinent information about the materials used will clarify the specifics for conducting these experiments.

BMH Competent Cell Preparation (for Electroporation)³

1. Inoculate a single colony of BMH into 5 mL LB + 40 ul of tetracycline at 37° C overnight.
2. Inoculate 5 mL culture in 1 L LB grow at 37° C until the OD at 600 nm. Reaches the interval [0.5 - 0.7].
3. Chill on ice for 10 minutes. Pour into a 1 L centrifuge bottle that has been pre-chilled.
4. Pellet 1 L of culture at 4000 g at 4° C for 15 minutes.
5. Resuspend in 1 L of sterile, cold H₂O. Pellet at 4000 g at 4° C for 15 minutes.
6. Resuspend in 500 mL of sterile, cold H₂O. Pellet at 4000 g at 4° C for 15 minutes.
7. Resuspend in 500 mL of sterile, cold H₂O. Pellet at 4000 g at 4° C for 15 minutes.
8. Resuspend in 20 mL of sterile, cold H₂O. Pellet at 4000 g at 4° C for 15 minutes.
9. Resuspend in 2 mL sterile, cold 10% glycerol (filter sterilized).
10. Aliquots may be frozen at -70° C.

² A detailed listing of the experimental procedures used, as well as a listing of the main materials, are available in appendix B.

³ Medical Biochemistry, Lab 207, June 14, 1998.

Dephosphorilation of the 5' End of the pET21d Vector⁴

1. Take the sample out of the -20° C storage and thaw.

Sample	l
pET21d vector, digested with NcoI and XhoI	40 ul
10X SAP buffer	4.5 ul
Shrimp Alkaline Phosphatase (1U/ul)	1 ul
Total volume	45.5 ul

2. Incubate the mixture for 60 minutes at 37° C.
3. Reload the same amount of enzyme after the first 60 minutes, and expose the mixture for another 60 minutes in the same conditions.
4. After 120 minutes of incubation, heat deactivate the shrimp alkaline phosphatase by introducing the mixture into a 65° C bath for 15 minutes.
5. Store the product at -20° C.

⁴ Medical Biochemistry, Lab 207, July 1, 1998.

Preparation of the pET21d Vector for the Later Sub-Cloning of the GAL4 Transcription Factors⁵

1. Digest the purified sample with NcoI and XhoI.

SAMPLE	l
pET21d (1.183 mg/ul)	3 ul
2.5 X 11.14 U of NcoI	3 ul
2.5 X 44.5 U of XhoI	7 ul
100 ug/ul of ABS	0.5 ul
10% total volume of buffer #2	5 ul
DD H ₂ O	31.5 ul
Total volume	50 ul

2. Incubate the mixture for 120 minutes at 37°C.
3. Purify the product by spin column.
4. 100 ul S-200 resin equilibrated with TE (centrifuge 5000-6000 rpm).
5. Check on 1.5% agarose + 1X TBE gel.
 - a) Uncut pET21d vector
 - b) Digested pET21d vector

⁵ Medical Biochemistry, Lab 207, May 24, 1998.

Transformation of Constructs into DH5alpha Competent Cells⁶

1. Set up the electro cell manipulator 600 as follows:

Choose mode: T 2.5KV/Resistance High Voltage (HV)

Set capacitance: C Not used in HV mode

Set resistance: R R5 (129 ohm)

BTX Disposable cuvette P/N 620 (2mm gap), place on
ice for five minutes

Set charging voltage: S 2.45 KV

Desired field strength: 12.25 KV/cm

Desired pulse length: I 5-6 mseconds

2. Remove 10 Eppendorf tubes of DH5alpha competent cells, prepared for electroporation and placed on ice.
3. Once the competent cells have thawed out, add the following DNA, to the competent cells:
 - a) 1 ul of GAL4-AD ligation 1' product
 - b) 1 ul of GAL4-D ligation 2' product
 - c) 1 ul of GAL4-AD ligation 1 product
 - d) 1 ul of GAL4-D ligation 2 product
 - e) 1 ul of vector pET21d ligation product
 - f) 1 ul of GAL4-AD insert ligation product
 - g) 1 ul of unligated pET21d vector
 - h) 1 ul of unligated GAL4-AD insert

⁶ Medical Biochemistry, Lab 207, July 17, 1998.

i) 1 ul of T.E. pH 8.0

4. Flick the Eppendorf tube to mix and settle the cell mixture by tapping the cuvette on the table.
5. Transfer the competent cells into the BTX cuvette, right after mixing with DNA, and put the cap on.
6. Press A to activate automatic charge and pulse sequence.
7. Immediately after green charging light goes out, add 1000 ul of 2xYT into the cuvette.
8. Briefly and gently pipette out the content from the cuvette into a sterile, small-size Falcon tube and vortex.
9. Incubate the Falcon tubes at 37° C for 60 minutes. (Shaking at 225 rpm will improve recovery.)
10. Plate 100 ul from each of the Falcon tubes onto LB agar plates containing 40 ug/ml of ampicilin.
11. Allow the plates to dry.
12. Incubate the plates at 37° C overnight.
13. Check for growth of individual colonies.

Preparation of the GAL4-AD and GAL4-D Inserts⁷

1. PCR amplify the GAL4-AD and GAL4-D fragments from the pGEX-cs vector.

SAMPLE	1	2
pGEX-cs vector (50ng/ul)	1 ul	1 ul
GAL4-AD 5' primer (50 pmol/ul)	2 ul	2 ul
GAL4-AD 3' primer (50pmol/ul)	2 ul	
GAL4-D 3' primer. (50pmol/ul.)		2 ul
DD H2O	76 ul	76 ul
10X Pfu buffer	10 ul	10 ul
2.5 mM dNTP	8 ul	8 ul
VORTEX SAMPLES		
2.5 U/ul Pfu DNA Polymerase.	1 ul	1 ul
Total volume	100 ul	100 ul

2. Program for PCR reaction:

25 cycles: 1 minute at 94° C, 1 minute at 50° C, 4 minutes at 72° C

1 cycle: 1 minute at 94° C, 1 minute at 50° C, 10 minutes at 72° C

3. Check results on a 0.5% agarose + 1X TBE gel.

⁷ Medical Biochemistry, Lab 207, July 7, 1998.

GAL4-VP16 Sequencing Reaction⁸

1. Preparation of the sequencing gel:

a) 8% Acrylamide

10 ml of 40% acrylamide

15 ml of E-pure H₂O

5 ml of 5 x TBE

25 g of urea

b) Mix gently until the urea is dissolved and filter through a 0.45 micron filter.

100 ul of TEMED

100 ul of APS

c) Pour the gel.

2. Sequencing reaction:

a) Denaturing reaction.

Sample.	1	2
GAL4-VP16 (bigger prep.)	8 ul	8 ul
1.0 M NaOH	2 ul	2 ul
PGEX-cs 3' (5pmol/ul)		1 ul
PGEX-cs down (5pmol/ul)	1 ul	

b) Mix thoroughly and incubate for 10' at 37 °C. Then place the mixture on ice and add to each of the tubes:

⁸ Medical Biochemistry, Lab 207, July 16, 1998.

Sample.	1	2
1.0 M HCL	2 ul	2 ul
Plasmid reaction buffer	2 ul	2 ul
Total volume	15 ul	15 ul

- c) Annealing reaction: Incubate the template/primer/buffer mixture at 37° C for 10 minutes, then chill on ice.
- d) While annealing, label, fill and cap tubes with 2.5 ul of each of the termination mixtures (G, A,T and C, red capped vials). Keep covered on ice for steps 5-7.
- e) If needed, dilute labeling mix 15-fold to working concentration. Retain for use in step 6.

Dilution	
Labeling mix	4 ul
DD H ₂ O	16 ul

- f) Pre-warm the termination tubes from step 2(d) in a water bath at 37° C.
- g) Labeling reaction: Add to ice-cold DNA from step 2(c). Mix and incubate at room temperature for 2-5 minutes.

Labeling	
DTT (0.1 M)	1 ul
Diluted labeling mix	2 ul
5 uCi ³⁵ dATP	0.5 ul
Sequenase polymerase	2 ul

3. Termination reaction: Transfer 4.5 ul of the labeling reaction into each of the termination tubes (G, A,T and C), mix and continue the incubation of the termination tubes at room temperature for 5 minutes.
 - a) Stop the reaction by adding 4 ul of stop solution.
 - b) Heat the samples to 75° C for 2 minutes immediately before loading onto the sequencing gel. Load 2-3 ul in each lane.

GAL4-VP16 Transcription Factor DNA Sequencing Reaction⁹

1. Preparation of the sequencing gel:

a) 8% Acrylamide

10 ml of 40% acrylamide

15 ml of E-pure H₂O

5 ml of 5 x TBE

25 g of urea

b) Mix gently until the urea is dissolved and filter through a 0.45 micron filter.

100 ul of TEMED

100 ul of APS

c) Pour the gel.

2. Sequencing reaction: Sequencing for GAL4-VP16 by Stratagene Cyclist Exo-Pfu DNA sequencing kit.

a) Add 3 ul of the appropriate ddNTPs to 4 termination tubes chilled on ice.

Sample	l
GAL4-VP16 (0.6 ug/ul)	2 ul
PGEX-cs down(5pmol/ul)	1 ul
10 X sequencing buffer	4 ul
S35-dATP (12.5uCi/ul)	1 ul
Exo-Pfu enzyme	1 ul
DD H ₂ O	17 ul
DMSO	4 ul
Total volume	30 ul

b) Aliquote 7 ul of the sequencing reaction mixture into each of 4 the termination tubes.

⁹ Medical Biochemistry Lab 207, May 19, 1998.

3. Program for PCR reaction:

1 cycle: 95° C for 30 minutes.

30 cycles: 95° C for 30 seconds. 60° C for 30 seconds. 72° C for 30 seconds.

a) Stop the reactions by adding 5 ul of the stop solution.

b) Heat samples at >80° C for 2-5 minutes. Then immediately load 2-4 ul onto sequencing gel.

BIGGERprep Plasmid DNA Preparation Procedure

1. Culture plasmid containing bacterial cells by inoculating 1 L of media of TB broth media with the appropriate concentration of antibiotics for 16-24 hours.
2. Pellet bacteria by centrifuging at 10,000 g for 3 minutes. Decant the supernatant.
3. Completely resuspend the bacterial pellet in 30 mL of solution I. Vortex to ensure pellet resuspension.
4. Lyse the bacteria by adding 30 mL of solution II to the resuspended bacteria. Mix very gently by rotating the centrifuge tube. Do not allow the solution to be active for more than 5 minutes. Lysate should be relatively clear.
5. Neutralize bacterial lysate by adding 30 mL of solution III. Immediately mix by extremely gentle inversion of the bottle for 5-10 seconds. A white material will appear. Increase the strength of the inversions, but do not vortex.
6. Pellet the bacterial cell wall by centrifuging at 16,000 g for 8 minutes. Transfer all the supernatant to a fresh centrifuge tube.
7. Bind the plasmid DNA by adding 80 mL of well-mixed BIGGERprep DNA Binding Matrix Suspension to supernatant and mix by vigorous inversion.
8. Pellet BIGGERprep DNA Binding Matrix and bound plasmid DNA at 16,000 g for 8 minutes. Pour off the supernatant making sure to keep the pellet.
9. Add 20 mL of diluted Purification Solution and mix by vigorous shaking. Purification solution is a 1:1 mixture of the reagent and 95% ethanol. Pour mixture into a BIGGERprep spin column.
10. Add 20 mL of the same mixture to rinse any possible residues from the centrifuge tube.

11. Purify the plasmid DNA by centrifuging at 1500 g for 7 minutes on a swinging-bucket centrifuge at room temperature. Pour off the filtrated liquid.
12. Repeat the same step for greater purity.
13. Transfer the BIGGERprep spin column to a fresh tube and elute the plasmid with 3 mL of 8.0 pH TE, preheated at 65-70° C.
14. Spin the column at 1500 g for 7 minutes. Transfer the elute plasmid DNA to a precipitation tube.
15. Precipitate the plasmid DNA by adding 140 ul of 5M NaCl, mix and then add 12 mL of 95%-100% ethanol at room temperature.
16. Mix well and pellet plasmid DNA at 16,000 g for 5 minutes at room temperature. Pour off supernatant.
17. Briefly wash with >70% ethanol. Wash twice and then let air dry.
18. Once the plasmid DNA is dry, resuspend in 500 ul of TE.
19. Store at -20° C.

GAL4-VP16 Transcription Factor Amplification (Taq Polymerase Protocol)

1. PCR amplify the GAL4-VP16 fragment from the pJL2 expression vector vector.

SAMPLE	1 and 2	3 and 4
pJL2 vector (50ng/ul)	1 ul	1 ul
50 mM MgCL ₂	1.5 ul	1.5 ul
GAL4-VP16 5' primer (50pmol/ul)	1 ul	1 ul
GAL4-VP16 3' primer. (50pmol/ul.)	1 ul	1 ul
DD H ₂ O	34.5 ul	34.5 ul
10X Taq extender buffer	5 ul	5 ul
10 mM dNTP	4 ul	4 ul
VORTEX SAMPLES		
2.5 U/ul Taq DNA Polymerase Exterder	1 ul	1 ul
2.5 U/ul Taq DNA Polymerase	1 ul	1 ul
Total volume	50 ul	50 ul

2. Program for PCR reaction:

1 cycle: 30 seconds at 94° C

35 cycles: 30 seconds at 94° C, 30 seconds at 55° C, 30 seconds at 72° C

1 cycle: 7 minutes at 72° C

3. Check results on a 1.5% agarose + 1X TBE gel.

GAL4-VP16 Transcription Factor Amplification (Pfu Turbo Protocol)

1. PCR amplify the GAL4-VP16 fragment from the pJL2 expression vector vector.

SAMPLE	1 and 2	3 and 4
pJL2 vector (50ng/ul)	1 ul	1 ul
GAL4-VP16 5' primer (50pmol/ul)	1 ul	1 ul
GAL4-VP16 3' primer. (50pmol/ul.)	1 ul	1 ul
DD H2O	31.5 ul	31.5 ul
10X Cloned Pfu buffer	5 ul	5 ul
2.5 mM dNTP	9 ul	9 ul
VORTEX SAMPLES		
2.5 U/ul Pfu Turbo DNA Polymerase	1.5 ul	1.5 ul
Total volume	50 ul	ul

2. Program for PCR reaction:

1 cycle: 30 seconds at 94° C

35 cycles: 30 seconds at 94° C, 30 seconds at 55° C, 30 seconds at 72° C

1 cycle: 7 minutes at 72° C

3. Check results on a 1.5% agarose + 1X TBE gel.

Ligation of the Dephosphorylated pET21d Vector and the GAL4-AD and GAL4-D***Transcription Factors.***

1. Preparation of the ligation reactions:

Sample	1 and 2	3 and 4	5	6
pET21d vector (100ng/ μ l)	2 μ l	2 μ l	2 μ l	2 μ l
GAL4-AD insert	2 μ l			2 μ l
GAL4-D insert		2 μ l		
10 X T4 DNA ligase buffer	1 μ l	1 μ l	1 μ l	1 μ l
T4 DNA Ligase	1 μ l	1 μ l	1 μ lv	1 μ lv
DD H ₂ O	4 μ l	4 μ l	4 μ l	4 μ l
TOTAL VOL.	10 μl	10 μl	10 μl	10 μl

2. Place the reaction mixture at a constant 16° C for an incubation period of 4 hours.
3. Store the ligation results at – 20° C.
4. Check the results on a 1.5% Agarose gel using a 1 Kb DNA ladder.

Digestion of the GAL4-AD and GAL4-D Inserts with NcoI and XhoI Restriction Enzymes.

1. Preparation of the digestion reactions of the GAL4-AD and GAL4-D inserts.
2. Extract the bands from the gel utilizing a gel extraction kit.

Sample	1	2
GAL4-AD insert	20 μ l	
GAL4-D insert		20 μ l
Buffer # 2	3.5 μ l	3.5 μ l
NcoI Restriction Enzyme	2.5 μ l	2.5 μ l
XhoI Restriction Enzyme	2.5 μ l	2.5 μ l
BSA (100mM)	3.5 μ l	3.5 μ l
DD H ₂ O	3 μ l	3 μ l
TOTAL VOL.	35 μ l	35 μ l

3. Incubate the reaction mixture at 37° C for four hours.
4. Inactivate the restriction enzymes by heating the reaction mixture at 65° C for 15 minutes
5. Check the results on a 1.5% Agarose gel.

TB Broth Media Preparation

1. Ingredients for preparation of 1 L of media:

Tryptophan	10 g
Yeast extract	15 g
NaCl	5 g
d H ₂ O	700 mL

2. Adjust to a final pH of 7.0 and equal volume to 1 L.
3. Sterilize by autoclaving for 20 minutes.
4. Add 100 ng/ul of ampiciline when the temperature is about 55° C.

2 x YT Media Preparation

1. Ingredients for preparation of 1 L of media:

Bacto-tryptone	16 g
Bacto-yeast extract	10 g
NaCL	5 g
dH ₂ O	900 mL

2. Adjust to a final pH of 7.0 and equal volume to 1 L.
3. Sterilize by autoclaving for 20 minutes.
4. Add 100 ng/ul of ampiciline when the temperature is about 55° C.

Experimental Results¹⁰

The restriction digestion of the expression vector pET21d, in which the desired constructs were inserted, showed a unique band on a 1.5% agarose gel, corresponding to a fragment of approximately 5350 ± 10 base pairs resulting from the digestion with the restriction enzymes: NcoI and XhoI. These restriction enzymes have specific cut sites at 1 base pair and 5365 base pairs respectively. The resulting fragment was then purified by spincolumn. As a comparison, a separate sample was run as a control, containing an undigested version of the vector. The results from this purification process can be seen in Figure #1, together with that of the pET21d expression vector digestion. To avoid further complications with the expression vector, this was dephosphorilated at its 5' end.

The preparation of the inserts by means of PCR amplification of the coding region for the transcription factors GAL4-AD and GAL4-D, utilizing the GAL4-AD 5', GAL4-AD 3' and GAL4-D 3' primers resulted in fragments of approximately 390 base pairs and 330 base pairs respectively, after digestion with restriction enzymes NcoI and XhoI, as shown in Figure #2. After the amplification, the working concentrations of both of the inserts are 0.97ug/ul for GAL4-AD and 0.93ug/gl for GAL4-D. 20 μ l from the amplification products were loaded into a 1.5% agarose gel from which they were later extracted and utilized for the digestion step. Figure #2 shows the gel from which they were extracted. The results from the ligation of the dephosphorilated pET21d expression vector and the GAL4-AD and Gal4-D constructs and the later electroporation are satisfactory. The growth of the transformants indicates a successful insertion of the plasmid. The controls used indicate growth of the transformant with the unligated plasmid but there is no growth of the bacteria transformed with the GAL4-

¹⁰ The results will be noted in chronological order of experiments.

AD insert alone, therefore resulting in a satisfactory ligation and transformation. The results from the ligation experiment can be seen in Figure #3. The restriction-digestion analysis from several of the transformed colonies indicates the presence of a band of approximately 5700 base pairs.

The results from the GAL4-VP16 sequencing have rendered an incomplete sequence for this transcription factor. The experimental sequence can be seen in appendix A. There has been no possibility of designing a DNA primer corresponding to the 5' end of the coding region of this transcription factor for the amplification procedure.

Using an alternative source of DNA containing the GAL4-VP16 insert, it has been possible to design the primers to be used in the amplification of this transcription factor. Also, the sequence for the GAL4-VP16 is available in Appendix A.

The bacterial culture containing the GAL4-VP16 insert has been grown and DNA extracted. The concentration of the working samples are 146 ng/ μ l and 176 ng/ μ l. The amplification of this transcription factor has been attempted using two different DNA polymerases. The first amplification has been performed using Taq polymerase and Taq extender. This amplification has resulted in good concentrations of amplified products, see Figure #4. The second amplification has been performed using Pfu Turbo as the DNA polymerase, the results from this latter amplification are more satisfactory in terms of yield and can be seen in Figure #5.

For a figure containing all the transcription factors together in an uncut GAL4-VP16 and digested versions of GAL4-AD and GAL4-D, refer to Figure #6.

Discussion

As a result of the satisfactory outcome obtained from the several experiments performed, a pET21d-GAL4 expression vector was obtained for each of the transcription factors (GAL4-AD and GAL4-D). The ultimate products from the transformation of the ligated plasmid-insert complex into *Escherichia coli* were kept as DNA stocks for future transformations. This method is preferred for DNA storage due to the relatively short life and much faster degradation rate of the bacterial cells. When transformed again into competent cells, they will enable an easy over-expression of the transcription factors in the future for their use in chromatin remodeling studies.

First of all, the resulting fragment from the pET21d expression vector digestion with NcoI and XhoI restriction enzymes, resulted in a unique band corresponding to a fragment of approximately 5400 base pairs approximately. This fragment is the expected outcome. The restriction enzyme NcoI cuts the expression vector at position 5365 and the restriction enzyme XhoI cuts at the expression vector at position 1. After digestion with both of these restriction enzymes, the expression vector is linearized in a fragment of 5364 base pairs, as indicated by experimental results, and there is a smaller fragment of approximately 75 base pairs. This fragment is too small to be detected with the size of the ladder used for the detection of our fragment of interest. The purification process as seen in Figure #1 maintains the size of the fragment. The fact that no other bands corresponding to fragments of similar sizes are apparent is another good indicator of the success of the digestion and purification processes. As a comparison tool, the sample has been run together with the undigested version of the vector that comprises 5440 base pairs in its entirety. A dephosphorilation step was performed on the purified sample of the pET21d expression vector with the sole intention of preventing

the vector from recircularizing, before the ligation experiment with the inserts was completed. The subsequent satisfactory ligation indicates that the concentration of the dephosphorylated vector was high enough to allow for a good ligation.

On the other hand, the amplification experiments have proved successful as well. The results from the amplification can be seen in Figure #2. We can see how the bands corresponding to the GAL4-AD transcription factor run about 390 base pairs, whereas those corresponding to the GAL4-D transcription factor are an average of 60 to 70 base pairs shorter, running about 320 base pairs. The reason for this is that the GAL4-D (domain) transcription factor lacks the last 60 base pairs that encode for the last 20 amino acids present in the GAL4-AD transcription factor sequence. As can be seen in Figure #3 the ligation products revealed the presence on a 1.5% agarose gel of fragments of two sizes. One is of approximately 5750 base pairs corresponding to the ligation of the pET21d expression vector and the GAL4-AD transcription factor insert. The other fragment is of approximately 5680 to 5690 base pairs corresponding to the ligation product of the expression vector and the GAL4-D transcription factor insert. The ligation experiment yielded a high enough concentration of ligated plasmid-insert complex to allow for a successful electroporation of the products into DH5 alpha competent cells. The controls used in the ligation experiment were also used in the transformation step. As expected, the negative control containing the products from the ligation of the GAL4-AD insert alone resulted in no growth on a LB broth plate containing ampicillin. The positive control consisted of the pET21d expression vector alone. After transformation the cells grew normally on the LB broth plates. The transformants resulting from the ligation of pET21d expression vector and the insert corresponding to the GAL4 series transcription factors all resulted in growth on the LB broth plates. These results are

satisfactory and the products have been kept as DNA stocks that can be used for later transformations into DH5 alpha competent cells for GAL4 proteins over-expression purposes. The purified version of these over-expressed proteins can be integrated into the study of chromatin remodeling.

The results from the sequencing of the GAL4-VP16 acidic transcription factor have provided only part of the sequence corresponding to the 3' end of the open reading frame containing this transcription factor. Therefore, no design of a primer covering the 5' region of the open reading frame was possible. The reason for this seems to be an advance stage of degradation of the DNA that has been used as a template for the sequencing experiments. The obtained partial sequence can be seen in Appendix A.

Utilizing a new *Escherichia coli* strain that contains the pJL2 expression vector, DNA has been isolated and the design of the necessary primers for the PCR amplification has been performed. To ensure a positive amplification of this transcription factor, two separate PCR reactions have been run. The first reaction consists of a combination of Taq polymerase and Taq polymerase extender. As mentioned in the results, the amplification obtained in this series of reactions is satisfactory. The second method is based on a PCR amplification using Pfu Turbo as the DNA polymerase and the results from this amplification are much more successful. The figures corresponding to these amplifications are Figure #4 and Figure #5 containing the Taq polymerase and the Pfu Turbo amplifications respectively.

In Figure #6 we can see the comparison of the different transcription factors in their digested version, with the exception of the GAL4-VP16 which is only present in its undigested version. As can be seen in the case of the GAL4-VP16 transcription factor there is more than one amplification product.

The project has rendered satisfactory results with respect to the cloning of the GAL4-AD and GAL4-D transcription factors. The obtained fragments matched the expected sizes after the PCR amplification step at approximately 390 base pairs for the GAL4-AD and 330 base pairs for the GAL4-D. Also, after the cloning step, the resulting fragments run about 5750 base pairs and 5680 base pairs respectively. These approximations calculated from the gels indicate that the cloning is acceptable. On the other hand, the GAL4-VP16 transcription factor amplification has shown positive results in both protocols that have been used for its amplification. The Taq polymerase amplified inserts are of the expected size of approximately 230 base pairs and the yield is considerably good, making it easier to digest and use in the cloning step. The second amplification performed with Pfu Turbo has resulted in extremely good amplification and there are different fragments. One of the fragments corresponds to the desired size while the others are slightly larger in size than expected as calculated from the genetic map of the pJL2 plasmid expression vector. These amplification products are excellent for the digestion protocol, and this step would allow for the isolation of the fragment that could be purified by spin column for a subsequent cloning step.


Appendix A

pGEX-cs Vector Linear Sequence

DNA sequence	5019 b.p.	ACGTTATCGACT	...	GCGTTGGAATT	circular	
	10	20	30	40	50	60
1	ACGTTATCGA	CTGCACGGTG	CACCAATGCT	TCTGGCGTCA	GCCAGCCATC	GGAAGCTGTG 60
61	GTATGGCTGT	GCAGGTCGTA	AATCACTGCA	TAATTCGTGT	CGCTCAAGGC	GCACTCCCGT 120
121	TCTGGATAAT	GTPTTTTGGC	CCGACATCAT	AACGGTTCTG	GCAAAATATC	TGAAATGAGC 180
181	TGTTGACAA	TAATCATCGG	CTCGTATAAT	GTGTGGAATT	GTGAGCGGAT	AACAATTTCA 240
241	CACAGGAAAC	AGTATTCATG	TCCCTATAC	TAGGTTATTG	GAAAATTAAG	GGCCTTGTCG 300
301	AACCCACTCG	ACTTCTTTTG	GAATATCTTG	AAGAAAAATA	TGAAGAGCAT	TTGTATGAGC 360
361	CGCATGAAGG	TGATAAATGG	CGAAACAAAA	AGTTTGAATT	GGGTTTGGAG	TTTCCCAATC 420
421	TTCCTTATTA	TAATGATGGT	GATGTTAAAT	TAACACAGTC	TATGGCCATC	ATACGTTATA 480
481	TAGCTGACAA	GCACAACATG	TTGGGTGGTT	GTCCAAAAGA	CCGTGCAGAG	ATTTCAATGC 540
541	TTGAAGGAGC	GGTPTTGGAT	ATTAGATACG	GTGTTTCGAG	AATTCGATAT	AGTAAAGACT 600
601	TTGAAACTCT	CAAAGTTGAT	TTTCTTAGCA	AGCTACCTGA	AATGCTGAAA	ATGTTCGAAG 660
661	ATCGTTTATG	TCATAAAACA	TATTTAAATG	GTGATCATGT	AACCCATCCT	GACTTCATGT 720
721	TGATGACCG	TCTTGATGTT	GTTTTATACA	TGGACCCAA	GTGCTGGAT	CGCTTCCCAA 780
781	AATTAGTTTG	TTTTAAAAAA	CGTATTGAAG	CTATCCACA	AATTGATAAG	TACTTGAAT 840
841	CGAGCAAGTA	TATAGCATGG	CCPTTGCAGG	GCTGGCAAGC	CACGTTTGGT	GGTGGCgacc 900
901	atcttcgagaa	ecttctatttc	cagggcccca	cggtactcat	ctttggcaca	gtcaacgcta 960
961	acatcccgaa	ggaaagtgtt	gggtggagctc	ggtagcccg	ggatccatccg	gaattcatccg 1020
1021	tgactgactg	acgatctgcc	tcggcggttt	cggtgatgac	gggtaaaacc	tctgacacat 1080
1081	gcagctcccg	gagacggcca	cagcttgcct	gtaagccgat	gcccgggagca	gacacggccc 1140
1141	tcagggcgcc	tcagccgggtg	ttggcgggtg	tcggggcgca	gccatgacc	agtcacgtag 1200
1201	cgatagccga	gtgtataatt	cttgaagacc	aaagggcctc	gtgatccgcc	tatttttata 1260
1261	ggtttaattg	atgataataa	tggtttctta	gacgtcaggt	ggcacttttc	ggggaaatgt 1320
1321	ggcgggaaac	cttatttgtt	tatttttcta	aatacattca	aatagtatc	cgctcatgag 1380
1381	acaataacc	tgataaatgc	ttcaataata	ttgaaaaagg	aagagtatga	gtattcaaca 1440
1441	ttcccgctg	gccctatttc	ctttttttgc	ggcattttgc	cttctgtttt	ttgtccacc 1500
1501	agaaaccgctg	gtgaaagtta	aagatgctga	agatcagttg	gggtccagag	tgggttacat 1560
1561	cgaaactggat	ctcaacagccg	gtaagatcct	tgagagtttt	cgccccgaag	aacgttttcc 1620
1621	aatgactgagc	acttttaaa	ttctgctatg	tgcccggtga	ttatcccggt	ttgacggccc 1680
1681	gcaagagcaca	ctcggctgcc	gcatacacta	ttctcagaat	gacttgggtg	agtactcacc 1740
1741	agtcacagaa	aagcatctta	cgatggcat	gacagtaaga	gaattatgca	gtgctgccat 1800
1801	aaccatgagt	gataacactg	cgcccaactt	acttctgaca	accatccgag	gaccgaagca 1860
1861	gctaaccgct	tttttcgaca	acatggggga	tcagttaact	cgctttgatc	gttgggaacc 1920
1921	ggagctgaat	gaagccatac	caaacgcaga	cgctgacacc	acgatgccctg	cagcaatggc 1980
1981	aacaacgttg	cgcaaaacta	taactggcga	actacttact	ctagctttccc	ggcaacaatt 2040
2041	aatgactgagc	atggagggcg	ataaagtgtc	aggaccactt	ctcgctccgg	cccttccggc 2100
2101	tgccctggtt	attgctgata	aatctggagc	cggtgagcgt	gggtctcggg	gtatcatgtc 2160
2161	agcaactggg	ccagatggtg	agccctcccg	tatcgtagtt	atctacacga	cggggagtca 2220
2221	ggcaactfatg	gatgaacgaa	atagacagat	cgctgagata	ggctcctcac	tgattaaagca 2280
2281	ttggtaactg	tcagaccaag	ttactcata	tatactttag	attgatttaa	aacttcattt 2340
2341	ttaatttaaa	aggatctagg	tgaagatcct	ttttgataat	ctcatgacca	aaatccctta 2400
2401	acgtgagttt	tcgttccact	gagctcaga	ccccgtagaa	aagatcaaa	gatcttcttg 2460
2461	agatcttttt	ttctccggcg	taactctctg	cttgcaaa	aaaaaacacc	cgctaccgca 2520
2521	ggtggtttgt	ttgcccgatc	aagagctacc	aaactttttt	ccgaaggtaa	ctggcttcag 2580
2581	cagagccgag	ataccaaata	ctgtccttct	aggtagccg	tagttagccc	accacttcaa 2640
2641	gaactctgta	gcaccgccta	cataacctgc	tctgctaatic	ctgttaccag	tggctgctgc 2700
2701	cagtgccgat	aagctggttc	ttaccgggtt	ggactcaaga	cgatagttac	ggataaagcc 2760
2761	gcagcggtgc	ggctgaaacg	ggggttcgtg	cacacagccc	agcttggagc	gaacgaccta 2820
2821	caccgaactg	agatacctac	agcctgagct	atgagaaacc	gccaccgttc	ccgaagggag 2880
2881	aaagccggac	aggtatcccg	taagccggag	ggtccgaa	ggagagccca	cgaggagact 2940
2941	tccaggggga	aagcctgggt	atctttatag	tcctgtcggg	tttccgacc	tctgacttga 3000
3001	ggctcgattt	ttgtgatgct	cgctcagggg	ggggagccta	tggaaaaacg	ccagcaacgc 3060
3061	ggccttttta	cggtttctgg	ctttttgctg	gcttttgcct	cacatgttct	ttcctgcttt 3120
3121	atccccgat	tctgttgata	accgtattac	gccccttgag	tgagctgata	cgctccggcc 3180
3181	cagccgaacg	accgagccga	gctgctcagt	gagccgaa	gcccgaagac	gcctgatcgc 3240
3241	gtatttttct	cttacgcatac	tgtgcccgtat	ttcacaccgc	ataaattccg	acaccatgca 3300
3301	atggtgcaaa	acctttccgg	gtatggcatg	atagcccccg	gaagagagtc	aattcaggg 3360
3361	gggtgaaatg	aaaccagtaa	cgttatacga	tgtccgagag	tatgcccggg	tctcttatca 3420
3421	gaccgtttcc	cgctggtgga	accagccag	ccacgtttct	gcaaaaaccg	gggaaaaagt 3480
3481	ggaagccggc	atggccgagc	tgaattacat	ttccaaaccg	gtggcaaac	aactgcccgg 3540
3541	caaacagctg	ttgctgattg	gcgttggcac	ctccagctgc	gcccctgacg	cgccctcgca 3600
3601	aattgtcggc	gcatataaat	ctcggccgca	tcacttgggt	gccagcgtgg	tgggtcgtat 3660
3661	ggtagAACGA	agccgctgctg	aagcctgtaa	agccgcccgtg	cacaatcttc	tcgcccAACG 3720
3721	cgctagTGGG	ctgatcattta	actatccgct	ggatgaccag	gatgccattg	ctgtggaaGC 3780
3781	tgccctgact	aatgttccgg	cgttatttct	tgatgtctct	gaccagacac	ccatcaaacg 3840
3841	tatttatttct	tcccatgaa	acggtaccgg	actggccgtg	gagcatctgg	tccgatpccg 3900
3901	tcaccagcaa	atccgctgtg	tagccggccc	atpaaagtct	gtctccggcc	gtctgctgct 3960
3961	ggctggctgc	cataaatatac	tcactccgaa	tcAAATTCAG	ccgatagccc	aacgggaaGG 4020
4021	cgactggag	gccatgtccg	gttttcaaca	aaccatgcaa	atgctgaaatg	agggcatcgt 4080
4081	tcccactgcg	atgctggttg	ccaacgatca	gatggccgtg	ggcccaatgc	gcgccattac 4140
4141	cgagtcggg	ctgcccgttg	gtccgatata	ctcgttagtg	ggataccgag	ataccgaaga 4200
4201	cagctcatgt	tatatccccc	cgftaacac	catcaaacag	gattttccgc	tgctggggca 4260
4261	aaccagctg	gaccgcttgc	tgcaactctc	tacggccag	gcccgtgaag	gcaatcagct 4320
4321	gttccccgct	tcactgggtg	aaagaaaaac	caccctggcc	cccaataccg	aaaccgctc 4380
4381	tcccctgcg	ttggccgatt	catataatgca	gctggcacga	caggtttccc	gactggaaag 4440
4441	cgggcagtg	gcccacgcga	atpaaatgta	gttagctcac	tcattagcca	cccaggctt 4500
4501	tacctttat	gcttccggct	cgatgttgt	gtgaaattct	gagccgata	caattttaca 4560
4561	caggaaacag	ctatgaccat	gattaccgat	tcactggccc	tcgttttaca	acgtcgtgac 4620
4621	tgggaaAAC	ctggccttac	ccaactpaa	cgccctggcag	cacatcccc	tttccgacg 4680
4681	tggcgtAATA	cgaaagagcc	ccgcaccgat	cgcccttccc	aacagttggc	cagcctgaa 4740
4741	ggcgaatggc	gctttgccc	gtttccggca	ccgaagccg	tgccggaaag	ctggctggag 4800
4801	tggcattctc	utgagccgca	tactgtcgtc	gtcccccaa	actggcagat	gcacggttac 4860
4861	gatgccccca	tctacaccaa	cgtaacctat	cccattaccg	tcaatccgcc	gtttgttccc 4920
4921	acggagaatc	cgaccgggtg	ttactcgcctc	acatttaattg	ttgatgaaag	ctggctacag 4980
4981	gaagccCAGA	cgccaattat	ttttgatggc	gttggaaatt		5019
	10	20	30	40	50	60

pGEX-cs Vector Genetic Map

DNA sequence 5019 b.p. ACGTTATCGACT ... GCGGTTGGAATT circular



pET21d Vector Linear Sequence


```

DNA sequence 5446 b.p. ATCCGGATATAG . TCCCATTCGCCA linear
      | 10 | 20 | 30 | 40 | 50 | 60
1  ATCCGGATAT AGTTCTCTCT TCCAGCAAAA AACCCCTCAA GACCCGTTTA GAGGCCCAAA 60
61 GGGGTTATGC TAGTTATTGC TCAGCGGTGG CAGCAGSCAA CTCAGCTTTC TTTCGGGCTT 120
121 TGTTAGCAGC CGJATCTCAG TGGTGGTGGT GGTGGTGGTC GAGTGGGGCC GCAAGCTTGT 180
181 CGACCGAGCT CGAATTCGGG TCCGACCCAT TTGCTGTCCA CCACTCATGC TAGCCATGGT 240
241 ATATCTCCTT CTTAAAGTTA AACAAAAATTA TTCTAGAGG GGAATTTGTA TCCGCTCACA 300
301 ATTCCTCTAT AGTGAGTGGT ATTAATTTCG CGGGATCGAG ATCTCGATCC TCTACGCCGG 360
361 ACGCATCGTG GCCGGCATCA CCGCGCCAC AGGTGGGGTT GCTGGGGCCT ATATCGCCGA 420
421 CATCACCGAT GGGGAAGATC GGGCTCGCCA TTTCGGGGTC ATGAGCGCTT GTTTCGGGCT 480
481 GGGTATGGTG CGAGCCCGCG TGGCCGGGGG ACTGTTGGGC GCCATCTCCT TGCATGCACC 540
541 ATTCCTTGGC GGGCGGGTGC TCAACGGGCT CAACCTACTA CTGGGTGGCT TCCATATGCA 600
601 GAGTCCGCAT AAGGGAGAGC GTCGAGATCC CGGACCCAT CGAATGGCC AAAACCTTTC 660
661 GCGGTATGGC ATGATAGCCG TGGGAAGAGA GTCAATTCAG GGTGGTGAAT GTGAAACCAG 720
721 TAACGTTATA CGATTCGCA GAGTATGCCG GTGCTCTTA TCAGACCCTT TCCCGCGTGG 780
781 TGAAACAGGC CAGCCAGCTT TCTCGAAAAA CCGGGAAAAA AGTGGAAAGC GCGATGGCGG 840
841 AGCTGAAATTA CAATCCCAAC CGGTGGCCAC AACAACTGCG GGGCAACAGC TCGTGTCTGA 900
901 TGGGCTTGGC CACTCTGAGT CTGGCCCTGC ACGGCTGTC GCAAAATGTC GCGCGGATTA 960
961 AATCTGGCGC CGATCAACTG GATGCCAGCG TGGTGGTGC GATGCTAGAA TGGTGGTGG 1020
1021 TCGAAGCCTG TAAAGCGCGC GTGCACAATC TTCTCCGCA ACGGCTCAGT GGGCTGATCA 1080
1081 TTAACATATC GCTGGATGAC TAGGATGCCA TTGCTGTGGA AGCTGCCCTG ACTAATGTTT 1140
1141 CGCGCTTATT TCTTGAATGC TCTGACCACA CACCATCAA CAGTATTATT TTCTCCCATG 1200
1201 AAGACGGTAC CGGACTGGGC GTGGAGCATC TGGTCCGATT GGGTCACCAG CAAATCGCGC 1260
1261 TGTTAGCGGG CCCATTAAGT TCTGTCTGG CGCGTCTGG TCTGGCTGGC TGGCATAAAT 1320
1321 ATCTCACCTG CAATCAAAAT CAGCCGATAG CGGAACGGGA AGCGGACTGG AGTGCCATGT 1380
1381 CCGGTTTTCA ACAAAACAATG CAAATGCTGA ATGAGGCGAT CGTTCACCAT CCGATGCTGG 1440
1441 TCGCAACGA TCAGATGGCG TCGGGCGCAA TCGCGCCAT TACCGAGTCC GGGCTGGCGG 1500
1501 TGGTTCGGGA TATCTGGTA GTGGGATAGC ACGATACCGA AGACAGCTCA TGTATATACC 1560
1561 CGCCGTTAAC CACCACTAAA CAGGATTTTC GCGTGTGGG GCAAAACAGC GTGACCCGCT 1620
1621 TGCTGCAACT CTCTCAGGGC CAGGCGGTGA AGGGCAATCA GCTGTGGCC GTCTCACTGG 1680
1681 TGAAGAAGAA AACCAACCTG GGGCCCAATA CGCAACCCGC CTCTCCCGCG GGTGTGGCGG 1740
1741 ATTCATTAAT CGACGTGGCA CGACAGGTTT CCGCACTGGA AAGCGGGCAG TGAGCCGAAC 1800
1801 GCAATTAATG TAAGTTAGCT CACTCATTAG GCACCCGGAT CTCGACCGAT GCCCTTGAGA 1860
1861 GCCTTCAACC CAGTCAGCTC CTTCGGGTGG CGCGGGGGA TGACTATGCT CGCCGCACTT 1920
1921 ATGACTGTCT TCTTATCAT GCAACTCGTA GGACAGGTGC CGGCAGCGCT CTGGGTCATT 1980
1981 TCGCGGAGG ACCGCTTCCG CTGGAGCGGG ACGATGATCG GCCTGTGCTG TGGGTATTTC 2040
2041 GGAATCTTGC ACGCCCTCGC TCAAGCCCTC GTCACTGCTC CGCCACCAA ACGTTCGGCG 2100
2101 JAGAAGCAGG CCATTAATGC CGGCATGGCG GCGCCACGGG TCGCATGATG CGTGCCTCTG 2160
2161 TCGTGGAGGA CCGCGCTAGG CTGGCGGGGT TCGCTTACTG GTTAGCAGAA TGAATCACCG 2220
2221 ATACCGGAGC GAACGTGAAG CGACTGCTGC TCGAAAACCT CTGGCACTG AGCAACAACA 2280
2281 TGAATGGTCT TCGGTTCCCG TGTTCGTA AAGTCTGAAA CGCGGAAGTC AGCGCCCTCG 2340
2341 ACCATTAATG TCGGGATCTG CATCGCAGGA TGCTGTGCGC TACCTGTGG AACACCTACA 2400
2401 TCTGTATTA CGAAGCGCTG GCATGACCC TGAGTGAATT TTCTCTGGTC CCGCGCATC 2460
2461 CATACCCGCA GTTGTATACC CTCACAACGT TCCAGTAACC GGGCATGTTT ATCATCAGTA 2520
2521 ACCCGTATCG TGAGCATCCT CTCTCGTTTC ATCGGTATCA TTACCCTCAT GAACAGAAAT 2580
2581 CCGCTTACA CCGAGGCATC AGTGACCAA CAGGAAAAAA CCGCCCTTAA CATGGCCCGC 2640
2641 TTTATCAGAA CCCAGACATT AACCTTCTG GAGAAACTCA ACGAGCTGGA CGCGGATGAA 2700
2701 CAGGCAGACA TCTGTGAATC GCTTACAGAC CACGCTGATG AGCTTTACCG CAGCTGCCTC 2760
2761 CGCGCTTTCG GTGATGACGG TGAAAACCTC TGACACATGC AGCTCCCGGA GACGGTCACA 2820
2821 GCTTCTCTGT AAGCGGATGC CGGGAGCAGA CAAGCCCTTC AGGGCCGCTC AGCGGGTGT 2880
2881 GCGCGGTGTC GGGCGCGAGC CATGACCAGC TCAGTAGGCG ATAGCGGAGT GTATACTGCG 2940
2941 TTAACATGCG GGCATCAGAG CAGATTGTAC TGAGAGTGA CCAATATGCG GGTGTGAAAT 3000
3001 ACCGCACAGA TCGCTAAGGA GAAAATACCG CATCAGGCGC TCTTCCGCTT CCTCGCTCAC 3060
3061 TGACTCGCTG CGCTCGCTCG TCGGCTGCG CCGAGGGAAG AACATGTGAG CAAAAGGCCA 3120
3121 AATCTGATTA TCCACAGAAAT CAGGGATAA CGCAGGAAAG AACATGTGAG CAAAAGGCCA 3180
3181 CGAAAAGGCC AGGAACCGTA AAAAGGCGCG GTTGTGCGCG TTTTTCATA GCTTCCGCCC 3240
3241 CCCTGACGAG CATCACAATA ATCGACGCTC AAGTCAGAGG TGGCGAAACC CGACAGGACT 3300
3301 ATAAAGATAC CAGGCGTTTC CCGCTGGAAG CTCCCTCTGT CGCTCTCTGT TTCCGACCTT 3360
3361 GCGCTTACC CGATACCTGT CCGCTTTCT CCGTTCGGGA AGCGTGGCCG TTTCTCATAG 3420
3421 CTCACGCTGT AGGTATCTCA GTTGGGTGA GGTGTCTGCG TCAAGCTGG GCTGTGTGCA 3480
3481 CGAACCCCCC GTTCAGCCCG ACCGCTGGCG CTTATCCGCT AACATATGTC TTGATGCCAA 3540
3541 CCGGTAAAGA CAGCACTTAT CGCCACTGGC AGCAGCCACT GGTAAACAGGA TTAGCAGAGC 3600
3601 GAGGTATGTA GCGCGTCTCA CAGAGTTCTT GAAGTGGTGG CCTAACTAAG GCTACACTAG 3660
3661 AAGGCAGATA TTTGGTATCT CGGCTCTGCT GAAGCCAGTT ACCTTCGGAA AAAGAGTTGG 3720
3721 TAGCTCTTGA TCGCGCAAC AAACCACCCG TGGTAGCGGT GGTTTTTTTT TTTGCAAGCA 3780
3781 GCAGATTCAG CGCAAAAAAA AAGGATCTCA AGAAGATCTT TTGATTTTTT CTACGGGCTG 3840
3841 TGACGCTCAG TCGAAGCAAA ACTCACGTTA AGGGATTTTG GTCATGAGAT TATCAAAAAA 3900
3901 GATCTTACC TAGATCTTCT TAAATTAATA ATGAAGTTTT AAATCAATCT AAAGTATATA 3960
3961 TGAGTAAACT TGGTCTGACA GTTACCAATG CTTAATCAGT GAGGCCACTA TCTCAGCGAT 4020
4021 CTGTCTATTT CGTTCATCCA TAGTTGCCGT ACTCCCGCTC GTGTAGATAA CTACGATAGC 4080
4081 GGAGGGCTTA CCACTTGGCC CCAATGCTGC AATGATACCG CGAGACCAC GCTCACCCGG 4140
4141 TCCAGATTA TCAGCAATAA ACCAGCCAGC CGGAAGGGCC GAGCCGAGAA GTGGTCTGCG 4200
4201 AACTTTATCC GCTTCATCC AGTCTATTA TTGTTCGGG GAAGCTAGAG TAAGTAGTTC 4260
4261 GCCAGTTAAT AGTTTGGCCA ACGTTGTGCG CATTGTCTGA GGCATCTGGT GTTCAACGCT 4320
4321 CTGCTTGGT ATGGCTTCA TCACTCCGG TTCCCAACGA TCAAGGCGAG TTACATGATC 4380
4381 CCCCATGTTG TCGAAAAAAG CCGTTAGCTC CTTCCGTCTC CAGTCTGTTG TCAGAAGTAA 4440
4441 GTTGGCGGCA GTTGTATCAC TCAATGTTAT GGCAGCACTG CATAATCTC TTACTGTGAT 4500
4501 CCGCATCGTA AGATGCTTCT CTGTGACTGG TGAGTACTCA ACCAAGTCAT TCTGCAAGTA 4560
4561 GTGATGCGG CGACCGAGTT GCTCTTGGCC GCGCTCAATA CGGGATAATA CCGCGCCACA 4620
4621 TAGCAGAACT TTAAGATGCG TCACTATTGG AAAACCTTCT TCGGGCGGAA AACTCTCAAG 4680
4681 GATCTTACCG CTGTTGAGAT CGAGTTOGAT GTAACCCACT CGTGCACCCA ACTGATCTTC 4740
4741 AGCATCTTCT ACTTTCACCA CGGTTCTCG GTGAGCAAAA ACAGGAAGGC AAAATGCCCG 4800
4801 AAAAAAGGGA ATAAGGCGGA CACGGAAATG TTGAATACTC ATACTCTTCC TTTTCAATA 4860
4861 TTAATGAAGC ATTTATCAGG GTTATTGTCT CATGAGCGGA TACATAATTT AATGATTTA 4920
4921 GAAAAATAAA CAAATAGGGG TCCCGCGCAC ATTTCCCGCA AAAGTGGCCAC CTGAATTTGT 4980
4981 AAGCTTTAAT ATTTTCTTAA AATTCGGCTT AAATTTTTGT TAAATAGCT CATTTTTTAA 5040
5041 TAAATAGGCC GAAATCGGCA AATCCCTTA TAAATCAAAA GAATAGACCG AGATAGGGTT 5100
5101 GAATGTTGTT CCAAGTTGGA ACAAGAGTCC ACTATTAAAG AACGTGGCT CCAACGTCAA 5160
5161 AGGCGAAAAA ACGCTTATC AGGCGGATGG CCGACTACGT GAACCATCAC CCTAATCAAG 5220
5221 TTTTTTGGCG TCGAGGTGCC GTAAGCACT AAATCUGAAC CCTAAAGGGA GCCCCGATT 5280
5281 TAGAGCTTGA CGGGAAAGC GGGCGAACGT GCGCAGAAAG GAAGGGAAGA AAGCGAAAG 5340
5341 AGCGGGGCT AGGGCGCTGG CAAGTGTAGC GGTACCGCTG CGGTAAACCA CCACACCCCG 5400
5401 CGGCTTAAAT CGCGGCTAC AGGGCGCTC CCATTGCCCA
      | 10 | 20 | 30 | 40 | 50 | 60

```

pET21d Vector Genetic Map

DNA sequence 5440 b.p. ATCCGGATATAG ... TCCCATTGCCA circular



GAL4-D 1-phase Translation

DNA sequence 324 b.p. atgaagctactg ... atcactacaggg linear


```

1 / 1 31 / 11
atg aag cta ctg tct tct atc gaa caa gca tgc gat att tgc cga ctt aaa aag ctc aag
met lys leu leu ser ser ile glu gln ala cys asp ile cys arg leu lys lys leu lys
61 / 21 91 / 31
tgc tcc aaa gaa aaa ccg aag tgc gcc aag tgt ctg aag aac aac tgg gag tgt cgc tac
cys ser lys glu lys pro lys cys ala lys cys leu lys asn asn trp glu cys arg tyr
121 / 41 151 / 51
tct ccc aaa acc aaa agg tct ccg ctg act agg gca cat ctg aca gaa gtg gaa tca agg
ser pro lys thr lys arg ser pro leu thr arg ala his leu thr glu val glu ser arg
181 / 61 211 / 71
cta gaa aga ctg gaa cag cta ttt cta ctg att ttt cct cga gaa gac ctt gac atg att
leu glu arg leu glu gln leu phe leu leu ile phe pro arg glu asp leu asp met ile
241 / 81 271 / 91
ttg aaa atg gat tct tta cag gat ata aaa gca ttg tta acc atg gac caa act gcg tat
leu lys met asp ser leu gln asp ile lys ala leu leu thr met asp gln thr ala tyr
301 / 101
aac gcg ttt gga atc act aca ggg
asn ala phe gly ile thr thr gly

```

GAL4-D Genetic Map

DNA sequence 324 b.p. atgaagctactg ... atcactacaggg circular



GAL4-AD 1-phase Translation

DNA sequence 384 b.p. atgaagctactg ... gatgaagataacc linear


```

1 / 1                               31 / 11
atg aag cta ctg tct tct atc gaa caa gca tgc gat att tgc cga ctt aaa aag ctc aag
met lys leu leu ser ser ile glu gln ala cys asp ile cys arg leu lys lys leu lys
61 / 21                               91 / 31
tgc tcc aaa gaa aaa ccg aag tgc gcc aag tgt ctg aag aac aac tgg gag tgt cgc tac
cys ser lys glu lys pro lys cys ala lys cys leu lys asn asn trp glu cys arg tyr
121 / 41                              151 / 51
tct ccc aaa acc aaa agg tct ccg ctg act agg gca cat ctg aca gaa gtg gaa tca agg
ser pro lys thr lys arg ser pro leu thr arg ala his leu thr glu val glu ser arg
181 / 61                              211 / 71
cta gaa aga ctg gaa cag cta ttt cta ctg att ttt cct cga gaa gac ctt gac atg att
leu glu arg leu glu gln leu phe leu leu ile phe pro arg glu asp leu asp met ile
241 / 81                              271 / 91
ttg aaa atg gat tct tta cag gat ata aaa gca ttg ttg acc atg gac caa act gcg tat
leu lys met asp ser leu gln asp ile lys ala leu leu thr met asp gln thr ala tyr
301 / 101                             331 / 111
aac gcg ttt gga atc act aca ggg atg ttt aat acc act aca atg gat gat gta tat aac
asn ala phe gly ile thr thr gly met phe asn thr thr thr met asp asp val tyr asn
361 / 121
tat cta ttc gat gat gaa gat acc
tyr leu phe asp asp glu asp thr

```

GAL4-AD Genetic Map

DNA sequence 384 b.p. atgaagctactg ... gatgaagatacc circular



GAL4-VP16 Experimental Linear Sequence

5' end – accatctgctaaactcgaagggtcggcccaattattcccataggaaggcgccgtagggcgagtcgtggggcgtaatcc
ggacccggggaatccccgtcccccaacatgtcagatcgaatcgtctagcgctcgcatgccatcgctcgtctagctggctccagtga
ctctgcatgggcattcctactcactttctact - 3' end.


*GAL4-VP16 Linear Sequence*¹¹

ATGAAGCTACTGTCTTCTATCGAACAAGCATGCGATATTTGCCGACTTAA
AAAGCTCAAGTGCTCCAAAGAAAAACCGAAGTGCGCCAAGTGTC'TGAAGA
ACAACTGGGAGTGTCGCTACTCTCCCAAACCAAAGGTCTCCGCTGACT
AGGGCACATCTGACAGAAGTGGAATCAAGGCTAGAAAGACTGGAACAGCT
ATTTCTACTGATTTTTCTCGAGAAGACCTTGACATGATTTTAAAAATGG
ATTCTTTACAGGATATAAAAGCATTGTTAACAGGATTATTTGTACAAGAT
AATGTGAATAAAGATGCCGTCACAGATAGATTGGCTTCAGTGGAGACTGA
TATGCCTCTAACATTGAGACAGCATAGAATAAGTGCGACATCATCATCGG
AAGAGAGTAGTAACAAAGGTCAAAGACAGTTGACTGTATCG

CCGGAATTCCTGGGGATCTGGG

GCCCCCCCACCGATGTCAGCCTGGGGGACGAGCTCCACTTAGAC
GGCGAGGACGTGGCGATGGCGCATGCCGACGCGCTAGACGATTTT
GATCTGGACATGTTGGGGGACGGGGATTCCCCGGGTCCGGGATTTACC
CCCCACGACTCCGCCCCCTACGGCGCTCTGGATATGGCCGACTTCGAG
TTTGAGCAGATGTTTACCGATGCCCTTGAATTGACGAGTACGGTGGGTAG

¹¹ Contributed by the Department of Biochemistry, Michigan State University.

pJL2 Bacterial Expression Vector

Plasmid name: pJL2

Plasmid size: 4256 bp

Constructed by: Janet Leatherwood

Construction date: 1987?

Comments/References: Expression of GAL4-VP16 fusion protein from ptac promoter.

Map constructed in 1993 by Steve Triezenberg. Some map positions, especially in P22-ant and ptac regions, are only approximate.

Appendix B

Materials:


1. pET21d plasmid (1997 Novagen Catalog, p. 122)
2. pGEX-cs plasmid (Novagen catalog)
3. GAL4-AD DNA, GAL4-D DNA
4. GAL4-VP16 DNA (Michigan State University)
5. PCR materials (Pfu, dNTP, PCR buffer, etc)
6. Shrimp alkaline phosphatase
7. T4 DNA ligase
8. Media components
9. DNA extraction kit (BIGGERprep Plasmid Preparation Kit)
10. Agarose (type)
11. Acrylamide gels
12. NcoI restriction enzyme (New England Biolabs Catalog, p. 46)
13. XhoI restriction enzyme (New England Biolabs Catalog, p. 60)
14. Resin for the micro column purification method

Methods:

1. Cell transformation by electroporation
2. DNA isolation kit. (Kiagen Biggerprep)
3. PCR amplification, insert construction.
4. Plasmid dephosphorylation
5. Plasmid-PCR constructs ligation


6. Competent cell production
7. pET21d plasmid digestion
8. pET21d plasmid purification by spincolumn procedure
9. Quantification of isolated DNA concentration by UV scan in spectrophotometer
10. GAL4-VP16 TF sequencing, method 1
11. GAL4-VP16 TF sequencing, method 2
12. Preparation of the media (recipes)
13. Agarose gels check for experimental results
14. Sequencing acrylamide gels

Appendix C

Figure 1: pET21d Digestion by *Xho*I and *Nco*I


1. 1 Kb DNA Ladder (loaded 0.5 μ l).
2. Undigested pET21d Vector (50ng/ μ l, loaded 3 μ l).
3. Digested pET21d Vector (50ng/ μ l, loaded 3 μ l).

Figure 2: GAL4-AD and GAL4-D Digested PCR Inserts



1. 50 bp DNA Ladder (loaded 0.5 μ l).
2. GAL4-AD Insert (loaded 20 μ l of the PCR reaction).
3. GAL4-D Insert (loaded 20 μ l of the PCR reaction).

Figure 3: GAL4-AD and GAL4-D Ligation Products



1. Low Mass DNA Ladder (loaded 2 μ l).
2. GAL4-AD Ligation Reaction (loaded 2 μ l).
3. GAL4-AD Ligation Reaction (loaded 2 μ l).
4. GAL4-D Ligation Reaction (loaded 2 μ l).
5. GAL4-D Ligation Reaction (loaded 2 μ l).
6. pET21d Ligation Reaction (loaded 2 μ l).
7. GAL4-AD Insert Ligation Reaction (loaded 2 μ l).
8. 1 Kb DNA Ladder (loaded 0.5 μ l).






Figure 4: Taq Polymerase Protocol Amplifications of GAL4-VP16




1. Low Mass DNA Ladder (loaded 2 μ l).
2. Taq Polymerase Amplified GAL4-VP16 Inserts (loaded 6 μ l from the PCR reaction).
3. Taq Polymerase Amplified GAL4-VP16 Inserts (loaded 6 μ l from the PCR reaction).
4. Taq Polymerase Amplified GAL4-VP16 Inserts (loaded 6 μ l from the PCR reaction). Taq Polymerase Amplified GAL4-VP16 Inserts (loaded 6 μ l from the PCR reaction).
5. Taq Polymerase Amplified GAL4-VP16 Inserts (loaded 6 μ l from the PCR reaction).
6. Taq Polymerase Amplified GAL4-VP16 Inserts (loaded 6 μ l from the PCR reaction).

Figure 5: Pfu Turbo Protocol Amplifications of GAL4-VP16



1. Low Mass DNA Ladder (loaded 2 μ l).
2. Pfu Turbo Amplified GAL4-VP16 Inserts (loaded 6 μ l from the PCR reaction).
3. Pfu Turbo Amplified GAL4-VP16 Inserts (loaded 6 μ l from the PCR reaction).
4. Pfu Turbo Amplified GAL4-VP16 Inserts (loaded 6 μ l from the PCR reaction).
5. Pfu Turbo Amplified GAL4-VP16 Inserts (loaded 6 μ l from the PCR reaction).

Figure 6: GAL4 Series Amplification Products



1. Low Mass DNA Ladder (loaded 2 μ l).
2. GAL4-AD Insert (loaded 6 μ l from digested insert).
3. GAL4-D Insert (loaded 6 μ l from digested insert).
4. GAL4-VP16 Insert (loaded 6 μ l from the PCR reaction).

Acknowledgements

I would like to thank my advisor on this project, Dr. Blaine Bartholomew, for this research opportunity and for his support and guidance. Also, I would like to thank Jim Persinger for all of his practical research advice.

I would like to express my gratitude to Søren Jaglo-Ottosen of Michigan State University's Department of Biochemistry, for providing with a vector containing GAL4-VP16.

Finally, I would like to thank my wife for her unconditional support and her assistance in the editing process of this thesis.

References

1. Beato, M. & K. Eisfeld. 1997. Transcription factor access to chromatin. *Nucleic Acids Res*, 25: 3559-3563.
2. Gadhavi P.L. 1997. An electrospray ionisation mass spectrometry (ESI-MS) study to probe the metal ion binding site in the DNA binding domain of the yeast transcriptional activator GAL4. *FEBS Lett*, 417(1): 145-149.
3. Lewin, B. 1997. Genes VI. Oxford Univ, New York. 1260 pp.
4. Liang S.D., R. Marmorstein, S.C. Harrison & M. Ptashine. 1996. DNA sequence preferences of GAL4 and PPR1: how a subset of Zn₂ Cys₆ binuclear cluster proteins recognize DNA. *Mol Cell Biol*, 16(7): 3773-3780.
5. Melcher K. & S.A. Johnston. 1995. GAL4 interacts with TATA-binding protein and coactivators. *Mol Cell Biol*, 15(5): 2839-2848.
6. Poch O. 1997. Conservation of a putative inhibitory domain in the GAL4 family members. *Gene*, 184(2), 229-235.
7. Xu H.E., T. Kodadek & S.A. Johnston. 1995. A single GAL4 dimer can maximally activate transcription under physiological conditions. *Proc Natl Acad Sci U S A*, 92(17): 7677-7680.
8. Xu M., R.T. Simpson & M.P. Kladde. 1998. GAL4p-mediated chromatin remodeling depends on binding site position in nucleosome but does not require DNA replication. *Mol Cel Biol*, 18(3): 1201-1212.