

Southern Illinois University Carbondale
OpenSIUC

Articles and Preprints

Department of Mathematics

2009

Multivariate Matching Polynomials of Cyclically Labelled Graphs

John P. McSorley

Southern Illinois University Carbondale, mcsorley60@hotmail.com

Philip Feinsilver

Southern Illinois University Carbondale

Follow this and additional works at: http://opensiuc.lib.siu.edu/math_articles

Published in McSorley, J. P., & Feinsilver, P. (2009). Multivariate matching polynomials of cyclically labelled graphs. *Discrete Mathematics*, 309(10), 3205-3218 .

Recommended Citation

McSorley, John P. and Feinsilver, Philip. "Multivariate Matching Polynomials of Cyclically Labelled Graphs." (Jan 2009).

This Article is brought to you for free and open access by the Department of Mathematics at OpenSIUC. It has been accepted for inclusion in Articles and Preprints by an authorized administrator of OpenSIUC. For more information, please contact opensiuc@lib.siu.edu.

Multivariate Matching Polynomials of Cyclically Labelled Graphs

John P. McSorley and Philip Feinsilver,
Department of Mathematics,
Mailcode 4408,
Southern Illinois University,
Carbondale. IL 62901-4408.
mcsorley60@hotmail.com
pfeinsil@math.siu.edu

Keywords

Matching, Matching Polynomial, Labelled, Graph, Recurrence,
MacMahon's Master Theorem

Abstract

We consider the matching polynomials of graphs whose edges have been cyclically labelled with the ordered set of t labels $\{x_1, \dots, x_t\}$.

We first work with the cyclically labelled path, with first edge label x_i , followed by N full cycles of labels $\{x_1, \dots, x_t\}$, and last edge label x_j . Let $\Phi_{i,Nt+j}$ denote the matching polynomial of this path. It satisfies the (τ, Δ) -recurrence: $\Phi_{i,Nt+j} = \tau \Phi_{i,(N-1)t+j} - \Delta \Phi_{i,(N-2)t+j}$, where τ is the sum of all non-consecutive cyclic monomials in the variables $\{x_1, \dots, x_t\}$ and $\Delta = (-1)^t x_1 \cdots x_t$. A combinatorial/algebraic proof and a matrix proof of this fact are given. Let G_N denote the first fundamental solution to the (τ, Δ) -recurrence. We express G_N (i) as a cyclic binomial using the Symmetric Representation of a matrix, (ii) in terms of Chebyshev polynomials of the second kind in the variables τ and Δ , and (iii) as a quotient of two matching polynomials. We extend our results from paths to cycles and rooted trees.

Introduction

The matching polynomial of a graph is defined in Farrell [1]. Often in pure mathematics and combinatorics it is interesting to consider cyclic structures, *eg.*, cyclic groups, cyclic designs, and circulant graphs. Here we consider the (multivariate) matching polynomial of a graph whose edges have been cyclically labelled.

We concentrate mainly on paths, cycles and trees. To cyclically label a path with the ordered set of t labels $\{x_1, \dots, x_t\}$, label the first edge with any x_i , the second with x_{i+1} , and so on until label x_t has been used, then start with x_1 , then x_2, \dots, x_t , then x_1 again \dots , repeating cyclically until all edges have been labelled, with the last edge receiving label x_j . Suppose that N full cycles of labels $\{x_1, \dots, x_t\}$ have been used. Call the matching polynomial of this labelled path $\Phi_{i,Nt+j}$. We show, for a fixed i and j , that $\Phi_{i,Nt+j}$ satisfies the following recurrence, the (τ, Δ) -recurrence:

$$\Phi_{i,Nt+j} = \tau \Phi_{i,(N-1)t+j} - \Delta \Phi_{i,(N-2)t+j},$$

where τ is the sum of all non-consecutive cyclic monomials in the variables $\{x_1, \dots, x_t\}$ (see Section 1), and $\Delta = (-1)^t x_1 \cdots x_t$. We give two different proofs of this fact. The first one is a combinatorial/algebraic proof in Section 2 that uses the following Theorem concerning decomposing the matching polynomial $\mathcal{M}(G, \mathbf{x})$ of a graph.

Theorem Let G be a labelled graph, H a subgraph of G , and M_H a matching of H , then

$$\mathcal{M}(G, \mathbf{x}) = \sum_{M_H} M_H(\mathbf{x}) \mathcal{M}(G - H - \overline{M}_H, \mathbf{x}),$$

where the summation is over every matching M_H of H . The second proof (Section 3) uses a matrix formulation of the recurrences that we develop.

Let G_N denote the first fundamental solution to the (τ, Δ) -recurrence; three different expressions for G_N are given in Section 4. The first expression is a sum of cyclic binomials and uses the Symmetric Representation of matrices from Section 3; the second involves Chebyshev polynomials of the second kind in the variables τ and Δ ; and the third is a quotient of two matching polynomials, see Theorem 4.5.

In Section 5 we extend our results from paths to cycles and rooted trees; we find explicit forms for the matching polynomial of a cyclically labelled cycle, and indicate how to find the matching polynomial of a cyclically labelled rooted tree, again using the decomposition Theorem stated above.

Many examples are given throughout the paper.

1 The multivariate matching polynomial of a graph, its decomposition; non-consecutive and non-consecutive cyclic functions

For a fixed $t \geq 1$ we use multi-index notations: $\mathbf{k} = (k_1, \dots, k_t)$, where each $k_s \geq 0$, $\mathbf{0} = (0, \dots, 0)$, and variables $\mathbf{x} = (x_1, \dots, x_t)$. The total degree of \mathbf{k} is denoted by $|\mathbf{k}| = k_1 + \dots + k_t$.

Let G be a finite simple graph with vertex set $V(G)$ where $|V(G)| \geq 1$, and edge set $E(G)$. We label these edges from the t commutative variables $\{x_1, \dots, x_t\}$, exactly one label per edge. A *matching* of G is a collection of edges, no two of which have a vertex in common. A *\mathbf{k} -matching* of G is a matching with exactly k_s edges with label x_s , for each s with $1 \leq s \leq t$. If M_G is a \mathbf{k} -matching of G we define its *weight* to be

$$M_G(\mathbf{x}) = x_1^{k_1} \cdots x_t^{k_t}.$$

The empty matching of G , which contains no edges, is denoted by M_\emptyset ; it is the unique $\mathbf{0}$ -matching and its weight is $M_\emptyset(\mathbf{x}) = 1$.

Define the *multivariate matching polynomial*, or simply, the *matching polynomial*, of G , by

$$\mathcal{M}(G, \mathbf{x}) = \sum_{M_G} M_G(\mathbf{x}),$$

where the summation is over every matching M_G of G .

Denote the number of \mathbf{k} -matchings of G by $a(G, \mathbf{k})$. Then an alternative definition of the multivariate matching polynomial of G is

$$\mathcal{M}(G, \mathbf{x}) = \sum_{(k_1, \dots, k_t)} a(G, \mathbf{k}) x_1^{k_1} \cdots x_t^{k_t}.$$

The multivariate matching polynomial is a natural extension of the matching polynomial of Farrell [1]. Indeed, here with $t = 1$ and in [1] with $w_1 = 1$ and $w_2 = x_1$, the polynomials are identical.

Let P_1 be the graph with one vertex and no edges, *i.e.*, an isolated vertex; we define $\mathcal{M}(P_1, \mathbf{x}) = 1$. Now suppose $G' = G \cup nP_1$, where $n \geq 1$, *i.e.*, G'

is the disjoint union of G and n isolated vertices, then we define $\mathcal{M}(G', \mathbf{x}) = \mathcal{M}(G, \mathbf{x})$.

For any edge $e \in E(G)$, let \bar{e} denote the set of edges that are incident to e ; and for any subgraph H of G , let $\bar{H} = \cup_{e \in E(H)} \bar{e}$. Define $\bar{M}_\emptyset = \emptyset$. Also let $G - H$ be the graph obtained from G when all the edges of H are removed, so $G - H$ has the same vertex set as G .

Now let H be a fixed subgraph of G and let M_H be a matching of H . In the following theorem we express $\mathcal{M}(G, \mathbf{x})$ as a sum of terms, each term containing the weight of a fixed matching, $M_H(\mathbf{x})$, of H ; we call this *decomposing* $\mathcal{M}(G, \mathbf{x})$ at H .

Theorem 1.1 *Let G be a graph labelled as above, H a fixed subgraph of G , and M_H a matching of H . Then*

$$\mathcal{M}(G, \mathbf{x}) = \sum_{M_H} M_H(\mathbf{x}) \mathcal{M}(G - H - \bar{M}_H, \mathbf{x}), \quad (1)$$

where the summation is over every matching M_H of H .

Proof. Let M_G be a matching of G which induces a (fixed) matching M_H on H , i.e., M_G contains exactly M_H and no other edges from H . Then $M_G(\mathbf{x}) = M_H(\mathbf{x}) M(\mathbf{x})$ where M is a matching of G with no edges in H , and also with no edges in \bar{M}_H or else M_G would not be a matching. Hence, M is a matching of $G - H - \bar{M}_H$, i.e., $M(\mathbf{x})$ is a term of $\mathcal{M}(G - H - \bar{M}_H, \mathbf{x})$. So $M_H(\mathbf{x}) \mathcal{M}(G - H - \bar{M}_H, \mathbf{x})$ is the sum of the weights of all the matchings in G which induce M_H on H .

Now every matching in G induces some matching on H , so we may sum over all matchings in H to give (1). \blacksquare

Theorem 1.1 extends known facts about matching polynomials, eg., see Theorem 1 of Farrell [1] for the case where H is a single edge. We have the corresponding:

Corollary 1.2 *Let G be a graph labelled as above, and let $H = e$ labelled with x be an edge of G . Then*

$$\mathcal{M}(G, \mathbf{x}) = \mathcal{M}(G - e, \mathbf{x}) + x \mathcal{M}(G - e - \bar{e}, \mathbf{x}). \quad (2)$$

Proof. The result comes from (1) since $H = e$ has just two matchings: the empty matching M_\emptyset with weight $M_\emptyset(\mathbf{x}) = 1$, and the matching e with weight $M_e(\mathbf{x}) = x$. ■

Notation Throughout this paper we use P_m to denote the path with m vertices and $m - 1$ edges.

Fix i and j where $1 \leq i \leq j \leq t$. Consider the path P_{j-i+2} with its $j-i+1$ edges labelled from the ordered set $\{x_i, \dots, x_j\}$, the first edge receiving label x_i , and the last x_j ; see Fig. 1.




Fig. 1: The labelled path P_{j-i+2} with matching polynomial $\phi_{i,j}$.

The pair $x_s x_{s+1}$ for any fixed s with $i \leq s \leq j - 1$ is called a *consecutive* pair. A monomial from the ordered set $\{x_i, \dots, x_j\}$ that contains no consecutive pairs is a *non-consecutive monomial*, a *nc-monomial*. Note that the empty monomial is a *nc-monomial* that we denote by 1.

Let $\phi_{i,j}$ be the sum of all *nc-monomials* in the ordered variables $\{x_i, \dots, x_j\}$. Then $\phi_{i,j} = \mathcal{M}(P_{j-i+2}, \mathbf{x})$ is the matching polynomial of the labelled path P_{j-i+2} . We call the functions $\phi_{i,j}$ *elementary non-consecutive functions*, and for any $i \geq 1$ define the initial values

$$\phi_{i,i-2} = \phi_{i,i-1} = 1. \quad (3)$$

These initial values ensure that the following recurrence is valid for any j with $i \leq j \leq t$.

Theorem 1.3 For a fixed i and j with $1 \leq i \leq j \leq t$ and the initial values in (3), we have

$$\phi_{i,j} = \phi_{i,j-1} + x_j \phi_{i,j-2}. \quad (4)$$

Proof. Let e be the rightmost edge of $G = P_{j-i+2}$ shown in Fig. 1, and apply (2). ■

Example 1 For arbitrary i we have

$$\begin{aligned}\phi_{i,i} &= 1 + x_i, & \phi_{i,i+1} &= 1 + x_i + x_{i+1}, \\ \phi_{i,i+2} &= 1 + x_i + x_{i+1} + x_{i+2} + x_i x_{i+2}, \\ \phi_{i,i+3} &= 1 + x_i + x_{i+1} + x_{i+2} + x_{i+3} + x_i x_{i+2} + x_i x_{i+3} + x_{i+1} x_{i+3}.\end{aligned}$$

Example 2 For arbitrary i , putting $j = i - 1$ and $j = i - 2$ in Recurrence (4) and using (3) give

$$\phi_{i,i-3} = 0 \quad \text{and} \quad \phi_{i,i-4} = \frac{1}{x_{i-2}}.$$

In the second equation, if $i = 1$ we replace x_{-1} by x_{t-1} , and if $i = 2$ we replace x_0 by x_t .

Consider Recurrence (4). It is convenient to work with a basis of solutions to this recurrence. Denote the first fundamental solution by $f_{i,j}$ and the second by $g_{i,j}$, with initial values

$$f_{i,i-2} = 0, f_{i,i-1} = 1 \quad \text{and} \quad g_{i,i-2} = 1, g_{i,i-1} = 0. \quad (5)$$

So

$$\phi_{i,i-2} = f_{i,i-2} + g_{i,i-2} \quad \text{and} \quad \phi_{i,i-1} = f_{i,i-1} + g_{i,i-1}.$$

Now, from Recurrence (4) and strong induction on j , we have (6) below for all j with $i \leq j \leq t$

$$\phi_{i,j} = f_{i,j} + g_{i,j}. \quad (6)$$

$$\phi_{i,j} = \phi_{i+1,j} + x_i \phi_{i+2,j}. \quad (7)$$

Equation (7) comes from decomposing $\phi_{i,j}$ at the leftmost edge of P_{j-i+2} , whose label is x_i , *i.e.*, decomposing $\phi_{i,j}$ at x_i ; see Corollary 1.2. These two equations suggest that the fundamental solutions are given by

$$f_{i,j} = \phi_{i+1,j} \quad \text{and} \quad g_{i,j} = x_i \phi_{i+2,j}.$$

This is indeed the case:

Lemma 1.4 *For any j with $i \leq j \leq t$ we have*

- (i) $f_{i,j} = \phi_{i+1,j}$,
- (ii) $g_{i,j} = x_i \phi_{i+2,j}$.

Proof. We need only prove (i) because of (6) and (7) above.

From (5) we have $f_{i,i-2} = 0$ and from Example 2 we have $\phi_{i+1,i-2} = 0$; thus $f_{i,i-2} = \phi_{i+1,i-2}$. Similarly, from (5) and (3), we have $f_{i,i-1} = \phi_{i+1,i-1}$. So both $f_{i,j}$ and $\phi_{i+1,j}$ have the same initial values at $j = i - 2$ and $j = i - 1$ and they both satisfy Recurrence (4), so they are equal for any j with $i \leq j \leq t$. ■

Thus we know combinatorially what the two fundamental solutions to Recurrence (4) are. The first, $f_{i,j}$, is the matching polynomial of the path shown in Fig. 2(a); the second, $g_{i,j}$, is $x_i \times$ the matching polynomial of the path in Fig. 2(b).




Fig. 2 (a) The labelled path with matching polynomial $f_{i,j}$.

(b) The labelled path with matching polynomial $\frac{g_{i,j}}{x_i}$.

Example 3 For arbitrary i we have

$$\begin{aligned} f_{i,i} &= 1, & g_{i,i} &= x_i, \\ f_{i,i+1} &= 1 + x_{i+1}, & g_{i,i+1} &= x_i, \\ f_{i,i+2} &= 1 + x_{i+1} + x_{i+2}, & g_{i,i+2} &= x_i + x_i x_{i+2}. \end{aligned}$$

Now arrange the variables $\{x_i, \dots, x_j\}$ clockwise around a circle. Thus x_i and x_j are consecutive. Call a pair $x_s x_{s'}$ *consecutive cyclic* if x_s and $x_{s'}$ are consecutive on this circle. Call a monomial from $\{x_i, \dots, x_j\}$ a *non-consecutive cyclic* monomial — *ncc*-monomial — if it contains no consecutive cyclic pairs. The empty monomial is a *ncc*-monomial that we denote by 1.

Let $\tau_{i,j}$ be the sum of all *ncc*-monomials in the variables $\{x_i, \dots, x_j\}$. Then, for $j \geq i + 2$, $\tau_{i,j} = \mathcal{M}(C_{j-i+1}, \mathbf{x})$ is the matching polynomial of the

labelled cycle C_{j-i+1} with $j - i + 1$ edges and $j - i + 1$ vertices, shown in Fig. 3; the cycle starts at the large vertex, and proceeds clockwise.




Fig. 3: The labelled cycle C_{j-i+1} with matching polynomial $\tau_{i,j}$.

For initial values let

$$\tau_{i,i-1} = 2, \quad \tau_{i,i} = 1, \quad \text{and} \quad \tau_{i,i+1} = 1 + x_i + x_{i+1}. \quad (8)$$

Lemma 1.5 *For any j with $i \leq j \leq t$ we have*

- (i) $\tau_{i,j} = f_{i,j} + g_{i,j-1}$,
- (ii) $\phi_{i,j} - \tau_{i,j} = x_i x_j \phi_{i+2,j-2}$.

Proof. (i) We check this equality at $j = i$ and $j = i + 1$ using (5), Example 3, and (8). For $j \geq i + 2$ we decompose $\tau_{i,j}$ at x_i yielding $\tau_{i,j} = \phi_{i+1,j} + x_i \phi_{i+2,j-1}$, which gives (i) via Lemma 1.4.

(ii) We check at $j = i$ and $j = i + 1$ using Examples 1 and 2, and (8). For $j \geq i + 2$ the difference $\phi_{i,j} - \tau_{i,j}$ consists of all nc -monomials that contain the consecutive cyclic pair $x_i x_j$; clearly this is $x_i x_j \times$ the sum of all nc -monomials on $\{x_{i+2}, \dots, x_{j-2}\}$, *i.e.*, $x_i x_j \phi_{i+2,j-2}$. ■

Example 4 For arbitrary i we have

$$\begin{aligned} \tau_{i,i+2} &= 1 + x_i + x_{i+1} + x_{i+2}, \\ \tau_{i,i+3} &= 1 + x_i + x_{i+1} + x_{i+2} + x_{i+3} + x_i x_{i+2} + x_{i+1} x_{i+3}. \end{aligned}$$

2 Cyclically labelled paths; $\Phi_{i,Nt+j}$ and the (τ, Δ) -recurrence

Consider a path P and the ordered set of t labels $\{x_1, \dots, x_t\}$. For a fixed i , where $1 \leq i \leq t$, and moving from left to right, label the first edge of P with x_i , the second with x_{i+1} , and so on until label x_t has been used; so the $(t-i+1)$ -th edge receives label x_t . Then label edge $t-i+2$ with x_1 , and edge $t-i+3$ with x_2 , and so on \dots , labelling cyclically with $\{x_1, \dots, x_t\}$ until all edges have been labelled. Let the last edge receive label x_j , where $1 \leq j \leq t$. Suppose that $N \geq 0$ full cycles of labels $\{x_1, \dots, x_t\}$ have been used beginning at edge $t-i+2$. Then if $j = t$ we call this path $P(i, Nt)$, or if $1 \leq j < t$ we call it $P(i, Nt+j)$. This labelling is a *cyclic labelling*. The cyclically labelled path $P(i, Nt+j)$ is shown in Fig. 4. Let $\Phi_{i,Nt+j}(\mathbf{x}) = \Phi_{i,Nt+j}$ denote the matching polynomial of the path $P(i, Nt+j)$.




Fig. 4: The cyclically labelled path $P(i, Nt+j)$ with matching polynomial $\Phi_{i,Nt+j}$.

We define the initial conditions for $\Phi_{i,Nt+j}$ as

$$\begin{aligned} N = -1 : \quad & \Phi_{i,-t+j} = \phi_{i,j}, \quad \text{for all } j \text{ with } 0 \leq j \leq t, \\ \text{also } N = 0 : \quad & \Phi_{i,0t+j} = \Phi_{i,j}. \end{aligned} \quad (9)$$

In order to find $\phi_{i,j}$ if $j < i$ we use the initial values for $\phi_{i,j}$ from (3) and push back Recurrence (4), as shown in Example 2.

Now $\Phi_{i,Nt+j}$ satisfies the same recurrence as that of $\phi_{i,j}$, Recurrence (4); the proof is similar, noting that x_0 must be replaced by x_t , and considering $Nt-1$ as $(N-1)t+t-1$, etc.

Lemma 2.1 *For any $N \geq -1$ and j with $0 \leq j \leq t$ we have*

$$\Phi_{i,Nt+j} = \Phi_{i,Nt+j-1} + x_j \Phi_{i,Nt+j-2}. \quad (10)$$

■

Notation For $i = 1$ we write $\phi_{i,j} = \phi_{1,j} = \phi_j$ and $\phi_t = \phi$, also $\tau_{1,j} = \tau_j$ and $\tau_t = \tau$, and $f_{1,j} = f_j$, etc. Also let $\Delta = (-1)^t x_1 \cdots x_t$.

Lemma 2.2 For any $N \geq 0$ and any j with $0 \leq j \leq t$ we have

$$\Phi_{i,Nt+j} = \Phi_{i,Nt} f_j + \Phi_{i,Nt-1} g_j. \quad (11)$$

Proof. With $N = 0$ and $j = 0$ Equation (11) is true using the initial values $f_0 = 1$ and $g_0 = 0$ of (5) with $i = 1$. Otherwise, consider the path $P(i, Nt + j)$ of Fig. 4 and decompose its matching polynomial, $\Phi_{i,Nt+j}$, at the edge labelled x_1 marked with a $*$. This gives

$$\begin{aligned} \Phi_{i,Nt+j} &= \Phi_{i,Nt} \phi_{2,j} + x_1 \Phi_{i,Nt-1} \phi_{3,j} \\ &= \Phi_{i,Nt} f_j + \Phi_{i,Nt-1} g_j, \end{aligned}$$

using Lemma 1.4. ■

Now we define the second order (τ, Δ) -recurrence

$$\Theta_N = \tau \Theta_{N-1} - \Delta \Theta_{N-2}. \quad (12)$$

Let $G_N(\mathbf{x}) = G_N$ denote the first fundamental solution to this recurrence. We will evaluate G_N in Section 4.

In Theorem 2.4 below we show that, for a fixed i and j , $\Phi_{i,Nt+j}$ satisfies the (τ, Δ) -recurrence. First:

Lemma 2.3 For any $N \geq 1$ we have

$$\begin{aligned} (i) \quad &\Phi_{i,Nt-1} f_t - \Phi_{i,Nt} f_{t-1} = \Delta \Phi_{i,(N-1)t-1}, \\ (ii) \quad &\Phi_{i,Nt-1} g_t - \Phi_{i,Nt} g_{t-1} = -\Delta \Phi_{i,(N-1)t}. \end{aligned} \quad (13)$$

Proof. (i) Using Recurrence (4) on f_t and on $\Phi_{i,Nt}$ (see Lemma 2.1), the left-hand side of (13) becomes

$$\Phi_{i,Nt-1} \{f_{t-1} + x_t f_{t-2}\} - \{\Phi_{i,Nt-1} + x_t \Phi_{i,Nt-2}\} f_{t-1} = -x_t \{\Phi_{i,Nt-2} f_{t-1} - \Phi_{i,Nt-1} f_{t-2}\}.$$

The second factor in the right-hand side of this equation is the left-hand side of (13) with subscripts shifted down by 1. After t such iterations the left-hand side of (13) becomes

$$(-x_t)(-x_{t-1}) \cdots (-x_1) \{\Phi_{i,(N-1)t-1} f_0 - \Phi_{i,(N-1)t} f_{-1}\} = \Delta \Phi_{i,(N-1)t-1},$$

using the initial values $f_0 = 1$ and $f_{-1} = 0$. The proof of (ii) is similar. ■

Now a main result: $\Phi_{i,Nt+j}$ satisfies the (τ, Δ) -recurrence.

Theorem 2.4 *For any $N \geq 1$, and any fixed i with $1 \leq i \leq t$, and any fixed j with $0 \leq j \leq t$, we have*

$$\Phi_{i,Nt+j} = \tau \Phi_{i,(N-1)t+j} - \Delta \Phi_{i,(N-2)t+j}. \quad (14)$$

Proof. Due to Recurrences (4) and (10) we need only show that (14) is true when $j = t$ and $t - 1$. It will then be true for all j with $0 \leq j \leq t$ by pushing back Recurrence (10).

With $N \geq 1$ and $j = t$, Equation (11) gives

$$\begin{aligned} \Phi_{i,Nt+t} &= \Phi_{i,Nt} f_t + \Phi_{i,Nt-1} g_t \\ &= \Phi_{i,Nt} f_t + \Phi_{i,Nt-1} g_t + \Phi_{i,Nt} g_{t-1} - \Phi_{i,Nt} g_{t-1} \\ &= \Phi_{i,Nt} f_t + \Phi_{i,Nt} g_{t-1} + \Phi_{i,Nt-1} g_t - \Phi_{i,Nt} g_{t-1} \\ &= \tau \Phi_{i,Nt} - \Delta \Phi_{i,(N-1)t}, \\ &= \tau \Phi_{i,(N-1)t+t} - \Delta \Phi_{i,(N-2)t+t}, \end{aligned}$$

using $\tau = \tau_t = f_t + g_{t-1}$ from Lemma 1.5(i), and Lemma 2.3(ii) at the fourth line. For $j = t - 1$ the proof is similar using Lemma 2.3(i). ■

3 Matrix formulation of recurrences

Here we use matrices to give another proof that $\Phi_{i,Nt+j}$ satisfies the (τ, Δ) -recurrence, and prepare for the evaluation of G_N in Section 4.

Recall from Section 1 that $f_{i,j}$ and $g_{i,j}$ are the 2 fundamental solutions to Recurrence (4). Now define the matrix

$$X_{i,j} = \begin{pmatrix} g_{i,j-1} & f_{i,j-1} \\ g_{i,j} & f_{i,j} \end{pmatrix}.$$

Then the recurrences for $f_{i,j}$ and $g_{i,j}$ can be written as:

$$X_{i,j} = \begin{pmatrix} 0 & 1 \\ x_j & 1 \end{pmatrix} \begin{pmatrix} g_{i,j-2} & f_{i,j-2} \\ g_{i,j-1} & f_{i,j-1} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ x_j & 1 \end{pmatrix} X_{i,j-1}. \quad (15)$$

Consistent with (5) we have $X_{i,i-1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = I$, the 2×2 identity matrix.

Thus, for $j \geq i$, we have

$$X_{i,j} = \begin{pmatrix} 0 & 1 \\ x_j & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ x_{j-1} & 1 \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ x_i & 1 \end{pmatrix}. \quad (16)$$

Let $\mathbf{1} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ and $\mathbf{e} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$, and let $\langle \cdot, \cdot \rangle$ denote the usual inner product. Then for $j \geq i$, and using (6),

$$\phi_{i,j} = \langle X_{i,j} \mathbf{1}, \mathbf{e} \rangle. \quad (17)$$

As before if $i = 1$ we let $X_{1,j} = X_j$ and if $j = t$ we let $X = X_t$, in particular,

$$X = \begin{pmatrix} g_{t-1} & f_{t-1} \\ g_t & f_t \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ x_t & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ x_{t-1} & 1 \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ x_1 & 1 \end{pmatrix}. \quad (18)$$

For $N \geq 0$, from (10) we may also write

$$\begin{pmatrix} \Phi_{i,Nt+j-1} \\ \Phi_{i,Nt+j} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ x_j & 1 \end{pmatrix} \begin{pmatrix} \Phi_{i,Nt+j-2} \\ \Phi_{i,Nt+j-1} \end{pmatrix},$$

and then repeated use of (15) gives

$$\Phi_{i,Nt+j} = \langle X_j X^N X_{i,t} \mathbf{1}, \mathbf{e} \rangle. \quad (19)$$

Now using (16) and (18) we see that $X_j X^{-1} X_{i,t} = X_{i,j}$. So, using (17) and (9), we have

$$\langle X_j X^{-1} X_{i,t} \mathbf{1}, \mathbf{e} \rangle = \langle X_{i,j} \mathbf{1}, \mathbf{e} \rangle = \phi_{i,j} = \Phi_{i,-t+j},$$

thus (19) is true for $N = -1$ also.

Theorem 3.1 For $N \geq -1$ we have

$$\Phi_{i,Nt+j} = \langle X_j X^N X_{i,t} \mathbf{1}, \mathbf{e} \rangle.$$

■

From Lemma 1.5(i) and (16) we have the following forms for the trace and determinant of matrix $X_{i,j}$

$$\text{tr}(X_{i,j}) = \tau_{i,j} \quad \text{and} \quad \det(X_{i,j}) = (-1)^{j-i+1} x_i \cdots x_j.$$

In particular, for matrix X from (18), we have

$$\text{tr}(X) = \tau \quad \text{and} \quad \det(X) = \Delta. \quad (20)$$

Now let Z be any invertible 2×2 matrix with trace $\text{tr}(Z)$ and determinant $\det(Z)$, and let T denote transpose. Then the Cayley-Hamilton theorem says that $Z^2 = \text{tr}(Z)Z - \det(Z)I$, so $Z^N = \text{tr}(Z)Z^{N-1} - \det(Z)Z^{N-2}$, for $N \geq 1$. Let \mathbf{u} and $\mathbf{v} \in \mathbf{R}^2$ and, for $N \geq -1$, define $\Psi_N = \langle Z^N \mathbf{u}, \mathbf{v} \rangle$. Then

Lemma 3.2 For $N \geq 1$, Ψ_N satisfies the recurrence

$$\Psi_N = \text{tr}(Z) \Psi_{N-1} - \det(Z) \Psi_{N-2},$$

with initial conditions $\Psi_{-1} = \langle Z^{-1} \mathbf{u}, \mathbf{v} \rangle$ and $\Psi_0 = \langle Z^0 \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle$.

■

Now for $N \geq -1$,

$$\Phi_{i,Nt+j} = \langle X_j X^N X_{i,t} \mathbf{1}, \mathbf{e} \rangle = \langle X^N X_{i,t} \mathbf{1}, X_j^T \mathbf{e} \rangle.$$

So, for $N \geq 1$, Lemma 3.2 with $Z = X$, $\mathbf{u} = X_{i,t} \mathbf{1}$, and $\mathbf{v} = X_j^T \mathbf{e}$, and (20), gives,

$$\Phi_{i,Nt+j} = \tau \Phi_{i,(N-1)t+j} - \Delta \Phi_{i,(N-2)t+j}.$$

This is a second proof that $\Phi_{i,Nt+j}$ satisfies the (τ, Δ) -recurrence.

4 The Symmetric Representation, MacMahon's Master Theorem, three expressions for G_N

Consider polynomials in the variables u_1, \dots, u_d . We will work with the vector space whose basis elements are the homogeneous polynomials of degree N in these variables, *i.e.*, with

$$\{u_1^{n_1} \cdots u_d^{n_d} \mid n_1 + \cdots + n_d = N, \text{ each } n_\ell \geq 0\},$$

this vector space has dimension $\binom{N+d-1}{N}$.

The symmetric representation of a $d \times d$ matrix $A = (a_{\ell\ell'})$ is the action on polynomials induced by:

$$u_1^{n_1} \cdots u_d^{n_d} \rightarrow v_1^{n_1} \cdots v_d^{n_d},$$

where

$$v_\ell = \sum_{\ell'} a_{\ell\ell'} u_{\ell'}$$

or, more compactly, $v = Au$. That is, define the matrix element $\left\langle \begin{matrix} m_1, \dots, m_d \\ n_1, \dots, n_d \end{matrix} \right\rangle_A$ to be the coefficient of $u_1^{n_1} \cdots u_d^{n_d}$ in $v_1^{m_1} \cdots v_d^{m_d}$. Then, for a fixed (m_1, \dots, m_d) , we have

$$v_1^{m_1} \cdots v_d^{m_d} = \sum_{(n_1, \dots, n_d)} \left\langle \begin{matrix} m_1, \dots, m_d \\ n_1, \dots, n_d \end{matrix} \right\rangle_A u_1^{n_1} \cdots u_d^{n_d}. \quad (21)$$

Observe that the total degree $N = |n| = \sum n_\ell = |m| = \sum m_\ell$, *i.e.*, homogeneity of degree N is preserved. We use multi-indices: $m = (m_1, \dots, m_d)$ and $n = (n_1, \dots, n_d)$. Then, for a fixed m , (21) becomes

$$v^m = \sum_n \left\langle \begin{matrix} m \\ n \end{matrix} \right\rangle_A u^n.$$

Successive application of B then A shows that this is a homomorphism of the multiplicative semi-group of square $d \times d$ matrices into the multiplicative semi-group of square $\binom{N+d-1}{N} \times \binom{N+d-1}{N}$ matrices.

Proposition 4.1 *Matrix elements satisfy the homomorphism property*

$$\left\langle \begin{matrix} m \\ n \end{matrix} \right\rangle_{AB} = \sum_k \left\langle \begin{matrix} m \\ k \end{matrix} \right\rangle_A \left\langle \begin{matrix} k \\ n \end{matrix} \right\rangle_B.$$

Proof. Let $v = (AB)u$ and $w = Bu$. Then,

$$\begin{aligned} v^m &= \sum_n \left\langle \begin{matrix} m \\ n \end{matrix} \right\rangle_{AB} u^n \\ &= \sum_k \left\langle \begin{matrix} m \\ k \end{matrix} \right\rangle_A w^k \\ &= \sum_n \sum_k \left\langle \begin{matrix} m \\ k \end{matrix} \right\rangle_A \left\langle \begin{matrix} k \\ n \end{matrix} \right\rangle_B u^n. \end{aligned}$$

■

Definition Fix the degree $N = \sum n_\ell = \sum m_\ell$. Define $\text{tr}_{\text{Sym}}^N(A)$, the *symmetric trace* of A in degree N , as the sum of the diagonal elements $\left\langle \begin{matrix} m \\ n \end{matrix} \right\rangle_A$, i.e.,

$$\text{tr}_{\text{Sym}}^N(A) = \sum_m \left\langle \begin{matrix} m \\ m \end{matrix} \right\rangle_A.$$

Equality such as $\text{tr}_{\text{Sym}}(A) = \text{tr}_{\text{Sym}}(B)$ means that the symmetric traces are equal in every degree $N \geq 0$.

Remark The action defined here on polynomials is equivalent to the action on symmetric tensor powers, see Fulton and Harris [2], pp. 472-5.

Now it is straightforward to see directly (cf. the diagonal case shown in the Corollary below) that if A is upper-triangular, with eigenvalues $\lambda_1, \dots, \lambda_d$, then $\text{tr}_{\text{Sym}}^N(A) = h_N(\lambda_1, \dots, \lambda_d)$, the N^{th} homogeneous symmetric function. The homomorphism property, Proposition 4.1, shows that $\text{tr}_{\text{Sym}}^N(AB) = \text{tr}_{\text{Sym}}^N(BA)$, and that similar matrices have the same trace. Again by the homomorphism property, if two $d \times d$ matrices are similar, $A = MBM^{-1}$, then that relation extends to their respective symmetric representations in every degree. Recall that any matrix is similar to an upper-triangular one with the same eigenvalues. Thus,

Theorem 4.2 Symmetric Trace Theorem (see pp. 51-2 of Springer [5]).

We have

$$\frac{1}{\det(I - cA)} = \sum_{N=0}^{\infty} c^N \text{tr}_{\text{Sym}}^N(A).$$

Proof. With λ_ℓ denoting the eigenvalues of A ,

$$\frac{1}{\det(I - cA)} = \prod_{\ell} \frac{1}{1 - c\lambda_\ell}$$

$$\begin{aligned}
&= \sum_{N=0}^{\infty} c^N h_N(\lambda_1, \dots, \lambda_d) \\
&= \sum_{N=0}^{\infty} c^N \text{tr}_{\text{Sym}}^N(A).
\end{aligned}$$

■

As a Corollary we have MacMahon's Master Theorem, which we express in the above terminology.

Corollary 4.3 MacMahon's Master Theorem.

The diagonal matrix element $\left\langle \begin{smallmatrix} m \\ m \end{smallmatrix} \right\rangle_A$ is the coefficient of $u^m = u_1^{m_1} \dots u_d^{m_d}$ in the expansion of $\det(I - UA)^{-1}$ where $U = \text{diag}(u_1, \dots, u_d)$ is the diagonal matrix with entries u_1, \dots, u_d on the diagonal.

Proof. From Theorem 4.2, with $c = 1$, we want to calculate the symmetric trace of UA . By the homomorphism property,

$$\begin{aligned}
\text{tr}_{\text{Sym}}^N(UA) &= \sum_m \left\langle \begin{smallmatrix} m \\ m \end{smallmatrix} \right\rangle_{UA} \\
&= \sum_m \sum_k \left\langle \begin{smallmatrix} m \\ k \end{smallmatrix} \right\rangle_U \left\langle \begin{smallmatrix} k \\ m \end{smallmatrix} \right\rangle_A.
\end{aligned}$$

Now, with $v = Uw$ and $v_\ell = u_\ell w_\ell$, then

$$v^m = (u_1 w_1)^{m_1} \dots (u_d w_d)^{m_d} = u^m w^m = \sum_k \left\langle \begin{smallmatrix} m \\ k \end{smallmatrix} \right\rangle_U w^k,$$

i.e.,

$$\left\langle \begin{smallmatrix} m \\ k \end{smallmatrix} \right\rangle_U = u_1^{m_1} \dots u_d^{m_d} \delta_{k_1 m_1} \dots \delta_{k_d m_d}$$

so that

$$\text{tr}_{\text{Sym}}^N(UA) = \sum_m \left\langle \begin{smallmatrix} m \\ m \end{smallmatrix} \right\rangle_A u^m.$$

■

Now we restrict ourselves to $d = 2$, and return to the (τ, Δ) -recurrence.

Recall, from (18), the 2×2 matrix

$$\begin{aligned} X &= \begin{pmatrix} 0 & 1 \\ x_t & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ x_{t-1} & 1 \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ x_1 & 1 \end{pmatrix} \\ &= \xi_t \xi_{t-1} \cdots \xi_1, \end{aligned}$$

where $\xi_s = \begin{pmatrix} 0 & 1 \\ x_s & 1 \end{pmatrix}$ for $1 \leq s \leq t$. Let us modify ξ_s slightly by defining $\alpha_s = \begin{pmatrix} 0 & 1 \\ x_s & a_s \end{pmatrix}$ for $1 \leq s \leq t$, and calling

$$\begin{aligned} \bar{X} &= \begin{pmatrix} 0 & 1 \\ x_t & a_t \end{pmatrix} \begin{pmatrix} 0 & 1 \\ x_{t-1} & a_{t-1} \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ x_1 & a_1 \end{pmatrix} \\ &= \alpha_t \alpha_{t-1} \cdots \alpha_1. \end{aligned}$$

Let

$$\text{tr}(\bar{X}) = \bar{\tau} \quad \text{and} \quad \det(\bar{X}) = \bar{\Delta},$$

and let \bar{G}_N be the first fundamental solution to the $(\bar{\tau}, \bar{\Delta})$ -recurrence:

$$\Theta_N = \bar{\tau} \Theta_{N-1} - \bar{\Delta} \Theta_{N-2}. \quad (22)$$

Then

$$\begin{aligned} \sum_{N=0}^{\infty} c^N \bar{G}_N &= \frac{1}{1 - \bar{\tau}c + \bar{\Delta}c^2} \\ &= \frac{1}{\det(I - c\bar{X})} \\ &= \sum_{N=0}^{\infty} c^N \text{tr}_{\text{Sym}}^N(\bar{X}). \end{aligned}$$

So

$$\bar{G}_N = \text{tr}_{\text{Sym}}^N(\bar{X}) = \sum_m \langle m \rangle_{\bar{X}} = \sum_m \langle m \rangle_{\alpha_t \alpha_{t-1} \cdots \alpha_1}.$$

We need to calculate the symmetric trace of \bar{X} and so identify \bar{G}_N . By the homomorphism property, we need only find the matrix elements for each matrix α_s , multiply together and take the trace.

For $\alpha_s = \begin{pmatrix} 0 & 1 \\ x_s & a_s \end{pmatrix}$ the mapping induced on polynomials is

$$v_1 = u_2, \quad v_2 = x_s u_1 + a_s u_2. \quad (23)$$

For any integer $N \geq 0$, the expansion of $v_1^m v_2^{N-m}$ in powers of u_1 and u_2 is of the form

$$v_1^m v_2^{N-m} = \sum_n \langle m \rangle_n \alpha_s u_1^n u_2^{N-n}, \quad (24)$$

with the notation for the matrix elements abbreviated accordingly. From (23) and (24), the binomial theorem yields

$$\langle m \rangle_n \alpha_s = \binom{N-m}{n} x_s^n a_s^{N-m-n}.$$

For example, when $t = 3$, the product $\bar{X} = \alpha_3 \alpha_2 \alpha_1$ gives the matrix elements, for homogeneity of degree N ,

$$\begin{aligned} \langle m \rangle_n \bar{X} &= \sum_{(k_2, k_3)} \langle m \rangle_{k_3} \alpha_3 \langle k_3 \rangle_{k_2} \alpha_2 \langle k_2 \rangle_n \alpha_1 \\ &= \sum_{(k_2, k_3)} \binom{N-m}{k_3} \binom{N-k_3}{k_2} \binom{N-k_2}{n} x_1^n x_2^{k_2} x_3^{k_3} a_1^{N-k_2-n} a_2^{N-k_3-k_2} a_3^{N-k_3-m}. \end{aligned}$$

Thus, the symmetric trace $\text{tr}_{\text{Sym}}^N(\bar{X}) = \sum_m \langle m \rangle_m \bar{X}$ is

$$\sum_{(k_1, k_2, k_3)} \binom{N-k_2}{k_1} \binom{N-k_3}{k_2} \binom{N-k_1}{k_3} x_1^{k_1} x_2^{k_2} x_3^{k_3} a_1^{N-k_1-k_2} a_2^{N-k_2-k_3} a_3^{N-k_3-k_1},$$

a *cyclic binomial*. In general, for a product of arbitrary length, the symmetric trace is given by the corresponding cyclic binomial.

Recall the recurrence

$$S_N(x) = 2x S_{N-1}(x) - S_{N-2}(x), \quad (25)$$

for $N \geq 1$. The Chebyshev polynomials of the first kind, $T_N = T_N(x)$, are solutions of this recurrence with initial conditions $T_{-1} = x$ and $T_0 = 1$, and the Chebyshev polynomials of the second kind, $U_N = U_N(x)$, are solutions with $U_{-1} = 0$ and $U_0 = 1$.

Combining these observations yields the main identities:

Theorem 4.4 Let $\overline{X} = \alpha_t \alpha_{t-1} \cdots \alpha_1$, with $\alpha_s = \begin{pmatrix} 0 & 1 \\ x_s & a_s \end{pmatrix}$ for $1 \leq s \leq t$, and let $\overline{\tau} = \text{tr}(\overline{X})$ and $\overline{\Delta} = \det(\overline{X})$. Let \overline{G}_N denote the first fundamental solution to the $(\overline{\tau}, \overline{\Delta})$ -recurrence (22).

Then we have the **cyclic binomial identity**

$$\begin{aligned} \overline{G}_N &= \sum_{(k_1, \dots, k_t)} \binom{N-k_2}{k_1} \binom{N-k_3}{k_2} \cdots \binom{N-k_1}{k_t} x_1^{k_1} x_2^{k_2} \cdots x_t^{k_t} a_1^{N-k_1-k_2} a_2^{N-k_2-k_3} \cdots a_t^{N-k_t-k_1} \\ &= \overline{\Delta}^{N/2} U_N \left(\frac{\overline{\tau}}{2\sqrt{\overline{\Delta}}} \right) \\ &= \sum_{k=0}^{\lfloor N/2 \rfloor} \binom{N-k}{k} \overline{\tau}^{N-2k} (-\overline{\Delta})^k, \end{aligned}$$

where U_N denotes the Chebyshev polynomial of the second kind.

Proof. The first equality follows by computing the symmetric trace for arbitrary t as indicated above. The second follows by induction on N using initial conditions $\overline{G}_{-1} = 0$ and $\overline{G}_0 = 1$, the $(\overline{\tau}, \overline{\Delta})$ -recurrence (22) and the Chebyshev recurrence (25). The third follows from the second by the Symmetric Trace Theorem applied to $\overline{X} = \begin{pmatrix} 0 & 1 \\ -\overline{\Delta} & \overline{\tau} \end{pmatrix}$, the shift matrix for the $(\overline{\tau}, \overline{\Delta})$ -recurrence. \blacksquare

Note that $G_{-1} = 0$ and $G_0 = 1$, so $G_1 = \tau$ using the (τ, Δ) -recurrence. This also follows directly from the condition $k_{s-1} + k_s \leq 1$ for non-zero terms in the cyclic binomial summation above. Note also that setting all $a_s = 1$ above gives explicit expressions for G_N .

Example 5 Here $N = 2$ and $t = 3$. Let $A^{\text{Sym}(N)}$ denote the symmetric representation in degree N of the matrix A . From the above we have

$$\begin{aligned} G_2 &= \sum_{(k_1, k_2, k_3)} \binom{2-k_2}{k_1} \binom{2-k_3}{k_2} \binom{2-k_1}{k_3} x_1^{k_1} x_2^{k_2} x_3^{k_3} \\ &= 1 + 2x_1 + 2x_2 + 2x_3 + x_1^2 + 2x_1x_2 + 2x_1x_3 + x_2^2 + 2x_2x_3 + x_3^2 + x_1x_2x_3. \end{aligned}$$

Also $d = 2$, so $\binom{N+d-1}{N} = 3$, and $\xi_s = \begin{pmatrix} 0 & 1 \\ x_s & 1 \end{pmatrix}$ for $1 \leq i \leq 3$, thus

$$X = \xi_3 \xi_2 \xi_1 = \begin{pmatrix} x_1 & x_2 + 1 \\ x_1 x_3 + x_1 & x_2 + x_3 + 1 \end{pmatrix}.$$

Now $\xi_s^{Sym(2)} = \begin{pmatrix} 0 & 0 & 1 \\ 0 & x_s & 1 \\ x_s^2 & 2x_s & 1 \end{pmatrix}$ for $1 \leq s \leq 3$, and so

$$\begin{aligned} X^{Sym(2)} &= \xi_3^{Sym(2)} \xi_2^{Sym(2)} \xi_1^{Sym(2)} \\ &= \begin{pmatrix} x_1^2 & 2x_1x_2 + 2x_1 & x_2^2 + 2x_2 + 1 \\ x_1^2x_3 + x_1^2 & x_1x_2x_3 + 2x_1x_2 + 2x_1x_3 + 2x_1 & x_2^2 + x_2x_3 + 2x_2 + x_3 + 1 \\ x_1^2x_3^2 + 2x_1^2x_3 + x_1^2 & 2x_1x_2x_3 + 2x_1x_3^2 + 2x_1x_2 + 4x_1x_3 + 2x_1 & x_3^2 + 2x_2x_3 + x_2^2 + 2x_2 + 2x_3 + 1 \end{pmatrix}. \end{aligned}$$

We check that $G_2 = \text{tr}(X^{Sym(2)})$, as indicated above.

We now give an expression for G_N as a quotient of two matching polynomials; this requires (29) from the next section.

Theorem 4.5 *For $N \geq 0$ we have*

$$G_N = \frac{\Phi_{1,Nt-2}}{\phi_{t-2}}.$$

Proof. Equation (29) is

$$\Phi_{i,Nt+j} = \Phi_{i,j} G_N - \Delta \phi_{i,j} G_{N-1}, \quad (26)$$

and from Example 2 we have $\phi_{i,i-3} = 0$. So (26) with $j = i - 3$ gives

$$G_N = \frac{\Phi_{i,Nt+i-3}}{\Phi_{i,i-3}} = \frac{\Phi_{1,Nt-2}}{\phi_{t-2}}, \quad (27)$$

the second equality comes from putting $i = 1$ in the first and then using (9) in the denominator. \blacksquare

Finally, consider the Fibonacci sequence $\{F_m \mid m \geq 1\} = \{1, 1, 2, 3, 5, 8, 13, 21, \dots\}$. It is straightforward to show that the number of matchings in the path P_m with $m-1$ edges is F_{m+1} . Now $\Phi_{1, Nt-2}$ is the matching polynomial of the path $P(1, Nt-2)$ which has $(N+1)t-2$ edges and so has $F_{(N+1)t}$ matchings. Similarly, the path whose matching polynomial is ϕ_{t-2} has F_t matchings. Now, evaluating (27) above with $N = N-1$ and $x_s = 1$ for all $1 \leq s \leq t$, gives $F_t | F_{Nt}$, a well-known result on Fibonacci numbers, see pp. 148-9, Hardy and Wright [4]. Furthermore, we have

$$\frac{F_{(N+1)t}}{F_t} = \sum_{(k_1, \dots, k_t)} \binom{N-k_2}{k_1} \binom{N-k_3}{k_2} \cdots \binom{N-k_t}{k_{t-1}}.$$

5 Examples: Paths, Cycles, Trees

In this Section we express the matching polynomial of some well-known graphs in terms of the fundamental solutions to the (τ, Δ) -recurrence (12).

G_N is the first fundamental solution to the (τ, Δ) -recurrence, so the initial values for G_N are

$$G_{-2} = \frac{-1}{\Delta}, \quad G_{-1} = 0, \quad G_0 = 1, \quad (\text{and } G_1 = \tau). \quad (28)$$

The second fundamental solution is $-\Delta G_{N-1}$.

5.1 Paths

$\Phi_{i, Nt+j}$ satisfies the (τ, Δ) -recurrence whose fundamental solutions are G_N and $-\Delta G_{N-1}$, thus $\Phi_{i, Nt+j} = a G_N + b(-\Delta G_{N-1})$ for some a and b . The initial conditions for $\Phi_{i, Nt+j}$ from (9) and for G_N from (28) give $a = \Phi_{i,j}$ and $b = \Phi_{i, -t+j} = \phi_{i,j}$. Hence for $N \geq -1$,

$$\Phi_{i, Nt+j} = \Phi_{i,j} G_N - \Delta \phi_{i,j} G_{N-1}. \quad (29)$$

Example 6 Here $i = 2$ and $t = 3$,

$$\begin{aligned} N = -1 & \quad \phi_{2,2} = 1 + x_2, \\ N = 0 & \quad \phi_{2,3} = 1 + x_2 + x_3, \\ N = 0 & \quad \Phi_{2,1} = 1 + x_1 + x_2 + x_3 + x_1 x_2, \\ N = 0 & \quad \Phi_{2,2} = 1 + x_1 + 2x_2 + x_3 + x_1 x_2 + x_2^2 + x_2 x_3, \\ N = 1 & \quad \Phi_{2,3} = 1 + x_1 + 2x_2 + 2x_3 + x_1 x_2 + x_1 x_3 + x_2^2 + 2x_2 x_3 + x_3^2 + x_1 x_2 x_3. \end{aligned}$$


For $N \geq 1$ let $P_{Nt+j+1} = P(1, (N-1)t+j)$ be the path with $Nt+j+1$ vertices and $Nt+j$ edges, cyclically labelled starting with label x_1 . Let $\mathcal{P}_{Nt+j+1}(\mathbf{x}) = \mathcal{P}_{Nt+j+1} = \Phi_{1, (N-1)t+j}$ be its matching polynomial. With this notation any subscript on a P , \mathcal{P} , C , or \mathcal{C} refers to the number of vertices in the appropriate graph.

Theorem 5.1 *For any $N \geq 1$ we have*


$$\begin{aligned} (i) \quad \mathcal{P}_{Nt+j+1} &= \Phi_{1,j} G_{N-1} - \Delta \phi_j G_{N-2}, \\ (ii) \quad \mathcal{P}_{Nt+1} &= G_N + (\phi - \tau) G_{N-1}. \end{aligned}$$

Proof. The proof of (i) is clear using (29) with $i = 1$ and $N = N - 1$. So (i) with $j = 0$ gives $\mathcal{P}_{Nt+1} = \Phi_{1,0} G_{N-1} - \Delta \phi_0 G_{N-2}$. But $\Phi_{1,0} = \Phi_{1,-t+t} = \phi_{1,t} = \phi_t = \phi$ and $\phi_0 = \phi_{1,0} = 1$, and then using the (τ, Δ) -recurrence for G_N gives (ii). \blacksquare

Example 7 Here $t = 3$,



$$\mathcal{P}_{1.3+2+1} = 1 + 2x_1 + 2x_2 + x_3 + x_1^2 + 2x_1x_2 + x_1x_3 + x_2^2 + x_2x_3 + x_1x_2x_3.$$



$$\mathcal{P}_{2.3+1} = 1 + 2x_1 + 2x_2 + 2x_3 + x_1^2 + 2x_1x_2 + 3x_1x_3 + x_2^2 + 2x_2x_3 + x_3^2 + x_1^2x_3 + 2x_1x_2x_3 + x_1x_3^2.$$

5.2 Cycles

Now we identify the first and the last vertices of the path $P(i, Nt + j)$ to form the cyclically labelled cycle $C(i, Nt + j)$ with matching polynomial $\Gamma_{i, Nt+j}(\mathbf{x}) = \Gamma_{i, Nt+j}$.

By decomposing $\Gamma_{i, Nt+j}$ at the ‘first’ edge labelled x_i we see that, *cf.* (29),

$$\begin{aligned}
\Gamma_{i, Nt+j} &= \Phi_{i+1, Nt+j} + x_i \Phi_{i+2, Nt+j-1}, \\
&= \Phi_{i+1, j} G_N - \Delta \phi_{i+1, j} G_{N-1} + x_i \{ \Phi_{i+2, j-1} G_N - \Delta \phi_{i+2, j-1} G_{N-1} \}, \\
&= \{ \Phi_{i+1, j} + x_i \Phi_{i+2, j-1} \} G_N - \Delta \{ \phi_{i+1, j} + x_i \phi_{i+2, j-1} \} G_{N-1}, \\
&= \Gamma_{i, j} G_N - \Delta \tau_{i, j} G_{N-1}, \tag{30}
\end{aligned}$$

using (29) at the second line, and decomposing $\Gamma_{i, j}$ and $\tau_{i, j}$ at the first edge x_i at the fourth line. Also, defining $\Gamma_{i, -t+j} = \tau_{i, j}$ ensures that (30) is true for all $N \geq -1$.

Example 8 Here $i = 2$ and $t = 3$ again,

$$\begin{aligned}
N = -1 & \quad \tau_{2, 2} = 1, \\
N = 0 & \quad \tau_{2, 3} = 1 + x_2 + x_3, \\
N = 0 & \quad \Gamma_{2, 1} = 1 + x_1 + x_2 + x_3, \\
N = 0 & \quad \Gamma_{2, 2} = 1 + x_1 + 2x_2 + x_3 + x_1x_2 + x_2x_3, \\
N = 1 & \quad \Gamma_{2, 3} = 1 + x_1 + 2x_2 + 2x_3 + x_1x_2 + x_1x_3 + x_2^2 + x_2x_3 + x_3^2.
\end{aligned}$$


Let $C_{Nt+j} = C(1, (N-1)t + j)$ be the cycle with $Nt + j$ vertices and $Nt + j$ edges in which labelling has started with x_1 , and let $\mathcal{C}_{Nt+j}(\mathbf{x}) = \mathcal{C}_{Nt+j} = \Gamma_{1, (N-1)t+j}$ be its matching polynomial. Compare with Theorem 5.1,

Theorem 5.2 *For any $N \geq 1$ we have*


$$\begin{aligned}
(i) \quad \mathcal{C}_{Nt+j} &= \Gamma_{1, j} G_{N-1} - \Delta \tau_j G_{N-2}, \\
(ii) \quad \mathcal{C}_{Nt} &= G_N - \Delta G_{N-2}.
\end{aligned}$$

Proof. The proof of (i) is clear from (30). Part (i) with $j = 0$ gives (ii), using $\Gamma_{1, 0} = \tau$, and $\tau_0 = 2$ from (8). \blacksquare

Example 9 Here $t = 3$ again, the cycle starts at the large vertex and proceeds clockwise,



$$C_{1,3+2} = 1 + 2x_1 + 2x_2 + x_3 + x_1^2 + x_1x_2 + x_1x_3 + x_2^2 + x_2x_3$$



$$C_{2,3} = 1 + 2x_1 + 2x_2 + 2x_3 + x_1^2 + 2x_1x_2 + 2x_1x_3 + x_2^2 + 2x_2x_3 + x_3^2 + 2x_1x_2x_3.$$

For a fixed $t \geq 1$ write $\widehat{\mathcal{P}}_N = \mathcal{P}_{Nt+1}$ and $\widehat{\mathcal{C}}_N = \mathcal{C}_{Nt}$. We now express G_N , $\widehat{\mathcal{P}}_N$, and $\widehat{\mathcal{C}}_N$ in terms of Chebyshev polynomials.

It is well-known that, in one variable x , the matching polynomial of the path P_{2m} is related to U_{2m} as follows

$$\mathcal{M}(P_{2m}, x) = (-1)^m x^m U_{2m} \left(\frac{i}{2\sqrt{x}} \right),$$

and, for P_{2m-1} we have

$$\mathcal{M}(P_{2m-1}, x) = (-1)^m x^m \left[U_{2m} \left(\frac{i}{2\sqrt{x}} \right) + U_{2m-2} \left(\frac{i}{2\sqrt{x}} \right) \right],$$

where $i = \sqrt{-1}$. Also, for the matching polynomials $\mathcal{M}(C_{2m})$ and $\mathcal{M}(C_{2m-1})$ of the cycles C_{2m} and C_{2m-1} there are similar formulas but with a factor of 2 on the right-hand side where U is replaced by T . See Theorem 3 of Godsil and Gutman [3], and Theorems 9 and 11 of Farrell [1].

Now Theorem 4.4 modified for G_N gives

$$G_N = \Delta^{N/2} U_N \left(\frac{\tau}{2\sqrt{\Delta}} \right). \quad (31)$$

Formulas for $\widehat{\mathcal{P}}_N$ and $\widehat{\mathcal{C}}_N$ in terms of U_N and T_N are given below, where the variable t is suppressed.

Theorem 5.3 *For any $N \geq 1$ we have*

$$(i) \quad \widehat{\mathcal{P}}_N = \Delta^{N/2} \left\{ U_N \left(\frac{\tau}{2\sqrt{\Delta}} \right) + \left(\frac{\phi - \tau}{\sqrt{\Delta}} \right) U_{N-1} \left(\frac{\tau}{2\sqrt{\Delta}} \right) \right\},$$

$$(ii) \quad \widehat{\mathcal{C}}_N = 2\Delta^{N/2} T_N \left(\frac{\tau}{2\sqrt{\Delta}} \right).$$

Proof. (i) This follows from Theorem 5.1(ii) and (31).

(ii) From Theorem 5.2(ii) we have $\widehat{\mathcal{C}}_N = G_N - \Delta G_{N-2}$, and now the well-known relation $2T_N = U_N - U_{N-2}$ between the two types of Chebyshev polynomials and (31) gives the result. \blacksquare

Expressions for G_N , $\widehat{\mathcal{P}}_N$, and $\widehat{\mathcal{C}}_N$ for $N = 0, 1, 2, 3$, and 4 are given below

$$\begin{array}{lll} G_0 = 1 & \widehat{\mathcal{P}}_0 = 1 & \widehat{\mathcal{C}}_0 = 2 \\ G_1 = \tau & \widehat{\mathcal{P}}_1 = \phi & \widehat{\mathcal{C}}_1 = \tau \\ G_2 = \tau^2 - \Delta & \widehat{\mathcal{P}}_2 = \phi\tau - \Delta & \widehat{\mathcal{C}}_2 = \tau^2 - 2\Delta \\ G_3 = \tau^3 - 2\tau\Delta & \widehat{\mathcal{P}}_3 = \phi\tau^2 - \phi\Delta - \tau\Delta & \widehat{\mathcal{C}}_3 = \tau^3 - 3\tau\Delta \\ G_4 = \tau^4 - 3\tau^2\Delta + \Delta^2 & \widehat{\mathcal{P}}_4 = \phi\tau^3 - 2\phi\tau\Delta - \tau^2\Delta + \Delta^2 & \widehat{\mathcal{C}}_4 = \tau^4 - 4\tau^2\Delta + 2\Delta^2. \end{array}$$

5.3 Trees

Here we consider cyclically labelled trees.

First let us extend the definition of a cyclically labelled path to include the path of Fig. 1, and the graph P_1 with one vertex and no edges.

A tree is a connected simple graph with no cycles, and a rooted tree is a tree in which some vertex of degree 1 has been specified to be the root, r . Given any rooted tree, let us label its edges by first labelling the edge incident to r with x_i . Then label all edges incident to this edge with x_{i+1} , then label all edges incident to these edges with x_{i+2} , and so on until label x_t has been used. Then label with the ordered set $\{x_1, \dots, x_t\}$ in a similar

manner to before, repeating cyclically until all edges have been labelled,..., and so on. Let T denote such a cyclically labelled tree, see Fig. 5 for an example with $i = 2$ and $t = 3$.




Fig. 5: A cyclically labelled tree with $i = 2$ and $t = 3$.

We may draw any such T with r as the leftmost vertex. Then we place the other vertices of T from ‘left to right’ according to their distance from r , *i.e.*, if a vertex v_1 is at distance d_1 from r and vertex v_2 is at distance d_2 from r where $d_2 > d_1$, then v_2 is placed to the right of v_1 .

Paths in T are of two types: (I) A path that always moves from left to right (a path that always moves from right to left can be thought of one that always moves from left to right): such a path is clearly cyclically labelled; or (II) a path that moves first from right to left and then from left to right; such a path must pass through at least one vertex of degree ≥ 3 , *i.e.*, a vertex where T ‘branches’.

Let V denote the set of vertices of degree ≥ 3 in T , and let $v \in V$ be arbitrary of degree $\deg(v)$. Vertex v has 1 edge to its left and $\deg(v) - 1 \geq 2$ edges to its right. Let H_v be the subgraph of T that consists of the ‘last’ $\deg(v) - 2 \geq 1$ edges as we rotate clockwise around v . Thus H_v is the star $K_{1, \deg(v)-2}$ centered at v . Set $H = \cup_{v \in V} H_v$.

Lemma 5.4 *The forest $T - H$ is a union of cyclically labelled paths.*

Proof. We show that $T - H$ does not contain a path of type (II). Suppose it does contain a path of type (II), then this path must pass through some

vertex $v \in V$. So 2 edges incident to v and to the right of v lie in this path and so lie in $T - H$, a contradiction because $T - H$ contains only 1 edge incident to v and to the right of v . Thus $T - H$ is a union of paths of type (I), each of which is a cyclically labelled path. ■

Thus $T - H$ is a union of cyclically labelled paths, and so $T - H - \overline{M}_H$ is also, for every matching M_H of H . We know the matching polynomial of any cyclically labelled path, so we can decompose the matching polynomial of T , $\mathcal{M}(T, \mathbf{x})$, at H , according to Theorem 1.1,

$$\mathcal{M}(T, \mathbf{x}) = \sum_{M_H} M_H(\mathbf{x}) \mathcal{M}(T - H - \overline{M}_H, \mathbf{x}),$$

where the summation is over every matching M_H of H .

Example 10 See Fig. 5.



H has 6 matchings with weights: $1, x_1, x_1, x_3, x_1x_3$, and x_1x_3 . Thus there are 6 terms in the decomposition, and $\mathcal{M}(T, \mathbf{x})$ is the sum of the following 6 terms:

$$\begin{aligned} & 1 \cdot \phi_1 \phi_{2,3} \Phi_{2,1} + x_1 \cdot \phi_1 \phi_{2,2} \phi_{2,3} + x_1 \cdot \phi_1 \phi_{2,2} \phi_{3,3} + x_3 \cdot \phi_1 \phi_{2,3} + x_1 x_3 \cdot \phi_{2,3} + x_1 x_3 \cdot \phi_{3,3} \\ & = 1 + 4x_1 + 2x_2 + 3x_3 + 3x_1^2 + 7x_1x_2 + 8x_1x_3 + x_2^2 + 3x_2x_3 + 2x_3^2 \\ & \quad + 5x_1^2x_2 + 3x_1^2x_3 + 3x_1x_2^2 + 7x_1x_2x_3 + 4x_1x_3^2 + 2x_1^2x_2^2 + 3x_1^2x_2x_3. \end{aligned}$$

Acknowledgement: We thank the referees for some helpful comments.

References

- [1] E.J. Farrell, *An Introduction to Matching Polynomials*, Jour. Comb. Theory, Series B, **27**, pp. 75–86, 1979.
- [2] W. Fulton, J. Harris, *Representation Theory, A First Course*, Graduate Texts in Mathematics, **129**, Springer-Verlag, 1991.

- [3] C.D. Godsil, I. Gutman, *On the Theory of the Matching Polynomial*, Jour. Graph. Theory, **5**, pp. 137–144, 1981.
- [4] G.H. Hardy, E.M. Wright, *An Introduction to the Theory of Numbers*, 5th Edition, Oxford University Press, 1979.
- [5] T.A. Springer, *Invariant Theory*, Lecture Notes in Mathematics, **585**, Springer-Verlag, 1977.