



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

**Parallel detection of bacteria with culture methods and 16S rRNA gene analysis:  
Preliminary results from a prospective cohort study of prosthetic joint infections**

Larsen, Lone Heimann; Xu, Yijuan; Thomsen, Trine Rolighed; Schønheyder, Henrik Carl

*Publication date:*  
2012

*Document Version*  
Early version, also known as pre-print

[Link to publication from Aalborg University](#)

*Citation for published version (APA):*

Larsen, L. H., Xu, Y., Thomsen, T. R., & Schønheyder, H. C. (2012). *Parallel detection of bacteria with culture methods and 16S rRNA gene analysis: Preliminary results from a prospective cohort study of prosthetic joint infections*. Poster presented at Symposium of The Danish Microbiological Society, Copenhagen, Denmark.

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

**Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

# Parallel detection of bacteria with culture methods and 16S *rRNA* gene analysis: Preliminary results from a prospective cohort study of prosthetic joint infections

Lone Heimann Larsen <sup>1,2</sup>, Yijuan Xu <sup>2</sup>, Trine Rolighed Thomsen <sup>2,3</sup>, Henrik C. Schönheyder <sup>1</sup> and the PRIS Study Group

<sup>1</sup>Department of Clinical Microbiology, Aalborg Hospital, Aarhus University Hospital

<sup>2</sup>Section for Biotechnology, Institute for Chemistry and Biotechnology, Aalborg University

<sup>3</sup>Section for Medical Biotechnology, Danish Technology Institute, Aarhus

## Introduction

'Artificial joints' have become indispensable treatment for osteoarthritis. Approximately 15,000 primary arthroplastic procedures and 2,000 revisions are performed each year in Denmark (1). The number of patients with prosthetic implants will continue to grow mainly due to an aging population.

However, the implants can lead to serious complications, one of which is bacterial infection. The risk is estimated to be 0.5-2% for primary arthroplasties and even higher for revision surgeries. The infections are difficult to diagnose due to vague symptoms and a modest inflammatory response. Routine culture-based methods failure to demonstrate a bacterial agent in a significant proportion of cases. The failure may be due


to inappropriate culture conditions, prior antibiotic treatment, fastidious or atypical organisms, and the biofilm mode of growth.

The successful diagnosis and treatment of prosthetic joint infections (PJIs) require a close cooperation between surgeons, radiologists, specialists in nuclear medicine, clinical microbiologists and molecular biologists. At Aalborg Hospital we have established a unique diagnostic algorithm within the framework of the PRIS Innovation project. In this presentation the objective was to compare positivity rates of extended cultures for the presence of bacteria with the positive rates of detection of 16S *rRNA* genes in parallel samples. We report preliminary results for the first 65 patients recruited consecutively December 2011 through June 2012.

## M&M

The study comprised patients seeking medical attention with a painful hip or knee prosthesis. Twenty five had PJI according to the physician's judgement; the remaining 40 patients had an aseptic loosening (AL). Fifty four underwent surgical revision, seven had a joint puncture, and four had two procedures (joint puncture+revision or revision+revision).

The prosthetic components were processed by submerging into isotonic buffer (1XPBS) for sonication (40 kHz) and aliquots of the buffer were centrifuged to concentrate the bacterial content and used for further processing (2). Bacteriological cultures were cultivated on agar plates and in broth medium under aerobic and anaerobic conditions for 14 days. For full length 16S *rRNA* gene analysis DNA were extracted and screened by endpoint PCR.



**Flow diagram:** Joint puncture; 3 samples were obtained, 2 for culture (1 aerobic blood culture vial and 1 for bacterial culturing) and 1 for 16S *rRNA* gene analysis. During revision up to 15 samples were obtained; 5 for 16S *rRNA* analysis, 5 for bacterial culturing and five for other purposes. Prior to this study, bone biopsies, prosthetic components, and biofilm swabs were not available for microbiological diagnosis of PJI.

## Results

**Surgical revisions (table).** In the PJI group (n=18), a total of 74 samples were positive by culture and 58 by 16S *rRNA* gene analysis. We observed a high level of concordance between culture-based and molecular methods with specimens of sonication fluid and synovial fluid whereas results were less consistent with synovial biopsies, bone biopsies and biofilm swabs (table 1). The highest numbers of positive specimens were

obtained with sonication fluid, followed by synovial fluid and tissue biopsies. In the AL group of patients (n= 40) one sample was positive only by culture and 14 by 16S *rRNA* gene analysis.

**Joint puncture.** One sample of synovial fluid for one patient was positive both by culture and 16S *rRNA* gene analysis, and samples from six other patients were negative by both culture and 16S *rRNA* gene analysis.

## Discussion

Prosthesis components show promise as a new sample type for the diagnosis of PJI. Using the established sonication procedure (2,3,4) and optimized culture methods we obtained agreement with 16S *rRNA* gene analysis in 57 procedures in 56 patients. Sonication has also previously been shown to be valuable for culturing of prosthetic components and to increase rates of bacterial growth (5).

At the current time we interpret the positive results in the AL group as spurious, but a conclusion cannot be reached before sequencing has been undertaken. Our results are promising for in depth exploration of the microbiology of prosthetic

joint infections. Parallel sampling for culture and 16S *rRNA* gene analysis allow an extensive evaluation of the diagnostic accuracy of different sample types and methods. A high concordance between clinical assessment and results obtained by optimized culture methods is indicated in several reports (2,3,4) and with the large patient series in the PRIS project these methods can be thoroughly tested against each other with different sample types.

We expect that the results from the ongoing PRIS project can contribute to the rational foundation for diagnosis of suspected prosthetic joint infections.

		Surgical revision no.																																								
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	18	19	20	21																					
Infections (PJI)	Joint Fluid	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red																					
	Prosthesis Swab	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red																					
	Synovialis	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red																					
	Bone	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red	Red																					
		Surgical revision no.																																								
		21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61
Aseptic Loosening (AL)	Joint Fluid	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green																					
	Prosthesis Swab	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green																					
	Synovialis	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green																					
	Bone	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green	Green																					

**Table 1:** Concordance between 16S *rRNA* gene analysis (PCR) and optimized culture methods. Red: PCR+Culture positive. Blue: PCR positive. Purple: Culture positive. Green PCR+Culture negative. White: No sample available. Two patients have two surgical procedures in the Infection Group (PJI).

## Acknowledgement


This study is part of the Danish 'Prosthetic-Related Infection and Pain' Innovation Project (Danish acronym PRIS), supported by a grant from the Danish Agency for Science, Technology and Innovation, Danish Ministry of Science, Innovation and Higher Education

[www.joint-prosthesis-infection-pain.dk](http://www.joint-prosthesis-infection-pain.dk)

## Reference

- (1) Danish Knee Arthroplasty Register and the Danish Hip Arthroplasty Register ([www.dkar.dk](http://www.dkar.dk) and [www.dhr.dk](http://www.dhr.dk))
- (2) Larsen LH., Lange J., Xu Y., Schönheyder HC. (2012). Optimizing culture methods for diagnosis of prosthetic joint infections: a summary of modifications and improvements reported since 1995. *J Med Microbiol* 61, 309-316
- (3) Trampuz A., Piper KE., Jacobson MJ., Hanssen AD., Unni KK., Osmon DR., Mandrekar JN., Cockerill FR., Seckelberg, JM. & et al. (2007). Sonication of removed hip and knee prostheses for diagnosis of infection. *N Engl J Med* 357, 654-663.
- (4) Gomez E. & Patel R. (2011). Laboratory diagnosis of prosthetic joint infection, Part I. *Clin Microbiol Newsl* 33, 55-60. Gomez E. & Patel R. (2011). Laboratory diagnosis of prosthetic joint infection, Part II. *Clin Microbiol Newsl* 33, 63-70.
- (5) Pitt WG., Ross A. (2003). Ultrasound Increases the rate of bacterial cell growth. *Biotechnol Prog* 19, 1038-1044

AALBORG HOSPITAL  
AARHUS UNIVERSITY HOSPITAL



DANISH  
TECHNOLOGICAL  
INSTITUTE



Environmental Biotechnology  
Aalborg University