



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Optimizing the identification of citrullinated peptides by mass spectrometry

Utilizing the inability of trypsin to cleave after citrullinated amino acids

Bennike, Tue; Lauridsen, Kasper; Olesen, Michael Kruse; Andersen, Vibeke; Birkelund, Svend; Stensballe, Allan

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Bennike, T., Lauridsen, K., Olesen, M. K., Andersen, V., Birkelund, S., & Stensballe, A. (2014). *Optimizing the identification of citrullinated peptides by mass spectrometry: Utilizing the inability of trypsin to cleave after citrullinated amino acids*. Poster presented at 19th Lorne Proteomics Symposium, Melbourne, Australia.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Optimizing the Identification of Citrullinated Peptides by Mass Spectrometry


Tue Bennike¹, Kasper B. Lauridsen¹, Michael Kruse Olesen², Vibeke Andersen³, Svend Birkelund¹, Allan Stensballe¹

¹Department of Health Science and Technology, Aalborg University, Denmark

²Department of Rheumatology, and Center for Clinical Research, Vendsyssel Teaching Hospital, Denmark

³Institute of Regional Health Services Research, University of Southern Denmark

AALBORG UNIVERSITY
AALBORG · ESBJERG · COPENHAGEN



AIM


- Investigate the cleavage properties of trypsin after a citrulline residue.
- Investigate the behavior of citrullinated peptides by reversed phase chromatography.
- Propose a verification strategy for detected citrullinated peptides in a MS workflow.

Conclusion

- Our results clearly demonstrate the inability of trypsin to cleave after citrulline residues. As a result, a miscleavage can be used to distinguish a citrullination from a deamidation of asparagine or glutamine.
- The shift in retention time was, for 22 of 24 peptides large enough to ensure that both peptides could be identified.

Introduction

Citrullination is a PAD-enzyme catalyzed deimination of arginine, yielding the non-standard amino acid citrulline.¹



Protein citrullination has been associated with several diseases and auto-antibodies against citrullinated proteins are today used as an important clinical biomarker in rheumatoid arthritis.^{2,7} The site-specific characterization of citrullination using mass spectrometry remains problematic, especially as citrullination and deamidation of asparagine or glutamine results in the same mass increase of +0.984016 Da. The verification, therefore, often relies on a tryptic miscleavage after citrulline.³ Furthermore, the mass increase is close to that of a neutron, +1.08665 Da.

However, tryptic cleavage after citrulline has in some cases been reported, so we here investigate the cleavage properties of trypsin after a citrulline residue.

Method

24 synthetic peptide sets containing either arginine or citrulline were analyzed (JPT Peptide Technologies GmbH). The peptide sequences originated from disease-associated *in vivo* citrullinated proteins. In-solution tryptic digestion was performed with sequencing grade trypsin (Promega). 1 pmol sample was analysed using ESI LC-MS/MS in positive ion mode, on a hybrid microQTOF mass spectrometer (Bruker). The peptides were separated using an in-house packed 10 cm reversed phase C18 column (Dr. Maisch; repositil-pur C18-AQ) with acetonitrile.

Extracted ion chromatograms (XIC) were constructed in Bruker Daltonics DataAnalysis v 3.4, with all predicted tryptic peptides +/- m/z 0.01, under the assumption that trypsin cleaves after arginine, lysine and citrulline.


References

- W. R. Fearon, "The carbamido diacetyl reaction: a test for citrulline.", *Biochem J.* 33, 902-907 (1939).
- G. A. Schellekens, et al., "Citrulline is an essential constituent of antigenic determinants recognized by rheumatoid arthritis-specific autoantibodies", *J. Clin. Invest.* 101, 273 (1998).
- M. De Ceuleneer, et al., "In vivo relevance of citrullinated proteins and the challenges in their detection", *PROTEOMICS* 12, 752-760 (2012).
- P. K. Gregersen, et al., "Circulating immune complexes contain citrullinated fibrinogen in rheumatoid arthritis", *Arthritis Res Ther.* 10, 94 (2008).
- G. Hao, et al., "Neutral loss of isocyanic acid in peptide CID spectra: a novel diagnostic marker for mass spectrometric identification of protein citrullination", *J. Am. Soc. Mass* 20, 4, 723-727 (2009).
- T. Bennike, et al., "Optimizing the Identification of Citrullinated Peptides by Mass Spectrometry: Utilizing the Inability of Trypsin to Cleave after Citrullinated Amino Acids", *Journal of Proteomics & Bioinformatics* 06, (2013)
- T. Bennike, et al., "Biomarkers in inflammatory bowel diseases: Current status and proteomics identification strategies. *World J. Gastroenterol.* 20, 3231-3244 (2014).

| SP # | Sequence | Protein ID | % RT shift |
|------|--|-------------|------------|
| 1 | DS R/Cit GNPTVEVDLFTSKGLFR | P06733.2 | 2.4 |
| 2 | DPS R/Cit YISPDQLADLYKSEIK | P06733.2 | 3.0 |
| 3 | EELGSKAKFAG R/Cit NF R/Cit NPLAK | P06733.2 | 1.6 |
| 4 | VTTSTRTYSLGSAL R/Cit PSTSR | AAH66956 | 1.4 |
| 5 | EQLKGGCKS R/Cit LGDLYEEMR | AAH66956 | 2.6 |
| 6 | NMKEEMARHL R/Cit EYQDLLNVK | AAH66956 | 1.4 |
| 7 | NMKEEMA R/Cit HLREYQDLLNVK | AAH66956 | 2.0 |
| 8 | LHVA R/Cit SEMDKV R/Cit VFQAT R/Cit GK | NP_036519.2 | 3.6 |
| 9 | GLKEFFIK R/Cit VMGPDFGYVTR | NP_036519 | 3.4 |
| 10 | PAPDRKGFRLLLASP R/Cit SCYK | NP_036519 | 2.0 |
| 11 | LS R/Cit TVRCTCISISNPQVNR | P02778.2 | 1.6 |
| 12 | EMHGKNWSKLC R/Cit DCQVIDGR | ACB10579 | 1.6 |
| 13 | SGVTKAISSTVS R/Cit LTDTTK | ACB10579 | 1.4 |
| 14 | AEGGGV R/Cit GPRVVE R/Cit HQSACK | P02671 | 0 |
| 15 | SHHPGIAEPPS R/Cit GKSSSYSK | P02671 | 0.4 |
| 16 | FTSSTSYN R/Cit GDSFESKSYK | P02671 | 1.0 |
| 17 | A R/Cit HGFLP R/Cit HRDTGILDSIGR | P02686.3 | 2.6 |
| 18 | LSRFSWGAEGQ R/Cit PGFGYGGK | P02686.3 | 2.0 |
| 19 | PGFGYGG R/Cit ASDYKSAHKGFK | P02686.3 | 0.6 |
| 20 | LSKIFKLG R/Cit DSRSGSPMAR | P02686.3 | 0.6 |
| 21 | Y R/Cit VYCDMNTENGWTVIQNR | P02675 | 0 |
| 22 | MYLIQPDSSVKPY R/Cit VYCDNR | P02675 | 3.4 |
| 23 | EAPSL R/Cit PAPPPISGGGYRAR | P02675 | 2.0 |
| 24 | SI R/Cit YLQEIYNSNNQKIVNLK | P02679.3 | 3.0 |

Results and Discussion

In situ digestion results were compared to the empirical data. For SP 10, prior to digestion only the synthetic peptide is detected and after digestion, peptides corresponding to PAPDR, LLLASPR and SCYK are detected, corresponding to a successful complete cleavage after 495_{Arg}. This is not the case after digestion of SP 10_{Cit}, where peptides corresponding to PAPDR and LLLASPCitSCYK are detected. All investigated peptides demonstrate this behavior.



Our results clearly demonstrate the inability of trypsin to cleave after citrulline residues. Hence, a miscleavage indicates the presence of the PTM. Furthermore, the shift in retention time between the citrulline and arginine peptides was large enough for 22 of the 24 peptides to ensure that coelution is not occurring to a detectable extent, ensuring that both peptides can be identified.

Download Paper

