

Southern Illinois University Carbondale  
**OpenSIUC**

---

Articles and Preprints

Department of Mathematics

---

2009

# Neighborhood Champions in Regular Graphs

John McSorley

*Southern Illinois University Carbondale*, [jmcsorley@math.siu.edu](mailto:jmcsorley@math.siu.edu)

Walter D. Wallis

*Southern Illinois University Carbondale*

Follow this and additional works at: [http://opensiuc.lib.siu.edu/math\\_articles](http://opensiuc.lib.siu.edu/math_articles)

---

## Recommended Citation

McSorley, John and Wallis, Walter D. "Neighborhood Champions in Regular Graphs." *Journal of Combinatorial Mathematics and Combinatorial Computing* 70 (Jan 2009): 49-56.

This Article is brought to you for free and open access by the Department of Mathematics at OpenSIUC. It has been accepted for inclusion in Articles and Preprints by an authorized administrator of OpenSIUC. For more information, please contact [opensiuc@lib.siu.edu](mailto:opensiuc@lib.siu.edu).

# Neighborhood Champions in Regular Graphs

J. P. McSorley and W. D. Wallis,  
Department of Mathematics,  
Southern Illinois University,  
Carbondale, IL 62901-4408. USA.

## Abstract

For a vertex  $x$  in a graph  $G$  we define  $\Psi_1(x)$  to be the number of edges in the closed neighborhood of  $x$ . Vertex  $x^*$  is a neighborhood *champion* if  $\Psi_1(x^*) > \Psi_1(x)$  for all  $x \neq x^*$ . We also refer to such an  $x^*$  as a *unique champion*. For  $d \geq 4$  let  $n_0(1, d)$  be the smallest number such that for every  $n \geq n_0(1, d)$  there exists a  $n$  vertex  $d$ -regular graph with a unique champion. Our main result is that  $n_0(1, d)$  satisfies  $d+3 \leq n_0(1, d) \leq 3d+1$ . We also observe that there can be no unique champion vertex when  $d = 3$ .

## 1 Introduction

We assume the standard ideas of graph theory. All graphs considered in this paper will be connected. We sometimes specify the vertex-set  $V$  and edge-set  $E$  of a graph  $G$  by denoting the graph  $G(V, E)$ .  $|V|$  and  $|E|$  are respectively the *order* and *size* of  $G$ . The *distance*  $d(v, u)$  between two vertices  $v$  and  $u$  in a graph is the number of edges in a shortest path from  $v$  to  $u$ . The *closed  $k$ -neighborhood* of a vertex  $v$  is defined as

$$N_k[v] = \{u \in V : d(v, u) \leq k\}.$$

When  $k = 1$  we simply use the term “neighborhood”.

The scale- $k$  locality statistic  $\Psi_k(x)$  of vertex  $x$  in a connected graph  $G(V, E)$  was defined in [4] to be the size of the subgraph induced by the closed  $k$ -neighborhood of  $x$ . (This statistic is also called the number of edges *seen* by  $x$ .) The scale- $k$  scan statistic  $M_k(G)$  of  $G$  was then defined to be the maximum over  $x \in V$  of the scale- $k$  locality statistics:

$$M_k(G) = \max_{x \in V} \Psi_k(x).$$

In a mild abuse of notation, we define  $\Psi_0(x)$  to be the degree of vertex  $x$  in  $G$ , and  $M_0(G)$  to be the maximum degree in  $G$ .

A *champion* for scale  $k$  is a vertex  $x^*$  such that  $\Psi_k(x^*) > \Psi_k(x)$  for all  $x \neq x^*$ . The reasons for studying champions are discussed in [4]; in that paper the authors discussed *dramatic champions*, for which  $\Psi_k(x^*) \gg \Psi_k(x)$  for all  $x \neq x^*$ . In [3] we construct some families of dramatic *neighborhood champions*, or champions for scale 1.

In this paper we focus on the pure graph theory of the situation, and consider the existence of neighborhood champions for scale 1 in connected regular graphs. We shall sometimes discuss graphs in which more than one vertex attains the maximum value  $M_1(G)$ . We shall use the word “co-champion” to denote these vertices.

## 1.1 Groupies

A related, but different, concept to a champion in  $G$  is a *groupie* in  $G$ . No confusion should arise between the two: A vertex in  $G$  is a groupie if the average degree of its neighbors is greater than or equal to the average degree of  $G$ . The concept was introduced in [1] (see also [2]).

## 2 Cubic graphs: $d = 3$

**Theorem 1.** *For  $d = 1, 2$ , and  $3$  there are no  $d$ -regular graphs with a unique neighborhood champion.*

**Proof** Clearly regular graphs of degrees  $d = 1$  or  $2$  have no champions.

Now suppose  $G$  is a cubic graph: If  $M_1(G) = 3$  then *every* vertex attains  $M_1(G)$ . If  $M_1(G) = 4$ , then any vertex  $x$  with  $\Psi_1(x) = 4$  lies in exactly one triangle, and the other vertices of the triangle are co-champions; so  $G$  contains at least three co-champions. If  $M_1(G) = 5$  and vertex  $x$  sees five edges, the configuration must be as shown in Figure 1, where  $y$  is a co-champion. And if  $M_1(G) = 6$  (the maximum) we have  $G = K_4$ , and every vertex is a co-champion. Thus no cubic graph has a champion.




Figure 1: A cubic configuration

■

However, one can construct cubic graphs with precisely two co-champions, or *twin champions*, for every even number  $n \geq 10$  of vertices. From above we must have  $M_1(G) = 5$ .

A short exhaustive search shows that this is impossible for fewer than 10 vertices (the graphs may be found on page 127 of [5]).

For every even  $n \geq 10$  we construct a cubic graph  $G$  on  $n$  vertices for which  $\Psi_1(x) = M_1(G) = 5$  for precisely two vertices. Our technique is to implant the graph shown in Figure 2 as a subgraph of a host graph. The implant graph  $H$  has six vertices  $a, b, p, q, y, z$  and adjacencies  $ap, bq, yp, yq, zp, zq, yz$ .




Figure 2: The graph  $H$  to be implanted

**Construction** Select any triangle-free cubic graph on  $n - 4$  vertices and choose any edge  $ab$  in that graph. Delete this edge. Then identify vertices  $a, b$  with the vertices  $a, b$  of  $H$ . See Figure 3 for an example of this; the value of  $\Psi_1(x)$  is shown on each vertex and the champion is emphasized.

$$\Psi_1(y) = \Psi_1(z) = 5, \Psi_1(p) = \Psi_1(q) = 4 \text{ and } \Psi_1(x) = 3 \text{ otherwise.}$$

To show that the construction is always possible for  $n \geq 10$ , we observe

**Lemma 1.** *If  $n - 4 = 2s \geq 6$ , there is a triangle-free cubic graph on  $n - 4$  vertices.*

**Proof** Take the integers modulo  $2s$  as vertices. For each  $i$ , let vertex  $i$  be adjacent to vertices  $i - 1, i + 1$ , and  $i + s$  (modulo  $2s$ ). ■

(This graph is called a *Möbius ladder* [5, p263].)

So we have

**Theorem 2.** *For every even  $n \geq 10$  there exists a cubic graph on  $n$  vertices with precisely two co-champions.* ■




Figure 3: Example for 10 vertices

### 3 General Constructions: $d \geq 4$

For even  $d \geq 4$  let  $n_0(c, d)$  be the smallest number such that for every  $n \geq n_0(c, d)$  there exists a  $n$  vertex  $d$ -regular graph with precisely  $c$  neighborhood co-champions; for odd  $d$  we require existence for even  $n \geq n_0(c, d)$  only.

In this section we discuss  $n_0(1, d)$ .

The complete graph on  $n$  vertices is denoted  $K_n$ , while  $K_{m,n}$  denotes the complete bipartite graph with vertex-sets of sizes  $m$  and  $n$ . A *one-factor* is a graph consisting of disjoint edges; in particular, given two ordered sets of vertices  $Y = \{y_0, y_1, \dots, y_{n-1}\}$  and  $Z = \{z_0, z_1, \dots, z_{n-1}\}$ , we define the one-factor  $F_j^n(Y, Z)$  to consist of the edges

$$y_0 z_j, y_1 z_{1+j}, \dots, y_{n-1} z_{n-1+j},$$

where subscripts are reduced modulo  $n$ . Then  $K_{n,n}$  can be represented as

$$F_0^n(Y, Z) \cup F_1^n(Y, Z) \cup \dots \cup F_{n-1}^n(Y, Z).$$

**Lemma 2.** *Suppose  $d \geq 4$ . For every  $t \geq 0$  there exists a  $d$ -regular graph, with a neighborhood champion, on  $n = 3d + 2t + 1$  vertices.*

**Proof** Let  $H$  represent the complete graph on the  $d+1$  vertices  $x_0, x_1, \dots, x_d$  with the  $d$  edges of the cycle  $x_0 x_1 x_2 \dots x_{d-1}$  deleted. Take a copy of  $F_0^n(Y, Z) \cup$

$F_1^n(Y, Z) \cup \dots \cup F_{d-1}^n(Y, Z)$ , where  $n = d+t$ , and delete the edges  $y_0z_0, y_1z_1, \dots, y_{d-1}z_{d-1}$ . Adjoin this to  $H$  by adding edges  $x_0y_0, x_0z_0, x_1y_1, x_1z_1, \dots, x_{d-1}y_{d-1}, x_{d-1}z_{d-1}$ .

In this graph,

$$\begin{aligned}\Psi_1(x_d) &= d(d-1)/2, \\ \Psi_1(x_i) &= (d^2 - 5d + 14)/2, \text{ for } 0 \leq i \leq d-1, \\ \Psi_1(y_j) &= \Psi_1(z_j) = d, \text{ for } 0 \leq j \leq n-1.\end{aligned}$$

Then  $x_d$  is a champion provided  $d(d-1)/2 > (d^2 - 5d + 14)/2$ , that is  $d \geq 4$ . ■

The above construction gives graphs whose order is of opposite parity to  $d$ . When  $d$  is odd, this provides all possible orders from some point on, because regular graphs of odd degree must have even order. However, for even degree, another construction is needed for even orders.

Suppose  $G$  is the graph of Lemma 2 in the case where  $d \geq 4$  is even. We modify  $G$  to form  $\hat{G}$  as follows: Add a vertex  $\hat{x}$ . Delete the  $d/2$  edges  $x_0y_0, x_2y_2, x_4y_4, \dots, x_{d-2}y_{d-2}$ , and add the  $d$  edges  $\hat{x}x_0, \hat{x}y_0, \hat{x}x_2, \hat{x}y_2, \hat{x}x_4, \hat{x}y_4, \dots, \hat{x}x_{d-2}, \hat{x}y_{d-2}$ .

The  $\Psi_1$  values of all vertices of  $G$  are unchanged. We have  $\Psi_1(\hat{x}) = d + \binom{d/2}{2} = (d^2 + 6d)/8$ . Thus vertex  $x_d$  is still the champion, and we have

**Lemma 3.** *Suppose  $d \geq 4$  is even. For every  $t \geq 0$  there exists a  $d$ -regular graph, with a neighborhood champion, on  $n = 3d + 2t + 2$  vertices.* ■

**Theorem 3.** *Suppose  $d \geq 4$ . Then*

$$d + 3 \leq n_0(1, d) \leq 3d + 1.$$


**Proof** For any  $d \geq 4$  the only  $d$ -regular graph with  $d+1$  vertices is  $K_{d+1}$ , which clearly doesn't have a unique champion. And for odd  $d \geq 4$  there is no  $d$ -regular graph with  $d+2$  vertices, so  $n_0(1, d) \geq d+3$ . And for even  $d \geq 4$  the only  $d$ -regular graph with  $d+2$  vertices is  $K_{d+2}$  minus a one-factor, which again doesn't have a unique champion; so  $n_0(1, d) \geq d+3$  here also. Hence, for any  $d \geq 4$ , we have  $n_0(1, d) \geq d+3$ . The upper bound  $n_0(1, d) \leq 3d+1$  comes from Lemmas 2 and 3. ■

## 4 Small degrees: $d = 4, 5$


By inspection, there are no 4-regular (quartic) graphs on  $n = 7$  or 8 vertices with a unique champion (see [5, p145]). From Theorem 3 and the examples for orders  $n = 9, 10, 11$  and 12 shown in Figure 4 we see that  $n_0(1, 4) = 9$ , *i.e.*, there is a 4-regular graph, with a neighborhood champion, on  $n$  vertices whenever  $n \geq 9$ .

Similarly, at degree 5, inspection shows (see [5, p154]) there are no quintic graphs on  $n = 6$  or 8 vertices with a unique champion. We present examples on 10, 12 and 14 vertices in Figure 5, showing that  $n_0(1, 5) = 10$ . Thus there is a 5-regular graph, with a neighborhood champion, on  $n$  vertices for every even  $n \geq 10$ . So the cases of  $d = 4$  or 5 are completely solved.







9 vertices



10 vertices




11 vertices




12 vertices

Figure 4: Small quartic graphs, each with a champion

10 vertices



12 vertices



14 vertices




Figure 5: Small quintic graphs, each with a champion

## 5 Directions for future work

In our constructions above of a  $d$ -regular graph with a unique champion we allowed the function  $\Psi_1$  to take on as many values as the construction demanded. In some applications it is desirable to restrict the range of  $\Psi_1$  to just two values, namely  $\Psi_1(x^*)$ , the value of  $\Psi_1$  at the champion vertex  $x^*$ , and  $\Psi_1(x)$  for all other vertices  $x \neq x^*$ . Some graphs with this property were constructed in [4], and the existence of further graphs is an area for future research.

Another problem is the existence of regular graphs with precisely two co-champions. We discussed the  $d = 3$  case earlier; some further results have been obtained and will be the subject of a future paper.

## References

- [1] M. Ajtai, J. Komlós and E. Szemerédi, A note on Ramsey numbers. *J. Combinatorial Theory* **29A** (1980), 354-360.
- [2] E. Bertram, P. Erdős, P. Horák, J. Širáň and Zs. Tuza, Local and global average degree in graphs and multigraphs. *J. Graph Theory* **18** (1994), 647-661.
- [3] J. P. McSorley, C. E. Priebe and W. D. Wallis, Neighborhood Champions. (In preparation)
- [4] C. E. Priebe and W. D. Wallis, On the Anomalous Behaviour of a Class of Locality Statistics. *Discrete Math.* **308** no.10 (2008), 2034-2037.
- [5] R. C. Read and R. J. Wilson, *An Atlas of Graphs* (Oxford U.P., 1999).