



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Migraine, body weight and psychological factors in children and adolescents

Tarantino, Samuela; De Ranieri, Cristiana; D'Ambrosio, Monica; Capuano, Alessandro; Frusciante, Roberto; Vigevano, Federico; Gentile, Simonetta; Valeriani, Massimiliano

Published in:

Journal of Headache and Pain

DOI (link to publication from Publisher):

[10.1186/1129-2377-16-S1-A153](https://doi.org/10.1186/1129-2377-16-S1-A153)

Publication date:

2015

Document Version

Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Tarantino, S., De Ranieri, C., D'Ambrosio, M., Capuano, A., Frusciante, R., Vigevano, F., ... Valeriani, M. (2015). Migraine, body weight and psychological factors in children and adolescents. *Journal of Headache and Pain*, 16(Suppl. 1), No. A153. DOI: 10.1186/1129-2377-16-S1-A153

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

POSTER PRESENTATION

Open Access

P029. Migraine, body weight and psychological factors in children and adolescents

Samuela Tarantino^{1*}, Cristiana De Ranieri², Monica D'Ambrosio², Alessandro Capuano¹, Roberto Frusciante¹, Federico Vigevano¹, Simonetta Gentile², Massimiliano Valeriani^{1,3}

From Abstracts from the 1st Joint ANIRCEF-SISC Congress
Rome, Italy. 29-31 October 2015

Background

Several studies have assessed the associations between adult migraine and underweight, pre-obesity or obesity. Prevalence, frequency, and severity of migraine appear to increase in relation to the body mass index, although this evidence is not supported by all the studies examined. The link between body weight and headache has hardly been examined in children. Data on the possible association between the body mass index (BMI) and the psychological profile in migraine children are sparse.

Objectives

Aims of the present study were: 1) to study the prevalence of pre-obesity and obesity in migraineur children/adolescents; 2) to analyze the possible relationship between frequency and severity of migraine and overweight; 3) to explore the role of anxiety and somatization on BMI in migraine patients.

Methods

We studied 92 migraineurs (mean age 11.3±2.3 years; 43 M and 49 F). Patients were divided into 2 groups according to headache attack frequency: 1) high frequency (HF) patients, having from weekly to daily episodes, and 2) low frequency (LF) patients, showing ≤ 3 episodes per month. Pain intensity was rated on a 3-level graduated scale (mild, moderate and severe pain). Given the low frequencies, the “moderate” and the “mild” intensity were collapsed into the same category. The psychological profile was assessed by SAFA Anxiety and Somatization scales. BMI was calculated as the

weight in kilograms divided by the height in meters squared.

Results

Among our patients, fifty-seven (62.0%) were classified as “normal weight”, 15.2% were obese and 17.4% pre-obese (both collapsed into the “overweight” group). Due to their low frequencies, “underweight” children/adolescents (5.4%) were eliminated from our subsequent analysis. The weight (“normal weight” or “overweight”) did not correlate with migraine frequency and intensity (respectively: $\chi^2 = 0.6853$, $p = 0.41$; $\chi^2 = 0.0058$; $p = 0.94$). Compared to normal weight children, overweight patients showed a significant higher score in “Separation anxiety” subscale. In the “overweight” patients, the BMI showed a positive and significant correlation with Anxiety (SAFA-A Total, $p = 0.000$) and with Somatization (SAFA-S Total, $p = 0.000$).

Conclusions

Our results suggest that, in young patients, there is an association between migraine, weight and psychological symptoms. Overweight migraineur patients are more prone to “separation anxiety”. In particular, we can hypothesize that overweight in migraine children may be related to anxiety and somatization symptoms.

Written informed consent to publish was obtained from the patient(s).

Authors' details

¹Headache Center, Division of Neurology, Pediatrico Bambino Gesù, IRCCS, Rome, Italy. ²Unit of Clinical Psychology, Ospedale Pediatrico Bambino Gesù, IRCCS, Rome, Italy. ³Center for Sensory-Motor Interaction, Aalborg University, Aalborg, Denmark.

* Correspondence: samuela.tarantino@opbg.net

¹Headache Center, Division of Neurology, Pediatrico Bambino Gesù, IRCCS, Rome, Italy

Full list of author information is available at the end of the article

Published: 28 September 2015

doi:10.1186/1129-2377-16-S1-A153

Cite this article as: Tarantino *et al.*: P029. Migraine, body weight and psychological factors in children and adolescents. *The Journal of Headache and Pain* 2015 **16**(Suppl 1):A153.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
