



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Does migraine follow benign paroxysmal torticollis?

Bernucci, Maria Chiara; Frusciante, Roberto; Capuano, Alessandro; Tarantino, Samuela; Vigevano, Federico; Valeriani, Massimiliano

Published in:

Journal of Headache and Pain

DOI (link to publication from Publisher):

[10.1186/1129-2377-16-S1-A143](https://doi.org/10.1186/1129-2377-16-S1-A143)

Publication date:

2015

Document Version

Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Bernucci, M. C., Frusciante, R., Capuano, A., Tarantino, S., Vigevano, F., & Valeriani, M. (2015). Does migraine follow benign paroxysmal torticollis? *Journal of Headache and Pain*, 16(Suppl. 1), No. A143. DOI: 10.1186/1129-2377-16-S1-A143

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

ORAL PRESENTATION

Open Access

O016. Does migraine follow benign paroxysmal torticollis?

Maria Chiara Bernucci¹, Roberto Frusciante¹, Alessandro Capuano¹, Samuela Tarantino¹, Federico Vigeveno¹, Massimiliano Valeriani^{1,2*}

From Abstracts from the 1st Joint ANIRCEF-SISC Congress
Rome, Italy. 29-31 October 2015

Background

Migraine equivalents are clinical conditions which often involve children who do not complain of headache. They include abdominal migraine, motion sickness, limb pain, cyclical vomiting, benign paroxysmal vertigo, and benign paroxysmal torticollis (BPT). The aim of our study was to investigate whether children referred to us for BPT have developed migraine at a distance from our first observation.

Methods

Forty-one children were included in the study. Only 36 families could be contacted by phone, but 2 of them refused to answer our questionnaire. Therefore, the present results were obtained from 34 children (22 girls and 12 boys).

Results

Migraine could be diagnosed in 14 children (41%), while the remaining 20 patients (59%) did not complain of headache. At the moment of our interview, children who had developed migraine had a mean age of 5 years, while the mean age of non-migrainous children was 3.5 years. Among migraine children, 43% developed it when they were 4 years, 21% at the age of 3, and 14% when they were 7 years old. The last three migraineurs developed migraine at the age of 5 years, 13 years and 18 months, respectively. Moreover, 55% of patients had developed other migraine equivalents. In particular, 73% children had abdominal migraine, 55% vertigo, 45% limb pain, 27% motion sickness, 27% cyclical vomiting. As for the paroxysmal torticollis time course, two children have had only one event, one child had been still

presenting episodes of torticollis, while in the remaining patients the torticollis events had not occurred for some years.

Conclusions

Our findings suggest that migraine follows BPT in approximately half of the children within the age of 13 years. Moreover, BPT is often associated to other migraine equivalents. Considered all together, all these periodic syndromes increase the risk of developing migraine.

Written informed consent to publish was obtained from the patient(s).

Authors' details

¹Headache Centre, Ospedale Bambino Gesù, IRCCS, Rome, Italy. ²Center for Sensory-Motor Interaction, Aalborg University, Aalborg, Denmark.

Published: 28 September 2015

doi:10.1186/1129-2377-16-S1-A143

Cite this article as: Bernucci et al.: O016. Does migraine follow benign paroxysmal torticollis? *The Journal of Headache and Pain* 2015 **16**(Suppl 1):A143.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com

* Correspondence: m.valeriani@tiscali.it

¹Headache Centre, Ospedale Bambino Gesù, IRCCS, Rome, Italy
Full list of author information is available at the end of the article