



Aalborg Universitet

AALBORG UNIVERSITY  
DENMARK

## Antipruritic effect of cold-induced and transient receptor potential-agonist-induced counter-irritation on histaminergic itch in humans

Andersen, Hjalte Holm; Melholt, Camilla; Hilborg, Sigurd Due; Jerwiarcz , Anne ; Randers, Amalie; Simoni , Amalie Højmoose; Elberling, Jesper; Arendt-Nielsen, Lars

*Published in:*  
Acta Dermato-Venereologica

*DOI (link to publication from Publisher):*  
[10.2340/00015555-2447](https://doi.org/10.2340/00015555-2447)

*Publication date:*  
2017

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication from Aalborg University](#)

### *Citation for published version (APA):*

Andersen, H. H., Melholt, C., Hilborg, S. D., Jerwiarcz , A., Randers, A., Simoni , A. H., ... Arendt-Nielsen, L. (2017). Antipruritic effect of cold-induced and transient receptor potential-agonist-induced counter-irritation on histaminergic itch in humans. *Acta Dermato-Venereologica*, 97(1), 63-70. DOI: 10.2340/00015555-2447

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

### **Take down policy**

If you believe that this document breaches copyright please contact us at [vbn@aub.aau.dk](mailto:vbn@aub.aau.dk) providing details, and we will remove access to the work immediately and investigate your claim.

# Antipruritic Effect of Cold-induced and Transient Receptor Potential-agonist-induced Counter-irritation on Histaminergic Itch in Humans\*

Hjalte H. ANDERSEN<sup>1</sup>, Camilla MELHOLT<sup>1</sup>, Sigurd D. HILBORG<sup>1#</sup>, Anne JERWIARZ<sup>1#</sup>, Amalie RANDERS<sup>1#</sup>, Amalie SIMONI<sup>1#</sup>, Jesper ELBERLING<sup>2</sup> and Lars ARENDT-NIELSEN<sup>1</sup>

<sup>1</sup>Laboratory of Experimental Cutaneous Pain Research, SMI®, Department of Health Science and Technology, Faculty of Medicine, Aalborg University, Aalborg, and <sup>2</sup>The Allergy Clinic, Copenhagen University Hospital, Gentofte, Copenhagen, Denmark

\*These authors contributed equally.

**A frequent empirical observation is that cold-induced counter-irritation may attenuate itch. The aim of this randomized, single-blinded, exploratory study was to evaluate the counter-irritation effects of cold-stimulation and topical application of transient receptor potential TRPA1/M8-agonists (trans-cinnamaldehyde/L-menthol, respectively), on histamine-induced itch, wheals and neurogenic inflammation in 13 healthy volunteers. Histamine 1% was applied to the volar forearms using skin prick-test lancets. Recorded outcome-parameters were itch intensity, wheal reactions, and neurogenic inflammation (measured by laser-speckle perfusion-imaging). Homotopic thermal counter-irritation was performed with 6 temperatures, ranging from 4°C to 37°C, using a 3 × 3-cm thermal stimulator. Chemical “cold-like” counter-irritation was conducted with 40% L-menthol and 10% trans-cinnamaldehyde, while 5% doxepin was used as a positive antipruritic control/comparator. Cold counter-irritation stimuli from 4°C to 22°C inhibited itch in a stimulus-intensity-dependent manner ( $p < 0.05$ ) and, to a lesser extent, also wheal reactions and neurogenic inflammation. Chemical “cold-like” counter-irritation with both L-menthol and trans-cinnamaldehyde had antipruritic efficacy similar to doxepin ( $p < 0.05$ ). Cold-induced counter-irritation had an inhibitory effect on histaminergic itch, suggesting that agonists of cold transduction receptors could be of potential antipruritic value.**

*Key words:* itch; histamine; cold stimulation; TRPs; counter-irritation; TRPM8; TRPA1.

Accepted Apr 28, 2016; Epub ahead of print May 3, 2016

Acta Derm Venereol 2017; 97: 63–70.

*Corr:* Lars Arendt-Nielsen, Fredrik Bajers Vej 7, Bld. D3, DK-9220 Aalborg E, Denmark. E-mail: LAN@hst.aau.dk

Itch is an unpleasant skin sensation that evokes scratching and frequently manifests as a chronic distressing symptom in association with diseases such as urticaria, atopic dermatitis, psoriasis, and uraemia (1, 2). Chronic

itch profoundly impacts on quality of life in affected patients and, due to a point prevalence of 8–13.5%, in combination with largely suboptimal treatments; it represents a substantial socioeconomic burden (3–5). Itch is transmitted in 2 separate pruritic pathways with distinct primary afferent populations (6–9). A subdivision of pruriceptive mechano-insensitive C-fibres (CMi) convey histaminergic itch (induced in the present study), whereas a subdivision of pruriceptive polymodal C-fibres (PmC) transmit so-called non-histaminergic itch (10). The peripheral fibres synapse with central projections in the superficial layer of the spinal dorsal horn, from where the pruriceptive signals are transmitted to the thalamus and parabrachial nuclei (11–13).

While scratching usually provides transient itch relief, it is thought to prolong exacerbations and be involved in chronification of itch by secondary mechanisms such as xerosis, inflammation, and infection, a phenomenon dubbed the vicious itch-scratch-itch cycle (14–16). However, the transient relief of itch achievable by introducing a mild painful or non-painful counter-irritation, such as scratching, a mechanism mediated by central segmental inhibition (17) via bhlhb5-interneurons in the spinal dorsal horn (18), signifies the existence of an effective endogenous mechanism capable of greatly attenuating itch. It has been shown experimentally that controlled thermal and mechanical painful stimuli inhibit itch and development of dysesthesias, such as alloknesis (10, 15, 19–21). In addition, patients with itch frequently report that it is reduced by cold-stimulation (11, 22–25). This has led to the idea that counter-irritation with non-painful/painful cooling or equivalent chemical agonists of cold thermosensory receptors, e.g. L-menthol, which is already in limited use, might be a feasible approach to alleviate chronic itch (12, 24). This approach is known from research into similar mechanisms related to pain-processing, where counter-irritation has been widely studied (13, 26, 27). A frequently used counter-irritation modality is homotopically-applied cold, which is often effective in relieving inflammatory pain, a phenomenon dubbed cold analgesia. However, evidence from psychophysical studies in patients with itch or using surrogate models of itch in healthy volunteers on the effect of cold or chemical “cold-like” counter-irritation is limited and contradictory (21, 24, 28, 29). For instance,

\*Subsets of the data within this paper were presented as a poster at the IFSI 2015 WCI in Nara Japan and the pertaining abstract was published in Acta Derm Venereol, 2015; 95: 875-912.

Bromm et al. (29) showed a strong antipruritic effect of mild non-noxious cooling and topical L-menthol, but aspects of these results were not reproduced in a later study with similar methodology (24, 30). The receptors conveying thermal sensations are known as transient receptor potential (TRP) channels and can be activated by a range of thermal and chemical stimuli (31–33). For cold thermosensation, TRPM8 is activated at  $\leq 23$ – $27^\circ\text{C}$  and TRPA1 is activated at  $\leq 15$ – $17^\circ\text{C}$ . Moreover, activation of TRPM8 and TRPA1 can be achieved by the use of their natural agonists L-menthol and trans-cinnamaldehyde (CA), respectively (27, 34–38), which provides an opportunity to study the potential gating of itch-signalling induced by homotopic cold counter-irritation using thermal and chemical stimuli comparatively.

The aim of the present exploratory study was to quantify the dose-response relationship between homotopic cold stimulations or chemical “cold-like” stimulations and their counter-irritation effect on experimentally induced histaminergic itch, wheals, and neurogenic inflammation in healthy volunteers. The antihistamine doxepin (5%, topical pretreatment) was used as a comparator.

## METHODS (for complete details see Appendix S1<sup>1</sup>)

### Test subjects and study design

Thirteen healthy subjects (mean age  $22.8 \pm 3$  years; 8 males, 5 females) participated in and completed the study after providing informed consent. The regional ethics committee of Northern Jutland approved the study protocol (N-20140078). All subjects participated in 2 90-min sessions with a minimum 24 h interval between sessions. The study was conducted in a randomized and single-blinded manner with the order of experimental interventions

and arm dominance randomized. Thus, histamine challenges and homotopic counter-irritation stimuli always alternated between the right and left arm. In total, 6 histaminergic itch provocations and 6 randomized counter-irritation stimuli (thermal/chemical) or control conditions were performed in each of the 2 experimental sessions (Fig. 1).

### Thermal sensory thresholds, itch induction, itch assessment, and counter-stimulation


Prior to induction of itch in the first of the 2 experimental sessions, individual cold sensitivity was established using quantitative sensory testing (QST). The cold detection threshold (CDT) and cold pain threshold (CPT) were assessed with a baseline temperature of  $32^\circ\text{C}$  and ramp stimuli decreasing at a rate of  $1^\circ\text{C}/\text{s}$ . When the subjects pressed a button denoting CDT or CPT, the temperature returned to the baseline temperature at a rate of  $5^\circ\text{C}/\text{s}$ .

Itch was induced using a solution of 1.0% histamine dihydrochloride (Allergopharma, Reinbek, Germany) and a standard shouldered 1-mm tip skin prick test (SPT) lancet. A drop of histamine solution was applied to the centre of a  $3 \times 3$  cm square on the skin of the forearm and a custom-made SPT lancet-mount (Aalborg University, Denmark) delivered  $\approx 200$  g of pressure for 2–3 s (41).

The subjects continuously expressed itch intensity on a 10-cm visual analogue scale (VAS<sub>0–10</sub>) after each itch provocation, with 0 being labelled as “no itch” and 10 being labelled as “worst imaginable itch”. The VAS was sampled 4 times/min for the first 5 min of each application. Area under the curve (AUC) was calculated for the 0–2 min (before thermode-induced cold counter-irritation) and at 2–5 min (during cold-counter stimulation). At 3 or 5 min after each interventional condition, the subjects were asked to rate the mean pain experienced as a consequence of the counter-irritation on a VAS<sub>0–10</sub>, with 0 being “no pain” and 10 “worst imaginable pain”.

Thermal counter-irritation stimuli and thermal threshold assessments of cold sensitivity (CDT and CPT) were conducted using the same  $3 \times 3$ -cm Thermal Stimulator probe attached to a Medoc Main Station (MMS). Thermal counter-irritation was applied at 6 temperatures: 37, 32, 28, 22, 12, and  $4^\circ\text{C}$ . The counter-irritation was commenced 1 min and 55 s post-introduction of histamine and lasted for 3 min and 5 s.

<sup>1</sup><https://www.medicaljournals.se/acta/content/abstract/10.2340/00015555-2447>



**Fig. 1. Overview of the experimental set-up of sessions 1 and 2.** The illustration exemplifies the order of experimental procedures conducted in a single subject. Note that the actual order of thermal/chemical interventions was randomized under the condition that applications always switched from arm to arm. Green arrows: skin prick testing lancet application of histamine. Blue arrows: onset of the experimental thermal interventions. Wave shapes: temporal profiles of the itch intensity. L-menthol, CA, and doxepin were applied by pretreatment (see Methods). CA: trans-cinnamaldehyde, CDT: cold detection threshold; CPT: cold pain threshold; FLPI: full-field laser perfusion imaging; VAS: visual analogue scale.

Chemical counter-irritation was induced with 40% L-menthol (>99.9%, TRPM8-agonist) and 10% CA (>99%, TRPA1-agonist), (both Sigma Aldrich, Broendby, Denmark). These were dissolved in 96% ethanol and topically applied to the skin using a 3 × 3 cm cotton pad on a plastic sheet attached to the arm by medical tape. The substances were applied to the skin 7 min before itch induction to achieve a high level of counter-irritation during the application period. Lastly, 5% doxepin (Region Hovedstadens Apotek, Copenhagen, Denmark) was applied to the skin under occlusion at least 60 min before the histamine application, as a comparator.

#### *Neurogenic inflammation (superficial blood flow) and wheal reactions*

To assess neurogenic inflammation, speckle contrast imaging (Moor FLPI, Moor Instruments, Axminster, UK) was recorded 6 min after each histamine application. The imager was placed with a 50-cm distance to the application area, and the pictures were analysed with the appertaining proprietary software. Infrared thermography (A40, FLIR systems, Wilsonville, OR, USA) was conducted to ensure that the control condition of 32°C was within an appropriate temperature range, i.e. close to the normal physiological temperature of the subject. Wheals were measured horizontally and vertically 9 min after every application.

#### *Statistical analysis*

All statistical analyses were performed using SPSS version 22 (IBM, New York, USA). The primary outcome was considered to be the intensity of itch 2–5 or 0–5 min after 1% histamine injections with/without counter-stimuli or control interventions and analysed with a repeated measures analysis of variance (RM-ANOVA) test using the Sidak *post hoc* correction for multiple paired comparisons. Comparisons were made between experimental interventions and their respective control; in particular, the thermal counter-stimuli were statistically compared with the 32°C condition. Secondary outcome parameters were wheal and neurogenic inflammation, which were analysed similarly to the primary outcome. Tests were considered significant at  $p < 0.05$ , and highly significant at  $p < 0.01$ . All results are presented as the arithmetic means ± standard error of the mean (SEM).

## RESULTS

The 13 healthy test subjects recruited for the experiment all completed both sessions of the study without experiencing any adverse reactions during or after the study.

#### *Quantitative sensory thermal tests*

The mean CDT was measured to  $27.89 \pm 1.05^\circ\text{C}$ , i.e. very close to the least intense cold counter-irritation condition at  $28^\circ\text{C}$ , while the mean CPT was measured to  $6.36 \pm 1.87^\circ\text{C}$ , and generally exhibited high inter-individual variation.

#### *Efficacy of counter-irritation*

The temporal development of itch in response to 1% histamine was highly reduced by the thermode-induced cold counter-irritation. Immediately after histamine injection, all test subjects rated rapidly increasing itch intensity,



reaching a mean of  $3.3 \pm 1.2$  (VAS<sub>0–10</sub>), immediately before application of cold-stimuli. Statistical analysis of the AUC after cold stimulation revealed significant reductions in itch intensity for all temperatures ( $p < 0.05$  or  $< 0.01$ ), except for  $32^\circ\text{C}$  and  $37^\circ\text{C}$ , which both caused insignificant reductions in itch. VAS scores after the histamine application without interventions remained between  $3.3 \pm 1.1$  and  $3.8 \pm 1.3$ , at 2 and 4 min, respectively (Fig. 2a). There were no significant differences in the comparison between itch intensity AUC from 0–2 min post-histamine application ( $p > 0.6$ ), indicating lack of bias between conditions. Inter-individual variation in the intensity of the histamine-provoked itch was high, i.e. the highest peak itch intensity score of the baseline condition was VAS = 8.2 and the lowest was VAS = 1.0. Similarly, all chemical counter-irritations, L-menthol ( $p \leq 0.05$ ), CA ( $p \leq 0.01$ ) and doxepin ( $p < 0.01$ ) applied by pre-treatment, caused a significant and pronounced anti-pruritic effect in comparison with the baseline application of histamine (Fig. 2b). The anti-pruritic effect size of the chemical interventions varied between  $-48.5 \pm 12.1\%$  (for L-menthol) and  $-73.6 \pm 10.4\%$  (for CA), but no significant differences were found between effect sizes for any of the substances.

When comparing thermode-induced cold counter-irritation interventions with  $32^\circ\text{C}$ , adjusting for the mechanical pressure stimulation introduced by the weight of the probe, only cold stimulation at 22, 12, and  $4^\circ\text{C}$  caused significant decreases in itch intensity, signifying that both cold-stimuli in the innocuous and noxious range were sufficient to decrease histaminergic itch (see Fig. 2c). Several of the applied counter-irritations gave rise to mild pain, with a mean VAS =  $0.3 \pm 0.2$  for all interventions. CA resulted in a pain score of VAS =  $1.7 \pm 0.5$ ,  $4^\circ\text{C}$  resulted in  $1.2 \pm 0.3$ , and  $12^\circ\text{C}$  stimulation resulted in VAS =  $0.5 \pm 0.3$ . All other interventions were rated VAS  $\leq 0.5$ . This is well aligned with the fact that CA,  $4^\circ\text{C}$ , and  $12^\circ\text{C}$  also gave rise to the most profound anti-pruritic effects.

#### *Neurogenic inflammation (superficial blood perfusion) and wheals after histamine application*

The mean skin temperature assessed by infrared thermography prior to any tests was  $32.67 \pm 0.33^\circ\text{C}$ , i.e. very close to the chosen control temperature of  $32^\circ\text{C}$  and with low inter-individual variability. The neurogenic inflammation, also known as axon-reflex-flare was, for all thermal conditions, most intense in the immediate injection area; 1–2 cm surrounding the injection site where a 3.2-fold increase in superficial skin perfusion was observed, while the flare progressively normalized approximately 4.0–4.5 cm away from the injection site. All thermal applications  $\leq 28^\circ\text{C}$  resulted in a significant decrease in skin perfusion compared with baseline, ( $p < 0.05$ ), Fig. 3a, but only  $22^\circ\text{C}$  and  $12^\circ\text{C}$  stimuli reduced the





neurogenic flare significantly compared with the 32°C control condition ( $p < 0.05$ ). For the chemical counter-irritation, the TRPA1-agonist CA expectedly resulted in a very large neurogenic inflammatory reaction (Fig. 3b),  $p < 0.001$ , as described previously (27, 36, 44). Doxepin and L-menthol both reduced the neurogenic inflammation by a moderate, but significant, extent ( $p < 0.05$ ). The intensity and dispersion of the neurogenic inflammation assessed by speckle contrast imaging is shown in Figs 3a and b. Wheal reactions occurred under all experimental conditions, but were significantly decreased by thermal counter-irritation compared with the 32°C control condition ( $p < 0.01$ ), with the exception of the 28°C stimulation (Fig. 3c). Both at 32°C and without any counter-irritation, the wheals were measured to  $0.22 \pm 0.01$  cm<sup>2</sup> on average. The decreases during thermal counter-irritation varied; from the lowest  $-0.06 \pm 0.01$  cm<sup>2</sup> at 37°C, to the highest  $-0.20 \pm 0.01$  cm<sup>2</sup> at 4°C, representing the almost total abolition of the reaction. For the chemical counter-irritations, both 40% L-menthol ( $p < 0.05$ ) and, to a much greater extent, 10% CA, and 5% doxepin ( $p < 0.001$ ), reduced the wheal reactions (Fig. 3d).

#### Correlational analyses

Correlational relationships were assessed between the magnitude of experienced anti-pruritic effect for any of the thermal or chemical counter-irritations and the QST parameters assessing individual cold hypersensitivity, i.e. CDT and CPT, as well as between the neurogenic flare and the itch intensity. There were no significant correlations between these groups of parameters; however, a nearly significant positive association was found between CPT and itch inhibition at 12°C ( $p = 0.061$ ,  $n = 12$ ), indicating that individuals exhibiting high cold pain sensitivity, i.e. high absolute CPT values, experienced more itch relief from cold counter-irritation near their pain threshold.

## DISCUSSION

The present study investigated the inhibitory effect of homotopic, monophasic cold-stimuli, and TRP-agonist-induced activity on histaminergic itch and vascular responses. For thermode-induced cold counter-irritation, a stimulus-response relationship was established for itch inhibition, starting at 22°C and increasing stepwise towards 4°C, while stimulation at 28°C and 37°C did not significantly inhibit histamine-induced itch perception compared with the control condition at 32°C. The chemical counter-irritations resulted in significant itch inhibition, the most effective being 10% CA followed by 40% L-menthol. The 5% topical doxepin applied as antipruritic control and comparator gave rise to inhibition of itch at a magnitude similar to 10% CA and 40% L-menthol. Listed in order of highest antipruritic effect

the counter-irritations: 4°C, 12°C, 10% CA, 22°C, 40% L-menthol, as well as the positive control, 5% doxepin, all gave rise to a total reduction in itch of more than 40% compared with respective control conditions. In addition, stimuli with 22°C, 12°C, but not 4°C, in addition to doxepin and L-menthol reduced the local neurogenic inflammatory histamine-evoked response.

#### Antipruritic effect of thermode-evoked cold and cold-like chemical stimulations

Few studies have attempted to assess or quantify the antipruritic effect of cold stimulation at different temperatures or chemical “cold-like” stimulation, despite the fact that cold-induced alleviation of itch is commonly reported in patients (4, 45). While it has frequently been observed that painful cold (or heat) stimulation results in inhibition of itch (46–48), the effect of non-painful thermode-induced cold counter-irritation or TRP-modulation by, for example, L-menthol, is less clear. Bromm et al. (29) found a significant decrease mediated by 1% L-menthol on histamine-induced itch; however, this finding was not reproduced in a subsequent study despite a 10-fold increase in L-menthol concentration. Similarly, studies have reported that both homotopic and heterotopic (in the same dermatome) innocuous cold-stimuli significantly attenuate histaminergic itch (21, 29), while 2 other studies, applying similar methodology, have reported that itch is aggravated by innocuous cold stimulations at 25°C (28, 30). The present data supports the prevailing notion that homotopic innocuous cold is sufficient to significantly reduce experimentally evoked itch. While cold analgesia of cutaneous and deep tissue pain is thought to rely on a combination of anti-nociceptive gating, decreased nerve conductivity, and diminished oedema, the antipruritic effect of cold-stimulation is probably less multifactorial. Since L-menthol does not change the actual temperature of the skin (required to affect neuronal conductivity), but decreases itch by a magnitude similar to thermal stimuli at 22°C and 12°C, it is likely that lateral inhibition from cold-encoding A $\delta$ -fibres is the primary mechanism behind the observed antipruritic effect. This proposition is also well supported by the fact that antipruritic efficacy slightly weaker than presented here has been observed when applying the thermal counter-stimulation 3-cm distally to the site of itch induction, hereby assuring limited interference from potential local inhibitory mechanisms (21).

In the present study, counter-irritation with 4°C and 10% CA were both found to evoke mild pain during the application period; an effect likely to be mediated by counter-irritancy through TRPA1-positive cold C-nociceptors (36, 49, 50). This is well in line with previous studies on the effect of painful counter-irritation of experimentally induced itch, wherein particularly painful mechanical stimuli (i.e. controlled scratching) and noxious

heat stimuli have been shown to significantly reduce itch (15, 20, 21). The mean of the observed CDT was on par with previously reported data for healthy controls (27, 40), while the mean CPT at 6.4°C was slightly below the usual reports of 10–13°C (27, 40). This is probably a consequence of CPT being a QST measure characterized by very high inter-individual variability (40) in combination with a relatively small sample size. A tendency towards a positive correlation was found between CPT and itch inhibition at 12°C ( $p=0.061$ ), indicating that individuals with high cold pain sensitivity, might be more susceptible to effective cold-induced itch relief.

#### *Vasomotor manifestations of thermode-evoked cold and cold-like stimulations*

For the thermal stimulations, a significant inhibitory effect on the neurogenic inflammatory response to histamine was found for stimulation at 22°C and 12°C, but not 4°C. Cutaneous cold-stimuli are well known to cause local vasoconstriction and, if sufficiently strong and persistent, subsequent transient paradoxical vasodilation after 3–4 min of stimulation. This, and the short delay between removal of the thermal probe and laser speckle contrast image recordings, could explain why cold-stimuli below the painful threshold (22°C and 12°C) appear to inhibit neurogenic inflammation, while stimulation above the pain threshold at 4°C, did not cause a significant decrease in neurogenic inflammation. In addition, painful cold stimulation is thought to entail TRPA1 activation, although this remains disputed (51), and TRPA1 exhibits co-expression with TRPV1 on peptidergic nociceptive C-fibres capable of generating axon-reflex flare (52, 53). This means that the paradoxical lack of inhibition of neurogenic inflammation at 4°C could be a consequence of responses being masked by additional recruitment of nociceptive C-units, beyond those CMi-fibres activated by histamine-introduction. Thermal stimulations  $\leq 22^\circ\text{C}$  stimulus-intensity-dependently decreased wheal reactions compared with the 32°C control condition. Wheal reactivity has not previously been assessed for a range of cold stimulations. Mechanistically, the observed vasoconstriction is probably mediated by a direct cold-induced smooth muscle contraction in the most superficial dermal capillaries (54).

As shown previously, the applied control, doxepin, drastically reduced neurogenic inflammation and wheal reactions, conceivably by direct inhibition of H1R on CMi-fibres (6, 43, 55, 56). L-menthol was also observed to reduce neurogenic inflammation and wheal formation, probably either via TRPM8-induced capillary constriction (57–59) or via potential TRPV1-cross-desensitization (60–63), given that TRPV1 is highly involved in downstream processing of histaminergic signalling (64). Interestingly, topical L-menthol, occasionally in much lower concentrations (above or equal to 1%), has

previously been found to reduce histaminergic and CA-induced neurogenic flare (27, 29); however, contradictory evidence does exist (24). The concentration of 10% CA used in the current study induced a stronger neurogenic inflammatory response than histamine, through direct activation of TRPA1-positive nociceptors (36, 65). On the other hand, CA profoundly reduced wheal formation and, in fact, abolished wheal formation completely in 5 subjects. The wheal reaction is based on localized dermal protein extravasation and is likely to wane when the applied histamine and extravasation products are sufficiently diluted by cutaneous blood perfusion (8). This process was conceivably augmented by the 5-fold increase in superficial blood perfusion induced by CA pre-treatment and a similar mechanism could be relevant for the slight, but significant, decrease in wheal size observed at 37°C.

#### *Study limitations*

High concentrations of topically applied ethanol are suspected to entail minor somatosensory, but not neurogenic, inflammatory changes, although previously published results are conflicting (27, 36, 66) and the supposed mechanism is unclear (67). The present study did not include a vehicle condition for application of the chemical substances, for the following reasons: (i) even in the few studies that do find somatosensory changes following ethanol application, the effect is very subtle (36, 68); (ii) there is no spontaneous sensation associated with ethanol when applied as in the present study (12, 68); and (iii) a previous study failed to find any effect of 80% ethanol on histaminergic itch (24). Nonetheless, it cannot be excluded that topical ethanol and the vehicle cream for application of doxepin have minor antipruritic effects, especially since ethanol is a TRPV1 agonist (69). Similarly, although the 32°C stimulation was chosen because it is known to be an approximation of the physiologically normal skin temperature, a significant decrease in itch was observed compared with histamine application without thermal counter-irritation, presumably due to the ongoing mechanical counter-stimulus from the thermal stimulator probe. Ideally, cold counter-irritation should be applied without concomitant activation of mechanoreceptive A $\beta$ -fibres, but this remains a technological challenge. CA produced spontaneous burning/pricking pain and extensive neurogenic inflammation in a number of subjects, which is known to drastically reduce homotopic itch. Future studies should include a titration of CA to a concentration at which it provides somatosensory counter-irritation without provoking spontaneous pain and neurogenic inflammation. In this context, it is a concern that lower concentrations of topical CA alone have previously been shown to have pruritogenic effects in humans (12). It should be kept in mind that itch and pain sensations were not quantified similarly in the present study; while itch was scored continuously as it occurred, subjects had to



recall the pain intensity of the previously applied counter-irritation stimuli, immediately after the stimuli had ended. As a result, comparisons can be made within the resulting itch and pain scores, but not across the sensory modalities. Lastly, the present study applied a histaminergic human surrogate model of itch conveyed by a subset of pruriceptive CMI-fibres. Future studies should ascertain whether itch induced by a non-histaminergic model, such as the mucunain-model (cowhage) (8, 12), and thus activating PmC-fibres, also known to transmit pain under certain circumstances (70), would be susceptible to effective cold-induced itch suppression.

### Conclusion

The thermal cold-stimuli applied in this study inhibited histaminergic itch in a stimulus-intensity-dependent manner of which the painful thermode-provoked stimulations had the highest antipruritic effect. A similar inhibitory tendency was observed on wheal reactions and neurogenic inflammation. In addition, the applied chemical proxies of cold-stimuli, L-menthol and trans-cinnamaldehyde, had significant inhibitory effects on histaminergic itch with an antipruritic effect-size comparable to that of 5% topical doxepin. Thermal counter-stimuli or selective and potent TRPA1-/TRPM8-agonists, titrated to concentrations evoking counter-irritation at sub-pain threshold levels could be potential antipruritic treatment modalities, particularly for those chronic itch patients reporting itch alleviation by cooling.

### ACKNOWLEDGEMENTS

The authors would like to thank Jacob B. Pedersen for assistance in the experimental procedures, and all volunteer participants.

*The authors declare no conflicts of interest.*

### REFERENCES

1. Ständer S, Weisshaar E, Mettang T, Szepietowski J, Carstens E, Ikoma A, et al. Clinical classification of itch: a position paper of the International Forum for the Study of Itch. *Acta Derm Venereol* 2007; 87: 291–294.
2. Weisshaar E, Gieler U, Kupfer J, Furue M, Saeki H, Yosipovitch G. Questionnaires to assess chronic itch: a consensus paper of the Special Interest Group of the International Forum on the Study of Itch. *Acta Derm Venereol* 2012; 92: 493–496.
3. Misery L, Brenaut E, Le Garrec R, Abasq C, Genestet S, Marcorelles P, et al. Neuropathic pruritus. *Nat Rev Neurol* 2014; 10: 408–416.
4. Patel T, Yosipovitch G. Therapy of pruritus. *Expert Opin Pharmacother* 2010; 11: 1673–1682.
5. Mattered U, Apfelbacher CJ, Loerbroks A, Schwarzer T, Büttner M, Ofenloch R, et al. Prevalence, correlates and characteristics of chronic pruritus: a population-based cross-sectional study. *Acta Derm Venereol* 2011; 91: 674–679.
6. Johaneck LM, Meyer RA, Hartke T, Hobelmann JG, Maine DN, LaMotte RH, et al. Psychophysical and physiological evidence for parallel afferent pathways mediating the sensation of itch. *J Neurosci* 2007; 27: 7490–7497.
7. Sun Y-G, Chen Z-F. A gastrin-releasing peptide receptor mediates the itch sensation in the spinal cord. *Nature* 2007; 448: 700–703.
8. Andersen HH, Elberling J, Arendt-Nielsen L. Human surrogate models of histaminergic and non-histaminergic itch. *Acta Derm Venereol* 2015; 95: 771–777.
9. Papoiu ADP, Coghill RC, Kraft RA, Wang H, Yosipovitch G. A tale of two itches. Common features and notable differences in brain activation evoked by cowhage and histamine induced itch. *Neuroimage* 2012; 59: 3611–3623.
10. Simone D a, Alreja M, LaMotte RH. Psychophysical studies of the itch sensation and itchy skin (“alloknesis”) produced by intracutaneous injection of histamine. *Somatosens Mot Res* 1991; 8: 271–279.
11. Darsow U, Scharein E, Simon D, Walter G, Bromm B, Ring J. New aspects of itch pathophysiology: component analysis of atopic itch using the “Eppendorf Itch Questionnaire.” *Int Arch Allergy Immunol* 2001; 124: 326–331.
12. Højland CR, Andersen HH, Poulsen JN, Arendt-nielsen L, Gazerani P. A human surrogate model of itch utilizing the TRPA1 agonist trans-cinnamaldehyde. *Acta Derm Venereol* 2015; 95: 798–803.
13. Ernst E, Fialka V. Ice freezes pain? A review of the clinical effectiveness of analgesic cold therapy. *J Pain Symptom Manage* 1994; 9: 56–59.
14. Leknes SG, Bantick S, Willis CM, Wilkinson JD, Wise RG, Tracey I. Itch and motivation to scratch: an investigation of the central and peripheral correlates of allergen- and histamine-induced itch in humans. *J Neurophysiol* 2007; 97: 415–422.
15. Kosteletzky F, Namer B, Forster C, Handwerker H. Impact of scratching on itch and sympathetic reflexes induced by cowhage (*Mucuna pruriens*) and histamine. *Acta Derm Venereol* 2009; 89: 271–277.
16. Heyer G, Ulmer FJ, Schmitz J, Handwerker HO. Histamine-induced itch and aloknesis (itchy skin) in atopic eczema patients and controls. *Acta Derm Venereol* 1995; 75: 348–352.
17. Ma Q. Population coding of somatic sensations. *Neurosci Bull* 2012; 28: 91–99.
18. Dhand A, Aminoff MJ. The neurology of itch. *Brain* 2013; 137: 313–322.
19. Pall PS, Hurwitz OE, King BA, LaMotte RH. Psychophysical measurements of itch and nociceptive sensations in an experimental model of allergic contact dermatitis. *J Pain* 2015; 16: 741–749.
20. Yosipovitch G, Fast K, Bernhard JD. Noxious heat and scratching decrease histamine-induced itch and skin blood flow. *J Invest Dermatol* 2005; 125: 1268–1272.
21. Yosipovitch G, Duque MI, Fast K, Dawn a. G, Coghill RC. Scratching and noxious heat stimuli inhibit itch in humans: a psychophysical study. *Br J Dermatol* 2007; 156: 629–634.
22. Brenaut E, Garlantezec R, Talour K, Misery L. Itch characteristics in five dermatoses: Non-atopic eczema, atopic dermatitis, urticaria, psoriasis and scabies. *Acta Derm Venereol* 2013; 93: 573–574.
23. Brenaut E, Marcorelles P, Genestet S, Ménard D, Misery L. Pruritus: an underrecognized symptom of small-fiber neuropathies. *J Am Acad Dermatol* 2015; 72: 328–332.
24. Yosipovitch G, Szolar C, Hui XY, Maibach H. Effect of topically applied menthol on thermal, pain and itch sensations and biophysical properties of the skin. *Arch Dermatol Res* 1996; 288: 245–248.
25. O’Neill JL, Chan YH, Rapp SR, Yosipovitch G. Differences in itch characteristics between psoriasis and atopic dermatitis patients: results of a web-based questionnaire. *Acta Derm Venereol* 2011; 91: 537–540.
26. Proudfoot CJ, Garry EM, Cottrell DF, Rosie R, Anderson H, Robertson DC, et al. Analgesia mediated by the TRPM8 cold receptor in chronic neuropathic pain. *Curr Biol* 2006; 16: 1591–1605.
27. Olsen RV, Andersen HH, Møller HG, Eskelund PW, Arendt-Nielsen L. Somatosensory and vasomotor manifestations of individual and combined stimulation of TRPM8 and TRPA1 using topical L-menthol and trans-cinnamaldehyde in healthy volunteers. *Eur J Pain* 2014; 18: 1333–1342.
28. Valet M, Pfab F, Sprenger T, Wöller A, Zimmer C, Behrendt H, et al. Cerebral processing of histamine-induced itch using



- short-term alternating temperature modulation – an fMRI study. *J Invest Dermatol* 2008; 128: 426–433.
29. Bromm B, Scharein E, Darsow U, Ring J. Effects of menthol and cold on histamine-induced itch and skin reactions in man. *Neurosci Lett* 1995; 187: 157–160.
  30. Pfab F, Valet M, Sprenger T, Toelle TR, Athanasiadis GI, Behrendt H, et al. Short-term alternating temperature enhances histamine-induced itch: a biphasic stimulus model. *J Invest Dermatol* 2006; 126: 2673–2678.
  31. Zhang X. Targeting TRP ion channels for itch relief. *Naunyn Schmiedeberg Arch Pharmacol* 2015; 388: 389–399.
  32. Dhaka A, Viswanath V, Patapoutian A. Trp ion channels and temperature sensation. *Annu Rev Neurosci* 2006; 29: 135–161.
  33. Zheng J. Molecular mechanism of TRP channels. *Compr Physiol* 2013; 3: 221–242.
  34. Lucaciu OC, Connell GP. Itch sensation through transient receptor potential channels: a systematic review and relevance to manual therapy. *J Manipulative Physiol Ther* 2013; 36: 385–393.
  35. Wei JJ, Fu T, Yang T, Liu Y, Wang GR. A TRPA1 channel that senses thermal stimulus and irritating chemicals in *Helicoverpa armigera*. *Insect Mol Biol* 2015; 24: 412–421.
  36. Namer B, Seifert F, Handwerker H, Maihöfner C. TRPA1 and TRPM8 activation in humans: effects of cinnamaldehyde and menthol. *Neuroreport* 2005; 16: 955–959.
  37. Macpherson LJ, Hwang SW, Miyamoto T, Dubin AE, Patapoutian A, Story GM. More than cool: promiscuous relationships of menthol and other sensory compounds. *Mol Cell Neurosci* 2006; 32: 335–343.
  38. Peier AM, Moqrich A, Hergarden AC, Reeve AJ, Andersson DA, Story GM, et al. A TRP channel that senses cold stimuli and menthol. *Cell* 2002; 108: 705–715.
  39. World Medical A. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA* 2013; 310: 2191–2194.
  40. Rolke R, Baron R, Maier C, Tölle TR, Treede RD, Beyer A, et al. Quantitative sensory testing in the German Research Network on Neuropathic Pain (DFNS): Standardized protocol and reference values. *Pain* 2006; 123: 231–243.
  41. Andersen HH, Sørensen A-KR, Nielsen GAR, Mølgård MS, Stilling P, Elberling J, et al. A reliability assessment of standardized human surrogate models of histaminergic and non-histaminergic itch using histamine and cowhage spicules. *Acta Derm Venereol* 2015; 95: 906.
  42. Gazerani P, Pedersen NS, Drewes AM, Arendt-Nielsen L. Botulinum toxin type A reduces histamine-induced itch and vasomotor responses in human skin. *Br J Dermatol* 2009; 161: 737–745.
  43. Sikand P, Dong X, LaMotte RH. BAM8-22 peptide produces itch and nociceptive sensations in humans independent of histamine release. *J Neurosci* 2011; 31: 7563–7567.
  44. Namer B, Carr R, Johaneck LM, Schmelz M, Handwerker HO, Ringkamp M. Separate peripheral pathways for pruritus in man. *J Neurophysiol* 2008; 100: 2062–2069.
  45. Yosipovitch G. Pruritus. *Curr Probl Dermatol* 2003; 15: 143–164.
  46. Fruhstorfer H, Hermanns M, Latzke L. The effects of thermal stimulation on clinical and experimental itch. *Pain* 1986; 24: 259–269.
  47. Murray FS, Weaver MM. Effects of ipsilateral and contralateral counterirritation on experimentally produced itch in human beings. *J Comp Physiol Psychol* 1975; 89: 819–826.
  48. Ward L, Wright E, McMahon SB. A comparison of the effects of noxious and innocuous counterstimuli on experimentally induced itch and pain. *Pain* 1996; 64: 129–138.
  49. Andersen HH, Olsen RV, Møller HG, Eskelund PW, Gazerani P, Arendt-Nielsen L. A review of topical high-concentration L-menthol as a translational model of cold allodynia and hyperalgesia. *Eur J Pain* 2013; 18: 315–325.
  50. McKemy DD. How cold is it? TRPM8 and TRPA1 in the molecular logic of cold sensation. *Mol Pain* 2005; 1: 16.
  51. McKemy DD. The molecular and cellular basis of cold sensation. *ACS Chem Neurosci* 2013; 4: 238–247.
  52. Schmelz M, Michael K, Weidner C, Torebjörk H, Handwerker H. Which nerve fibers mediate the axon reflex flare in human skin? *Neuroreport* 2000; 11: 645–648.
  53. Zylka MJ, Rice FL, Anderson DJ. Topographically distinct epidermal nociceptive circuits revealed by axonal tracers targeted to Mrgprd. *Neuron* 2005; 45: 17–25.
  54. Thompson-Torgerson CS, Holowatz LA, Flavahan NA, Larry Kenney W. Rho kinase-mediated local cold-induced cutaneous vasoconstriction is augmented in aged human skin. *AJP Hear Circ Physiol* 2007; 293: 30–36.
  55. Abila B, Ezeamuzie IC, Igbigbi PS, Ambakederemo AW, Asomugha L. Effects of two antihistamines on chloroquine and histamine induced weal and flare in healthy African volunteers. *Afr J Med Med Sci* 1994; 23: 139–142.
  56. Hägermark O. Influence of antihistamines, sedatives, and aspirin on experimental itch. *Acta Derm Venereol* 1973; 53: 363–368.
  57. Yang XR, Lin MJ, McIntosh LS, Sham JSK. Functional expression of transient receptor potential melastatin- and vanilloid-related channels in pulmonary arterial and aortic smooth muscle. *Am J Physiol Lung Cell Mol Physiol* 2006; 290: 1267–1276.
  58. Johnson CD, Melanaphy D, Purse A, Stokesberry SA, Dickson P, Zholos A V. Transient receptor potential melastatin 8 channel involvement in the regulation of vascular tone. *AJP Hear Circ Physiol* 2009; 296: 1868–1877.
  59. Topp R, Winchester L, Mink AM, Kaufman JS, Jacks DE. Comparison of the effects of ice and 3.5% menthol gel on blood flow and muscle strength of the lower arm. *J Sport Rehabil* 2011; 20: 355–366.
  60. Dessirier JM, O'Mahony M, Carstens E. Oral irritant properties of menthol: sensitizing and desensitizing effects of repeated application and cross-desensitization to nicotine. *Physiol Behav* 2001; 73: 25–36.
  61. Albin KC, Carstens MI, Carstens E. Modulation of oral heat and cold pain by irritant chemicals. *Chem Senses* 2008; 33: 3–15.
  62. Cliff MA, Green BG. Sensitization and desensitization to capsaicin and menthol in the oral cavity: interactions and individual differences. *Physiol Behav* 1996; 59: 487–494.
  63. Green BG, McAuliffe BL. Menthol desensitization of capsaicin irritation. Evidence of a short-term anti-nociceptive effect. *Physiol Behav* 2000; 68: 631–639.
  64. Shim WS, Tak MH, Lee MH, Kim M, Koo JY, et al. TRPV1 mediates histamine-induced itching via the activation of phospholipase A2 and 12-lipoxygenase. *J Neurosci* 2007; 27: 2331–2337.
  65. Eid SR, Crown ED, Moore EL, Liang HA, Choong K-C, Dima S, et al. HC-030031, a TRPA1 selective antagonist, attenuates inflammatory- and neuropathy-induced mechanical hypersensitivity. *Mol Pain* 2008; 4: 48.
  66. Wasner G, Schattschneider J, Binder A, Baron R. Topical menthol – a human model for cold pain by activation and sensitization of C nociceptors. *Brain* 2004; 127: 1159–1171.
  67. Löffler H, Kampf G. Hand disinfection: how irritant are alcohols? *J Hosp Infect* 2008; 70 Suppl 1: 44–48.
  68. Namer B, Kleggetveit IP, Handwerker H, Schmelz M, Jorum E. Role of TRPM8 and TRPA1 for cold allodynia in patients with cold injury. *Pain* 2008; 139: 63–72.
  69. Trevisani M, Smart D, Gunthorpe MJ, Tognetto M, Barbieri M, Campi B, et al. Ethanol elicits and potentiates nociceptor responses via the vanilloid receptor-1. *Nat Neurosci* 2002; 5: 546–551.
  70. LaMotte RH, Dong X, Ringkamp M. Sensory neurons and circuits mediating itch. *Nat Rev Neurosci* 2014; 15: 19–31.