



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Preoperative pain mechanisms assessed by cuff algometry are associated with chronic postoperative pain relief after total knee replacement

Petersen, Kristian Kjær; Graven-Nielsen, Thomas; Simonsen, Ole; Laursen, Mogens Berg; Arendt-Nielsen, Lars

Published in:
Pain

DOI (link to publication from Publisher):
[10.1097/j.pain.0000000000000531](https://doi.org/10.1097/j.pain.0000000000000531)

Publication date:
2016

Document Version
Peer reviewed version

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Petersen, K. K., Graven-Nielsen, T., Simonsen, O., Laursen, M. B., & Arendt-Nielsen, L. (2016). Preoperative pain mechanisms assessed by cuff algometry are associated with chronic postoperative pain relief after total knee replacement. *Pain*, 157(7), 1400-1406. DOI: 10.1097/j.pain.0000000000000531

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- ? Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- ? You may not further distribute the material or use it for any profit-making activity or commercial gain
- ? You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

Chronic postoperative pain following total knee replacement (TKR) in knee osteoarthritis (KOA) implies clinical challenges. Widespread hyperalgesia, facilitated temporal summation of pain (TSP), and impaired conditioning pain modulation (CPM) have been found in painful KOA. **This exploratory study investigated postoperative pain relief 12 months after TKR in 4 sub-groups of patients preoperatively profiled by mechanistic quantitative sensory testing.**

In 103 KOA patients pressure pain detection and tolerance thresholds (PDT, PTT) were assessed at the lower leg using cuff algometry. TSP was measured as an increase in pain intensity scores during 10 repeated (2 seconds intervals) painful cuff stimuli. CPM was calculated as the relative increase in PDT during painful conditioning stimulation. The grand averages of TSP and CPM were calculated and values below or above were used for sub-grouping. **Facilitated TSP/impaired CPM (Group-A, N=16), facilitated TSP/normal CPM (Group-B, N=15), normal TSP/impaired CPM (Group-C, N=44), and normal TSP/normal CPM (Group-D, N=28).** Clinical VAS pain intensity score were collected before and 12 months after TKR-surgery **and the pain relief calculated. Less pain relief was found in Group A (52.0±14.0% pain relief) compared with Group B (81.1±3.5%, P=0.023) and Group C (79.6±4.4%, P=0.007), but not Group D (69.4±7.9%, P=0.087). Low preoperative PDT was associated with a less postoperative pain relief (R=-0.222, P=0.034) whereas TSP or CPM alone showed no associations with postoperative pain relief.**

This explorative study indicated that OA patients with facilitated TSP together with impaired CPM are more vulnerable to experience less pain relief after TKR.

1 **Preoperative pain mechanisms assessed by cuff algometry are associated with chronic**
2
3
4 **postoperative pain relief after total knee replacement**
5
6

7 Kristian Kjær Petersen¹, Thomas Graven-Nielsen², Ole Simonsen³, Mogens Berg Laursen³, Lars
8
9 Arendt-Nielsen^{1*}
10

11
12 (1) SMI, Department of Health Science and Technology, Faculty of Medicine, Aalborg University,
13
14 Aalborg, Denmark
15

16
17 (2) Center for Neuroplasticity and Pain (CNAP), SMI, Department of Health Science and
18
19 Technology, Faculty of Medicine, Aalborg University, Aalborg, Denmark
20

21 (3) Aalborg University Hospital, Orthopaedic Surgery Research Unit, Aalborg, Denmark
22
23
24

25 **Original paper for: Pain**
26

27 **Article type: Clinical/Basic Science Research Report**
28

29 **Number of Figures and Tables: 2 figures and 2 tables**
30

31 **Number of Pages: 26 (27 including abstract)**
32
33

34 **Keywords:** Osteoarthritis, conditioned pain modulation, temporal summation, chronic
35
36 postoperative pain relief, total knee replacement.
37
38
39
40
41
42

43 ***Corresponding Author:**
44

45 Prof. Lars Arendt-Nielsen, Ph.D., DMSc.
46

47 SMI
48

49 Department of Health Science and Technology
50

51 Faculty of Medicine, Aalborg University
52

53 Fredrik Bajers Vej 7 D3, DK-9220 Aalborg, Denmark
54

55 Phone:+45 9940 8831, Fax:+45 9815 4008, E-mail: LAN@HST.AAU.DK
56
57
58
59
60
61
62

1. Introduction

Patients with osteoarthritis (OA) may eventually suffer from chronic pain leading to disabilities of the individual and associated costs for public health care systems [20]. OA is the most frequent painful musculoskeletal condition in the elderly population [26]. The end-stage treatment of knee OA (KOA) is total knee replacement (TKR) of which approximately 500,000 are performed in the US every year, and this number is estimated to increase sevenfold by 2030 [41]. Evidence suggests that around 20% of KOA patients will develop chronic postoperative pain after TKR [7,29].

Quantitative sensory testing (QST) is widely used to characterise the underlying mechanisms of KOA pain [3,4,13,30]. KOA patients have decreased pressure pain thresholds (PPTs) around the knee and at extra-segmental sites compared with healthy controls [3,4,13] known as local and widespread hyperalgesia [14]. Patients with chronic e.g. OA [3,4,13], fibromyalgia [15], or low back pain [19] show facilitated pain responses to repeated painful stimulation (temporal summation of pain:TSP) compared with controls. This indicates that the central integrative mechanisms are facilitated potentially due to sensitization processes. A measure for the status of the descending pain control system is conditioned pain modulation (CPM) [5] which is impaired in patients with, e.g. KOA [3,4,13], fibromyalgia [21,23], chronic tension-type headache [36], or chronic pancreatitis [27].

Cuff algometry is widely used for assessment of the deep somatic tissue pain sensitivity [26] in addition to single-point algometry which stimulates more localized superficial somatic structures [12]. Computer controlled cuff algometry has previously been used to assess PPT, TSP, and CPM [4,13,33,40] and offers the advantage of being user independent [33] and has a good test-retest reliability [16].

1 Accumulated findings suggest that preoperative pain and sensitization play a role in the
2
3 development of chronic postoperative pain. Impaired CPM before thoracotomy [49] and
4
5 abdominal [47] surgery has shown to be predictive for the risk of developing chronic postoperative
6
7 pain. Recently, it was demonstrated that patients who developed severe chronic postoperative
8
9 pain after TKR had a fourfold increase in preoperative TSP compared with patients who did not
10
11 develop chronic pain [30]. Another study showed that the degree of widespread hyperalgesia
12
13 predicted postoperative pain in patients after total hip replacement [48]. Subgrouping KOA
14
15 patients based on radiographic characterization and pain reveals different sub-groups [10]
16
17 emphasizing that subgrouping of patients may play a major role in understanding the involvement
18
19 of different pain mechanisms in KOA and hence possible how vulnerable they are to develop
20
21 chronic pain after surgery. No studies have investigated the possible value of combined
22
23 preoperative **mechanistic QST parameters (e.g. TSP and CPM)** in KOA patients before TKR and the
24
25 associations to chronic postoperative pain outcomes.
26
27
28
29
30
31
32
33
34

35 The aims of this exploratory study were (1) preoperative profiling of KOA patients based on
36
37 spreading hyperalgesia, TSP, and CPM characteristics, (2) to investigate **4 sub-groups of patients**
38
39 **preoperatively profiled by mechanistic quantitative sensory testing (TSP, and CPM)** and the
40
41 relation to chronic postoperative **pain relief 12 months after TKR**, and (3) to investigate the
42
43 associations between preoperative widespread hyperalgesia and the development of the chronic
44
45 postoperative **pain relief**.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2. Methods

2.1 Patients

KOA patients (N=135) scheduled for unilateral TKR from the outpatient clinic at Hospital Vendsyssel, Frederikshavn, Denmark, were invited to join the study. Radiological KOA progression was evaluated using the Kellgren & Lawrence (KL) score [22]. Patients with other diagnosed pain problems (e.g., hip OA, rheumatoid arthritis, fibromyalgia, neuropathic pain), sensory dysfunction, or mental impairment were excluded from the study. The study was approved by The North Denmark Region Committee on Health Research Ethics (N-20120015) and conducted in accordance with the Helsinki Declaration. All patients read and signed an informed consent. This cohort of patients has not previous been included in any other scientific publications.

2.2 Protocol

The peak pain intensity within the last 24 hours (visual analogue scale, VAS), body mass index (BMI), and QST recordings (pressure pain sensitivity, TSP and CPM) were collected before surgery by handheld pressure algometry and cuff algometry. The patients were asked not to take any analgesic medication 24 hours prior to the examination. Moreover, the patients were contacted 12 months after surgery to collect the VAS score of the peak pain intensity within the last 24 hours (VAS). Pain relief was calculated as the percentage difference between pre- and postoperative VAS and used as the main outcome measure.

2.3 Cuff Algometry

Deep-tissue pain sensitivity was evaluated by cuff pressure stimuli using a computer-controlled cuff algometer (NociTech and Aalborg University, Denmark) including a 13-cm wide tourniquet cuff (VBM, Sulz, Germany) and an electronic VAS (Aalborg University, Denmark) for recording of

1 the pain intensity. The cuff was placed at the level of the head of the gastrocnemius muscle of the
2
3 leg most affected by KOA. The electronic continuous VAS (sliding resistor) was 10 cm long and
4
5 sampled at 10 Hz; 0 cm indicated “no pain” and 10 cm indicated “maximum pain”.
6
7

8 9 ***Pressure Detection and Tolerance Threshold***

10
11
12 The pressure was increased by 1 kPa/s and the patient was instructed to rate the pain intensity
13
14 continuously on the electronic VAS until the tolerance level was reached and, further, the patient
15
16 was instructed to press a stop button after which the pressure was released immediately. The
17
18 pressure pain detection threshold (PDT) was defined as the pressure where the VAS score
19
20 exceeded 2 cm^[43] and the pain tolerance threshold (PTT) was defined when the patient pressed
21
22 the stop button. The measures were repeated three times and the average was used for further
23
24 analysis.
25
26
27
28
29
30

31 32 ***Temporal Summation of Pain***

33
34
35 The concept of temporal summation is an increase in the pain ratings during a series of e.g. 10
36
37 identical painful stimuli delivered at a rate of e.g. 0.5 Hz. The automatic cuff algometer was used
38
39 to elicit temporal summation [16].
40
41
42

43
44 A total of 10 repeated mechanical pressure stimuli at the average of the PDT and PTT levels were
45
46 delivered at 0.5 Hz (1 s stimulus duration and 1 s interval between stimuli) to the lower leg. A
47
48 constant pressure between the individual pressure stimuli of 1 kPa was applied to avoid
49
50 movement of the cuff. During the 10 repeated stimuli, the patients rated continuously the pain
51
52 intensity on a 10 cm continuous VAS (sliding resistor) (“0” represented “no pain”, and “10”
53
54 represented “maximal pain”).
55
56
57
58
59
60
61

1 For analysis of TSP, the mean VAS score was calculated in the interval from the first to the end of
2
3 the 4th stimulus (VAS-I) and in the interval from the 8th to the end of the 10th stimulus (VAS-II). TSP
4
5 was defined as the difference between VAS-I and VAS-II (i.e. VAS-II minus VAS-I) [44].
6
7

8 9 **Conditioned Pain Modulation**

10
11
12
13 The concept of CPM is that a tonic painful stimulus (conditioning stimulus) will inhibit pain evoked
14
15 simultaneously from another site (test stimulus).
16

17
18
19 The painful conditioned stimulus was initially set to the level of 60 kPa as this value was found
20
21 equivalent to a general pain perception of 5 cm on the VAS in a preliminary assessment. If not
22
23 tolerated, the conditioned stimulus was reduced to 30 kPa. Simultaneously, assessment of PDT
24
25 was performed using a single chamber cuff on the ipsilateral lower leg (test stimulus). The
26
27 conditioned stimulus was terminated right after the PDT was assessed. CPM was defined as the
28
29 difference between PDT during and before conditioned pain (i.e. “during” minus “before”).
30
31
32
33
34

35 **2.4 Handheld Algometry**

36
37
38 A handheld algometer (Somedic AB, Sweden) was used for measuring PPT. A 1-cm² probe was
39
40 used and placed perpendicularly to the skin. The pressure was applied at 30 kPa/s until the patient
41
42 identified the pressure as pain and pressed a button. The algometry was performed on 7 sites at
43
44 the most affected knee with two distant sites at the tibialis anterior muscle (TA, 5 cm distal to the
45
46 tibial tuberosity) and at the extensor carpi radialis longus muscle (arm, 5 cm distal to the lateral
47
48 epicondyle of humerus). The sites in the peripatellar regions were: 2 cm distal to the inferior
49
50 medial edge of patella (Site 1); 2 cm distal to the inferior lateral edge of patella (Site 2); 3 cm
51
52 lateral to the midpoint of the lateral edge of patella (Site 3); 2 cm proximal to the superior lateral
53
54 edge of patella (Site 4); 2 cm proximal to the superior edge of patella (Site 5); 2 cm proximal to the
55
56
57
58
59
60
61
62
63
64
65

1 superior medial edge of patella (Site 6); 3 cm medial to the midpoint of the medial edge of patella
2
3
4 (Site 7). An average of the seven peripatellar PPTs was used for further analysis [3].
5
6

7 **2.5 Statistics**

8
9
10 To phenotype patients with a high degree of central sensitization, i.e., facilitated TSP and impaired
11
12 CPM, the patients were arbitrarily divided into four groups based on a mean cut-off splits of
13
14 preoperative TSP and CPM defined by facilitated TSP and impaired CPM (Group A), facilitated TSP
15
16 and normal (defined as the average of the group) CPM (Group B), normal TSP and impaired CPM
17
18 (Group C), and normal TSP and normal CPM (Group D). Later studies should be designed to
19
20 optimize these cut-off splits. The data are presented as mean and standard error of the mean
21
22 (SEM) if not otherwise stated. A one-way analysis of variance (ANOVA) was performed to compare
23
24 preoperative parameters and the postoperative pain relief for the four groups. The Fisher post-hoc
25
26 test was used in case of significant differences. Pearson's correlation was used for correlation
27
28 analysis. Linear regression, including significant correlating parameters from the Pearson's
29
30 correlations, was used to categorize independent parameters. A previous study showed that cuff
31
32 measurement in KOA patients is age-dependent, for which reason all analyses were adjusted for
33
34 age [11]. $P < 0.05$ was considered significant.
35
36
37
38
39
40
41
42
43

44 **3. Results**

45 **3.1 Demographics**

46
47
48 Of the 135 patients recruited, 32 patients (24%) were lost at the 12 month follow-up where three
49
50 patients could not be reached, two patients had undergone major surgery within the last month,
51
52 one patient had developed Alzheimer, and 26 patients did not attend the follow-up despite
53
54
55
56
57
58
59
60
61
62
63
64
65

1 several phone calls. In total 103 patients were included in the current analysis. All patients had
2
3 technically successful TKRs at follow-up.
4
5

6
7 Four groups were initially suggested based on the arbitrary QST cut-off levels below or above the
8
9 grand average of TSP (1.55 ± 0.17 cm) and CPM (5.40 ± 1.05 kPa), respectively.
10
11

12
13 Group A: facilitated TSP and impaired CPM (N=16).
14
15

16
17 Group B: facilitated TSP and normal CPM (N=15).
18
19

20
21 Group C: normal TSP and impaired CPM (N=44).
22
23

24
25 Group D: normal TSP and normal CPM (N=28).
26
27

28
29 Group B showed higher BMI compared with Group D (ANOVA: $F=4.772$, $P<0.002$, Fisher: $P=0.020$)
30 (Table 1). Otherwise the groups were identical.
31
32

33 **3.2 Postoperative Pain Relief**

34
35

36 No difference between preoperative VAS scores was found among the four groups (Table 1;
37
38 ANOVA: $F=1.86$, $P=0.13$). Group A had significantly (Table 1; ANOVA: $F=2.55$, $P<0.045$) less
39
40 postoperative pain relief as compared with Group B (Fisher: $P=0.023$) and C (Fisher: $P=0.007$) but
41
42 not with Group D (Fisher: $P=0.087$).
43
44
45

46 **3.3 Cuff Algometry**

47
48

49 *Pressure Pain Detection and Tolerance Thresholds*

50
51

52 Group D showed significantly higher PDT values compared with Group A and Group B (Figure 1;
53
54 ANOVA: $F=3.62$, $P<0.005$, Fisher: $P<0.016$) and higher PTT values compared with Group A (ANOVA:
55
56 $F=2.86$, $P<0.028$, Fisher: $P<0.027$).
57
58
59
60
61

Temporal Summation of Pain

A significant interaction was found between groups and TSP (ANOVA: $F=52.53$, $P<0.001$) showing less TSP in Group B (Fishers, $P=0.002$), Group C (Fishers, $P<0.001$) and Group D (Fishers, $P<0.001$) compared with Group A, and less TSP in Group C (Fishers, $P<0.001$) and Group D (Fishers, $P<0.001$) compared with Group B (Figure 1). **TSP alone did not predict postoperative pain relief.**

Conditioned Pain Modulation

Group A and Group C showed significantly impaired CPM compared with Group B and Group D (ANOVA: $F=25.6$, $P<0.001$, Fisher: $P<0.001$), but no significant difference was found between the groups with facilitated and normal TSP (i.e., Group A and Group C or Group B and Group D) (

Figure 1. **CPM alone did not predict postoperative pain relief.**

3.4 Handheld Pressure Algometry

No significant difference in PPTs was found between the groups in the peripatellar area (Figure 2; ANOVA: $F=0.59$, $P=0.67$), TA (ANOVA: $F=1.41$, $P=0.24$), or on the arm (ANOVA: $F=0.49$, $P=0.75$).

PPTs alone did not predict postoperative pain relief.

3.5 Correlations

A positive correlation was found between postoperative **pain relief** and preoperative pain ($R=0.241$, $P=0.009$). A significant negative correlation was found between postoperative **pain relief** and preoperative PDT ($R=-0.216$, $P=0.021$). A linear stepwise regression analysis, including preoperative VAS and PDT (Table 2), was performed to investigate the possible prediction of postoperative **pain relief**. The analysis showed that PDT was an independent parameter ($R=-0.222$, $P=0.034$) and a trend towards an independent parameter was found for preoperative pain

1 (R=0.263, P=0.080). As such PDT, independently of preoperative pain, could predict postoperative
2
3
4 **pain relief.**
5

6
7 **No significant correlations between the postoperative pain relief and the preoperative TSP, CPM,**
8
9 **or handheld algometry, respectively, were found.**
10

11 12 13 14 **4. Discussion**

15 This **preliminary** study showed that preoperative subgrouping of KOA patients based on TSP and
16
17
18 CPM can identify vulnerable groups experiencing less postoperative **pain relief** 12 months after
19
20
21 TKR. Reduced preoperative pressure pain detection thresholds at the lower leg were as a single
22
23
24 parameter associated with **pain relief** 12 months after TKR.
25

26 27 **4.1 Preoperative Pain Biomarkers and associations for Postoperative Pain Relief**

28
29 Sensitization in OA is often characterized by widespread hyperalgesia, impaired descending pain
30
31 control, and increased facilitation of temporal summation [6,25,42]. As in the present study
32
33
34 preoperative widespread hyperalgesia has been shown to predict postoperative pain outcome
35
36
37 after total hip replacement [48]. Yarnitsky et al. [49] suggested that impaired CPM was associated
38
39
40 with a higher risk of chronic pain after thoracotomy and Wilder-Smith et al. [47] showed similar
41
42
43 results in patients after abdominal surgery. Preoperatively facilitated TSP has been shown to
44
45
46 predict chronic postoperative pain after abdominal surgery [46] and, recently also after TKR [30].
47
48
49 An increasing number of studies suggest that subgrouping of KOA patients based on clinical and
50
51
52 experimental pain assessment parameters can reveal groups of specifically sensitized patients
53
54
55 [2,3,10] which could be the patients more vulnerable to experience less pain relief after TKR. The
56
57
58 present study confirmed this suggestion and found that subgrouping KOA patients based on
59
60
61 preoperatively facilitated TSP and impaired CPM revealed a sub-group of KOA patients with the
62
63
64
65

1 least pain relief after TKR. This group had less postoperative pain relief as compared with patients
2
3 showing either showing 1) facilitated TSP but normal CPM or 2) impaired CPM but normal TSP.
4
5 This indicates that a preoperative multimodal pain mechanistic QST approach is important for
6
7 predicting the chronic postoperative pain outcome. Drugs designed to block the N-methyl-D-
8
9 aspartate (NMDA) receptors have shown to reduce TSP [34,45], and drugs promoting the re-
10
11 uptake of serotonin and norepinephrine have shown to strengthen CPM [32] suggesting that
12
13 such drugs in combination may be used for selected vulnerable KOA patients before TKR, but
14
15 future randomized control trials are needed to confirm the hypothesis of such a targeted
16
17 individualized treatment regime.
18
19
20
21
22
23
24

25 It is well known that pain in KOA varies throughout the day (e.g. during walking, while climbing
26
27 stairs, or when the patient is at rest)[4,29]. To avoid misinterpretation of an e.g. lower current
28
29 pain while the patients were examined in the laboratory, the peak pain intensity within the last 24
30
31 hours is, as in the present study, often applied to assess the knee OA pain [2-4,30,31,35,37,38].
32
33 Pain ratings during rest, during the last 24 hours, or during the last 8 days are strongly correlated
34
35 in patients with hip and knee OA [28].
36
37
38
39
40
41

42 The present was not designed to predict if patients developed chronic postoperative pain or not
43
44 and hence the sub-grouping of patients were not optimized for this purpose. However the
45
46 combination of mechanistic QST parameters for assessing central pain processing may be
47
48 indicative of possibly biologically important contributions for the postoperative outcome. In case
49
50 of specific sensitive patients (group A) red flags should be raised prior to surgery to indicate that a
51
52 given patient could be more vulnerable for developing less postoperative pain relief.
53
54
55
56
57
58
59
60
61
62
63
64
65

4.2 Temporal Summation of Pain and Conditioned Pain Modulation

Previous studies have shown that patients with KOA have enhanced temporal summation compared with healthy controls [4] and that patients with pain after revision TKR have continued enhanced temporal summation as compared with patients without pain [39]. Preoperative temporal summation has been associated with postoperative chronic pain after TKR [30]. The current study defined two groups with facilitated preoperative TSP, and the group with the most facilitated TSP showed the least pain relief after TKR surgery. Patients with KOA have shown a less efficient CPM system compared with healthy controls [4] but the CPM normalises after pain free recovery after TKR surgery [13]. Dysfunctional preoperative CPM has been suggested to have a predictive value for which patients may develop chronic postoperative pain [47,49], for which reason the current study focused on combining these two apparently important pain mechanisms. It should be stated that neither TSP nor CPM individually as parameters could be associated with postoperative outcomes. The present preliminary study arbitrarily selected cut-off levels of the different QST parameters used for sub-grouping patients and further studies should investigate if these cut-off levels could be further optimised and hence increase the ability to predict postoperative outcomes.

Cuff algometry has previously been widely used to study pain mechanisms in KOA, such as pressure pain thresholds, TSP, and CPM [13,39,40], and offers the advantage of being user-independent [33]. Pain sensitivity has been shown to be related with age for which reason the analysis in the present study was adjusted for age [11]. In addition, studies have shown that increased BMI is associated with increased inflammation [24] and increased inflammation is associated sensitization of the peripheral nociceptors [1], why BMI is important to adjust for in the statistical analysis as done in the present study.

4.3 Widespread Hyperalgesia

Cuff assessment performed on the lower part of the leg in KOA is suggested a proxy for widespread hyperalgesia. The current study showed that KOA patients with preoperatively facilitated TSP and impaired CPM and patients with facilitated TSP had a decreased pressure detection threshold measured by cuff algometry compared with the other groups. Patients with normal CPM combined with normal TSP (normal defined as the grand average in the present population) did not show widespread hyperalgesia compared with the other groups which may suggest that facilitated TSP and widespread hyperalgesia somehow reflect the same pain mechanisms.

Preoperative widespread hyperalgesia measured by handheld pressure algometry has been shown in KOA patients compared with healthy controls [4,8,9,13,17,18,30]. A recent study found that preoperative widespread hyperalgesia measured by pressure algometry did not, as a single parameter, predict postoperative pain after total knee replacement [48]. The present study did confirm that preoperative widespread hyperalgesia as assessed by handheld algometry is not capable of predicting the postoperative **pain relief** after TKR.

In addition, this study illustrated that widespread hyperalgesia measured by cuff and handheld algometry yielded different sensory profiles which had different predictive values for postoperative KOA patients and hence argue for the value of multi-modal, mechanism based sensory testing.

4.4 Limitations

This study presented a 24% dropout at follow-up limiting the study results by the low sample size of the groups, but these patients are most likely the patients with no pain. **This lowers the**

1 sensitivity of the current study, and the findings should be replicated in large studies before the
2
3 clinical implications can be concluded. The study was designed as an explorative study and as such
4
5 no firm statistical plan was formulated *à priori* which further highlight the preliminary nature at
6
7 the study. The findings should therefore be replicated in future studies before the findings can be
8
9 suggested as indicative for diagnosis and treatment of patients before TKR.
10
11
12
13

14 The groups were initially established based on arbitrary selected cut-off limits of the mechanistic
15
16 QST parameters which should be optimised to enhance the sensitivity of the sub-grouping. The
17
18 current study did not include a healthy control group, and therefore, the terms “normal” TSP and
19
20 “normal” CPM is used in contrast to “facilitated” TSP and “impaired” CPM.
21
22
23
24

25 The present preliminary study can be indicative of how possibly biologically important
26
27 sensitization processes may contribute to the continued postoperative pain problems in KOA
28
29 patients and hence provide clinically important pre-operative indications for specifically vulnerable
30
31 patients.
32
33
34
35

36 The clinical outcome parameter in the present study was based on the peak pain intensity in the last 24
37
38 hours as this has been shown as a reliable clinical parameter in characterizing KOA pain. If this is the most
39
40 sensitive parameter to be associated with mechanistic QST parameters is not known and should be further
41
42 investigated.
43
44
45
46

47 This study was conducted in a single hospital and a multi-centre study should confirm the findings
48
49 of this study. In addition, the patients were end-stage KOA patients and the criteria for eligibility
50
51 for surgery may vary from country to country.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

5. Conclusions

KOA patients with preoperatively facilitated TSP and impaired CPM constitute a group of vulnerable patients who experience **less postoperative pain relief 12 months** after TKR as compared with patients with either only facilitated TSP or impaired CPM alone. Widespread hyperalgesia assessed by cuff algometry may be a predictive preoperative mechanistic QST parameter for the postoperative **pain relief** after TKR.

Conflicts of Interest

NociTech is partly owned by Aalborg University and KKP is partly employed by NociTech.

Acknowledgements

The authors thank The Danish Rheumatism Association, The Innovation Fund Denmark (j.no. 136-2014-5) and The Shionogi Science Program for providing the opportunity to conduct the study.

TGN is a part of the Center for Neuroplasticity and Pain (CNAP) which is supported by the Danish National Research Foundation (DNRF121). Stud. Ph.D. Hjalte Holm Andersen is acknowledged for assistance with the artwork.

References

- [1] Andrew D, Greenspan JD. Mechanical and heat sensitization of cutaneous nociceptors after peripheral inflammation in the rat. *J Neurophysiol* 1999;82:2649-2656.
- [2] Arendt-Nielsen L, Egsgaard L, Petersen K, Eskehave T, Graven-Nielsen T, Hoeck H, Simonsen O. A mechanism-based pain sensitivity index to characterize knee osteoarthritis patients with different disease stages and pain levels. *Eur J Pain* 2015;19(10):1406-1417.
- [3] Arendt-Nielsen L, Eskehave TN, Egsgaard LL, Petersen KK, Graven-Nielsen T, Hoeck HC, Simonsen O, Siebuhr AS, Karsdal M, Bay-Jensen AC. Association Between Experimental Pain Biomarkers and Serologic Markers in Patients With Different Degrees of Painful Knee Osteoarthritis. *Arthritis Rheumatol* 2014;66:3317-3326.
- [4] Arendt-Nielsen L, Nie H, Laursen MB, Laursen BS, Madeleine P, Simonsen OH, Graven-Nielsen T. Sensitization in patients with painful knee osteoarthritis. *Pain* 2010;149:573-581.
- [5] Arendt-Nielsen L, Graven-Nielsen T. Translational musculoskeletal pain research. *Best Prac Res Clin Rheumatol* 2011;25:209-226.
- [6] Arendt-Nielsen L, Skou ST, Nielsen TA, Petersen KK. Altered Central Sensitization and Pain Modulation in the CNS in Chronic Joint Pain. *Curr Osteoporos Rep.* 2015;13(4):225-234.
- [7] Beswick AD, Wylde V, Gooberman-Hill R, Blom A, Dieppe P. What proportion of patients report long-term pain after total hip or knee replacement for osteoarthritis? A systematic review of prospective studies in unselected patients. *BMJ Open* 2012;2:e000435-2011-000435.
- [8] Felson DT, Anderson JJ, Naimark A, Walker AM, Meenan RF. Obesity and knee osteoarthritis. *Ann Intern Med* 1988;109:18-24.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

[9] Felson DT, Zhang Y, Anthony JM, Naimark A, Anderson JJ. Weight loss reduces the risk for symptomatic knee osteoarthritis in women. The Framingham Study. *Ann Intern Med* 1992;116:535-539.

[10] Finan PH, Buenaver LF, Bounds SC, Hussain S, Park RJ, Haque UJ, Campbell CM, Haythornthwaite JA, Edwards RR, Smith MT. Discordance between pain and radiographic severity in knee osteoarthritis: findings from quantitative sensory testing of central sensitization. *Arthritis Rheum* 2013;65:363-372.

[11] Finocchietti S, Arendt-Nielsen L, Petersen KK, Simonsen O, Graven-Nielsen T. Cuff algometry to assess sensitization in elderly patients with severe knee osteoarthritis compared with controls. The 15th World Congress on Pain, 6-11 October 2014, Buenos Aires, Argentina. International Association for the Study of Pain/IASP Press, 2014. s. No. PT232

[12] Finocchietti S, Takahashi K, Okada K, Watanabe Y, Graven-Nielsen T, Mizumura K. Deformation and pressure propagation in deep tissue during mechanical painful pressure stimulation. *Med Biol Eng Comput* 2013;51:113-122.

[13] Graven-Nielsen T, Wodehouse T, Langford R, Arendt-Nielsen L, Kidd B. Normalisation of widespread hyperesthesia and facilitated spatial summation of deep-tissue pain in knee osteoarthritis patients after knee replacement. *Arthritis Rheum* 2012;64:2907-2916.

[14] Graven-Nielsen T, Arendt-Nielsen L. Assessment of mechanisms in localized and widespread musculoskeletal pain. *Nat Rev Rheumatol* 2010;6:599-606.

[15] Graven-Nielsen T, Aspegren Kendall S, Henriksson KG, Bengtsson M, Sörensen J, Johnson A, Gerdle B, Arendt-Nielsen L. Ketamine reduces muscle pain, temporal summation, and referred pain in fibromyalgia patients. *Pain* 2000;85:483-491.

- 1 [16] Graven-Nielsen T, Vaegter HB, Finocchietti S, Handberg G, Arendt-Nielsen L. Assessment of
2 musculoskeletal pain sensitivity and temporal summation by cuff pressure algometry: A reliability study.
3 Pain 2015;156(11):2193-202.
4
5
6
7
8
9 [17] Harden RN, Bruehl S, Stanos S, Brander V, Chung OY, Saltz S, Adams A, Stulberg SD. Prospective
10 examination of pain-related and psychological predictors of CRPS-like phenomena following total knee
11 arthroplasty: a preliminary study. Pain 2003;106:393-400.
12
13
14
15
16
17 [18] Hawker G, Davis A, French M, Cibere J, Jordan J, March L, Suarez-Almazor M, Katz J, Dieppe P.
18 Development and preliminary psychometric testing of a new OA pain measure—an OARSI/OMERACT
19 initiative. Osteoarthritis Cartilage 2008;16:409-414.
20
21
22
23
24
25 [19] Hübscher M, Moloney N, Leaver A, Rebeck T, McAuley JH, Refshauge KM. Relationship between
26 quantitative sensory testing and pain or disability in people with spinal pain—A systematic review and
27 meta-analysis. Pain 2013;154:1497-1504.
28
29
30
31
32
33 [20] Jinks C, Jordan K, Croft P. Osteoarthritis as a public health problem: the impact of developing knee pain
34 on physical function in adults living in the community: (KNEST 3). Rheumatology (Oxford) 2007;46:877-881.
35
36
37
38
39 [21] Julien N, Goffaux P, Arsenault P, Marchand S. Widespread pain in fibromyalgia is related to a deficit of
40 endogenous pain inhibition. Pain 2005;114:295-302.
41
42
43
44
45 [22] Kellgren J, Lawrence J. Radiological assessment of osteo-arthrosis. Ann Rheum Dis 1957;16:494.
46
47
48
49 [23] Kosek E, Hansson P. Modulatory influence on somatosensory perception from vibration and
50 heterotopic noxious conditioning stimulation (HNCS) in fibromyalgia patients and healthy subjects. Pain
51 1997;70:41-51.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 [24] Lee YC, Lu B, Bathon JM, Haythornthwaite JA, Smith MT, Page GG, Edwards RR. Pain sensitivity and
2 pain reactivity in osteoarthritis. *Arthritis Care Res* 2011;63:320-327.
3
4

5
6
7 [25] Lluch E, Torres R, Nijs J, Van Oosterwijck J. Evidence for central sensitization in patients with
8 osteoarthritis pain: A systematic literature review. *Eur J Pain* 2014;18(10):1367-1375.
9

10
11
12 [26] Manafi-Khanian B, Arendt-Nielsen L, Frøkjær J, Graven-Nielsen T. Deformation and pressure
13 propagation in deep somatic tissue during painful cuff algometry. *Eur J Pain* 2015;19(10):1456-1466.
14
15

16
17
18 [27] Olesen SS, Brock C, Krarup AL, Funch-Jensen P, Arendt-Nielsen L, Wilder-Smith OH, Drewes AM.
19 Descending inhibitory pain modulation is impaired in patients with chronic pancreatitis. *Clin Gastroenterol*
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

[28] Perrot S, Poiraudou S, Kabir-Ahmadi M, Rannou F. Correlates of pain intensity in men and women
with hip and knee osteoarthritis. Results of a national survey: The French ARTHRIX study. *Clin J Pain*
2009;25:767-772.

[29] Petersen KK, Simonsen O, Laursen MB, Nielsen TA, Rasmussen S, Arendt-Nielsen L. Chronic
Postoperative Pain After Primary and Revision Total Knee Arthroplasty. *Clin J Pain* 2015;31:1-6.

[30] Petersen KK, Arendt-Nielsen L, Simonsen O, Wilder-Smith O, Laursen MB. Presurgical assessment of
temporal summation of pain predicts the development of chronic postoperative pain 12 months after total
knee replacement. *Pain* 2015;156:55-61.

[31] Petersen KK, Siebuhr AS, Graven-Nielsen T, Simonsen O, Boesen M, Gudbergesen H, Karsdal M, Bay-
Jensen AC, Arendt-Nielsen L. Sensitization and Serological Biomarkers in Knee Osteoarthritis Patients With
Different Degrees of Synovitis. *Clin J Pain* 2015 [Epub ahead of print].

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

[32] Phillips K, Clauw DJ. Central pain mechanisms in chronic pain states—maybe it is all in their head. *Best Prac Res Clin Rheumatol* 2011;25:141-154.

[33] Polianskis R, Graven-Nielsen T, Arendt-Nielsen L. Computer-controlled pneumatic pressure algometry—a new technique for quantitative sensory testing. *Eur J Pain* 2001;5:267-277.

[34] Price DD, Mao J, Frenk H, Mayer DJ. The N-methyl-D-aspartate receptor antagonist dextromethorphan selectively reduces temporal summation of second pain in man. *Pain* 1994;59:165-174.

[35] Rathleff MS, Petersen KK, Arendt-Nielsen L, Thorborg K, Graven-Nielsen T. Impaired Conditioned Pain Modulation in Young Female Adults with Long-Standing Patellofemoral Pain: A Single Blinded Cross-Sectional Study. *Pain Medicine* 2015 [Epub ahead of print].

[36] Sandrini G, Rossi P, Milanov I, Serrao M, Cecchini A, Nappi G. Abnormal modulatory influence of diffuse noxious inhibitory controls in migraine and chronic tension-type headache patients. *Cephalalgia* 2006;26:782-789.

[37] Siebuhr AS, Bay-Jensen AC, Leeming DJ, Plat A, Byrjalsen I, Christiansen C, van de Heijde D, Karsdal MA. Serological identification of fast progressors of structural damage with rheumatoid arthritis. *Arthritis Res Ther* 2013;14;15(4):R86

[38] Siebuhr A, Petersen KK, Arendt-Nielsen L, Egsgaard LL, Eskehave T, Christiansen C, Simonsen O, Hoek HC, Karsdal MA, Bay-Jensen AC. Identification and characterisation of osteoarthritis patients with inflammation derived tissue turnover. *Osteoarthritis Cartilage* 2014;22(1):44-50.

[39] Skou ST, Graven-Nielsen T, Rasmussen S, Simonsen O, Laursen MB, Arendt-Nielsen L. Widespread sensitization in patients with chronic pain after revision total knee arthroplasty. *Pain* 2013;154(9):1588-1594.

1 [40] Skou S, Graven-Nielsen T, Rasmussen S, Simonsen O, Laursen M, Arendt-Nielsen L. Facilitation of pain
2
3 sensitization in knee osteoarthritis and persistent post-operative pain: A cross-sectional study. Eur J Pain
4
5 2014;18(7):1024-1031
6

7
8
9 [41] Kurtz S, Ong K, Lau E, Mowat F, Halpern M. Projections of primary and revision hip and knee
10
11 arthroplasty in the United States from 2005 to 2030. J Bone Joint Surg Am 2007;89:780-785.
12
13

14
15 [42] Suokas A, Walsh D, McWilliams D, Condon L, Moreton B, Wylde V, Arendt-Nielsen L, Zhang W.
16
17 Quantitative sensory testing in painful osteoarthritis: a systematic review and meta-analysis. Osteoarthritis
18
19 Cartilage 2012;20(10):1075-1085.
20
21

22
23 [43] Uchida Y, Petersen K, Graven-Nielsen T, Arendt-Nielsen L. Within and between daily session reliability
24
25 of computer-controlled pressure cuff algometry. Abstract book of the 9th Congress of the European Pain
26
27 Federation, EFIC vol. PO28 No. 363. pp. EFIC5-0373.
28
29

30
31 [44] Vægter HB, Handberg G, Graven-Nielsen T. Isometric exercises reduce temporal summation of
32
33 pressure pain in humans. Eur J Pain 2015;19(7):973-983.
34
35

36
37 [45] Vierck CJ,Jr, Cannon RL, Fry G, Maixner W, Whitsel BL. Characteristics of temporal summation of
38
39 second pain sensations elicited by brief contact of glabrous skin by a preheated thermode. J Neurophysiol
40
41 1997;78:992-1002.
42
43

44
45 [46] Weissman-Fogel I, Granovsky Y, Crispel Y, Ben-Nun A, Best LA, Yarnitsky D, Granot M. Enhanced
46
47 presurgical pain temporal summation response predicts post-thoracotomy pain intensity during the acute
48
49 postoperative phase. J Pain 2009;10:628-636.
50
51

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

[47] Wilder-Smith OH, Schreyer T, Scheffer GJ, Arendt-Nielsen L. Patients with chronic pain after abdominal surgery show less preoperative endogenous pain inhibition and more postoperative hyperalgesia: a pilot study. *J Pain Palliat Care Pharmacother* 2010;24:119-128.

[48] Wylde V, Sayers A, Lenguerrand E, Gooberman-Hill R, Pyke M, Beswick AD, Dieppe P, Blom AW. Preoperative widespread pain sensitization and chronic pain after hip and knee replacement: a cohort analysis. *Pain* 2015;156(1):47-54.

[49] Yarnitsky D, Crispel Y, Eisenberg E, Granovsky Y, Ben-Nun A, Sprecher E, Best L, Granot M. Prediction of chronic post-operative pain: pre-operative DNIC testing identifies patients at risk. *Pain* 2008;138:22-28.

1 **Figure Legends**


2
3
4
5
6
7
8 Figure 1. A: Mean (\pm SEM) preoperative pressure detection thresholds (PDTs, red bars, [kPa]), pressure pain
9 tolerance thresholds (PTTs, yellow bars, [kPa]), conditioned pain modulation (CPM, green bars, [kPa]) and
10 temporal summation of pain (TSP, blue bars, [VAS]) were measured using cuff algometry at the lower leg.
11
12 Groups are defined by: facilitated TSP and impaired CPM (Group A, N=16), facilitated TSP and normal CPM
13 (Group B, N=15), normal TSP and impaired CPM (Group C, N=44), and normal TSP and normal CPM (Group
14 D, N=28). B: Mean (\pm SEM) pain relief from the four groups after total knee replacement. * indicates
15 significant difference ($P<0.05$) compared with group A, and # indicates significant difference ($P<0.05$)
16 compared with group B.
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 Figure 2. Mean (\pm SEM) preoperative pressure pain thresholds (PPTs) measured using handheld pressure
32 algometry at the peripatellar area (knee), the tibialis anterior muscle (TA), and the extensor carpi radialis
33 longus muscle (Arm). No significant differences were found between groups.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Summary

Knee osteoarthritis patients with low conditioned pain modulation and high temporal summation have more pain 12 months after total knee replacement surgery.






Table 1: Demographic characteristics of the subgrouped KOA patients

	Group A	Group B	Group C	Group D
Number (N) of patients	N=16	N=15	N=44	N=28
Preoperative pain VAS scores (cm, mean \pm SEM)	6.6 \pm 0.5	6.8 \pm 0.5	6.8 \pm 0.3	5.9 \pm 0.4
BMI (kg/m ² , mean \pm SEM)	28.5 \pm 1.2	32.2 \pm 1.7	30.4 \pm 0.8	28.6 \pm 1.1*
Age (years, mean \pm SEM)	73.2 \pm 2.1	69.6 \pm 2.2	66.8 \pm 1.7	67.0 \pm 1.7
Sex (% female)	73.3%	57.1%	70.7%	51.9%
KL (mean (range))	3.7 (2-4)	3.9 (1-4)	3.7 (3-4)	3.8 (3-4)
Pain relief (% VAS reduction, mean \pm SEM)	52.0 \pm 14.0	81.1 \pm 3.5 [#]	79.6 \pm 4.4 [#]	69.4 \pm 7.9

BMI: Body Mass Index. KL: Kellgren & Lawrence radiological scores. Significant difference compared with

Group B (*, ANOVA: P = 0.020) or Group A (#; ANOVA: P < 0.023).

Table 2. The crude coefficient shows the result from the univariate logistic regression analysis between the postoperative pain relief and the preoperative pain and pressure detection threshold (PDT) measured by cuff algometry. The adjusted coefficient shows the results from the multivariate stepwise logistic regression analysis.

	Crude coefficient	P-value	Adjusted coefficient	P-value
Preoperative pain	0.263	0.005	0.183	0.080
PDT	-0.196	0.042	-0.222	0.034