

POSTER

LEUKOCYTE ANTIGENS AS CANDIDATE GENES TO IMPROVE
THE IMMUNORESPONSE IN SWINEANTÍGENOS LEUCOCITARIOS COMO GENES CANDIDATOS PARA MEJORAR LA
RESPUESTA INMUNE EN CERDOS

Moreno, A., L. Morera, J.J. Garrido, A. Jiménez, G. Estesó, G. Paños, N. Yubero,
M. Barbancho and D. Llanes

Department of Genetics. Veterinary Faculty. Molecular Markers of Domestic Animals Group. UCO-CSIC.
University of Cordoba. Edificio C-5. Campus de Rabanales. 14071 Córdoba. Spain.

ADDITIONAL KEYWORDS

Immunogenetics. Leukocyte antigens. SNP.

PALABRAS CLAVE ADICIONALES

Inmunogenética. Antígenos leucocitarios. SNP.

SUMMARY

Leukocyte antigens (CD) have functions related to immune response and are of interest as *classical* candidate genes for health. Polymorphisms (e.g. SNPs) in these genes may be associated with variation in the immune response and consequently in disease response. This approach is being taken in search of susceptibility genes for swine disease. In addition, these genes may vary between populations, especially where specific adaptation to pathogens has occurred, and are of potential interest in characterising pig biodiversity.

RESUMEN

Los antígenos leucocitarios (CD) tienen funciones relacionadas con la respuesta inmune y son de interés con genes candidatos *clásicos* para la salud. Los Polimorfismos (ej. SNPs) en estos genes pueden estar asociados con variaciones en la respuesta inmune y consecuentemente con la respuesta a la enfermedad. Este ensayo se está desarrollando en la búsqueda de

susceptibilidades genéticas a enfermedades porcinas.

Adicionalmente, estos genes pueden variar entre poblaciones, especialmente en la que han ocurrido adaptaciones a patógenos, y suponen un interés potencial para la caracterización de la biodiversidad porcina.

In the last twenty years the number of leukocyte receptors discovered in humans and domestic animals have been increasing enormously. The first International Workshop and conference on Human Leukocyte differentiation antigens (HLDA), which was held in Paris in 1982, proposed the *Cluster Differentiation* (CD) nomenclature for the leukocyte surface antigens recognized by at least two independent antibodies. The number of defined CD has been growing since then with each

Arch. Zootec. 52: 249-253. 2003.

workshop; 151 CD clusters in the 5th Boston workshop in 1993 and 240 in the 7th Harrogate workshop in 2000. The same nomenclature was adopted for leukocyte antigens in different animal species. In swine 38 CD clusters were described at the last workshop held in Amsterdam in 1999 (Haverson *et al.*, 2001).

Many different biological functions have been described for these receptors that include adhesion structures, B cells receptors, cytokine receptors, endothelial cells receptors, myeloid cells receptors, NK antigens, non-lineage antigens, platelet receptors and T cell receptors. All of them are of fundamental importance in the control of the immune response. In certain cases changes in these receptors have severe consequences for an individual, e.g. CD18 (α2 integrin) deficiency in humans causes a severe granulocyte disorder with susceptibility to bacterial infections and high morbidity and mortality. This deficiency causes leukocyte adhesion deficiency disease (BLAD) in the bovine specie and CLAD in the canine specie (Jeyaseelan *et al.*, 2000). In swine the complex CD11/CD18 mediates the cellular adhesion of PRV (pseudorabies virus) infected monocytes to endothelial cells and subsequently the virus transmission to endothelial cells (Van de Walle *et al.*, 2003).

Recent advances in DNA technology have shown that a large repertoire of functional variants exists in CD receptor genes as is mentioned by Iida and Nakamura (2003), Iida *et al.* (2003), Mackelprang *et al.* (2002), Meller *et al.* (2001) and Tomer *et al.* (2002).

Most of the CD receptor studied are polymorphic at the cDNA level, showing single nucleotide polymorphisms (SNPs) and some of them also present variability at the protein level (coding SNPs). This polymorphism may influence CD gene expression resulting in a change in the immune response and consequently in the disease response

Studies using single nucleotide polymorphisms (SNPs) or microsatellite polymorphisms have now become technologically realistic and our group proposes to use CD gene variants in the *classical* candidate gene approach in search of susceptibility genes to disease and other related characters such as fertility in swine populations (Tsuchiya *et al.*, 2002).

Our group is specialized in the study of differentiation antigens of the immune system of domestic animals, especially in swine. The first step in our goal has been to characterize functionally and phenotypically the cells by means of monoclonal antibodies from the immune system of these animals. So during the past ten years we have characterized many swine leucocytes antigens, among them CD5; CD11; CD18; CD29; MHC Class I; CD41/61; CD46; CD230 and Fibrinogen.

In a second step we have characterized the genes that controlled swine CD receptors using a general protocol presented in **figure 1**.

Following this protocol swine CD9, (accession number Genbank, AY072785) CD29, (accession number Genbank, AF192528) CD51 (accession number Genbank, pending insertion), CD61 (accession number Genbank,

LEUKOCYTE ANTIGENS AS CANDIDATE GENES

AF282890) and CD97 (accession number Genbank, AF413207) have been characterized in our laboratory.

At this moment we are obtaining our first results about CD gene polymorphism. We have described (Morera *et al.*, 2002) a microsatellite in the 3'-UTR region of the swine CD61. Two SNP have been detected after a partial screening of the coding

and regulatory regions of the swine CD61 gene. Our group is carrying out more studies to detect variability in the coding and regulatory regions of the previously characterized CD9, CD29 and CD51 genes.

These markers may be a valuable tool to be used in case control association analyses in various swine populations, e.g. in the case of CD9




Figure 1. Protocol to clone candidate genes for swine leukocyte antigens. (Protocolo seguido para la clonación de un gen candidato presente en leucocitos porcinos).

gene for fertility and with different resistance to various infectious diseases in the case of CD29, CD51 and CD61 genes.

In addition, such polymorphism may be useful in characterising diversity within pig populations (see Ciobanu *et*

al., 2001; Blott *et al.*, 2003; Rothschild, 2003). Of particular interest in such studies is adaptation to local conditions, including disease agents that are geographically restricted, therefore the analysis of variation in CD genes may be an important component of this research.

REFERENCES

- Blott, S., L. Andersson, M. Groenen, M. San Cristobal, C. Chevalet, R. Cardellino, N. Li, L. Huang, K. Li, G. Plastow and C. Haley. 2003. Characterisation of genetic variation in the pig breeds of China and Europe-the PigBioDiv2 project. *Arch. Zootec.*, 52: 207-217.
- Ciobanu, D.C., A.E. Day, A. Nagy, R. Wales, M.F. Rothschild and G.S. Plastow. 2001. Genetic variation in two conserved local Romanian pig breeds using type I DNA markers. *Genet. Sel. Evol.*, 33: 417-432.
- Haverson, K., A. Saalmuller, B. Alvarez, F. Alonso, M. Bailey, A.T. Bianchi, W.J. Boersma, Z. Chen, W.C. Davis, J. Dominguez, H. Engelhardt, A. Ezquerro, L.S. Grosmaire, M.J. Hamilton, E. Hollemweguer, C.A. Huang, K.V. Khanna, G. Kuebart, G. Lackovic, J.A. Ledbetter, R. Lee, D. Llanes, J.K. Lunney, K.C. McCullough, T. Molitor, J. Nielsen, T.A. Niewold, M.D. Pescovitz, J.M. de la Lastra, Z. Rehakova, H. Salmon, W.M. Schnitzlein, J. Seebach, A. Simon, J. Sinkora, M. Sinkora, C.R. Stokes, A. Summerfield, L. Sver, E. Thacker, I. Valpotic, H. Yang, F.A. Zuckermann and R. Zwart. 2001. Overview of the third international workshop on swine leukocyte differentiation antigen. *Veterinary Immunology Immunopathology*, 80: 5-23.
- Iida, A. and Y. Nakamura. 2003. High-resolution SNP map in the 55-kb region containing the selectin gene family on chromosome 1q24-q25. *J. Human Genetics*, 48: 150-4.
- Iida, A., S. Saito, A. Sekine, C. Mishima, Y. Kitamura, K. Kondo, S. Harigae, S. Osawa and Nakamura. 2003. Catalog of 668 SNPs detected among 31 genes encoding potential drug targets on the cell surface. *J. Human Genetics*, 48: 23-46.
- Jeyaseelan, S., S.L. Hsuan, M.S. Kannan, B. Walcheck, J.F. Wang, M.E. Kehrl, E.T. Lally, G.C. Sieck and S.K. Maheswaran. 2000. Lymphocyte function-associated antigen 1 is a receptor for Pasteurella haemolytica leukotoxin in bovine leukocytes. *Infection Immunity*, 68: 72-9.
- Mackelprang, R., C.S. Carlson, L. Subrahmanyam, R.J. Livingston, M.A. Eberle and D.A. Nickerson. 2002. Sequence variation in the human T-cell receptor loci. *Immunological Review*, 190: 26-39.
- Meller, S., P. Jagiello, S. Borgmann, H. Fricke, J.T. Epplen and M. Gencik. 2001. Novel SNPs in the CD18 gene validate the association with MPO-ANCA+ vasculitis. *Genes Immunology*, 2: 269-72.
- Morera, L., A. Jiménez-Marín, M. Yerle, D. Llanes, M.J. Barbancho and J.J. Garrido. 2002. A polymorphic microsatellite located on pig chromosome band 12p11-2/3p13, within the 3'-UTR of the ITGB3 gene. *Animal Genetics*, 33: 224-248.
- Rothschild, M. F. 2003. Approaches and Challenges in Measuring Genetic Diversity in Pigs. *Arch. Zootec.*, 52: 129-135.
- Tomer, Y., E. Concepcion and D.A. Greenberg. 2002. A C/T Single-Nucleotide Polymorphism in the Region of the CD40 Gene is Associated with Graves' Disease. *Thyroid*, 12: 1129-35.

LEUKOCYTE ANTIGENS AS CANDIDATE GENES

Tsuchiya, N., J. Ohashi and K. Tokunaga. 2002. Variations in immune response genes and their associations with multifactorial immune disorders. *Immunological Rev.*, 190: 169-81.
Van de Walle, G.R., H.W. Favoreel, H.J.

Nauwynck, T.C. Mettenleiter and M.B. Pensaert. 2003. Transmission of pseudorabies virus from immune-masked blood monocytes to endothelial cells. *J. General Virology*, 84: 629-37.