

1           **Heavy metals in an impacted wetland system: a typical case from southwestern China**

2

3   Xiangyang Bi<sup>a, b</sup>, Xinbin Feng<sup>a, \*</sup>, Yuangen Yang<sup>a</sup>, Guangle Qiu<sup>a</sup>, Xiaoli Qian<sup>a, c</sup>, Feili Li<sup>a, d</sup>, Taoze Liu<sup>a, b</sup>,

4           Tianrong He<sup>a, b</sup>, Ping Li<sup>a, b</sup>, Zhiyou Fu<sup>a, b</sup>, Xiangdong Li<sup>e</sup> and Grace Shin<sup>e</sup>

5

6   <sup>a</sup>State Key Laboratory of Environmental Geochemistry, Institute of Geochemistry, Chinese Academy

7           of Sciences, Guiyang 550002, P. R. China;

8   <sup>b</sup>Graduate School of the Chinese Academy of Sciences, Beijing 100039, P. R. China;

9   <sup>c</sup>Guizhou University, Guiyang 550025, P. R. China;

10 <sup>d</sup>College of Biological and Environmental Engineering, Zhejiang University of Technology, Hangzhou

11           310032, P. R. China

12 <sup>e</sup>Department of Civil and Structural Engineering, The Hong Kong Polytechnic University, Hung Hom,

13           Kowloon, Hong Kong

14

15           \* Corresponding author. Tel.: +86-851-5891356; fax: +86-851-5891609.

16           E-mail address: [fengxinbin@vip.skleg.cn](mailto:fengxinbin@vip.skleg.cn)

17

18 **Abstract**

19 Historic zinc smelting in Hezhang, southwestern China, has resulted in significant heavy metal  
20 contamination to the surrounding ecosystems. The Caohai wetland system, an important national nature  
21 reserve close to the Hezhang zinc smelting area, was investigated in the present study. Results showed  
22 that sediments from the Caohai wetland system have been seriously contaminated by Cd, Pb and Zn  
23 with the highest enrichment factors (*EF*) up to 91, 16 and 41, respectively. The heavy metals in the  
24 sediments were mainly presented in the organic and residual fractions, both of which can be changed  
25 into Fe-Mn oxide fraction in a more oxidized condition induced by the aquatic plants. Pb isotopic  
26 compositions in the sediments indicated a ternary mixing model with ore Pb, gasoline Pb and geogenic  
27 Pb as the three end members/sources which accounted for about 60, 26, and 14%, respectively, of the  
28 inventory of Pb in the sediments. Heavy metal contamination in aquatic plants was also studied and the  
29 results indicated that heavy metals accumulated by plants may pose a potential threat to the higher  
30 trophic-level organisms, even humans.

31

32 *Keywords:* Heavy metals; Pb isotopes; Sequential chemical fraction; Aquatic plant; Zinc smelting;

33 Caohai wetland; China

34

## 35 **1 Introduction**

36 Sediments are recognized as an important sink for heavy metals emitted from anthropogenic sources,  
37 as well as a potential non-point pollution source which may directly affect overlying waters and the  
38 aquatic organisms. It is well accepted that the total concentrations of heavy metals in sediments can not  
39 provide sufficient information about their impacts to the ecosystem because the mobility,  
40 bioavailability and toxicity of metals depends not only on their total concentrations but also on the  
41 geochemical fractions in which they occur. Thus, various extraction procedures have been developed to  
42 assess the partitioning of sediment/soil-associated metals among various geochemical fractions and to  
43 evaluate the metal mobility and bioavailability (Tessier et al., 1979; Kersten and Förstner, 1986; Ure et  
44 al., 1993). The Tessier sequential extraction procedure is among the most common used methods,  
45 although the disadvantages of this extraction scheme have been well recognized (e.g., non-specificity  
46 of extraction and re-sorption of metals) (Nirel and Morel, 1990; Gomez-Ariza et al., 1999; Rapin et al.,  
47 1986).

48 Metal fractionations in sediment are strongly depended on the substrata geochemical conditions,  
49 such as the pH and redox potential (Tessier et al., 1979). Many studies (e.g., Jacob and Otte, 2004a, b;  
50 Almeida et al., 2004) found that aquatic plants can alter the chemistry of the sediment (for instance,  
51 causing changes in pH and redox potential), modifying the metal fractionations in the sediment. For  
52 example, Almeida et al. (2004) reported that metals associated with the residual fraction were  
53 significantly lower in the rhizosediment than in the sediment. On the other hand, metals settled in the  
54 sediment can be accumulated by aquatic plants. Many studies reported that high metal concentrations  
55 in sediments often result in high levels of metals in the onsite vegetations (Deng et al., 2004; Hozhina  
56 et al., 2001; Cymerman-Samecka and Kempers, 2004). Therefore, the investigations of metal

57 accumulations in plants can provide vital information on what extents the contamination has impacted  
58 on an aquatic ecosystem.

59 In addition to the quantitative measurement of heavy metals, Pb isotopic compositions can be very  
60 useful for identifying sources and contributions of anthropogenic Pb and have been widely used in a  
61 wide range of environmental studies (Mukai et al., 1993, 2001; Gélinas and Schmit, 1997; Zhu et al.,  
62 2001; Wong and Li, 2004; Lee et al., 2006; Gallon et al., 2006). As a general rule, anthropogenic Pb  
63 derived from an ore is less radiogenic than the geogenic Pb. For example, the  $^{206}\text{Pb}/^{207}\text{Pb}$  ratios of Pb  
64 used to manufacture gasoline additives in south China are generally lower than 1.17 (Zhu et al., 2001).  
65 In comparison, the  $^{206}\text{Pb}/^{207}\text{Pb}$  ratios of naturally derived Pb are often higher than 1.20 (Wong and Li,  
66 2004; Lee et al., 2006).

67 Zinc smelting activities have caused many serious environmental problems in China in terms of  
68 heavy metal pollution due to the fact that not only China is the largest zinc producer in the world, but  
69 also to the fact that most zinc industry in China has used the old technique in smelting without the  
70 necessary pollution control measures. However, this kind of pollution has not been well studied so far.  
71 Hezhang, located in southwest China, is a typical region which had been seriously impacted by  
72 historical zinc smelting activities (Feng et al., 2004, 2006; Bi et al., 2006a, b; Yang et al., 2006). In this  
73 study, the impact of the deposits from the Hezhang zinc smelting emissions on a natural freshwater  
74 wetland (Caohai) was investigated. The objectives of the present research are (1) to evaluate the  
75 metal enrichment characteristics in the wetland sediments; (2) to determine the geochemical  
76 fractionations of heavy metals in the sediments using Tessier sequential extraction method for their  
77 mobility and bioavailability; (3) to identify the potential origins of the sediment Pb by lead isotopes;  
78 and (4) to delineate the extent to which these metals accumulate in the onsite vegetations.

79 **2. Materials and methods**

80 *2.1. Study site*

81 The Caohai natural wetland (104°12'-104°18' E, 26°49'-26°53' N), an important national natural  
82 reserve for migratory birds in China, is situated in northwestern Guizhou, approximately 350 km west  
83 of Guiyang, the capital of Guizhou Province. It lies on the Yunnan-Guizhou Plateau with an altitude of  
84 about 2100 m above sea level. Its climate represents a typical subtropical humid monsoon type with an  
85 average temperature of 10.6°C and an average annual rainfall of 951 mm. The Caohai wetland is  
86 underlain mainly by Carboniferous limestone and dolomite with water surface area about 25 km<sup>2</sup> and  
87 depth of 1-5 m. The major source of water in the Caohai wetland is precipitation. The Hezhang zinc  
88 smelting area is located approximately 15 km northeast of the Caohai wetland (Fig. 1)

89 *2.2. Sample collection and analysis*

90 Field sampling at the Caohai wetland was carried out in September 2005 and June 2006. Ten surface  
91 sediment samples (from the top 5 cm) were collected by grab sampler for heavy metal screening. In  
92 order to develop profiles of heavy metals with respect to sediment depths, two cores were also  
93 collected by gravity sampler. Both core and grab samples were air dried and ground < 100 mesh prior  
94 to the chemical analysis. For the total metal concentration analysis, sediment samples (0.5g) were  
95 digested by a 10 mL HNO<sub>3</sub> (65%, v/v), HCl (30%, v/v) and HF (40%, v/v) mixture in a sealed Teflon  
96 vessel. Metal fractionations were performed using a modified version of the method described by  
97 Tessier et al. (1979). The sequential extraction procedure consisted of extractions in the following order  
98 with associated chemical reagents and conditions: exchangeable fraction (1 M MgCl<sub>2</sub>, pH 7.0, for 1h),  
99 carbonate fraction (1 M NaOAc, pH 5.0, for 5h), Fe-Mn oxide bound fraction (0.04 M NH<sub>2</sub>OH·HCl in

100 25% (v/v) HOAc at 96°C for 6h), organic fraction (30% H<sub>2</sub>O<sub>2</sub> and 0.02 M HNO<sub>3</sub> for 2h, second 30%  
101 H<sub>2</sub>O<sub>2</sub> at 85°C for 3h), and residual fraction (total digestion with concentrated HNO<sub>3</sub>-HCl-HF acid  
102 mixture). After each extraction step was completed, the samples were centrifuged at 3000 g for 15  
103 minutes at room temperature, and the resulting supernatant was filtered through a 0.45 µm filter and  
104 saved for analysis. Between each reaction step, samples were rinsed with 8 mL of deionized water, and  
105 the wash solution was discarded after centrifuging for 15 minutes.

106 Plant samples, including emergent, submerged and floating aquatic vegetations, were taken from a  
107 range of species present at the Caohai wetland. The submerged species, primarily *C. demersum*, *M.*  
108 *spicatum*. and *P. malainus*, were the most common type of aquatic macrophyte. They grow in most  
109 parts of the wetland, especially in the deep water region. Whereas the emergent and floating species are  
110 generally found in the shallow water near the edge of the wetland. The above and under water stem and  
111 leaves (aboveground tissues) of plants were collected by hand with gloves. Each sample consisted of  
112 5-10 subsamples collected from about 20 m<sup>2</sup> area. Plant samples were thoroughly washed with tap  
113 water and de-ionized water to remove soil particles before being oven-dried at about 60 °C. The dried  
114 plant samples were then cut into pieces and ground to powder. The prepared plant samples (0.5g) were  
115 digested by a 10 mL HNO<sub>3</sub> (65%) and H<sub>2</sub>O<sub>2</sub> (30%) solution in a sealed Teflon vessel for total metal  
116 analysis.

117 Metal (Cd, Cu, Pb and Zn) concentrations in all prepared solutions were determined using ICP-OES  
118 (Vista MPX, Varian Inc.). Quality assurance and quality control of metal analysis were assessed using  
119 duplicates, method blanks and standard reference materials (SRM2710, SMR8704 and GBW07603).  
120 Reliability of the sequential extraction procedure was determining by comparison of the sum of  
121 fractions with independent total metal digestion concentrations. In most cases, agreement was within

122 20% indicating the extraction results were reliable and accurate.

123 The Pb isotopic composition analysis was performed on selected sediment samples by ICP-MS  
124 (Perkin-Elmer Elan 6100 DRC<sup>plus</sup>). The details of the procedure were reported by Lee et al. (2006). The  
125 analytical parameters were set as 190 sweeps/reading, one reading/replicate, and 10 replicates per  
126 sample solution. Dwell times of 40, 25, 25, and 25 ms were used for <sup>204</sup>Pb, <sup>206</sup>Pb, <sup>207</sup>Pb, and <sup>208</sup>Pb,  
127 respectively. The Pb counts of the procedural blank were below 0.5% of the samples. The relative  
128 standard deviations (RSD) of the 10 replicates were generally below 0.5%. A standard reference  
129 material (NIST SRM981 Common Pb Isotopic Standard) was used for quality control. The measured  
130 Pb ratios of <sup>204</sup>Pb/<sup>207</sup>Pb, <sup>206</sup>Pb/<sup>207</sup>Pb, and <sup>208</sup>Pb/<sup>207</sup>Pb were 0.0645±0.0001, 1.0938±0.0011, and  
131 2.3710±0.0030, which were in good agreement with the standard reference values of 0.0645, 1.0933,  
132 and 2.3704, respectively.

133 Analysis of organic matter in the sediments was performed using the potassium  
134 dichromate-volumetric method (Liu, 1999).

### 135 **3. Results**

#### 136 *3.1. Total metals in sediments*

137 The total concentrations of Cd, Pb, Zn and Cu in the surface and core sediment samples collected  
138 from the Caihao wetland are shown in Table 1. Metal concentrations in the surface sediments exhibited  
139 relatively wide ranges of 2.6-71 µg g<sup>-1</sup> for Cd, 66-160 µg g<sup>-1</sup> for Pb and 140-1100 µg g<sup>-1</sup> for Zn. The  
140 relatively high levels of metals in the surface sediments were generally found in the north sampling  
141 sites (e.g., site 1, 3 and 4), and metal concentrations decreased gradually along the transect from north  
142 to south, except for Cu, probably reflecting the dispersion of the emissions from the zinc smelting area

143 located at the northeast part of Caohai. The metal profiles in the two sediment cores showed the similar  
144 trends with significant elevated Cd ( $10\text{-}81\ \mu\text{g g}^{-1}$ ), Pb ( $68\text{-}200\ \mu\text{g g}^{-1}$ ) and Zn ( $320\text{-}1500\ \mu\text{g g}^{-1}$ )  
145 concentrations in the upper layers ( $<20\ \text{cm}$ ), which was consistent with the intensified zinc smelting  
146 activities in the Hezhang area since the last few decades. It was therefore reasonable to assume from  
147 the metal concentrations and the spatial-temporal distribution patterns of heavy metals in the sediments  
148 that the metal contamination was derived principally from past smelting operations in this area.

### 149 *3.2. Metal fractionation in sediments*

150 The sequential chemical fraction of Cd, Pb, Zn and Cu in the selected sediment (surface sediments  
151 and Core 2 sediments) samples is depicted in Fig. 2. In the surface sediments, Cd and Pb were mainly  
152 present in the organic fraction (Cd, 31-84%; Pb, 34-82%), while the residual fraction was second  
153 dominant (Cd, 4.0-55%; Pb, 14-39%). As for Zn and Cu, the residual fraction became more important  
154 (Zn, 17-69%; Cu, 50-75%) than the organic fraction (Zn, 3.8-46%; Cu, 6.6-49%). In general, heavy  
155 metals in the exchangeable, carbonate and Fe-Mn oxide fractions were less important for most samples  
156 (generally less than 10%), except for two, i.e., Sample 1 and 3, which exhibited abnormally high  
157 percentage of Fe-Mn oxide fraction for Cd (63-76%), Pb (42-47%) and Zn (62-63%).

158 The distribution of metal fractions in the sediment core (especially the upper layers) was similar to  
159 those of the surface sediments with organic and residual fractions being dominant phases. But their  
160 distribution trends varied with depth (Fig. 2b). In general, the percentage of Cd, Pb and Zn in the  
161 organic fraction decreased and the residual fraction increased with depth. The metals in the sediment  
162 core were also slightly associated with the Fe-Mn oxide fraction with a rapid increase in the lowermost  
163 profile. This was probably due to a Fe-Mn oxide boundary existing in these layers. A slightly increase  
164 of the exchangeable and carbonate fractions for Cd and Pb were also observed in the deeper sediments.



165 In contrast, the chemical fractions for Cu were evenly distributed in the profile.

### 166 3.3. *Pb isotope ratios*

167 Pb isotopic analysis results of the selected sediment (the upper layers ( $\leq 20$  cm) of the Core 2), local  
168 background soil and the feeding ore samples are given in Table 2. Generally, the local background soils  
169 have relatively high  $^{206}\text{Pb}/^{207}\text{Pb}$  (1.2441-1.2489) ratios. In contrast, the  $^{206}\text{Pb}/^{207}\text{Pb}$  ratios of the upper  
170 core sediments fell within a relatively low range of 1.1778-1.2016, which increased with depth,  
171 suggesting a mixing of anthropogenic sources and geogenic Pb. The zinc smelting in Hezhang mainly  
172 used the native ores. The  $^{206}\text{Pb}/^{207}\text{Pb}$  ratios of the feeding ores were 1.1756-1.1877, which  
173 corresponded to the previous study (1.1736-1.1870) (Zheng, 1994). These values for ores were  
174 comparable to those measured in the sediments, thus indicating their predominant contribution to the  
175 Pb burden in the sediments.

176 However, the sediment samples also showed slightly lower  $^{208}\text{Pb}/^{207}\text{Pb}$  (2.4611-2.4732) ratios than  
177 either the background soils (2.4771-2.4782) or the ores (2.4837-2.4987). This result suggested that  
178 other unknown anthropogenic sources with low  $^{208}\text{Pb}/^{207}\text{Pb}$  ratios (at least should be lower than those of  
179 the sediments) must be involved. It is reasonable to assume that the coal combustion and gasoline Pb  
180 might be the possible important sources of Pb due to the lack of other large anthropogenic emissions in  
181 the study area.

### 182 3.4. *Metals in plants*

183 Cd, Pb, Zn and Cu concentrations in the aquatic plants (aboveground tissues) collected from the  
184 Caohai wetland are shown in Table 3. The data show that the metal concentrations in the plants varied  
185 among species, indicating their different capacities for metal uptake. Generally, the submerged species

186 *C. demersum*. accumulated the highest concentrations of Cd (8.2  $\mu\text{g g}^{-1}$ ), Pb (22  $\mu\text{g g}^{-1}$ ) and Zn (200  $\mu\text{g}$   
187  $\text{g}^{-1}$ ), and the floating species (*Azolla imbricata* and *Spirogyra communis*) contained the second highest  
188 values of Cd (4.4  $\mu\text{g g}^{-1}$ ) and Pb (20  $\mu\text{g g}^{-1}$ ). While the lowest metal concentrations (Cd, 0.4  $\mu\text{g g}^{-1}$ ; Pb,  
189 1.4  $\mu\text{g g}^{-1}$ ; Zn, 34  $\mu\text{g g}^{-1}$ ; and Cu, 12  $\mu\text{g g}^{-1}$ ) were found in the emergent species *S. triqueter*. In  
190 summary, all the metals (especially Cd, Pb and Zn) found in the plants had much lower concentrations  
191 than those measured in the sediments.

## 192 **4. Discussion**

### 193 *4.1. Enrichment factors in sediments*

194 In this study, enrichment factor (*EF*) was used to assess the anthropogenic contribution of metals in  
195 the sediments. For a given sediment, the metal concentration was normalized to that of Al, a  
196 conservative element relative to silicate crustal materials, to calculate the enrichment factors according  
197 to the following equation:

$$198 \quad EF = \frac{[M / Al]_{sample}}{[M / Al]_{background}} \quad (1)$$

199 where *EF* is the enrichment factor,  $[M/Al]_{sample}$  is the ratio of given metal and Al concentration of the  
200 sample, and  $[M/Al]_{background}$  is the ratio of given metal and Al concentration of a background. In this  
201 study, the relatively low metal concentrations from the deeper sediment layers (35-50cm) of Core 2  
202 was used as the less impacted background values ( $\mu\text{g g}^{-1}$ ): Cd,  $2.1 \pm 0.53$ ; Pb,  $29 \pm 2.5$ ; Zn,  $82 \pm 23$ ; and  
203 Cu,  $21 \pm 1.6$  (Fig. 2). These results are comparable to the background values in the local soil, except for  
204 Cd, which is some what elevated (China Environmental Monitoring Station, 1990).

205 *EF* values were interpreted as suggested by Birch (2003) where  $EF < 1$  indicates no enrichment;  $< 3$  is  
206 minor; 3~5 is moderate; 5~10 is moderately severe; 10~25 is severe; 25~50 is very severe; and  $> 50$  is

207 extremely severe. In this case, only the surface and upper core (<20 cm) sediment samples would  
208 reflect the anthropogenic impacts. In most samples, Cd has the highest *EF* values varying from  
209 moderately severe (5.5) to extremely severe (91) enrichment, followed by Zn (2.7-41) and Pb (2.8-16),  
210 while Cu only shown minor anthropogenic enrichment (generally *EF*<3) (Table 1). These results were  
211 comparable to the previous study which reported that moss samples collected close to the same  
212 smelting area were seriously contaminated by Cd, Pb and Zn, but less impacted by Cu (Bi et al., 2006b).  
213 However, the much lower *EF* value for Pb relative to Cd and Zn in the sediments suggested that the  
214 sediments received less Pb pollutant, which is quite different from that in the moss samples. Pb in the  
215 moss samples exhibited equal or even higher contamination level compared to Cd and Zn (Bi et al.,  
216 2006b). Considering metal pollutants in the Caohai wetland are derived mainly from the emissions of  
217 the zinc smelting, and the wetland is much farther away from the smelting area (~15 km) than the sites  
218 (~2 km) where the moss samples were collected, this result thus indicates that smelting emitted Pb  
219 might prefer to deposit in the close vicinity of the smelter. Telmer et al. (2004) also found a higher  
220 enrichment of Pb compared to Cd and Zn in the near surrounding area (< 5 km) of a smelter.

#### 221 *4.2. Metal motility in sediments*

222 Sequential extraction procedure despite known limitations does provide some insight information on  
223 the potential mobility and redox sensitively of metals in sediments. In a given environment, different  
224 chemical fractions of metals may reveal various degrees of potential mobility and bioavailability  
225 depending on the substrata geochemical conditions and the nature of the metals. Exchangeable and  
226 carbonate fractions are considered to be weakly bonded metals which may equilibrate with the aqueous  
227 phase and thus become more rapidly bioavailable (Pardo et al., 1990). The low percentages of these  
228 fractions of metals in the Caohai sediments thus indicated the low mobility and bioavailability of

229 metals in the sediments. Typically, the Fe-Mn oxide bound fraction are thermodynamically unstable  
230 under anoxic conditions (i.e., low redox potential), while under oxidized conditions, organic matter can  
231 be degraded, leading to a release of trace metals from the organic bound metal fraction (Tessier et al.,  
232 1979).

233 Heavy metals in the sediments were strongly associated with the organic fraction. This could be  
234 related to the high OM levels (14-45%) of the sediments. Previous studies also reported high  
235 percentages of metals bound to the organic fraction with the high OM concentrations in sediments  
236 (Dollae et al., 2001; Olivares-Rieumont et al., 2005). Stoepler (1992) demonstrated that major factors,  
237 such as metal uptake by plants and subsequent decomposition, metal complex formation with soluble  
238 OM, and metal absorption onto settable minerals, were all possible vehicles that might contribute to  
239 metal accumulation in the organic fraction.

240 However, in some cases, i.e., the surface sample 1 and 3, the Fe-Mn oxide fraction of Cd, Pb and Zn  
241 was dominant instead of the organic fraction (Fig. 2a). This was not due to the relatively low OM  
242 concentration of these two samples (Table 1), but a reflection of the influences of the aquatic plants.  
243 The fact is that the sites from which the samples were collected have much more biomass of the  
244 macrophytes than other sites. Many studies reported that living plants can induce sediment oxidation  
245 (higher redox potential) through increased root radial oxygen loss (Madureira et al., 1997; Jacob and  
246 Otte, 2004a, b; Almeida et al., 2004), by which may lead to an increase of metals bound to the Fe-Mn  
247 oxide. Indeed, Carroll et al. (2002) also demonstrated that the sulfides of Cd and Zn in sediments can  
248 dissolve under oxidized conditions and the majorities of these dissolved concentrations will eventually  
249 undergo sorption to or co-precipitation with an iron (oxy)hydroxide.

250 The oxidized conditions induced by plants can simultaneously lead to a decay of the organic matter

251 (Tessier et al., 1979), therefore, the lower OM level of the Sample 1 and 3 was not a cause, but a  
252 result/reflection of the influences of the plants.

253 It is clear that the increased metals in the Fe-Mn oxide fraction must be derived from the  
254 transformation from other fractions, such as the organic and residual fractions. By comparing the  
255 difference of metal fractionations between sample1, 3 and others (especially sample 2 and 4) (Fig. 2a),  
256 it was reasonable to assume that the increase of the Fe-Mn oxide fraction for Cd and Zn in the  
257 sediments was attributed to both the organic and residual fraction conversion, while for Pb the increase  
258 was principally derived from the organic fraction change.

259 In comparison with other metals, the Cu, representing a less impacted element in the sediments, had  
260 the chemical fractions seemed to be independent to the substrata conditions such as the redox potential.

261 The sediments from the deeper layers of the core seemed to be less impacted by human activities and  
262 their metals derived mainly from the geogenic sources. The increasing trend of the residual fraction of  
263 Cd, Pb and Zn with depth in the sediment core thus indicated that the geogenic originated metals tend  
264 to be associated with the residual fraction, i.e., were retained in the crystal lattice of minerals. In  
265 contrast, anthropogenic inputted metals prefer to bind to the non-residual fractions, especially the  
266 organic fraction, in the Caohai sediments, posing a potential hazard on the ecosystem.

#### 267 4.3. Source attribution of Pb in sediments

268 The Pb isotopic compositions indicated that the Pb burden in the sediments were a mixing of  
269 geogenic Pb and a variety of anthropogenic sources, including the ore Pb and other sources with low  
270  $^{208}\text{Pb}/^{207}\text{Pb}$  ratios. It is well known that gasoline Pb has been an important source of Pb to the  
271 atmosphere in the last few decades, in addition, coal combustion has also contributed significantly to  
272 the atmospheric Pb in southwestern China (Mukai et al., 1993, 2001). Besides, no other large

273 anthropogenic emissions were present in the study region. Although no isotopic data are available for  
274 coal in southwestern China, the values from the aerosol can be used because aerosol Pb in southwestern  
275 China was derived principally from the coal combustion (Mukai et al., 1993, 2001). The Pb isotope  
276 ratios in the aerosol from Guiyang were 1.170-1.205 and 2.444-2.476 for  $^{206}\text{Pb}/^{207}\text{Pb}$  and  $^{208}\text{Pb}/^{207}\text{Pb}$ ,  
277 respectively (calculated from Mukai et al. (2001)). The tetraethyl Pb additives in China gasoline were  
278 mainly from the Octel Inc. (UN). According to Zhu et al. (2001), the  $^{206}\text{Pb}/^{207}\text{Pb}$  and  $^{208}\text{Pb}/^{207}\text{Pb}$  ratios  
279 in south China gasoline were 1.1562-1.1632 and 2.4182-2.4260, respectively.

280 Therefore, all the measured and literature isotopic data for various samples in the  $^{206}\text{Pb}/^{207}\text{Pb}$  vs  
281  $^{208}\text{Pb}/^{207}\text{Pb}$  diagram is plotted in Fig. 3. It is clear to note that all the data from the Caohai sediments  
282 fall within a triangle region with ore, gasoline and background soil as the three end members. This  
283 suggests that Pb in the Caohai sediments derived mainly from these three sources: the geogenic Pb, the  
284 gasoline Pb, and of course, the ore Pb from the smelting operations. In contrast, as shown in this Figure,  
285 the coal Pb may not be regarded as a possible end member due to its similar or even higher isotope  
286 ratios compared to those of the sediments.

287 If the Pb in the Caohai sediments was from just these three sources, it is possible to calculate the  
288 relative amounts of the three sources according to a ternary mixing model:

$$289 \quad X = \frac{R_{\text{sample}} - (R_{\text{gasoline}} - R_{\text{geogenic}})Y - R_{\text{geogenic}}}{R_{\text{ore}} - R_{\text{geogenic}}} \quad (2)$$

290 where  $X$  and  $Y$  are the proportion of ore Pb and gasoline Pb in the sediments and  $R_{\text{sample}}$ ,  $R_{\text{ore}}$ ,  $R_{\text{gasoline}}$   
291 and  $R_{\text{geogenic}}$  are the isotope ratios ( $^{206}\text{Pb}/^{207}\text{Pb}$  or  $^{208}\text{Pb}/^{207}\text{Pb}$ ) in the sediment samples, ore, gasoline and  
292 local background soils, respectively. The values of  $^{206}\text{Pb}/^{207}\text{Pb}$  and  $^{208}\text{Pb}/^{207}\text{Pb}$  ratios for background  
293 soils, ore and gasoline are the average of the measured or reported ratios for these groups of samples.  
294 Using the above equation, it was estimated that on the average 60% (47-68%) and 26% (15-37%) of the

295 inventory of Pb in the Caohai sediments originated from the ores used in the smelting and from the  
296 gasoline, respectively. The remaining 14% (5-38%) derived from the geogenic Pb.

#### 297 4.4 Metal accumulations in plants

298 Plant species differ widely in their ability to accumulate heavy metals. Our study showed that the  
299 emergent plants accumulated lower amounts of metals than submerged and floating aquatic vegetation.  
300 This is consistent with previous observations (Albers and Camardese, 1993; Yurukova and Kochev,  
301 1994; Rai et al., 1995). In order to determine whether significant bioaccumulation occurred in the  
302 Caohai vegetations, it is necessary to compare our data with those of previous studies. Outridge and  
303 Noller (1991) reported that Cd, Pb, Zn and Cu concentrations in normal plants were 1.9, 6.3-9.9, 66 and  
304 37  $\mu\text{g g}^{-1}$ , respectively. Baldantoni et al. (2005) observed that Cd, Pb, Zn and Cu concentrations in  
305 plants *Najas marina* shoots and *P. Pectinatus* leaves from a less impacted site on Lake Averno, Naples,  
306 were generally lower than 0.4, 4.0, 60 and 5.0  $\mu\text{g g}^{-1}$ , respectively. Hozhina et al. (2001) investigated  
307 aquatic plants *Equisetum fluviatile*, *Typha latifolia* and *Scirpus sylvaticus* in the Kemerovo region,  
308 Russia, and found that the aboveground tissues of these plants from the control area contained  
309 0.023-1.8  $\mu\text{g g}^{-1}$  of Cd, 0.89-23  $\mu\text{g g}^{-1}$  of Pb, 26-84  $\mu\text{g g}^{-1}$  of Zn and 4.7-39  $\mu\text{g g}^{-1}$  of Cu. Our data  
310 indicated that Cd, Zn and Pb in most plants had higher concentrations than these thresholds, but were  
311 still within the range found in contaminated plants (Deng et al., 2004; Hozhina et al., 2001;  
312 Cymerman-Samecka and Kempers, 2004). In contrast, Cu values in our study were similar to the  
313 literature values mentioned above. The extents of metal accumulations in plants agreed well with the  
314 metal contamination conditions of the Caohai sediments and thus confirming that contamination on  
315 plants did occur.

316 However, the much lower metal concentrations found in the plants compared to those in the

317 sediments suggested the limited bioavailability of the metals in the sediments, which corresponds to the  
318 metal chemical fractionation results in sediments. Nevertheless, the elevated metals accumulated in the  
319 aquatic plants, especially Cd and Pb in *C. demersum*, *Azolla imbricate* and *Spirogyra communis*, would  
320 be a potential threat to a number of higher trophic-level organisms, especially the migratory birds, once  
321 they enter their food chain. In addition, the local people often take some of the aquatic plants, such as *C.*  
322 *demersum*, *P. malainus*. and *Azolla imbricata*, as the food for poultry and livestock, by which the  
323 metals accumulated in these plants could probably enter human body through food chain.

## 324 **5. Conclusions**

325 The present field investigation shows that Chaohai wetland has been seriously contaminated by Cd,  
326 Pb and Zn due to the historic zinc smelting activities in the Hezhang area. Contaminated metals in the  
327 wetland sediments were mainly presented in the organic and residual fractions. However, these  
328 fractions of metals were quite unstable and might be changed into other chemical forms, such as the  
329 Fe-Mn oxide fraction, if the substrata conditions change, such as the increase of the redox potential.  
330 The aquatic plants may act as an important trigger to alter the metal fractionations in the Caihao  
331 wetland. The elevated metal accumulations in the plants should deserve to be particularly concerned  
332 since these metals can enter into higher trophic-level organisms, even humans, through food chain.

## 333 **Acknowledgements**

334 The present study was supported by Chinese Academy of Sciences (CAS) through Key Innovation  
335 Project KZCX3-SW-443, the Natural Science Foundation of China (40473049) and the Western Light  
336 of CAS.

## 337 **References**



338 Albers, P.H., Camardese, M.B., 1993. Effects of Acidification on Metal Accumulation by Aquatic  
339 Plants and Invertebrates. 1. Constructed Wetlands. *Environmental Toxicology and Chemistry* 12:  
340 959-967.

341 Almeida, C.M.R., Mucha, A.P., Vasconcelos, M.T.S.D., 2004. Influence of the Sea Rush *Juncus*  
342 *maritimus* on Metal Concentration and Speciation in Estuarine Sediment Colonized by the Plant.  
343 *Environmental Science and Technology* 38: 3112-3118.

344 Baldantoni, D., Maisto, G., Bartoli, G., Alfani, A., 2005. Analyses of the Three Native Aquatic Plant  
345 Species to Assess Spatial Gradients of Lake Trace Element Contamination. *Aquatic Botany* 83:  
346 48-60.

347 Bi, X., Feng, X., Yang, Y., Qiu, G., Li, G., 2006a. Quantitative assessment of cadmium emission from  
348 zinc smelting and its influences on the surface soils and mosses in Hezhang County, Southwestern  
349 China. *Atmospheric Environment* 40, 4228-4233.

350 Bi, X., Feng, X., Yang, Y., Qiu, G., Li, G., Li, F., Liu, T., Fu, Z., Jin, Z., 2006b. Environmental  
351 contamination of heavy metals from zinc smelting areas in Hezhang County, western Guizhou,  
352 China. *Environmental International* 32, 883-890.

353 Birch, G. 2003. A scheme for assessing human impacts on coastal aquatic environments using  
354 sediments. In: Woodcoffe, C.D., Furness, R.A. (Eds.), *Coastal GIS 2003*. Wollongong University  
355 Papers in Center for Maritime Policy, 14, Australia.

356 Carroll, S., O'Day, P.A., Esser, B., Randall, S., 2002. Speciation and fate of trace metals in estuarine  
357 sediments under reduced and oxidized conditions, Seaplane Lagoon, Alameda Naval Air Station  
358 (USA). *Geochemical Transactions* 3, 81-101.

359 China Environmental Monitoring Station, 1990. Element background values of China's soil. China

360 Environment Science Press, Beijing.

361 Cymerman-Samecka, A., Kempers, A.J., 2004. Toxic metals in aquatic plants surviving in surface  
362 water polluted by copper mining industry. *Ecotoxicology and Environmental Safety* 59, 64-69.

363 Deng, H., Ye, Z.H., Wong, M.H., 2004. Accumulation of lead, zinc, copper and cadmium by 12  
364 wetland plant species thriving in metal-contaminated sites in China. *Environmental Pollution* 132,  
365 29-40.

366 Dollar, N.L., Souch, C.J., Filippelli, G.M., Mastalerz, M., 2001. Chemical fractionation of metals in  
367 wetland sediments: Indiana dunes national lakeshore. *Environmental Science and Technology* 35,  
368 3608-3615.

369 Feng, X., Li, G., Qiu, G., 2004. A preliminary study on mercury contamination to the environment  
370 from artisanal zinc smelting using indigenous method in Hezhang County, Guizhou, China: Part 1.  
371 Mercury emission from zinc smelting and its influences on the surface waters. *Atmospheric  
372 Environment* 38, 6223-6230.

373 Feng, X., Li, G., Qiu, G., 2006. A preliminary study on mercury contamination to the environment  
374 from artisanal zinc smelting using indigenous method in Hezhang County, Guizhou, China: Part 2.  
375 Mercury contaminations to soil and crop. *Science of the Total Environment* 368, 47-55.

376 Gallon, C., Tessier, A., Gobeil, C., 2006. History perspective of industrial lead emissions to the  
377 atmosphere from a Canadian smelter. *Environmental Science and Technology* 2006, 741-747.

378 Gélinas, Y., Schmit, J-P., 1997. Extending the use of the stable lead isotope ratios as a tracer in  
379 bioavailability studies. *Environmental Science and Technology* 1997, 1968-1972.

380 Gomez-Ariza, J.L., Giraldez, I., Sanchez-Rodas, D., Morales, E., 1999. Metal readsorption and  
381 redistribution during the analytical fractionation of trace elements in oxic estuarine sediments.

382 Analytica Chimica Acta 399, 295-307.

383 Hozhina, E.I., Khramov, A.A., Gerasimov, P.A., Kumarkov, A.A., 2001. Uptake of heavy metals,  
384 arsenic, and antimony by aquatic plants in the vicinity of ore mining and processing industries.  
385 Journal of Geochemical Exploration 74, 153-162.

386 Jacob, D.L., Otte, M.L., 2004a. Long-term effects of submergence and wetland vegetation on metals in  
387 a 90-year old abandoned Pb-Zn mine tailings pond. Environmental Pollution 130, 337-345.

388 Jacob, D.L., Otte, M.L., 2004b. Influence of *Typha latifolia* and fertilization on metal mobility in two  
389 different Pb-Zn mine tailing types. Science of the Total Environment 333, 9-24.

390 Kersten, M., Förstner, U., 1986. Chemical fractionation of heavy metals in anoxic estuarine and coastal  
391 sediments. Water Science and Technology 18, 121-130.

392 Lee, C.S., Li, X., Shi, W., Cheung, S.C., Thornton, I., 2006. Metal contamination in urban, suburban,  
393 and country park soils of Hong Kong: A study based on GIS and multivariate statistics. Science of  
394 the Total Environment 356, 45-61.

395 Lu, R.K., 1999. Analytical methods for soil and agricultural chemistry. Beijing: China Agricultural  
396 Science and Technology Press.

397 Madureira, M.J., Vale, C., Goncalves, M.L., 1997. Effect of plants on sulphur geochemistry in the  
398 Tagus salt-marshes sediments. Marine Chemistry 58, 27-37.

399 Mukai, H., Furuta, N., Fujii, T., Ambe, Y., Sakamoto, K., Hashimoto, Y., 1993. Characterization of  
400 sources of lead in the urban air of Asia using ratios of stable lead isotopes. Environmental Science  
401 and Technology 27, 1347-1356.

402 Mukai, H., Tanaka, A., Fujii, T., Zeng, Y., Hong, Y., Tang, J., Guo, S., Xue, H., Sun, Z., Zhou, J., Xue,  
403 D., Zhao, J., Zhai, G., Gu, J., Zhai, P., 2001. Regional characteristics of sulfur and lead isotope

404 ratios in the atmosphere at several Chinese urban sites. *Environmental Science and Technology* 35,  
405 1064-1071.

406 Nirel, P.M.V., Morel, F.M.M., 1990. Pitfalls of sequential extractions. *Water Research* 24, 1055-1056.

407 Olivares-Rieumont, S., Rosa, D., Lima, L., Graham, D.W., Alessandro, K.D., Borroto, J., Martínez, F.,  
408 Sánchez, J., 2005. Assessment of heavy metal levels in Almendares River sediments—Havana City,  
409 Cuba. *Water Research* 39, 3945-3953.

410 Outridge, P.M., Noller, B.N., 1991. Accumulation of toxic trace elements by freshwater vascular plants.  
411 *Reviews of Environmental Contamination and Toxicology* 121, 1-63.

412 Pardo, R., Barrado, E., Perez, L., Vega, M., 1990. Determining and association of heavy metals in  
413 sediments of the Pisucrga river. *Water Research*. 24, 373-379.

414 Rai, U.N., Sinha, S., Tripathi, R.D., Chandra, P., 1995. Wastewater treatability potential of some  
415 aquatic macrophytes: removal of heavy metals. *Ecological Engineering* 5, 5-12.

416 Rapin, F., Tessier, A., Campbell, P.G.C., Carignan, R., 1986. Potential artifacts in the determination of  
417 metal partitioning in sediments by a sequential extraction procedure. *Environmental Sciences and*  
418 *Technology* 20, 836-840.

419 Stoeppler, M., 1992. *Hazardous metals in environment*. Elsevier Press, Amsterdam.

420 Telmer, K., Bonham-Carter, G.F., Kliza, D.A., Hall, G.E.M., 2004. The atmospheric transport and  
421 deposition of smelter emissions: Evidence from the multi-element geochemistry of snow, Quebec,  
422 Canada. *Geochimica et Cosmochimica Acta* 68, 2961-2980.

423 Tessier, A., Campbell, P.G.C., Blsson, M., 1979. Sequential extraction procedure for the speciation of  
424 particulate trace metals. *Analytical Chemistry* 51, 844-851.

425 Ure, A.M., Quevauviller, Muntau, H., Griepink, B., 1993. Speciation of heavy metals in solids and

426 harmonization of extraction techniques undertaken under the auspices of the BCR of the  
427 Commission of the European Communities. *International Journal of Environmental Analytical*  
428 *Chemistry* 51, 135-151.

429 Wong, C.S.C., Li, X.D., 2004. Pb contamination and isotopic composition of urban soils in Hong Kong.  
430 *Science of the Total Environment* 2004, 185-195.

431 Yang, Y., Liu, C., Wu, P., Zhang, G., Zhu, W., 2006. Heavy metal accumulation from zinc smelters in  
432 a carbonate rock region in Hezhang County, Guizhou Province, China. *Water, Air, and Soil*  
433 *Pollution* 174, 321-339.

434 Yurukova, L., Kochev, K., 1994. Heavy metal concentrations in freshwater macrophytes from the  
435 Aldomirovsko Swamp in the Sofia District, Bulgaria. *Bulletin of Environmental Contamination and*  
436 *Toxicology* 52, 627-632.

437 Zheng, C., 1994. An approach on the source of ore-forming metals of lead-zinc deposits in  
438 northwestern part, Guizhou. *Journal of Guilin College of Geology* 14, 113-124 (in Chinese with  
439 English abstract).

440 Zhu, B., Chen, Y., Peng, J., 2001. Lead isotope geochemistry of the urban environment in the Pearl  
441 River Delta. *Applied Geochemistry* 16, 409-417.

442

443 Fig. 1

444 Study area (Caohai wetland) and sampling locations.

445 Fig. 2

446 Percentages of different chemical fractions of Cd, Pb, Zn and Cu for the surface (a) and core (b)

447 sediments collected from Caohai.

448 Fig. 3

449 A plot of  $^{206}\text{Pb}/^{207}\text{Pb}$  vs  $^{208}\text{Pb}/^{207}\text{Pb}$  for different samples.

450

451 Table 1

452 Metal concentrations ( $\mu\text{g g}^{-1}$ , dry weight), EF and organic matter (%) of sediments

Sample ID	Depth (cm)	OM	Cd	Pb	Zn	Cu	EFCd	EFPb	EFZn	EFCu
Surface sediment										
1	0-5	23	56	120	1100	25	61	9.4	31	2.7
2	0-5	29	29	84	630	17	14	2.8	7.5	0.8
3	0-5	20	71	160	1200	22	38	5.8	16	1.2
4	0-5	32	51	140	1000	30	41	7.6	21	2.4
5	0-5	44	16	120	430	33	7.6	3.8	5.0	1.5
6	0-5	38	9.6	76	190	28	5.5	3.1	2.7	1.6
7	0-5	33	16	92	380	28	9.8	4.0	5.9	1.7
8	0-5	24	6.0	69	160	25	6.9	5.6	4.7	2.8
9	0-5	32	3.0	70	140	29	1.7	2.7	1.9	1.6
10	0-5	14	2.6	66	130	28	1.2	2.2	1.6	1.3
Core 1										
	2	44	69	170	1200	28	91	16	41	3.6
	5	42	76	190	1400	32	89	16	40	3.8
	7	43	81	200	1500	32	91	16	42	3.6
	10	46	51	140	1100	34	48	9.5	27	3.2
	15	47	10	68	320	39	6.4	3.1	5.1	2.5
	20	32	2.1	62	190	36				
	25	28	1.7	57	170	39				
	30	29	2.2	58	160	31				
	35	14	2.8	38	130	16				
	40	14	3.1	40	130	17				
Core 2										
	2	30	51	140	1100	31	39	7.4	21	2.3
	5	31	52	130	1000	28	43	7.8	21	2.4
	7	31	51	140	1000	30	42	8.0	21	2.4
	10	33	50	110	1000	22	51	8.2	26	2.2
	15	25	25	71	660	22	21	4.1	13	1.8
	20	17	4.9	35	240	24				
	25	15	3.3	34	230	26				
	30	23	2.0	32	180	25				
	35	26	1.9	28	110	20				
	40	16	2.2	27	88	21				
	45	13	2.8	33	70	19				
	50	1.2	1.5	29	58	23				

453

454

455 Table 2

456 Pb isotope ratios measured in the selected sediments and the local background soils

Sample type	Depth (cm)	$^{204}\text{Pb}/^{207}\text{Pb}$	$^{206}\text{Pb}/^{207}\text{Pb}$	$^{208}\text{Pb}/^{207}\text{Pb}$
Sediment				
	2	0.0636	1.1796	2.4611
	5	0.0637	1.1791	2.4664
	7	0.0637	1.1778	2.4682
	10	0.0637	1.1788	2.4664
	15	0.0636	1.1855	2.4715
	20	0.0638	1.2016	2.4732
Local background soil				
		0.0635	1.2442	2.4776
		0.0635	1.2451	2.4771
		0.0635	1.2489	2.4781
Zinc ore				
smithsonite		0.0635	1.1760	2.4837
smithsonite		0.0635	1.1756	2.4987
sphalerite		0.0636	1.1877	2.4975

457

458




459 Table 3

460 Metal concentrations in different aquatic plant species collected from Caihai ( $\mu\text{g g}^{-1}$ , dry weight)

Species	N	Cd	Pb	Zn	Cu
Submerged					
<i>Ceratophyllum demersum</i>	2	8.2 (6.8-9.6)	22 (20-25)	200 (170-220)	18 (17-18)
<i>Myriophyllum spicatum</i>	4	2.8 (1.9-4.6)	14 (2.3-30)	120 (85-180)	17 (7.4-21)
<i>Potamogeton malainus</i>	3	1.6 (1.2-2.1)	2.4 (1.2-4.6)	62 (52-69)	14 (9.0-22)
<i>Potamogeton pectinatus</i>	3	3.0 (2.0-4.8)	8.2 (7.8-8.7)	90 (80-107)	16 (12-23)
<i>Potamogeton lucens</i>	1	3.2	13	120	20
<i>Potamogeton maackianus</i>	1	2.6	4.6	84	26
Floating					
<i>Azolla imbricata</i>	3	4.4 (4.2-4.7)	12 (9.4-17)	100 (95-110)	11 (7.2-15)
<i>Spirogyra communis</i>	3	4.2 (2.3-7.5)	20 (13-31)	99 (69-150)	16 (7.2-26)
Emergent					
<i>Alternanthera philoxeroides</i>	3	1.5 (1.1-1.7)	10 (2.7-22)	82 (78-88)	15 (13-16)
<i>Scirpus triqueter</i>	3	0.41 (0.35-0.46)	1.4 (0.44-3.3)	34 (19-43)	12 (11-13)
<i>Nymphoides peltata</i>	3	2.6 (2.1-2.9)	9.5 (5.0-14)	81 (71-92)	18 (14-25)

461 Metal concentrations are presented in mean and range values (in parenthesis); N, number of analyzed  
462 samples.


463




464

465 Fig. 1


466




467



468




469




470

471


Fig. 2a




472



473




474



475

476

Fig.2b



477

478 Fig. 3