

Functional and Radiologic Outcomes of Degenerative Versus Traumatic Full-Thickness Rotator Cuff Tears Involving the Supraspinatus Tendon

Cornelia Baum,* MD, Laurent Audigé, PhD, Thomas Stojanov, MSc, Sebastian A. Müller, MD, Christian Candrian, MD, ARCR_Pred Study Group, and Andreas M. Müller, MD
Investigation performed at University Hospital Basel, Basel and the Schulthess Klinik, Zurich, Switzerland

Background: Arthroscopic rotator cuff repair (ARCR) is among the most commonly performed orthopaedic procedures. Several factors—including age, sex, and tear severity—have been identified as predictors for outcome after repair. The influence of the tear etiology on functional and structural outcome remains controversial.

Purpose: To investigate the influence of tear etiology (degenerative vs traumatic) on functional and structural outcomes in patients with supraspinatus tendon tears.

Study Design: Cohort study; Level of evidence, 2.

Methods: Patients undergoing ARCR from 19 centers were prospectively enrolled between June 2020 and November 2021. Full-thickness, nonmassive tears involving the supraspinatus tendon were included. Tears were classified as degenerative (chronic shoulder pain, no history of trauma) or traumatic (acute, traumatic onset, no previous shoulder pain). Range of motion, strength, the Subjective Shoulder Value, the Oxford Shoulder Score (OSS), and the Constant-Murley Score (CMS) were assessed before (baseline) and 6 and 12 months after ARCR. The Subjective Shoulder Value and the OSS were also determined at the 24-month follow-up. Repair integrity after 12 months was documented, as well as additional surgeries up to the 24-month follow-up. Tear groups were compared using mixed models adjusted for potential confounding effects.

Results: From a cohort of 973 consecutive patients, 421 patients (degenerative tear, $n = 230$; traumatic tear, $n = 191$) met the inclusion criteria. The traumatic tear group had lower mean baseline OSS and CMS scores but significantly greater score changes 12 months after ARCR (OSS, 18 [SD, 8]; CMS, 34 [SD, 18] vs degenerative: OSS, 15 [SD, 8]; CMS, 22 [SD, 15]) ($P < .001$) and significantly higher 12-month overall scores (OSS, 44 [SD, 5]; CMS, 79 [SD, 9] vs degenerative: OSS, 42 [SD, 7]; CMS, 76 [SD, 12]) ($P \leq .006$). At the 24-month follow-up, neither the OSS (degenerative, 44 [SD, 6]; traumatic, 45 [SD, 6]; $P = .346$) nor the rates of repair failure (degenerative, 14 [6.1%]; traumatic 12 [6.3%]; $P = .934$) and additional surgeries (7 [3%]; 7 [3.7%]; $P = .723$) differed between groups.

Conclusion: Patients with degenerative and traumatic full-thickness supraspinatus tendon tears who had ARCR show satisfactory short-term functional results. Although patients with traumatic tears have lower baseline functional scores, they rehabilitate over time and show comparable clinical results 1 year after ARCR. Similarly, degenerative and traumatic rotator cuff tears show comparable structural outcomes, which suggests that degenerated tendons retain healing potential.

Keywords: degenerative rotator cuff tear; rotator cuff repair; shoulder arthroscopy; shoulder surgery outcome; traumatic rotator cuff tear

Arthroscopic rotator cuff repair (ARCR) is the most frequently performed shoulder surgery in patients aged 50 to 70 years.^{22,23} The incidence of degenerative rotator cuff tears is clearly age-related, but traumatic events

may cause cuff tears in all adult age groups.^{12,39,40} The distinction between degenerative and traumatic tear etiology is important because the success of tendon repair may be limited by the restricted regenerative properties of degenerated tendons.^{2,13,28} Nonetheless, these biological observations have limited translation to clinical healing, and the literature on the effect of tear etiology on structural and functional outcomes after ARCR is scarce.^{1,8,19,25,31,36} Previous studies comparing degenerative versus traumatic tears also lack the necessary adjustments in their analyses



to consider tear severity, even though it is one of the most relevant predictors of structural outcome after ARCR.^{24,26,27} Thus, they are prone to selection bias. By restricting analyses to more homogeneous patient groups, causal associations can be assessed more accurately. Therefore, this study aimed to compare functional and radiologic outcomes of comparable patient groups with isolated degenerative or traumatic full-thickness and non-massive tears involving the supraspinatus tendon. We hypothesized that patients with traumatic tears would have better functional and structural outcomes over those with degenerative rotator cuff tears and, therefore, would derive greater benefits from ARCR.

METHODS

Patient Cohort and Allocation to Degenerative Versus Traumatic Tear Groups

The patient sample was drawn from a prospective multi-center cohort implemented to assess the safety and effectiveness of ARCR, as described elsewhere.⁴

Briefly, after ethics approval was obtained, a cohort of 973 patients who had primary ARCR was prospectively enrolled at 19 orthopaedic centers, 18 Swiss and 1 German, between June 2020 and November 2021, and observed for 24 months postoperatively.⁴ Procedures were performed all-arthroscopically, with various surgical techniques used in both groups, including single row, conventional double row, or transosseous equivalent (with and without knots) (Table 1).²⁹

Likewise, postoperative rehabilitation protocols varied among treating surgeons and patients in both groups. Functional and structural outcomes, patient-reported outcome measures, and adverse events were recorded at the 6-week, 6-, 12-, and 24-month follow-ups. Patients with full-thickness rotator cuff tears involving the supraspinatus tendon were selected from this specific cohort. Massive tears—full-thickness tear of ≥ 2 tendons as defined by Gerber et al¹⁸—and partial tears were excluded to avoid any imbalances favoring the causes of traumatic tears (64%) and degenerative tears (80%), respectively.

Patients were allocated to 1 of the 2 groups based on the patient's medical history and the surgeons' assessment of the cause of the tear. Patients with chronic (ie, symptomatic for >3 months) shoulder pain with no history of trauma or previous shoulder surgery were included in

TABLE 1
Distribution of Repair Methods Used in the Degenerative and Traumatic Groups^a

Suture Technique SSP	Degenerative	Traumatic	StdDiff
Single row	42 (18)	21 (11)	0.237
Conventional double row	60 (26)	45 (24)	
TOE with knots	87 (38)	88 (47)	
TOE without knots	41 (18)	35 (19)	
Partial repair and debridement		2 (1)	

^aData are presented as n (%). SSP, supraspinatus; StdDiff, standardized difference; TOE, transosseous equivalent.

the degenerative group. Patients who recalled an acute onset of symptoms after a traumatic event, had no history of previous shoulder surgery or complaints, and underwent ARCR within 6 months after the traumatic event were allocated to the traumatic group.

Preoperative Baseline Characteristics

Age at surgery, sex, dominance of the affected shoulder, smoking status, body mass index, diabetes status, the American Society of Anesthesiologists physical status classification (I to IV), previous nonoperative interventions (physical therapy or steroid infiltrations), and the number of steroid infiltrations were recorded before surgery. In addition, the following were assessed by physical examination and patient questionnaire: (1) range of motion—that is, active shoulder flexion, abduction, external and internal rotation; (2) strength at 90° of abduction—measured with a handheld dynamometer; mean strength values of 3 consecutive measurements recorded for each patient; (3) pain levels measured based on the numeric rating scale (NRS), ranging from 0 (no pain) to 10 (highest possible pain); (4) the Subjective Shoulder Value (SSV); (5) the Constant-Murley Score (CMS); (6) the Oxford Shoulder Score (OSS); and (7) the EQ5D5L utility index. Osteoarthritis grade (Samilson Prieto classification³⁴), tear pattern adapted from the classification of Collin et al,¹⁰ supraspinatus tendon atrophy by the areal percentage,³⁸ fatty infiltration of cuff muscles (magnetic resonance imaging [MRI]—adapted Goutallier classification¹⁵), and tendon retraction (Patte classification³⁰) were recorded on preoperative baseline MRI and radiographs. Intraoperative findings were also recorded—such as superior labrum anterior to posterior lesions and humeral or glenoid cartilage lesions.

*Address correspondence to Cornelia Baum, MD, Department of Orthopaedic and Trauma Surgery, University Hospital Basel, Spitalstrasse 21, Basel, 4031, Switzerland. (email: cornelia.baum@usb.ch).

All authors are listed in the Authors section at the end of this article.

Submitted June 16, 2023; accepted October 2, 2023.

One or more of the authors has declared the following potential conflict of interest or source of funding: This project was funded by the Swiss National Science Foundation (SNF Project ID 320030_184959, <http://p3.snf.ch/project-184959>). Complementary grants were provided by Swiss Orthopedics to support project site documentation. The following sites are funding their participation in the project: Charité Medicine University, Berlin, Germany; Public Hospital Solothurn, Solothurn, Switzerland; Endoclinic, Zurich, Switzerland; Inselspital, Bern, Switzerland; and University Clinic Balgrist, Zurich, Switzerland. C.B. received funding for her research salary from the University Hospital Basel, Department of Surgery. AOSSM checks author disclosures against the Open Payments Database (OPD). AOSSM has not conducted an independent investigation on the OPD and disclaims any liability or responsibility relating thereto.

Outcome Parameters

Clinical outcome parameters—including active range of motion, strength at 90° of abduction, and CMS—were assessed 6 months and 1 year after surgery. All consultations with clinical examination were performed in a face-to-face study visit. Overall patient satisfaction, OSS, SSV, pain, and the EQ5D5L utility index were documented via online or paper-based patient questionnaires at the 6-, 12-, and 24-month follow-ups, or if we did not get a response, via phone calls. Adverse events were documented according to a standardized core set⁵ on separate adverse event forms throughout the 2-year postoperative period and reviewed for plausibility by 2 fellowship-trained orthopaedic surgeons (C.B. and S.A.M.). Adverse events included symptomatic recurrent rotator cuff defects diagnosed by ultrasound or MRI that led to a change in treatment. Patients also underwent a routine ultrasound check of their operated shoulder 1 year after surgery to document asymptomatic recurrent defects of the repaired tendons. A Delphi process with all ultrasound operators was conducted before the follow-up examinations, and training videos were provided to improve comparability among the ultrasound assessments. The ultrasound operators were radiologists with musculoskeletal specialty training ($n = 21$), experienced rheumatologists, or orthopaedic surgeons ($n = 8$). Of all sonographers, 72% had >10 years of work experience. Repair integrity was documented at 12 months via a composite outcome parameter combining the occurrence of a symptomatic rotator cuff adverse event ≤ 12 months after surgery and the diagnosis of a complete recurrent defect of any repaired tendon by ultrasound. The occurrence of a secondary surgery for any reason was also documented. Finally, patients were asked how satisfied they were with the overall result of their shoulder treatment—with the NRS, which ranges from 0 (not at all satisfied) to 10 (fully satisfied)—and they were asked if they would opt for ARCR again with this knowledge of satisfaction level (no; I do not know; yes).

Data Management and Statistical Analysis

Study data were managed using the REDCap Electronic Data Capture system²⁰ and exported for statistical analysis using Intercooled Stata Version 17 (StataCorp LP). Baseline patient parameters were tabulated separately per group using standard descriptive statistics and compared using clinical judgment and standardized differences (StdDiffs),⁶ where values closest to 0.10 indicate stronger group similarity.

We compared both groups using generalized linear mixed models to account for repeated measurements when outcome data were available at each follow-up time point, as applicable. All models were adjusted for sex and the number of preoperative steroid infiltrations. Regression coefficients (beta) were estimated along with their 95% CIs and the results of the adjusted t test. Dichotomous outcomes were tabulated by the study group with absolute and relative frequencies (risk) and compared between groups using adjusted logistic regression. Relative risks

(RRs) were calculated along with their 95% CIs. All analyses were explorative, with a significance level set at .05.

The sample size was not a priori estimated for this analysis, as all eligible patients were selected from the ARCR_Pred cohort. Nevertheless, considering a minimal important difference of 5 points on the CMS¹¹ and the observed variability of this score in our cohort (SD, 11), this analysis had a power of 99% to identify such a minimal important difference at the 12-month follow-up, with a significance level set at 5%.

RESULTS

Patient Group Baseline Characteristics

From the ARCR_Pred cohort of 973 patients, 421 patients met the inclusion criteria; degenerative rotator cuff tears were diagnosed in 230 patients (55%), and 191 patients (45%) had tears of traumatic origin (Figure 1). Follow-up rates were 90% (377/421) for the study visits at the 12-month follow-up and 95% (400/421) and 84% (223/267) for the patient-reported questionnaires at the 12- and 24-month follow-ups, respectively (Table 2).

The difference in age distribution between groups was negligible (StdDiff, 0.334), and there was a higher percentage of women in the degenerative compared with the traumatic group (StdDiff, 0.457) (Table 2). A higher proportion of patients with degenerative tears (76%) underwent non-operative interventions before surgery compared with the traumatic group (55%), which was associated with a higher number of steroid infiltrations administered to the degenerative group (StdDiff, 1.049).

Functional Outcome

A detailed overview of the range of motion data per tear group is shown in Appendix Table A1 (available in the online version of this article). Patients in the traumatic group had lower mean baseline values for flexion (109° vs 137°), abduction (96° vs 129°), and internal rotation (34° vs 40°) compared with those with degenerative tears. Even though the absolute values for flexion, abduction, and external rotation at 6 and 12 months after ARCR were similar between the groups, there were significant changes in the mean values of flexion (degenerative [$n = 19$] vs traumatic [$n = 48$]), abduction (degenerative [$n = 25$] vs traumatic [$n = 59$]), and internal rotation (degenerative [$n = 6$] vs traumatic [$n = 14$]) from baseline to the 12-month follow-up ($P \leq .007$). In addition, patients with traumatic tears achieved higher overall internal rotation at the 12-month follow-up over the patients with degenerative tears (49 vs 46, respectively; beta, 6.68 [95% CI, 1.17-12.2]; $P = .018$) (Figure 2).

The mean baseline CMS scores in the traumatic group were lower (44 vs 54). Yet, the score changes by 6 and 12 months after ARCR were significantly higher, with higher overall scores at the 12-month follow-up, when compared with the degenerative group ($P \leq .006$) (Figure 3 and see Appendix Table A2, available online). Similarly, the




Figure 1. Patient selection flowchart highlighting follow-up rates at 6, 12, and 24 months after ARCR. ARCR, arthroscopic rotator cuff repair; SSP, supraspinatus.

traumatic group showed greater improvements in strength at 90° of abduction from baseline to both the 6- and 12-month follow-ups ($P < .001$); nonetheless, no significant difference was observed in the 12-month strength values between groups ($P = .16$). For patients with traumatic tears, the OSS was significantly higher at the 6 and 12 month follow-ups ($P < .032$); nevertheless, this score did not significantly differ with that of the degenerative group by the 24 months follow-up ($P = .346$).

Structural Outcome, Secondary Surgeries, and Adverse Events

Ultrasound examinations at the 1-year follow-up revealed symptomatic recurrent defects in 14 of 230 patients (6.1%) with degenerative tears and 12 of 191 patients (6.3%) with traumatic tears (adjusted RR, 1.4 [95% CI, 0.56-3.5]; $P = .934$). No significant difference was found regarding the proportions of secondary surgeries for the degenerative (7/230 [3%]) and traumatic (7/191 [3.7%]) groups; 7 additional surgeries were completed in each group (adjusted RR, 0.94 [95% CI, 0.30-2.9]; $P = .723$). Reasons for these interventions included removal of a displaced anchor (degenerative, $n = 2$; traumatic, $n = 1$), intra-articular lavage and wound irrigation due to infection (traumatic, $n = 1$), additional biceps tenodesis for persistent biceps pain (degenerative, $n = 1$; traumatic, $n = 1$), or ruptured tendon and revision rotator cuff reconstruction with or without patch augmentation (degenerative, $n = 4$; traumatic, $n = 4$). Similarly, no significant difference was found between groups for any recorded adverse event (Table 3).

Quality of Life and Level of Satisfaction

The mean EQ5DL utility indices were similar for the degenerative and traumatic groups at 12 and 24 months

after ARCR ($P = .215$) (see Appendix Table A3, available online).

The mean overall levels of patient satisfaction at 12 months were 8.8 (SD, 1.8) and 9.1 (SD, 1.4) for the degenerative and traumatic groups, respectively ($P = .083$). By 24 months, the respective mean satisfaction levels were 9.2 (SD, 1.7) and 9.3 (SD, 1.6) ($P = .595$). At 12 months, most patients in the degenerative (84%) and traumatic (90%) groups stated that they would undergo the surgery again (adjusted $P = .185$). By the final follow-up, the proportions of patients with degenerative (92%) and traumatic tears (91%) opting for ARCR again remained high (adjusted $P = .777$).

DISCUSSION

In this analysis, we compared 2 homogeneous groups of patients with full-thickness supraspinatus tendon tears caused by either degenerative changes or traumatic events. Even though both types of rotator cuff tears can cause pain and disability, our findings suggest that their patterns of baseline characteristics and postoperative recovery of functional outcomes differ. Patients with traumatic tears had significantly lower subjective as well as objective shoulder function scores at baseline, yet with significantly greater recovery at the 1-year follow-up. One year after ARCR, patients with traumatic tears showed significantly better CMS scores; however, with minimal clinically important differences of 5 to 10 reported in the literature,¹¹ this difference may not be clinically relevant. Moreover, patient-reported outcomes—such as OSS, SSV, EQ5D5L, and overall patient satisfaction—were no longer significantly different by the 24-month follow-up, indicating that the tear etiology no longer affects outcome after this time.

TABLE 2
Baseline Descriptive Data and Comparison Between Degenerative and Traumatic Groups^a

Parameter	Degenerative (n = 230)		Traumatic (n = 191)		StdDiff	Diff (95% CI)
	n (%)	Mean (SD)	n (%)	Mean (SD)		
Age at surgery		59 (9)		57 (8)	0.334	-3 (-4 to -1)
Sex					0.457	
Female	119 (52)		57 (30)			
Male	111 (48)		134 (70)			
BBody mass index		27 (4)		27 (5)	0.039	0 (-1 to 1)
Current smoker					0.092	
No	189 (82)		150 (79)			
Yes	41 (18)		41 (21)			
ASA classification					0.286	
I	81 (35)		90 (47)			
II	128 (56)		93 (49)			
III	21 (9)		8 (4)			
Diabetes					0.185	
No	217 (94)		187 (98)			
Yes	13 (6)		4 (2)			
Surgery on the dominant arm					0.112	
No	56 (24)		56 (29)			
Yes	174 (76)		135 (71)			
Previous nonoperative interventions					0.434	
No	56 (24)		85 (45)			
Yes	174 (76)		106 (55)			
Physical therapy					0.490	
No	114 (50)		139 (73)			
Yes	116 (50)		52 (27)			
Steroid infiltrations					1.049	
None	122 (53)		179 (94)			
1	59 (26)		10 (5)			
2 +	49 (21)		2 (1)			
Fatty infiltration of SSP					0.334	
Stage 0	111 (48)		113 (59)			
Stage 1	88 (38)		68 (36)			
Stage 2	27 (12)		10 (5)			
Stage 3	4 (2)					
Fatty infiltration of ISP					0.228	
Stage 0	142 (62)		132 (69)			
Stage 1	76 (33)		48 (25)			
Stage 2	10 (4)		11 (6)			
Stage 3	2 (1)					
Fatty infiltration of SSC					0.193	
Stage 0	163 (71)		151 (79)			
Stage 1	57 (25)		35 (18)			
Stage 2	10 (4)		5 (3)			
SSP atrophy					0.206	
I = normal to slight atrophy	195 (85)		170 (89)			
II/III = medium to severe atrophy	35 (15)		21 (11)			
SSP tendon retraction					0.348	
No retraction	33 (14)		32 (17)			
Grade I	105 (46)		65 (34)			
Grade II	79 (34)		66 (35)			
Grade III	13 (6)		28 (15)			
Osteoarthritis grade					0.059	
Grade 0 (none)	193 (86)		165 (88)			
Grade 1 (mild)	32 (14)		23 (12)			
Intraoperative findings						
SLAP lesion	40 (17)		24 (13)		0.136	
Humeral cartilage lesion	12 (5)		6 (3)		0.104	
Glenoid cartilage lesion	4 (2)		5 (3)		0.06	
Tear profile					0.403	
SSP	95 (41)		56 (29)			
SSP and superior SSC	58 (25)		36 (19)			
SSP and complete SSC	5 (2)		13 (7)			
SSP and ISP	72 (31)		86 (45)			

^aASA classification, American Society of Anesthesiologists Physical Status classification system; Diff, difference; CI, confidence interval; StdDiff, standardized difference; ISP, infraspinatus; SLAP, superior labrum anterior to posterior; SSC, subscapularis; SSP, Supraspinatus.




Figure 2. Comparison of (A) abduction, (B) flexion, (C) internal rotation at 90° of abduction, and (D) external rotation at 0° of abduction at the baseline, 6, and 12 months of follow-up for patients with degenerative and traumatic tears. Means and SEMs are presented, where * indicates statistically significant differences between groups.

With our 2 sizable and comparable patient groups, this prospective study could address the question of whether the cause of tear influences functional and structural outcomes after ARCR. Previous studies investigating the difference between degenerative and traumatic tear repair used mixed definitions for the latter, included heterogeneous tear patterns, and were not adjusted for additional factors affecting the outcome of ARCR.

Despite its clinical relevance, tear etiology is not uniformly defined, as indicated by a more recent systematic literature review addressing the term definitions of “acute” and “traumatic.” For example, only 24 of 46 studies overall on traumatic rotator cuff tears reported that the injuries occurred in previously asymptomatic patients.³² It is important, however, to distinguish purely traumatic tears from acute on chronic tears because the latter occurs in predegenerated tendons with potentially reduced healing capacity.^{2,13,28} Therefore, the term traumatic, as implemented in our study and as suggested by Pogorzelski et al,³² should be restricted to patients with an acute onset of symptoms after a shoulder trauma and who have had no previous shoulder complaints.

The restriction of full-thickness tears involving the supraspinatus tendon in our study is of great importance. Within the initial cohort of 973 patients, a substantial proportion of partial tears were degenerative (80%), and a significant proportion of massive tears were of traumatic origin

(64%). The registry study of Kukkonen et al²⁵ included 306 patients and reported a higher rate of larger tear sizes in patients with traumatic rotator cuff tears, which is in line with our findings that massive tears are more likely to derive from trauma. Therefore, including massive and partial tears would inevitably introduce selection bias.

Because of previously unaddressed methodological issues, the literature comparing degenerative and traumatic tears has been inconsistent. Tan et al³⁶ investigated the influence of the cause of tear on outcome in a cohort of 1300 patients at a single center. They found a more restricted range of motion pre- and postoperatively in the traumatic group but no difference in retear rates between groups. Besides the retrospective design and the short follow-up duration of 6 months, a major limitation was the definition of traumatic tears. Patients were considered to have sustained a traumatic tear if a trauma event was recalled. No information was provided regarding previous shoulder complaints. Furthermore, patients in the traumatic group were included until 24 months after trauma. This duration alone can lead to tendon degeneration in the event of a persisting tear. Braune et al⁸ reported significantly higher CMS for patients with traumatic rotator cuff tears (94) when compared with those with nontraumatic, degenerative tears (75) at the mean post-ARCR follow-ups of 47 and 41 months, respectively. Besides the small sample size of 46 patients, the traumatic group was




Figure 3. (A) The Constant-Murley Score, (B) the pain level on the Numeric Rating Scale (NRS), (C) abduction strength at 90° of abduction, (D) the Subjective Shoulder Value, (E) the EQ5D5L utility index, and (F) the Oxford Shoulder Score for the degenerative and traumatic groups at 6-, 12-, and 24-month follow-ups. Means and SEMs are presented, where * indicates statistically significant differences.

younger, with a mean age of 34 years versus 54 years in the degenerative group. In contrast, our patient groups were very comparable with respect to the mean age. Kukkonen et al²⁵ reported significantly higher CMS in the degenerative group when compared to the traumatic group at baseline (52 vs 46) and 12 months postoperatively (77 vs 73), and both groups had comparable patient ages. Abechain et al¹ reported a lack of significant difference in functional outcomes between their degenerative and traumatic groups that shared similar mean ages of 59.9 and 59.0 years, respectively; the authors applied the University of California, Los Angeles, Shoulder Score tool over the

CMS and the study was limited by a small total sample size of 87 patients. Godshaw et al¹⁹ demonstrated higher preoperative functional deficits in the traumatic group but greater improvements in range of motion, strength, and shoulder function compared with their degenerative group, which mirrors our findings. However, it is a single-center study with a smaller sample size (73 traumatic versus 148 atraumatic tears).

It is generally believed that degenerative rotator cuff tears have poorer healing capacity over traumatic rotator cuff tears after repair, although the evidence to prove this theory is meager.^{1,19,25,31,36} Tan et al³⁶ assessed

TABLE 3
Distribution of Local Postoperative Adverse Event Categories and Comparison
Between Degenerative and Traumatic Groups^a

Adverse Event Category	Degenerative		Traumatic		RR (95% CI) ^b	P ^b
	n	Risk (%)	n	Risk (%)		
Any local adverse event	71	30.9	50	26.2	0.92 (0.65-1.3)	.740
Device events ^c	4	1.7	1	0.5		
Osteochondral	3	1.3	1	0.5		
Symptomatic AC arthritis	1		0			
Fracture	2		1			
Persisting or worsening pain	30	13	20	10.5	0.86 (0.47-1.6)	.620
Rotator cuff defect	9	3.9	8	4.2	1.4 (0.46-4.6)	.080
Peripheral neurological	4	1.7	1	0.5		
Nerve injury	1		1			
CRPS	3		0			
Surgical-site infection ^d	0	0	1	0.5		
Superficial soft tissue ^e	2	0.9	0	0		
Deep soft tissue	24	10.4	18	9.4	1.4 (0.46-4.6)	.080
Subacromial space	6		2			
Biceps	3		2			
Capsule (stiffness)	16	7	13	6.8	1.2 (0.59-2.3)	.504
Other deep soft tissue	0		1			
Other postoperative local	5		0			

^aAC, acromioclavicular; CRPS, complex regional pain syndrome; RR, relative risk.

^bRR, 95% CI, and P values—using binomial regression adjusted for sex and the number of preoperative steroid infiltrations—were only calculated if the total number of events was ≥ 10 .

^cAll device events were anchor displacements.

^dDeep surgical-site infection (incisional and organ/space).

^eThese 2 events were edema.

postoperative tendon integrity in correlation with tear size and did not find any significant differences in retear rates between the traumatic and degenerative groups for each tear size, but did report higher retear rates for larger tears. Similarly, Raman et al³³ found insufficient evidence for the influence of the cause of tear on postoperative tendon integrity in their systematic review summarizing 490 patients from 3 studies. In line with these published findings, we could not show any significant differences in retear or revision surgery rates between the 2 groups, which indicates that healing potential is preserved, even in the case of degenerated tendons.

The retear rates for our patient groups were surprisingly low at around 6% when compared with the reported rates of 5% to 92%.^{3,7,16,17,21,35,37} Repair failure may be considered a multifactorial issue because of the biological properties of the tendon and biomechanical factors, and expected healing after ARCR correlates with tear size.^{26,27,33} The retear rate of isolated full-thickness supraspinatus repairs was previously reported to be 13%.¹⁴ Moreover, the relatively younger mean age of our patient groups may also play a role in supporting the healing potential of affected tendons.

Strengths and Limitations

We acknowledge the following limitations of our study. Our classification of rotator cuff tears as traumatic was based on

surgeons' judgment, the history of trauma, and the absence of previous symptoms. Neither preoperative MRI findings nor the mechanism of injury were recognized and documented for these patients at the baseline patient assessments and cannot be accurately assessed at a later stage. Histopathologic differences between degenerative and traumatic tendons were also not evaluated because implementation was not clinically feasible. Thus, we cannot completely rule out that some primary tendon degeneration may be evident in patients classified with traumatic tears.

Another limitation is that repair integrity was measured by ultrasound in a multicenter setting, which may be associated with examiner-dependent variations in accuracy. In addition, patient recruitment was conducted during the COVID-19 pandemic. Thus, we cannot rule out that some enrolled patients experienced a delay in time to the surgical theater and had increased use of steroid infiltrations when compared with similar patient cohorts.

As a strength of our work, the follow-up time of 2 years adequately reflects functional results after ARCR, as it extends beyond the 1-year period in which functional recovery is known to improve after surgery and is followed by stabilization.⁹

CONCLUSION

Patients with degenerative and traumatic full-thickness tears involving the supraspinatus tendon both show

satisfactory functional results at the short-term follow-up after ARCR. Although patients with traumatic tears have lower baseline functional scores, such as the CMS, they rehabilitate over time and demonstrate significantly greater score changes up to the 6- and 12-month follow-ups. At 12 to 24 months after ARCR, clinical differences in functional and patient-reported outcomes are no longer evident. Likewise, structural outcomes are comparable between degenerative and traumatic rotator cuff tears, which indicates that degenerated tendons retain healing potential.

AUTHORS

Cornelia Baum, MD (Department of Orthopaedic and Trauma Surgery, University Hospital Basel, Basel, Switzerland); Laurent Audigé, PhD (Surgical Outcome Research Center, Department of Clinical Research, University Hospital of Basel, Basel, Switzerland; Research and Development, Shoulder and Elbow Surgery, Schulthess Klinik, Zurich, Switzerland); Thomas Stojanov, MSc (Department of Orthopaedic and Trauma Surgery, University Hospital Basel, Basel, Switzerland; Research and Development, Shoulder and Elbow Surgery, Schulthess Klinik, Zurich, Switzerland); Sebastian A. Müller, MD (Department of Orthopaedic and Trauma Surgery, Shoulder and Elbow, Cantonal Hospital Basel, Bruderholz, Switzerland; Faculty of Medicine, University of Basel, Basel, Switzerland); Christian Candrian, MD (Trauma and Ortho Unit, Ospedale Regionale di Lugano, Lugano, Switzerland); ARCR_Pred Study Group; and Andreas M. Müller, MD (Department of Orthopaedic and Trauma Surgery, University Hospital Basel, Basel, Switzerland).

Members of the ARCR_Pred Study Group are listed below per site and partner institution:

ARTHRO Medics, Basel, CH: Claudio Rosso (Principal Investigator [PI]), Lena Fankhauser, Gernot Willscheid; Charité Medicine University, Berlin, DE: Philipp Moroder (PI), Doruk Akgün, Victor Danzinger, Henry Gebauer, Jan-Philipp Imiolczyk, Katrin Karpinski, Lucca Lacheta, Marvin Minkus, Alp Paksoy, Eduardo Samaniego, Kathi Thiele, Isabella Weiss; Cantonal Hospital Baselland, Bruderholz, CH: Thomas Suter (PI), Julia Müller-Lebschi, Sebastian Mueller, Markus Saner, Claudia Haag-Schumacher, Giorgio Tamborrini-Schütz; Public Hospital Solothurn, Solothurn, CH: Mai Lan Dao Trong (PI), Carlos Buitrago-Tellez, Julian Hasler, Ulf Riede, Sandra Weber; Hôpital du Valais—Centre Hospitalier du Valais Romand, Martigny, CH: Beat Moor (PI), Matthias Biner, Sarah Fournier, Nicolas Gallusser, Deborah Marietan, Sebastien Pawlak; Endoclinic, Zurich, CH: Christoph Spormann (PI), Britta Hansen, Nadja Mamisch; Klinik Gut, St Moritz, CH: Holger Durchholz (PI), Jakob Bräm; Hirslanden Clinique la Colline, Geneva, CH: Gregory Cunningham (PI), Abed Kourhani, Sarah Ossipow, Patricia Simao; La Tour Hospital, Meyrin, CH: Alexandre Lädermann (PI); Inselspital, Bern, CH: Rainer Egli, Stephanie Erdbrink, Remy Flückiger, Paolo Lombardo, Tawan Pinworasarn, Philipp Scacchi, Johannes Weihs, Matthias Zumstein; In-Motion, Wallisellen, CH: Matthias Flury (PI), Ralph Berther, Christine Ehrmann, Larissa Hübscher; Institute of Social and Preventive Medicine, University of Bern, Bern, CH: David Schwappach; Cantonal Hospital of Baden, Baden, CH: Karim Eid (PI), Susanne Bensler, Yannick Fritz, Nisha Grünberger, Philipp Kriechling, Daniel Langthaler, Richard Niehaus, Raffaella Nobs; Cantonal Hospital of Winterthur, Winterthur, CH: Emanuel Benninger (PI), Quintin de Groot, Aleksis Doert, Sebastian Ebert, Philemon Grimm, Fabian Mottier, Markus Pisan, Jan Schätz, Ariane Schwank, Julian Wiedenbach; Schulthess Klinik, Zurich,

CH: Markus Scheibel (PI), Laurent Audigé, Frederik Bellmann, Daniela Brune, Marije de Jong, Stefan Diermayr, David Endell, Marco Etter, Florian Freislederer, Nikitas Gkikopoulos, Michael Glanzmann, Cécile Grobet, Christian Jung, Fabrizio Moro, Philipp Moroder, Ralph Ringer, Jan Schätz, Hans-Kaspar Schwyzer, Béatrice Weber, Martina Wehrli, Barbara Wirth, Manuela Nötzli, Anne Franz, Jörg Oswald, Birgit Steiger, Yacine Ameziane, Christopher Child, Giovanni Spagna; Ospedale Regionale di Lugano, Lugano, CH: Christian Candrian (PI), Filippo Del Grande, Pietro Feltri, Giuseppe Filardo, Francesco Marbach, Florian Schöneweger; Cantonal Hospital of St. Gallen, St. Gallen, CH: Bernhard Jost (PI), Michael Badulescu, Stephanie Lüscher, Fabian Napieralski, Lena Öhrström, Martin Olach, Jan Rechsteiner, Jörg Scheler, Christian Spross, Vilijam Zdravkovic; Orthopädie Sonnenhof, Bern, CH: Matthias A. Zumstein (PI), Adrian Chlasta, Kate Gerber, Annabel Hayoz, Julia Müller-Lebschi, Frederick Schuster; University Clinic Balgrist, Zurich, CH: Karl Wieser (PI), Paul Borbas, Samy Bouaicha, Roland Camenzind, Sabrina Catanzaro, Christian Gerber, Florian Grubhofer, Anita Hasler, Bettina Hochreiter, Roy Marcus, Farah Selman, Reto Sutter, Sabine Wyss; University Library Basel, University of Basel, Basel, CH: Christian Appenzeller-Herzog; University Hospital of Basel, Basel, CH: Soheila Aghmandi, Ilona Ahlborn, Cornelia Baum, Franziska Eckers, Kushtrim Grezda, Simone Hatz, Sabina Hunziker, Thomas Stojanov, Mohy Taha, Giorgio Tamborrini-Schütz, Andreas Marc Mueller (PI).

ACKNOWLEDGMENT

The authors thank M. Wilhelmi, PhD (medical writer at Schulthess Klinik), for language editing the final manuscript.

REFERENCES

1. Abechain JJK, Godinho GG, Matsunaga FT, Netto NA, Daou JP, Tamaoki MJS. Functional outcomes of traumatic and non-traumatic rotator cuff tears after arthroscopic repair. *World J Orthop.* 2017;8(8):631-637.
2. Andarawis-Puri N, Flatow EL, Soslowsky LJ. Tendon basic science: development, repair, regeneration, and healing. *J Orthop Res.* 2015;33(6):780-784.
3. Anderson K, Boothby M, Aschenbrenner D, van Holsbeeck M. Outcome and structural integrity after arthroscopic rotator cuff repair using 2 rows of fixation: minimum 2-year follow-up. *Am J Sports Med.* 2006;34(12):1899-1905.
4. Audige L, Bucher HCC, Aghmandi S, et al. Swiss-wide multicentre evaluation and prediction of core outcomes in arthroscopic rotator cuff repair: protocol for the ARCR_Pred cohort study. *BMJ Open.* 2021;11(4):e045702.
5. Audige L, Flury M, Muller AM, Panel ACC, Durchholz H. Complications associated with arthroscopic rotator cuff tear repair: definition of a core event set by Delphi consensus process. *J Shoulder Elbow Surg.* 2016;25(12):1907-1917.
6. Austin PC. Balance diagnostics for comparing the distribution of baseline covariates between treatment groups in propensity-score matched samples. *Stat Med.* 2009;28(25):3083-3107.
7. Boileau P, Brassart N, Watkinson DJ, Carles M, Hatzidakis AM, Krishnan SG. Arthroscopic repair of full-thickness tears of the supraspinatus: does the tendon really heal? *J Bone Joint Surg Am.* 2005;87(6):1229-1240.
8. Braune C, von Eisenhart-Rothe R, Welsch F, Teufel M, Jaeger A. Mid-term results and quantitative comparison of postoperative shoulder function in traumatic and non-traumatic rotator cuff tears. *Arch Orthop Trauma Surg.* 2003;123(8):419-424.

9. Charoussat C, Grimberg J, Duranthon LD, Bellaiche L, Petrover D, Kalra K. The time for functional recovery after arthroscopic rotator cuff repair: correlation with tendon healing controlled by computed tomography arthrography. *Arthroscopy*. 2008;24(1):25-33.
10. Collin P, Matsumura N, Lädermann A, Denard PJ, Walch G. Relationship between massive chronic rotator cuff tear pattern and loss of active shoulder range of motion. *J Shoulder Elbow Surg*. 2014;23(8):1195-1202.
11. Cvetanovich GL, Gowd AK, Liu JN, et al. Establishing clinically significant outcome after arthroscopic rotator cuff repair. *J Shoulder Elbow Surg*. 2019;28(5):939-948.
12. Diebold G, Lam P, Walton J, Murrell GAC. Relationship between age and rotator cuff retear: a study of 1,600 consecutive rotator cuff repairs. *J Bone Joint Surg Am*. 2017;99(14):1198-1205.
13. Fenwick SA, Hazleman BL, Riley GP. The vasculature and its role in the damaged and healing tendon. *Arthritis Res*. 2002;4(4):252-260.
14. Fuchs B, Gilbert MK, Hodler J, Gerber C. Clinical and structural results of open repair of an isolated one-tendon tear of the rotator cuff. *J Bone Joint Surg Am*. 2006;88(2):309-316.
15. Fuchs B, Weishaupt D, Zanetti M, Hodler J, Gerber C. Fatty degeneration of the muscles of the rotator cuff: assessment by computed tomography versus magnetic resonance imaging. *J Shoulder Elbow Surg*. 1999;8(6):599-605.
16. Galatz LM, Ball CM, Teefey SA, Middleton WD, Yamaguchi K. The outcome and repair integrity of completely arthroscopically repaired large and massive rotator cuff tears. *J Bone Joint Surg Am*. 2004;86(2):219-224.
17. Gazielly DF, Gleyze P, Montagnon C. Functional and anatomical results after rotator cuff repair. *Clin Orthop Relat Res*. 1994;304:43-53.
18. Gerber C, Fuchs B, Hodler J. The results of repair of massive tears of the rotator cuff. *J Bone Joint Surg Am*. 2000;82(4):505-515.
19. Godshaw BM, Hughes JD, Boden SA, Lin A, Lesniak BP. Comparison of functional outcomes after arthroscopic rotator cuff repair between patients with traumatic and atraumatic tears. *Orthop J Sports Med*. 2022;10(10):23259671221126551.
20. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42(2):377-381.
21. Harryman DT II, Mack LA, Wang KY, Jackins SE, Richardson ML, Matsen FA III. Repairs of the rotator cuff. Correlation of functional results with integrity of the cuff. *J Bone Joint Surg Am*. 1991;73(7):982-989.
22. Jain NB, Higgins LD, Losina E, Collins J, Blazar PE, Katz JN. Epidemiology of musculoskeletal upper extremity ambulatory surgery in the United States. *BMC Musculoskelet Disord*. 2014;15:4.
23. Karjalainen TV, Jain NB, Heikkinen J, Johnston RV, Page CM, Buchbinder R. Surgery for rotator cuff tears. *Cochrane Database Syst Rev*. 2019;12(12):CD013502.
24. Kim YK, Jung KH, Kim JW, Kim US, Hwang DH. Factors affecting rotator cuff integrity after arthroscopic repair for medium-sized or larger cuff tears: a retrospective cohort study. *J Shoulder Elbow Surg*. 2018;27(6):1012-1020.
25. Kukkonen J, Joukainen A, Itala A, Aarimaa V. Operatively treated traumatic versus non-traumatic rotator cuff ruptures: a registry study. *Ups J Med Sci*. 2013;118(1):29-34.
26. Le BT, Wu XL, Lam PH, Murrell GA. Factors predicting rotator cuff retears: an analysis of 1000 consecutive rotator cuff repairs. *Am J Sports Med*. 2014;42(5):1134-1142.
27. Lee YS, Jeong JY, Park CD, Kang SG, Yoo JC. Evaluation of the risk factors for a rotator cuff retear after repair surgery. *Am J Sports Med*. 2017;45(8):1755-1761.
28. Matthews TJ, Hand GC, Rees JL, Athanasou NA, Carr AJ. Pathology of the torn rotator cuff tendon: reduction in potential for repair as tear size increases. *J Bone Joint Surg Br*. 2006;88(4):489-495.
29. Park MC, Elattrache NS, Ahmad CS, Tibone JE. "Transosseous-equivalent" rotator cuff repair technique. *Arthroscopy*. 2006;22(12):1360.e1361-1365.
30. Patte D. Classification of rotator cuff lesions. *Clin Orthop Relat Res*. 1990;254:81-86.
31. Paul S, Yadav AK, Goyal T. Comparison of tear characteristics, outcome parameters and healing in traumatic and non-traumatic rotator cuff tear: a prospective cohort study. *Musculoskelet Surg*. 2022;106(4):433-440.
32. Pogorzelski J, Erber B, Themessl A, et al. Definition of the terms "acute" and "traumatic" in rotator cuff injuries: a systematic review and call for standardization in nomenclature. *Arch Orthop Trauma Surg*. 2021;141(1):75-91.
33. Raman J, Walton D, MacDermid JC, Athwal GS. Predictors of outcomes after rotator cuff repair—a meta-analysis. *J Hand Ther*. 2017;30(3):276-292.
34. Samilson RL, Prieto V. Dislocation arthropathy of the shoulder. *J Bone Joint Surg Am*. 1983;65(4):456-460.
35. Sugaya H, Maeda K, Matsuki K, Moriishi J. Repair integrity and functional outcome after arthroscopic double-row rotator cuff repair: a prospective outcome study. *J Bone Joint Surg Am*. 2007;89(5):953-960.
36. Tan M, Lam PH, Le BT, Murrell GA. Trauma versus no trauma: an analysis of the effect of tear mechanism on tendon healing in 1300 consecutive patients after arthroscopic rotator cuff repair. *J Shoulder Elbow Surg*. 2016;25(1):12-21.
37. Thomazeau H, Boukobza E, Morcet N, Chaperon J, Langlais F. Prediction of rotator cuff repair results by magnetic resonance imaging. *Clin Orthop Relat Res*. 1997;344:275-283.
38. Thomazeau H, Rolland Y, Lucas C, Duval JM, Langlais F. Atrophy of the supraspinatus belly: assessment by MRI in 55 patients with rotator cuff pathology. *Acta Orthop Scand*. 1996;67(3):264-268.
39. Yamaguchi K, Ditsios K, Middleton WD, Hildebolt CF, Galatz LM, Teefey SA. The demographic and morphological features of rotator cuff disease: a comparison of asymptomatic and symptomatic shoulders. *J Bone Joint Surg Am*. 2006;88(8):1699-1704.
40. Yamamoto A, Takagishi K, Osawa T, et al. Prevalence and risk factors of a rotator cuff tear in the general population. *J Shoulder Elbow Surg*. 2010;19(1):116-120.