

Isolation and Structure Determination of Drought-Induced Multihexose Benzoxazinoids from Maize (*Zea mays*)

Sylvain Sutour,* Van Cong Doan, Pierre Mateo, Tobias Züst, Ella Raymonde Hartmann, Gaétan Glauser, and Christelle Aurélie Maud Robert*




Cite This: <https://doi.org/10.1021/acs.jafc.3c09141>



Read Online

ACCESS |

 Metrics & More

 Article Recommendations

 Supporting Information

ABSTRACT: Benzoxazinoids (BXDs) are plant specialized metabolites exerting a pivotal role in plant nutrition, allelopathy, and defenses. Multihexose benzoxazinoids were previously observed in cereal-based food products such as whole-grain bread. However, their production in plants and exact structure have not been fully elucidated. In this study, we showed that drought induced the production of di-, tri-, and even tetrahexose BXDs in maize roots and leaves. We performed an extensive nuclear magnetic resonance study and elucidated the nature and linkage of the sugar units, which were identified as gentiobiose units β -linked ($1'' \rightarrow 6'$) for the dihexoses and ($1'' \rightarrow 6'$)/($1''' \rightarrow 6''$) for the trihexoses. Drought induced the production of DIMBOA-2Glc, DIMBOA-3Glc, HMBOA-2Glc, HMBOA-3Glc, and HDMBOA-2Glc. The induction was common among several maize lines and the strongest in seven-day-old seedlings. This work provides ground to further characterize the BXD synthetic pathway, its relevance in maize-environment interactions, and its impact on human health.

KEYWORDS: maize, drought, NMR, multihexose benzoxazinoids, DIMBOA dihexose, DIMBOA trihexose

1. INTRODUCTION

Benzoxazinoids (BXDs) are plant specialized metabolites that modulate plant nutrition, reproduction, interactions with microbes, and defenses against herbivores.¹ They are commonly found in cereals and frequently recorded in wholegrain cereal products.^{2–5} BXDs can have cascading effects on food quality and human health, and their consumption is associated with anti-inflammatory and anticancer effects.^{6,7} Understanding the different molecular structures of BXDs is crucial for improving the reliability and robustness of analytical methods, for identifying their underlying biosynthesis genes, and ultimately, for comprehending their potential influence on human health.^{8,9}

BXDs include benzoxazolinones with a 1,3-benzoxazol-2-one core structure and benzoxazinones, possessing a 1,4-benzoxazin-3-one scaffold. In rye, BXD species with no methoxy group on the aromatic ring, such as DIBOA and DIBOA-Glc, are dominant, whereas in maize, methylated forms such as DIMBOA and DIMBOA-Glc are predominant. In wheat, both species can be found in similar amounts.¹⁰ Multihexose BXDs, such as H(M)BOA and DI(M)BOA di-, tri-, and tetrahexoses, were reported in wheat, maize, and rye.^{4,10–16} The structures of dihexose BXDs were described using MS/MS fragmentation as H(M)BOA-Glc-Hex and DI(M)BOA-Glc-Hex.^{4,8,10–14} Consequently, the exact nature and configuration of the hexose units as well as the relative and absolute stereochemistry of multihexose BXDs have remained undefined.

BXDs can transition from plants to the human food supply, where they can impact human health. Food processing, such as watering/drying the seeds, or hydrothermal processing (HTP) of cereals were found to increase BXD levels,

including dihexose BXDs, in processed products.^{3,4,12} For instance, rye-based products contain approximately 122 $\mu\text{g/g}$ dry mass HBOA-Glc-Hex and 300 $\mu\text{g/g}$ dry mass DIBOA-Glc-Hex.⁴ The impact of multihexose BXDs on human health remains unexplored, hindered by several factors, including the unavailability of purified compounds (attributable to their limited presence in healthy plants), a lack of structural characterization, and their omission from analytical methodologies.

Here, we demonstrate that drought enhances the production of multihexose BXDs in maize root and shoot tissues. We conducted an extensive nuclear magnetic resonance (NMR) study to determine the nature and configuration of the multihexose chains. We then characterized their induction patterns and kinetics in different maize tissues. Understanding the structure of these multihexose BXDs is pivotal for enabling their inclusion in analytical methods, enhancing our ability to monitor their presence and impact on food quality and human nutrition.

2. MATERIALS AND METHODS

2.1. Biological Resources. Maize (*Zea mays*, varieties B73, CML277, Hp301, and Oh7B) were obtained from Maize GDB (https://maizegdb.org/data_center/phenotype#stocks) and bred in-house. The seeds were soaked in aqueous bleach (5%, Migros, CH)

Received: December 4, 2023

Revised: January 25, 2024

Accepted: January 26, 2024




Figure 1. Drought induced putative multihexose benzoxazinoids (BXDs) in maize leaves. Base peak intensity (BPI) chromatograms of control (A) and drought-stressed (B) maize leaves. Putative multihexose benzoxazinoids (BXDs) are labeled as numbers in the chromatograms. (1) Putative DIMBOA-dihexose (m/z 534.1459), (2) putative DIMBOA-trihexose (m/z 696.1987), (3) putative HMBOA-dihexose (m/z 518.1510), (4) putative HMBOA-trihexose (m/z 680.2038), and (6) putative HDMBOA-dihexose (m/z 594.1670). m/z : mass-to-charge ratio.

for 5 min, washed with distilled water, and then soaked in distilled water for one night before being placed in preweighed 100 mL plastic pots (4 cm diameter, 11.2 cm height; Platz GmbH Medizintechnik, Dorsten-Wulfen; DE) containing soil (40% sand, 35% silt, and 25% clay (Landerde, Ricoter, Aarberg; CHE)) with 23% moisture. The moisture level was controlled by drying soil in an oven (24 h, 75 °C) and adding the corresponding amount of water prior to planting. The plants were cultivated in a greenhouse with the following conditions: 14 h light (20 °C \pm 2 °C), 10 h dark (18 °C \pm 2 °C), 55% relative humidity, with 250 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ additional light supplied by Master GreenPower 600W 400 V E40 high-pressure sodium bulbs (Philips Lighting, CHE).

2.2. Drought Induction. After planting, the soil moisture was either kept at 23% v/v (ambient conditions) or left to decrease to 16% v/v (drought conditions, reached at day 4 after planting). Ambient and drought conditions were then maintained by weighing the pots and adding the required amounts of water daily. The drought moisture conditions were determined based on the projected RCP8.5 climate scenario, using a correlation between decrease precipitation levels and soil moisture levels as in refs 17–19. A 16.6% v/v moisture led to moderate drought symptoms in maize plants. All plants were harvested 10 days after sowing. Roots and shoots were harvested separately, flash-frozen in liquid nitrogen, and ground to a fine powder using a mortar and a pestle in the presence of liquid nitrogen.

2.3. Benzoxazinoid Analyses. The BXDs analysis was adapted from a previously described protocol.²⁰ Aliquots of powder samples (100 mg) were mixed with 1 mL of a 70:30 mixture of methanol (HPLC grade) and water (milli-Q) containing 0.1% formic acid (LC-MS grade). All extracts were vortexed for 20 s and centrifuged at 13000 rpm for 20 min at 10 °C, and the supernatants were collected for LC-MS analyses. Where necessary, samples were diluted 1:10 prior to analysis.

The BXD profiling was performed with an Acquity i-Class UHPLC system equipped with a photodiode array detector (PDA) and a single quadrupole detector (QDa) with an electrospray source (Waters, USA). Gradient elution was performed on an Acquity BEH C18 (1.7 μm , 2.1 \times 100 mm, Waters, USA) column maintained at 40 °C. The elution conditions were as follows: solvent A, H₂O (Milli-Q) + 0.1% formic acid (LC-MS grade); solvent B, ACN (LC-MS grade) + 0.1% formic acid (LC-MS grade); flow rate, 0.4 mL/min; 0–3 min, linear gradient from 2 to 20% B; 3–6 min, linear gradient to 100% B; 6–8 min, 100% B; 8–10 min, 2% B. Absolute

quantities of DIMBOA-2Glc, DIMBOA-3Glc and HMBOA-2Glc were determined through a semiquantitative method using standard curves of the corresponding pure monoglucoside DIMBOA-Glc for the two first ones and the aglucone HMBOA for the latter.

2.4. Crude Extract. Ten days after germination, root and shoot samples were gently washed in tap water, patted dry with paper, flash frozen in liquid nitrogen, and stored at –80 °C until further processing. The samples were ground manually with mortar and pestle in the presence of liquid nitrogen. Root and shoot samples were pooled to give 450 g of raw material. The extraction was performed in two equal batches (225 g each) as follows: the cold ground material was added to 1.5 L of MeOH and homogenized for 3 min with an immersion disperser (Polytron PT 10–35, Kinematica, CHE), followed by suction filtration through a P3 sintered glass filter equipped with two layers of filter paper. The filter cake was collected and suspended again in 1.5 L MeOH. After a second homogenization and a new filtration, the filtrates were combined and concentrated under reduced pressure on a rotary evaporator. Upon evaporation, a yellow-green sticky residue formed on the round-bottom flask walls. This residue was discarded, and the remaining aqueous phase was separated. The same procedure was repeated for the second batch of frozen ground material. The two aqueous residues obtained were pooled and lyophilized to provide a strong yellow crude dry extract. The presence of compounds 1–7 was verified by HPLC-MS prior to isolation.

2.5. Preparative Chromatography. The crude extract (1.8 g) was first fractionated by open column chromatography (CC) using 45 g of silica gel (63–200 μm , 60 Å) as the stationary phase. A step gradient elution consists of mixtures of dichloromethane-ethyl acetate, followed by ethyl acetate-methanol, and finally pure methanol yielded 11 fractions (F1–11). Compounds 1–7, which were all present in F9 (126.8 mg), were further purified by semipreparative HPLC using a LC-20AR Shimadzu pump (Japan) connected to a UV–vis detector (Shimadzu, model SPD-20A) and a fraction collector FRC-40 (Shimadzu). A detailed description of the semipreparative conditions for each molecule is provided in Supporting Information S11.

2.6. NMR Spectroscopy. NMR spectra were acquired on a Bruker Avance Neo Ascend 600 MHz (Bruker, Germany) in D₂O. Standard pulse sequences were used to acquire 1D and 2D NMR analyses. HSQC and HMBC analyses were recorded in nonuniform sampling (NUS) mode with a sampling amount of 12.5%. Acetone was used as calibration standard ($\delta^1\text{H}$ = 2.225 ppm; $\delta^{13}\text{C}$ = 30.5




Figure 2. Extracted ion chromatograms of multihexose benzoxazinoids detected in control (dashed line) and drought-stressed (solid line) maize leaves. (1) Putative DIMBOA-dihexose (m/z 534.1459), (2) putative DIMBOA-trihexose (m/z 696.1987), (3) putative HMBOA-dihexose (m/z 518.1510), (4) putative HMBOA-trihexose (m/z 680.2038), (5) putative HM₂BOA-dihexose (m/z 548.1615), (6) putative HDMBOA-dihexose (m/z 594.1670), (7) putative DIBOA-dihexose (m/z 504.1353), and (8) putative DIMBOA-tetrahexose (m/z 858.2515). cps: count per second. m/z : mass-to-charge ratio.




Figure 3. Structure of compounds 1 to 7.

ppm) except for compounds 6 and 7. In this case, spectra were calibrated on H-2' (3.25 ppm) and C2' (73.1 ppm). Acquired spectra were processed using the Mnova NMR software package (v.14.2.0, MestReLab Research S.L., Spain).

2.7. Statistical Analysis. All statistical analyses were conducted on Sigma Plot 14.5. The distribution and heteroscedasticity of the data residuals were assessed by using the Shapiro–Wilk and Brown–Forsythe tests. Two-way ANOVAs on ranks were conducted to assess the effects of drought and time on multihexose benzoxazinoid levels. Student *t*-tests and Mann–Whitney tests were performed to evaluate the impact of drought on individual compounds in maize.

3. RESULTS AND DISCUSSION

3.1. Drought Promotes the Production of Putative Multihexose Benzoxazinones in Maize. Although BXDs with multiple hexose units were previously reported,^{4,10–16,21} their exact structures remained unidentified, probably due to their low concentrations in plants. During our investigation

on stress-induced BXDs in maize, we observed that drought induced higher amounts of putative di-, tri-, and tetrahexose BXDs in maize (Figures 1 and 2).

Extracted ion chromatograms revealed drought-increased peaks for DIMBOA-dihexose (m/z 534.1459, 1), DIMBOA-trihexose (m/z 696.1987, 2), HMBOA-dihexose (m/z 518.1510, 3), HMBOA-trihexose (m/z 680.2038, 4) HM₂BOA-dihexose or HDMBOA-dihexose (m/z 548.1615 and 594.1670, 5 and 6), DIBOA-dihexose (m/z 504.1353, 7), and DIMBOA-tetrahexose (m/z 858.2515, 8) (Figure 2). As high-resolution mass spectrometry was unable to provide the exact nature and linkage of those hexose units as well as to distinguish between isomeric aglucones such as HM₂BOA and HDMBOA, we decided to purify these multihexose BXDs from large enough quantities of plant material to enable NMR analyses.

While certain dihexose BXDs such as DIMBOA-dihexose (1) or HDMBOA/HM₂BOA-dihexose (6) were produced in

Table 1. ^1H NMR (600 MHz, D_2O) Data for Compounds 1–7

no.	1	2	3	4	5	6	7
2	5.99	5.92	5.84	5.84	6.07	6.02	5.95
5	7.38	7.47	7.01	7.02	7.19	7.36	7.58
6	6.83	6.82	6.76	6.78	6.91	6.88	7.19
7							7.24
8	6.85	6.81	6.86	6.86		6.92	7.18
11	3.84	3.85	3.83	3.84	3.89	3.87	
12					3.92		
N-OMe						3.98	
1'	4.86	4.85	4.86	4.86	4.88	4.88	4.87
2'	3.25	3.26	3.25	3.27	3.25	3.25	3.25
3'	3.51	3.50	3.51	3.50	3.52	3.50	3.54
4'	3.49	3.49	3.50	3.49	3.51	3.49	3.50
5'	3.63	3.64	3.61	3.61	3.60	3.63	3.68
6'	a	3.92	3.94	3.91	3.96	3.93	3.90
	b	4.13	4.13	4.13	4.12	4.07	4.21
1''	4.40	4.37	4.41	4.35	4.39	4.41	4.48
2''	3.28	3.29	3.29	3.28	3.23	3.29	3.31
3''	3.44	3.48	3.45	3.43	3.42	3.46	3.47
4''	3.39	3.43	3.39	3.48	3.36	3.40	3.42
5''	3.27	3.30	3.27	3.32	3.23	3.28	3.34
6''	a	3.71	3.82	3.72	3.82	3.69	3.73
	b	3.87	4.14	3.88	4.15	3.84	3.92
1'''		4.49		4.49			
2'''		3.30		3.31			
3'''		3.49		3.49			
4'''		3.38		3.39			
5'''		3.43		3.43			
6'''	a	3.71		3.72			
	b	3.90		3.90			

fairly high concentrations as shown by their prominent peaks in the chromatogram, others were present in much lower amounts (e.g., HMBOA-trihexose (4), DIBOA-dihexose (7), Figures 1 and 2). Compound 8 was present in such low quantities that it could not be isolated from the plant material. To elucidate the structure of all detected di- and trihexose BXDs, we purified them from maize plants grown under drought for 10 days and fully characterized them by a combination of spectroscopic techniques.

3.2. Structural Elucidation of Multihexose Benzoxazinoids. A methanol extract from maize roots was first fractionated by open column chromatography using a step gradient elution. The fraction (F9) containing dihexose and trihexose BXDs was eluted with a 1/1 mixture of ethyl acetate and methanol. This fraction was submitted to multiple steps of semipreparative fractionation to obtain compounds 1–7. All of the purification steps were controlled by LC-MS and the purity of each isolated compound was finally checked by NMR spectroscopy. Compounds 1 to 7 were identified as multihexose BXDs (Figure 3).

In the following section, the structural elucidation of compounds 1 to 7, based on NMR, HRMS and circular dichroism (CD) data, is briefly described. All ^1H NMR chemical shifts are displayed in Table 1 and a detailed report of all obtained data is provided in Supporting Information S12–S18.

The elemental composition of 1 was deduced to be $\text{C}_{21}\text{H}_{29}\text{NO}_{15}$ based on the HRESIMS and NMR data (Table 1, Supporting Information S12). The chemical shifts,

multiplicity, and coupling constant of the aromatic protons were characteristic of a monosubstituted aromatic ring fused to 1,4-benzoxazine [(7.38 (d, $J = 8.9$ Hz), 6.85 (d, $J = 2.6$ Hz), 6.83 (dd, $J = 8.9, 2.6$ Hz)]. The proton H-11 of the methoxy group [δ_{C} 56.2 (CH_3), δ_{H} 3.80] showed a long-range HMBC cross-peak with the carbon C-7. Thus, the aglycone was DIMBOA (2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one). The remaining signals observed in the ^1H and ^{13}C NMR spectra exhibited resonance of two hexose units characterized by two anomeric methines [δ_{C} 102.5 (CH), δ_{H} 4.86 (d, 7.9 Hz)] and [δ_{C} 103.5 (CH), δ_{H} 4.40 (d, 7.8 Hz)], two methylenes [δ_{C} 68.9 (CH_2), δ_{H} 3.92/4.13] and [δ_{C} 60.9 (CH_2), δ_{H} 3.87/3.71] and two closely related series of 4 methines bonded with an oxygen. Long range HMBC spectrum displayed correlations from H-1' (δ_{H} 4.86) to C-2 (δ_{C} 97.7) and from both H-6'a (δ_{H} 4.13) and H-6'b (δ_{H} 3.92) to C-1'' (δ_{C} 103.5), which established the linkage between the aglycone part and the sugar units (1'→2) and between the two sugar units (1''→6'). Both glucose units exhibited β conformations [δ_{C} 102.5 (CH), δ_{H} 4.86 (d, 7.9 Hz)] and [δ_{C} 103.5 (CH), δ_{H} 4.40 (d, 7.8 Hz)]. The coupling constants $^1J_{(\text{C}-\text{H})}$ of 163.7 and 161.8 Hz of both anomeric carbons C1' and C1'' were further evidence of a β conformation. Selective 1D-TOCSY, 1D-NOESY together with 2D-band selective HSQC were employed to unambiguously assign chemical shifts and to determine the stereochemistry of the sugar units. Correlations between H2 (δ_{H} 5.99) and H-1' (δ_{H} 4.86) confirmed the β linkage between the DIMBOA aglycone and the sugar unit. Additional NOESY enhancement observed from H-1' to H-3' and to H-5' and in a similar way from H-1'' to H-3'' and to H-5'' indicated that they have a *syn* orientation. The (2R) configuration could be proven by means of the CD spectrum (Supporting Information S12), showing a positive Cotton effect at 231 nm and a negative one at 315 nm in aqueous solution. This feature is in coincidence with that of the closely related (2R)-2- β -D-glucopyranosyloxy-7-methoxy-2H-1,4-benzoxazin-3(4H)-one.²² Compound 1 was thus identified as DIMBOA- β -gentiobiose (DIMBOA-2Glc) exhibiting a β -D-glucopyranosyl-(1''→6')- β -D-glucopyranose sugar unit (Figure 3).

The elemental composition of 2 was deduced to be $\text{C}_{27}\text{H}_{39}\text{NO}_{20}$ based on the HRESIMS and NMR data (Table 1, Supporting Information S13). The aglycone part was identified as DIMBOA. Signals observed in the ^1H and ^{13}C NMR spectra of 2 exhibited an extra set of chemical shifts in the glucoside region of both spectra matching with the additional $\text{C}_6\text{H}_{10}\text{O}_5$ moiety compared to 1. Long range HMBC spectrum displaying correlations from H-6''a (δ_{H} 4.13) to C-1''' (δ_{C} 103.7) established the linkage between the second and the third sugar units (1'''→6''). The third glucose unit exhibited an β conformation [δ_{C} 103.7 (CH), δ_{H} 4.49 (d, 8.0 Hz)]. The coupling constants $^1J_{(\text{C}-\text{H})}$ of anomeric carbons C 1', C 1'' and C 1''', 162.0, 161.3, and 159.3 Hz, respectively, are characteristic of a β conformation. The (2R) configuration could be proven by means of the CD spectrum, showing a positive Cotton effect at 231 nm and a negative one at 282 nm in aqueous solution. This feature is in accordance with that of the closely related DIMBOA- β -gentiobiose. Compound 2 was thus identified as DIMBOA- β -D-glucopyranosyl-(1''→6')- β -D-glucopyranosyl-(1'''→6'')-D- β -glucopyranose (DIMBOA-3Glc, Figure 3).




Figure 4. Drought induced di- and trihexose benzoxazinoids (BXDs) in maize roots and shoots. BXD peak heights (Mean \pm sem) in B73 maize leaves (A) and roots (B) ($n = 4\text{--}5$ per treatment and tissue). Drought was established 4 days after sowing and all plants were harvested 6 days later. Gray bars: ambient conditions, black bars: drought conditions. cps: count per second. Student t tests and Mann–Whitney Rank Sum tests were conducted. Dots and stars indicate trends and significant differences respectively (\cdot : $p < 0.10$, * $p < 0.05$).

The elemental composition of **3** was deduced to be $C_{21}H_{29}NO_{14}$ based on the HRESIMS and NMR data (Table 1, Supporting Information S14). Compound **1** possessed extra oxygen compared to **3**. NMR data of **1** and **3** were closely related. The alcohol function located on the nitrogen induced a gamma-steric effect for compound **1**. This gamma-steric effect of the alcohol function located on the nitrogen induced a variation of the ^{13}C chemical shift value of $\Delta\delta = -5.1$ ppm on C-3 of **1** compared to **3** (Supporting Information S12–S14). NOESY correlations and the CD spectrum confirmed the structure of the dihexose unit as well as the (2R) configuration (Supporting Information S14). Compound **3** was identified as HMBOA- β -gentiobiose (HMBOA-2Glc, Figure 3).

The elemental composition of **4** was deduced to be $C_{27}H_{39}NO_{19}$ based on the HRESIMS and NMR data (Table 1, Supporting Information S15). Signals observed in the 1H and ^{13}C NMR spectra exhibited signals related to both the HMBOA aglycone and the trihexose unit previously described for **2**. CD spectrum confirmed the (2R) configuration. Compound **4** was identified as HMBOA- β -D-glucopyranosyl-(1'' \rightarrow 6')- β -D-glucopyranosyl-(1''' \rightarrow 6'')-D- β -glucopyranose (HMBOA-3Glc, Figure 3).

The elemental composition of **5** was deduced to be $C_{22}H_{31}NO_{15}$ based on the HRESIMS and NMR data (Table 1, Supporting Information S16). Signals observed in the 1H and ^{13}C NMR spectra of **5** exhibited resonances for two methoxy groups [δ_C 56.2 (CH₃), δ_H 3.89] and [δ_C 61.6

(CH₃), δ_H 3.92]. Protons H-11 and H-12 located on each of the methoxy groups showed long-range HMBC cross-peaks, respectively, with the carbons C-8 and C-7. The aglycone unit was identified as HM₂BOA. Chemical shifts of the two hexose units were closely related to both compounds **1** and **3** (Table 1). NOESY correlations and CD spectrum confirmed the structure of the dihexose unit as well as the (2R) configuration. Compound **5** was identified as HM₂BOA- β -gentiobiose (HM₂BOA-2Glc, Figure 3).

Compound **6** exhibited a prominent adduct $[M+HCOO]^-$ at m/z 594.1670 and a minor $[M-H]^-$ at m/z 548.1619 leading to the same elemental composition $C_{22}H_{31}NO_{15}$ as **5** based on the HRESIMS and NMR data (Table 1, Supporting Information S17). The proton NMR spectrum exhibited two singlet proton signals characteristic of methoxy groups [δ_C 56.3 (CH₃), δ_H 3.87] and [δ_C 63.6 (CH₃), δ_H 3.98]. Long-range HMBC showed cross-peaks between H-11 and C-7. Spectral data of the second methoxy group [δ_C 63.6 (CH₃), δ_H 3.98] were characteristic of an N-OMe group. The aglycone unit was identified as HDMBOA. Chemical shifts of the two hexose units were closely related to compounds **1**, **3**, and **5** (Table 1). NOESY correlations and the CD spectrum confirmed the structure of the dihexose unit as well as the (2R) configuration. Compound **6** was identified as HDMBOA- β -gentiobiose (HDMBOA-2Glc, Figure 3).

The elemental composition of **7** was deduced to be $C_{20}H_{27}NO_{14}$ based on the HRESIMS and NMR data (Table 1, Supporting Information S18). Signals observed in the 1H




Figure 5. Drought-mediated induction of double and triple hexoses was transient. DIMBOA-2Glc concentrations in maize shoots (A) and roots (B) under control and drought conditions. DIMBOA-3Glc concentrations in maize shoots (C) and roots (D) under control and drought conditions. HMBOA-2Glc concentrations in maize shoots (E) and roots (F) under control and drought conditions ($n = 8$ per treatment and time point). Control (green lines): 23% soil moisture (v/v); Drought (yellow lines): 16.6% soil moisture (v/v). tmt: treatment (control or drought). FW: fresh weight. Mean \pm SEM are shown. Two-way ANOVAs on ranks were conducted, followed by posthoc Holm–Sidak tests when relevant. Stars indicate significant differences within time points (\cdot : $p < 0.10$; $*$: $p < 0.05$; $**$: $p < 0.01$; $***$: $p < 0.001$).

spectrum were characteristic of an aromatic ring without any substituents fused to 1,4-benzoxazine [7.58 (dd, $J = 7.5, 2.5$ Hz), 7.24 (ddd, $J = 7.9, 6.5, 2.5$ Hz), 7.19 (dd, $J = 7.5, 6.5$ Hz), 7.18 (d, $J = 7.9$ Hz)]. Thus, the aglycone was DIBOA. Chemical shifts of the two hexose units were closely related to compounds 1, 3, 5 and 6 (Table 1). Compound 7 was identified as DIBOA- β -gentiobiose (DIBOA-2Glc, Figure 3).

3.3. Drought Profoundly Modifies Benzoxazinoid Profiles in Leaf and Root Tissues. Drought decreased DIMBOA-Glc and DIBOA-Glc levels but increased the levels of DIMBOA-2Glc, HMBOA-2Glc, DIMBOA-3Glc, and HDMBOA-2Glc in maize leaves. Drought further induced higher levels of DIMBOA-2Glc, HMBOA-2Glc, and HDMBOA-2Glc in roots (Figure 4). The production of di- and trihexose BXDs was previously observed in maize¹³ and was induced under stressful conditions.^{15,16}

3.4. Induction of Dihexose- and Trihexose Benzoxazinoids Is Transient. Under ambient conditions, the oligosaccharides DIMBOA-2Glc, and DIMBOA-3Glc could only be detected in four-day-old seedlings, as their levels dropped under limit of detection afterward (Figure 5A–D).

However, low levels of HMBOA-2Glc could be detected in roots over 24 days of growth (Figure 5E,F).

Under drought conditions, maize seedlings transiently produced higher concentrations of DIMBOA-2Glc, DIMBOA-3Glc, and HMBOA-2Glc in both shoot and root tissues (Figure 5). The production of the di- and trihexose BXDs was the highest in roots of 4-day-old seedlings and in shoots of 7-day-old seedlings (Figure 5). Whether these BXDs are produced in roots and subsequently transported to shoots remains to be investigated.

3.5. Drought-Induced Multihexose Benzoxazinoid Production Is Common in Maize. The drought-mediated induction of multihexose BXDs was found in leaves and roots of the four maize varieties, B73, CML277, Hp301, and Oh7B (Figure 6). Interestingly, the increase in multihexose BXDs was accompanied by a decrease in monoglucosylated BXDs, suggesting their use as precursors for multihexose BXDs (Figure 6). Some line-specific patterns were further observed, suggesting that genetic variability could be used to identify the biosynthesis pathways and glucosyl transferases involved in the production of multihexose BXDs.

3.6. Toward the Function(s) of Multihexose Benzoxazinoids. Overall, this study reports a method to produce maize extracts enriched in multihexose BXDs and provides the first characterization of the exact hexose nature and configuration. Glycosylation of a small organic compound can alter its solubility, stability, bioavailability, and bioactivity and modulate its storage, transport, and/or interactions with other organisms. Plant multihexose secondary metabolites have been previously reported in wheat, rye, tomato, daisy flowers, Madagascar periwinkle flowers, and Japanese morning glory.^{23–26} They include phenolic compounds such as quercetin 3-*O*-gentiobiosides,²⁶ and flavonoids, such as quercetin 3-*O*-gentiobiosides, gentiotriosides, and gentiotetrosides,²³ anthocyanidin 3-*O*-rutinosides,^{24,25} or 3-*O*-2''-*O*- β -glucuronosyl-6''-*O*-malonylglucoside.²⁴ Evidence suggests that the conversion of glycoside plant specialized metabolites into multihexoses can modulate plant volatile emissions²⁶ and flower coloration.²⁴

The role of multihexose BXDs still remains to be elucidated. Several, nonexclusive, hypotheses can be formulated. First, multihexose BXDs may act as osmoprotectants through maintaining cell turgor pressure and osmotic balance. Second, these BXDs may form, through the sugar moieties, hydrogen bonds with water molecules and reduce the water loss associated with transpiration. Third, the multihexose compounds may act as stronger antioxidants than their precursors and prevent cellular damage associated with reactive oxygen species (ROS). Fourth, multihexoses may be used to reallocate and store sugars as an energy source to tolerate drought and decreased photosynthesis. Fifth, because BXDs modulate the plant interactions with pathogens and herbivores, multihexose BXDs may be involved in protecting the plant from biotic stress under challenging abiotic conditions. Alternatively, the sugar-rich compounds may be exuded in the rhizosphere and promote beneficial microorganisms. While our understanding of the biosynthesis and ecological functions of multihexose plant specialized metabolites is still in its infancy, they represent a fascinating aspect of phytochemical diversity, calling for future research.




Figure 6. Multihexose benzoxazinoids were induced by drought in several maize varieties. Heatmap of benzoxazinoid (BXD) levels in drought-stressed plants (leaves and roots) compared to control plants in B73, CML277, Hp301, and Oh7B maize varieties ($n = 2-5$ per treatment and variety). The log fold (\log_{10}) changes between drought-stressed and control plants are shown. Green indicates a decrease in BXD levels upon drought. Red indicates an increase in BXD levels upon drought. Student t tests and Mann–Whitney Sum Rank tests were conducted. Dots and stars indicate trends and significant differences respectively ($\cdot: p < 0.10$, $*: p < 0.05$, $** : p < 0.01$, $***: p < 0.001$).

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.jafc.3c09141>.

(SI1) Preparative chromatography; (SI2) structure elucidation of DIMBOA-2Glc (1); (SI3) structure elucidation of DIMBOA-3Glc (2); (SI4) structure elucidation of HMBOA-2Glc (3); (SI5) structure elucidation of HMBOA-3Glc (4); (SI6) structure elucidation of HM₂BOA-2Glc (5); (SI7) structure elucidation of HDMBOA-2Glc (6); (SI8) structure elucidation of DIBOA-2Glc (7) (PDF)

AUTHOR INFORMATION

Corresponding Authors

Sylvain Sutour – Neuchâtel Platform of Analytical Chemistry, University of Neuchâtel, Neuchâtel 2000, Switzerland; orcid.org/0000-0001-9676-7123; Email: sylvain.sutour@unine.ch

Christelle Aurélie Maud Robert – Institute of Plant Sciences, University of Bern, Bern 3013, Switzerland; Oeschger Centre for Climate Change Research (OCCR), University of Bern, Bern 3012, Switzerland; orcid.org/0000-0003-3415-2371; Email: christelle.robert@unibe.ch

Authors

Van Cong Doan – Institute of Plant Sciences, University of Bern, Bern 3013, Switzerland; Oeschger Centre for Climate Change Research (OCCR), University of Bern, Bern 3012, Switzerland; Plant Physiology Unit, The Department of Life Sciences and Systems Biology of the University of Turin, Torino 10123, Italy

Pierre Mateo – Institute of Plant Sciences, University of Bern, Bern 3013, Switzerland; orcid.org/0000-0003-4338-3024

Tobias Züst – Department of Systematic and Evolutionary Botany, University of Zürich, Zürich 8008, Switzerland; orcid.org/0000-0001-7142-8731

Ella Raymonde Hartmann – Institute of Plant Sciences, University of Bern, Bern 3013, Switzerland

Gaëtan Glauser – Neuchâtel Platform of Analytical Chemistry, University of Neuchâtel, Neuchâtel 2000, Switzerland; orcid.org/0000-0002-0983-8614

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.jafc.3c09141>

Author Contributions

S.S. collected and analyzed data, and wrote the first draft of the manuscript; C.v.D. collected and analyzed data; T.Z. collected and analyzed data; P.M. collected data and wrote the first draft of the manuscript; E.R.H. collected data; G.G. collected and analyzed data, supervised the project, and wrote the first draft of the manuscript; C.A.M.R. conceived the study, secured funding, collected and analyzed data, supervised the project, and wrote the first draft of the manuscript. All authors have contributed to the final version of the manuscript.

Funding

The work of P.M., E.R.H., and C.A.M.R. was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program [ERC-2019-STG949595] and the Swiss National Science Foundation [310030–189071]. The work of C.v.D. was supported by the University of Bern (UniBe 2021) and the National Recovery and Resilience Plan (NRRP), Italy (DOAV_PNRR_YOUR_SOE_22_01). The work of T.Z. was supported by the Swiss National Science Foundation [PCEFP3_194590]. The work of S.S. and G.G. was supported by the University of Neuchâtel.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We are grateful to Thi My Duyen Nguyen for her technical help during drought experiments. We thank Thomas Degen and Ted Turlings for providing drawings of maize plants. This work was supported by the European Research Council

(ERC), the Swiss National Science Foundation, the Universities of Bern and Neuchâtel, and the Italian National Recovery and Resilience Plan.

ABBREVIATIONS USED

ACN: acetonitrile
 BPI: base peak intensity chromatogram
 BXD: benzoxazinoid
 CHCl₃: chloroform
 CH: Switzerland
 CLIP-HSQC: coupled heteronuclear single quantum coherence clean inphase multiplets
 COSY: correlation spectroscopy (NMR spectroscopy)
 DE: Germany
 DIBOA: 2,4-dihydroxy-1,4-benzoxazin-3-one
 DIMBOA: 2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one
 EtOH: ethanol
 Glc: glucose
 Hex: hexose
 HBOA: 2-hydroxy-1,4-benzoxazin-3-one
 HDMBOA: 2-hydroxy-4,7-dimethoxy-1,4-benzoxazin-3-one
 HMBC: heteronuclear multiple bond correlation (NMR spectroscopy)
 HMBOA: 2-hydroxy-4-methoxy-1,4-benzoxazin-3-one
 HM₂BOA: 2-hydroxy-4,7-dimethoxy-1,4-benzoxazin-3-one
 HPLC: high-performance liquid chromatography
 HTP: hydrothermal processing
 HRESIMS: high-resolution electrospray mass spectrometry
 HSQC: heteronuclear single quantum coherence (NMR spectroscopy)
 IPCC: Intergovernmental Panel on Climate Change
 LC: liquid chromatography
 MeOH: methanol
 MS: mass spectrometry
m/z: mass-to-charge ratio
 NMR: nuclear magnetic resonance
 NOESY: nuclear Overhauser effect spectroscopy (NMR spectroscopy)
 ppm: parts per million
 QTOF: quadrupole time-of-flight (mass spectrometry)
 RCP: representative concentration pathways (climate modeling)
 TOCSY: total correlation spectroscopy (NMR spectroscopy)
 UPLC: ultrahigh-performance liquid chromatography
 USA: United States of America

REFERENCES

- Robert, C. A. M.; Mateo, P. The Chemical Ecology of Benzoxazinoids. *Chimia (Aarau)* **2022**, *76* (11), 928.
- Gross, J. J.; Mateo, P.; Ramhold, D.; Kramer, E.; Erb, M.; Robert, C. A. M. Turnover of Benzoxazinoids during the Aerobic Deterioration of Maize Silage (*Zea Mays*). *J. Agric. Food Chem.* **2023**, *71* (5), 2370–2376.
- Pedersen, H. A.; Laursen, B.; Mortensen, A.; Fomsgaard, I. S. Bread from Common Cereal Cultivars Contains an Important Array of Neglected Bioactive Benzoxazinoids. *Food Chem.* **2011**, *127* (4), 1814–1820.
- Steffensen, S. K.; Adhikari, K. B.; Laursen, B. B.; Jensen, C.; Gregersen, P. L.; Bhattarai, B.; Marais, L. M.; Schnorr, H.; Jensen, B. M.; Poulsen, L. K.; Nielsen, C. H.; Borre, M.; Borre, M.; Høyer, S.; Fomsgaard, I. S. Bioactive Small Molecules in Commercially Available Cereal Food: Benzoxazinoids. *J. Food Compos. Anal.* **2017**, *64*, 213–222.

- Gross, J. J.; Mateo, P.; Schlaeppli, K.; Wyss, U.; Kramer, E.; Ramhold, D.; Erb, M.; Robert, C. A. M. Short Communication: Metabolization of Benzoxazinoids during Silage Fermentation of Maize and Their Effects on Silage Quality. *Anim. Feed Sci. Technol.* **2023**, *304*, No. 115748.

- Adhikari, K. B.; Tanwir, F.; Gregersen, P. L.; Steffensen, S. K.; Jensen, B. M.; Poulsen, L. K.; Nielsen, C. H.; Høyer, S.; Borre, M.; Fomsgaard, I. S. Benzoxazinoids: Cereal Phytochemicals with Putative Therapeutic and Health-Protecting Properties. *Mol. Nutr. Food Res.* **2015**, *59* (7), 1324–1338.

- Matos, P.; Figueirinha, A.; Paranhos, A.; Nunes, F.; Cruz, P.; Geraldes, C. F. G. C.; Cruz, M. T.; Batista, M. T. Bioactivity of *Acanthus Mollis* – Contribution of Benzoxazinoids and Phenylpropanoids. *J. Ethnopharmacol.* **2018**, *227*, 198–205.

- Bhattarai, B.; Steffensen, S. K.; Staerk, D.; Laursen, B. B.; Fomsgaard, I. S. Data-Dependent Acquisition-Mass Spectrometry Guided Isolation of New Benzoxazinoids from the Roots of *Acanthus Mollis* L. *Int. J. Mass Spectrom.* **2022**, *474*, No. 116815.

- Nordin, E.; Steffensen, S. K.; Laursen, B. B.; Andersson, S.-O.; Johansson, J.-E.; Åman, P.; Hallmans, G.; Borre, M.; Stærk, D.; Hanhineva, K.; Fomsgaard, I. S.; Landberg, R. An Inverse Association between Plasma Benzoxazinoid Metabolites and PSA after Rye Intake in Men with Prostate Cancer Revealed with a New Method. *Sci. Rep.* **2022**, *12* (1), 5260.

- Pihlava, J.-M.; Kurtelius, T. Determination of Benzoxazinoids in Wheat and Rye Beers by HPLC-DAD and UPLC-QTOF MS. *Food Chem.* **2016**, *204*, 400–408.

- Bhattarai, B.; Steffensen, S. K.; Gregersen, P. L.; Kristensen, H. L.; Fomsgaard, I. S. Stepwise Mass Spectrometry-Based Approach for Confirming the Presence of Benzoxazinoids in Herbs and Vegetables. *Phytochemical Analysis* **2021**, *32* (3), 283–297.

- Dihm, K.; Vendelbo Lind, M.; Sundén, H.; Ross, A.; Savolainen, O. Quantification of Benzoxazinoids and Their Metabolites in Nordic Breads. *Food Chem.* **2017**, *235*, 7–13.

- Fomsgaard, I. S.; Mortensen, A. G.; Holm, P. B.; Gregersen, P. L. Use of Benzoxazinoids-Containing Cereal Grain Products for Health-Improving Purposes. Patent application. EP 2 265 133 A1 2009.

- Hanhineva, K.; Rogachev, I.; Aura, A.-M.; Aharoni, A.; Poutanen, K.; Mykkänen, H. Qualitative Characterization of Benzoxazinoid Derivatives in Whole Grain Rye and Wheat by LC-MS Metabolite Profiling. *J. Agric. Food Chem.* **2011**, *59* (3), 921–927.

- Pedersen, H. A.; Heinrichson, K.; Fomsgaard, I. S. Alterations of the Benzoxazinoid Profiles of Uninjured Maize Seedlings during Freezing, Storage, and Lyophilization. *J. Agric. Food Chem.* **2017**, *65* (20), 4103–4110.

- Batyrshina, Z. S.; Shavit, R.; Yaakov, B.; Bocobza, S.; Tzin, V. The Transcription Factor TaMYB31 Regulates the Benzoxazinoid Biosynthetic Pathway in Wheat. *J. Exp. Bot.* **2022**, *73* (16), 5634–5649.

- van Doan, C.; Pfander, M.; Guyer, A. S.; Zhang, X.; Maurer, C.; Robert, C. A. M. Natural Enemies of Herbivores Maintain Their Biological Control Potential under Short-Term Exposure to Future CO₂, Temperature, and Precipitation Patterns. *Ecol. Evol.* **2021**, *11* (9), 4182–4192.

- Guyer, A.; van Doan, C.; Maurer, C.; Machado, R. A. R.; Mateo, P.; Steinauer, K.; Kesner, L.; Hoch, G.; Kahmen, A.; Erb, M.; Robert, C. A. M. Climate Change Modulates Multitrophic Interactions Between Maize, A Root Herbivore, and Its Enemies. *J. Chem. Ecol.* **2021**, *47* (10), 889–906.

- IPCC. *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*; IPCC, 2014.

- Robert, C. A. M.; Zhang, X.; Machado, R. A. R.; Schirmer, S.; Lori, M.; Mateo, P.; Erb, M.; Gershenzon, J. Sequestration and Activation of Plant Toxins Protect the Western Corn Rootworm from Enemies at Multiple Trophic Levels. *Elife* **2017**, *6*, No. e29307.

(21) Tanwir, F.; Fredholm, M.; Gregersen, P. L.; Fomsgaard, I. S. Comparison of the Levels of Bioactive Benzoxazinoids in Different Wheat and Rye Fractions and the Transformation of These Compounds in Homemade Foods. *Food Chem.* **2013**, *141* (1), 444–450.

(22) Nagao, T.; Otsuka, H.; Kohda, H.; Sato, T.; Yamasaki, K. Benzoxazinones from Coix Lachryma-Jobi Var. *Ma-Yuen*. *Phytochemistry* **1985**, *24* (12), 2959–2962.

(23) Masada, S.; Terasaka, K.; Oguchi, Y.; Okazaki, S.; Mizushima, T.; Mizukami, H. Functional and Structural Characterization of a Flavonoid Glucoside 1,6-Glucosyltransferase from *Catharanthus Roseus*. *Plant Cell Physiol* **2009**, *50* (8), 1401–1415.

(24) Sawada, S.; Suzuki, H.; Ichimaida, F.; Yamaguchi, M.; Iwashita, T.; Fukui, Y.; Hemmi, H.; Nishino, T.; Nakayama, T. UDP-Glucuronic Acid:Anthocyanin Glucuronosyltransferase from Red Daisy (*Bellis Perennis*) Flowers. Enzymology and Phylogenetics of a Novel Glucuronosyltransferase Involved in Flower Pigment Biosynthesis. *J. Biol. Chem.* **2005**, *280* (2), 899–906.

(25) Morita, Y.; Hoshino, A.; Kikuchi, Y.; Okuhara, H.; Ono, E.; Tanaka, Y.; Fukui, Y.; Saito, N.; Nitasaka, E.; Noguchi, H.; Iida, S. Japanese Morning Glory Dusky Mutants Displaying Reddish-Brown or Purplish-Gray Flowers Are Deficient in a Novel Glycosylation Enzyme for Anthocyanin Biosynthesis, UDP-Glucose:Anthocyanidin 3-O-Glucoside-2"-O-Glucosyltransferase, Due to 4-Bp Insertions in the Gene. *Plant Journal* **2005**, *42* (3), 353–363.

(26) Tikunov, Y. M.; Molthoff, J.; de Vos, R. C. H.; Beekwilder, J.; van Houwelingen, A.; van der Hooft, J. J. J.; Nijenhuis-de Vries, M.; Labrie, C. W.; Verkerke, W.; van de Geest, H.; Viquez Zamora, M.; Presa, S.; Rambla, J. L.; Granell, A.; Hall, R. D.; Bovy, A. G. NON-SMOKY GLYCOSYLTRANSFERASE1 Prevents the Release of Smoky Aroma from Tomato Fruit. *Plant Cell* **2013**, *25* (8), 3067–3078.