

Azithromycin resistance in *Escherichia coli* and *Salmonella* from food-producing animals and meat in Europe

Mirena Ivanova¹, Armen Ovsepiyan ^{1,2}, Pimlapas Leekitcharoenphon ³, Anne Mette Seyfarth¹, Hanne Mordhorst³, Saria Otani³, Sandra Koeberl-Jelovcan⁴, Mihail Milanov⁵, Gordan Kompes⁶, Maria Liapi⁷, Tomáš Černý⁸, Camilla Thougard Vester⁹, Agnès Perrin-Guyomard¹⁰, Jens A. Hammerl ¹¹, Mirjam Grobbel¹¹, Eleni Valkanou¹², Szilárd Jánosi¹³, Rosemarie Slowey¹⁴, Patricia Alba ¹⁵, Virginia Carfora ¹⁵, Jelena Avsejenko¹⁶, Asta Pereckiene¹⁷, Dominique Claude¹⁸, Renato Zerafa¹⁹, Kees T. Veldman²⁰, Cécile Boland²¹, Cristina Garcia-Graells²¹, Pierre Wattiau²¹, Patrick Butaye ^{22,23}, Magdalena Zajac ²⁴, Ana Amaro²⁵, Lurdes Clemente²⁵, Angela M. Vaduva²⁶, Luminita-Maria Romascu²⁷, Nicoleta-Manuela Milita²⁷, Andrea Mojžišová²⁸, Irena Zdovc²⁹, Maria Jesús Zamora Escribano³⁰, Cristina De Frutos Escobar³⁰, Gudrun Overesch³¹, Christopher Teale³², Guy H. Loneragan³³, Beatriz Guerra³⁴, Pierre Alexandre Beloeil³⁴, Amanda M. V. Brown³⁵, Rene S. Hendriksen¹, Valeria Bortolaia^{1,36}† and Jette Sejer Kjelsgaard^{1*†}

¹European Union Reference Laboratory for Antimicrobial Resistance (EURL-AR), Research Group for Global Capacity Building, Technical University of Denmark, Kongens Lyngby, Denmark; ²DIANA-Lab, Dept. of Computer Science and Biomedical Informatics, University of Thessaly, Lamia, Greece; ³Research Group for Genomic Epidemiology, Technical University of Denmark, Kongens Lyngby, Denmark; ⁴Austrian Agency for Health and Food Safety, Graz, Austria; ⁵National Diagnostic and Research Veterinary Institute, Sofia, Bulgaria; ⁶Croatian Veterinary Institute, Zagreb, Croatia; ⁷Bacteriology Serology Laboratory, Veterinary Services, Cyprus; ⁸State Veterinary Institute, Prague, Czech Republic; ⁹Danish Veterinary and Food Administration, Ringsted, Denmark; ¹⁰French Agency for Food, Environmental and Occupational Health & Safety, Maisons-Alfort, France; ¹¹German Federal Institute for Risk Assessment, Berlin, Germany; ¹²Veterinary Laboratory of Chalkis, Chalkis, Greece; ¹³National Food Chain Safety Office, Veterinary Diagnostic Directorate, Budapest, Hungary; ¹⁴Central Veterinary Research Laboratory, Kildare, Ireland; ¹⁵Istituto Zooprofilattico Sperimentale del Lazio e della Toscana 'M. Aleandri', Rome, Italy; ¹⁶Institute of Food Safety, Animal Health and Environment BIOR, Riga, Latvia; ¹⁷National Food and Veterinary Risk Assessment Institute, Vilnius, Lithuania; ¹⁸Laboratoire de Médecine Vétérinaire de l'État, Dudelange, Luxembourg; ¹⁹Public Health Laboratory, Valletta, Malta; ²⁰Wageningen Bioveterinary Research, Part of Wageningen University & Research, Lelystad, Netherlands; ²¹Sciensano, Brussels, Belgium; ²²Department of Pathobiology, Ghent University, Merelbeke, Belgium; ²³Jockey Club College of Veterinary Medicine and Life Sciences, Kowloon, Hong Kong; ²⁴National Veterinary Research Institute, Pulawy, Poland; ²⁵Instituto Nacional de Investigação Agrária e Veterinária, Oeiras, Portugal; ²⁶Institute for Hygiene and Veterinary Public Health, Bucharest, Romania; ²⁷Institute for Diagnosis and Animal Health, Bucharest, Romania; ²⁸State Veterinary and Food Institute, Dolny Kubin, Slovakia; ²⁹Institute for Microbiology and Parasitology, Ljubljana, Slovenia; ³⁰Spanish Agency for Food Safety and Nutrition, Madrid, Spain; ³¹Vetsuisse Faculty, Institute of Veterinary Bacteriology, University of Bern, Bern, Switzerland; ³²Animal & Plant Health Agency, Weybridge, UK; ³³School of Veterinary Medicine, Texas Tech University, Amarillo, TX, USA; ³⁴European Food Safety Authority, Parma, Italy; ³⁵Department of Biological Sciences, Texas Tech University, Lubbock, TX, USA; ³⁶Statens Serum Institut, Copenhagen, Denmark

*Corresponding author. E-mail: jetk@food.dtu.dk
†Equally contributing authors.

Received 12 July 2023; accepted 30 April 2024

Objectives: To characterize the genetic basis of azithromycin resistance in *Escherichia coli* and *Salmonella* collected within the EU harmonized antimicrobial resistance (AMR) surveillance programme in 2014–18 and the Danish AMR surveillance programme in 2016–19.

Methods: WGS data of 1007 *E. coli* [165 azithromycin resistant (MIC > 16 mg/L)] and 269 *Salmonella* [29 azithromycin resistant (MIC > 16 mg/L)] were screened for acquired macrolide resistance genes and mutations in *rplDV*, 23S rRNA and *acrB* genes using ResFinder v4.0, AMRFinder Plus and custom scripts. Genotype–phenotype concordance was determined for all isolates. Transferability of *mef*(C)-*mph*(G)-carrying plasmids was assessed by conjugation experiments.

Results: *mph(A)*, *mph(B)*, *mef(B)*, *erm(B)* and *mef(C)-mph(G)* were detected in *E. coli* and *Salmonella*, whereas *erm(C)*, *erm(42)*, *ere(A)* and *mph(E)-msr(E)* were detected in *E. coli* only. The presence of macrolide resistance genes, alone or in combination, was concordant with the azithromycin-resistant phenotype in 69% of isolates. Distinct *mph(A)* operon structures were observed in azithromycin-susceptible ($n=50$) and -resistant ($n=136$) isolates. *mef(C)-mph(G)* were detected in porcine and bovine *E. coli* and in porcine *Salmonella enterica* serovar Derby and *Salmonella enterica* 1,4, [5],12:-, flanked downstream by ISCR2 or TnAs1 and associated with IncIy and IncFII plasmids.

Conclusions: Diverse azithromycin resistance genes were detected in *E. coli* and *Salmonella* from food-producing animals and meat in Europe. Azithromycin resistance genes *mef(C)-mph(G)* and *erm(42)* appear to be emerging primarily in porcine *E. coli* isolates. The identification of distinct *mph(A)* operon structures in susceptible and resistant isolates increases the predictive power of WGS-based methods for *in silico* detection of azithromycin resistance in Enterobacterales.

Introduction

The macrolide azithromycin is a critically important clinical antimicrobial,¹ increasingly used as an alternative when typical first-line antimicrobials (e.g. quinolones) are no longer effective in the treatment of severe cases of bacterial gastrointestinal infections.^{2–5} Considering that azithromycin is one of the few available options for treatment of MDR bacteria and that the majority of the azithromycin resistance genes are acquired, the spread of azithromycin resistance could seriously decrease the options to fight life-threatening infections. Among the azithromycin resistance genes,⁶ *mph(A)* and *erm* genes, encoding macrolide 2'-phosphotransferase and rRNA methylases, respectively, are the two main mechanisms involved in high-level azithromycin resistance.^{7,8} Another recently identified resistance mechanism in *Escherichia coli* conferred by tandemly arranged plasmid-borne genes *mef(C)-mph(G)*, encoding an efflux pump and a phosphorylase, respectively, has been described to mediate high-level azithromycin resistance.⁹ Additionally, substitutions in the 50S ribosomal subunit proteins L4 (*rplD*) and L22 (*rplV*), in 23S rRNA (*rrlH*) and in the efflux pump AcrB (R717Q/L) also can lead to increased macrolide resistance.^{2,6} However, genes encoding efflux pumps, e.g. *msr(A)*, *msr(D)*, *mef(A)*, *mef(B)*, *ere* genes encoding macrolide esterases, and *mph(B)* encoding a macrolide phosphorylase appear to have no role or only marginal roles in azithromycin resistance in Enterobacterales.^{6–8}

Due to the transmission of antimicrobial-resistant bacteria between animals and humans, the EU has implemented harmonized monitoring and reporting of antimicrobial resistance (AMR) in zoonotic and commensal bacteria from food-producing animals and food since 2014.^{10,11} In this monitoring programme, azithromycin susceptibility of *E. coli* and *Salmonella enterica* is tested phenotypically by broth microdilution.

Phenotypic azithromycin susceptibility testing is technically challenging and presents reproducibility issues in classifying isolates consistently as susceptible or resistant, which could be overcome by using WGS methods. However, these methods perform with high accuracy only for well-studied AMR determinants. Therefore, elucidating AMR gene patterns to increase the accuracy and effectiveness of WGS approaches is of great importance.

Although the prevalence of azithromycin-resistant *E. coli* and *Salmonella* in the EU has generally been reported to be low depending on the country and isolation source,¹² a proper

assessment of the risk to humans posed by azithromycin-resistant bacteria from food animals and food requires knowledge of resistance determinants. Therefore, the objectives of this study were to elucidate the genetic basis of azithromycin resistance in *E. coli* and *Salmonella* from food-producing animals and meat from 27 European countries and explore the azithromycin genotype-phenotype correspondence. Furthermore, this study aimed to resolve the molecular basis of genotype-phenotype discordance previously observed for *mph(A)*, which is the most common azithromycin resistance gene in *E. coli*, and to characterize the genetic context and transferability of *mef(C)-mph(G)* genes, which are emerging in Enterobacterales.

Methods

Bacterial isolates

A total of 1276 isolates were examined in this study. These isolates comprised 1007 *E. coli* and 269 *Salmonella enterica* subsp. *enterica* serovars (Table S1, available as [Supplementary data](#) at JAC Online) collected within the EU harmonized monitoring of AMR in zoonotic and indicator bacteria from food-producing animals and food in 2014–18,^{10,13} and the Danish Programme for surveillance of antimicrobial consumption and resistance in bacteria from food animals and food (www.danmap.org) in 2017–19. The EU harmonized monitoring system involved biannual surveillance of different animal types. Isolates from 2014, 2016 and 2018 originated from poultry caeca and meat, whereas isolates from 2015 and 2017 and from the Danish AMR surveillance programme were recovered from porcine and bovine caeca and meat. For the Danish collection, only isolates with azithromycin MIC > 16 mg/L were included. Antimicrobial susceptibility testing was performed by broth microdilution using Sensititre MIC susceptibility plates (EUVSEC1 and EUVSEC2, Thermo Fisher Scientific) in duplicate, by the originating laboratory and at the EU Reference Laboratory – Antimicrobial Resistance (EURL-AR). In cases of more than one 2-fold dilution difference, the tests were repeated a third time. The azithromycin MIC values were interpreted in accordance with The European Food Safety Authority (EFSA)-defined surveillance breakpoint. Isolates with MIC ≤ 16 mg/L were classified as WT and isolates with MIC > 16 mg/L as non-WT.¹³

Identification of genes and chromosomal mutations associated with macrolide resistance

WGS of the isolates was performed using Illumina paired-end sequencing on MiSeq, HiSeq or NextSeq platforms (Illumina, Inc., San Diego, CA, USA). Sequencing reads were trimmed using Trimmomatic v0.38¹⁴ and

assembled by metaSPAdes v3.13.0.¹⁵ Assemblies were evaluated by Quast¹⁶ and genomes with ≤ 500 contigs were included in further analysis (Table S1). The ResFinder database v4.0¹⁷ incorporated into ABRicate v0.9.8 (<https://github.com/tseemann/abricate>) was used to detect acquired macrolide resistance genes with minimum identity and coverage thresholds of 90% and 60%, respectively. Chromosomal mutations in *acrB* and 23S rRNA genes were screened for using AMRFinder (<https://github.com/ncbi/amr/wiki>). The sequences of chromosomal *rplDV* genes were screened for mutations mediating azithromycin resistance^{7,18} using custom scripts. Sequence alignments of *rplDV* genes were performed using MAFFT in Geneious Prime v2020.0.4 (<https://www.geneious.com>) and compared with *rplDV* of *E. coli* ATCC 25922 (NC_000913.3) and *Salmonella enterica* serovar Typhimurium LT2 (NC_003197) as reference genes. MLST was performed using mlst v2.19.0 according to the Achtman schemes (<https://github.com/tseemann/mlst>). *Salmonella* serovar determination was carried out by SeqSero2 v1.1.1.¹⁹ Sequences from isolates positive for *mph(A)* and *mef(C)-mph(G)* were screened for plasmid replicon genes using PlasmidFinder (database version February 2020)²⁰ with minimum identity and coverage thresholds as above. IS and Tn were identified using ISfinder (<https://www-is.biotoul.fr>)²¹ and TnCentral (<https://tncentral.ncc.unesp.br/index.html>). Visualization of the genetic contexts of the macrolide resistance genes was performed by pyGenomeViz v0.4.3 (<https://github.com/moshi4/pyGenomeViz>). Graphs were created using 'ggplot2' and 'networkD3' packages in R v4.1.0.^{22,23} Raw sequence data were submitted to the European Nucleotide Archive (ENA) at EMBL-EBI under accession numbers PRJEB18618, PRJEB21546, PRJEB33169, PRJEB43436, PRJEB43584, PRJEB63535 and PRJEB63683 (Table S1).

Conjugation experiments and characterization of *mef(C)-mph(G)*-harbouring contigs

Isolates harbouring *mef(C)-mph(G)* (Table 1) were used as donors in filter-mating experiments with a rifampicin-resistant, lactose-negative *E. coli* J62-2 strain as recipient.²⁴ Transconjugants (TCs) were selected on MacConkey agar (Sigma, Denmark) supplemented with 16 mg/L azithromycin (Sigma, Denmark) and 50 mg/L rifampicin (Sigma, Denmark). TCs were initially verified by MIC determination as described for the original isolates and by colony PCR targeting *mef(C)-mph(G)* using primers F3220-5'-ATTGGCGGTGTCATCCTGAG-3' and R3221-5'-CGTGACTTGTGCA GTTGAC-3'. Plasmid DNA from TCs with azithromycin MIC > 16 mg/L and *mef(C)-mph(G)* positive by PCR was extracted using the QIAGEN Plasmid Midi kit (QIAGEN, Germany), prepared for sequencing using the Nextera XT DNA Library Preparation Kit and sequenced with a NextSeq 500/550 Mid Output v2.5 Kit (300 cycles) on a NextSeq 500 platform (Illumina). Raw plasmid sequence data were subjected to quality checking using FastQC v0.11.5,²⁵ trimmed with BBTools v.36.49²⁶ and assembled by SPAdes v3.15.3.²⁷ Assemblies were analysed by ResFinder and PlasmidFinder with thresholds as described above. Due to the common occurrence of co-extraction of chromosomal DNA during plasmid DNA extraction, seven-gene MLST analysis was successfully performed to distinguish between true TCs (having ST of the recipient strain) and mutated donor strains (having ST of the respective donor strain).

Results and discussion

Macrolide resistance genes in *E. coli* and *Salmonella* from food-producing animals and meat in the EU

We screened commensal *E. coli* and *Salmonella* isolates from food-producing animals and meat in Europe for the presence of macrolide resistance determinants and compared these genotypes with their respective azithromycin phenotypes. Eleven different macrolide resistance genes and five non-synonymous

rplDV mutations were detected in 248 *E. coli* and 26 *Salmonella* (*enterica* serovars *S. Rissen*, *S. Blockley*, *S. Typhimurium*, *S. Paratyphi-B-var-Java*, *S. Bredeney*, *S. Infantis*, *S. 1,4, [5],12:i:-*, *S. Derby* and *S. Dublin*) isolates (Tables S4, S5 and S6). The presence of one or more macrolide resistance gene(s) or known mutations was associated with the azithromycin resistance phenotype in 159 (66%) *E. coli* and 24 (92%) *Salmonella* isolates (Figure 1 and Table S4). No known genes or mutations mediating azithromycin resistance were detected in the remaining 1002 isolates, although 10 of these isolates exhibited azithromycin MIC > 16 mg/L (Table S4). Six genes, namely *mph(A)*, *mph(B)*, *mef(B)*, *erm(B)* and *mef(C)-mph(G)*, were detected in both *E. coli* and *Salmonella*, whereas *erm(C)*, *erm(42)*, *ere(A)* and *msr(E)-mph(E)* were detected in *E. coli* only (Table S4). Seven of all detected macrolide resistance genes [*mph(A)*, *mph(B)*, *mef(B)*, *erm(B)*, *erm(42)*, *msr(E)-mph(E)*] were found in isolates from all animal sources, while *mef(C)-mph(G)* were detected in isolates of bovine and porcine origin only, and *ere(A)* and *erm(C)* were harboured by isolates of porcine origin only (Figure 2, Table S4).

Regarding chromosomal mutations, five non-synonymous *rplDV* mutations were detected in seven *E. coli* isolates with MIC ≤ 16 mg/L (Table S4), while no *rplDV* mutations were detected in *Salmonella*. Mutations in 23S rRNA genes were not detected in any of the analysed genomes. Two azithromycin-resistant *Salmonella* isolates (serovars 1,4, [5],12:i:- and Dublin) with no azithromycin resistance gene harboured the AcrB-R717Q substitution. The 10 isolates with no identified mechanism of azithromycin resistance were screened for additional non-synonymous mutations in the *acrR* and *acrB* genes, and either no mutations were found, or the detected amino acid substitutions/deletions were also observed in susceptible isolates (Table S4).

Mph(A) operon structures in azithromycin-resistant and -susceptible isolates

One of the established macrolide resistance genes that confers high-level azithromycin resistance in Enterobacteriales is *mph(A)*.^{5,6,28} Here, *mph(A)* was detected in 15% of the isolates ($n=186$). The *E. coli* *mph(A)*-harbouring isolates ($n=175$) were genetically diverse, belonging to 77 STs, with ST744, ST1011, ST10 and ST410 being the most represented. The *mph(A)*-carrying *Salmonella* isolates ($n=11$) belonged to four STs (ST469, ST34, ST19 and ST52) (Table S4). The isolates harbouring *mph(A)* displayed a wide range of azithromycin MIC values ranging from 4 to >64 mg/L (Figure 2 and Table S4). Of these isolates, 27% ($n=50$) had a susceptible phenotype (MIC ≤ 16 mg/L). Previous studies have also shown that *mph(A)* can be present in azithromycin-susceptible isolates. For instance, Gomes *et al.*⁷ reported that 7% of the *mph(A)*-harbouring *E. coli* isolates in their study were susceptible, which suggests that the *mph(A)* gene does not always confer a resistant phenotype in Enterobacteriales.

The *mph(A)* gene encoding macrolide 2'-phosphotransferase I is part of an operon, *mph(A)-mrx-mphR(A)*, in which the downstream genes *mrx* and *mphR(A)* encode a protein with unknown function and a repressor that controls the inducible expression of *mph(A)*, respectively.²⁹ In this study, annotation of the complete *mph(A)* operon and analysis of the *mph(A)* promoter region, which overlaps with the *mphR(A)* binding site, revealed differences

Table 1. Characteristics of *E. coli* and *Salmonella* sp. isolates carrying *mef(C)-mph(G)* genes

Strain ID	Species	ENA run accession number	Host	Country of origin	Source of isolation	Year of isolation	MLST ST	Azithromycin MIC (mg/L)	Conjugative transfer of		Other AMR genes ^a
									<i>mef(C)-mph(G)</i>	Plasmid replicons ^a	
NRS-2015 ESBL-08-69	<i>E. coli</i>	ERR2019198	Porcine	Netherlands	Caecal content	2015	453	32	Yes	IncFIB, IncX1, IncIγ , IncFII, IncQ1, Col8282	tet(A) , <i>sul3</i> , sul2 , <i>ant(3')-Ia</i> , <i>cmiA1</i> , <i>aph(3')-Ib</i> , <i>aph(6)-Id</i> , <i>dfrA5</i> , bla_{TEM-52}
ZTA15:00420EB1	<i>E. coli</i>	ERR2019249	Porcine	Spain	Caecal content	2015	7456	32	No	ColE10, Col156, IncFII(pRSB107), Col(MG828), IncI γ , IncX4, IncHI2; IncHI2A, IncQ1	<i>tet(M)</i> , <i>mcr-4.1</i> , <i>aph(4)-Ia</i> , <i>aac(3)-Ia</i> , <i>tet(A)</i> , <i>bla_{CTX-M-14}</i> , <i>bla_{TEM-1B}</i> , <i>aph(3')-Ia</i> , <i>mcr-1.1</i> , <i>mph(B)</i> , <i>sul1</i> , <i>ant(3')-Ia</i> , <i>dfrA1</i> , <i>aph(6)-Id</i> , <i>aph(3')-Ib</i> , <i>sul2</i> , <i>aadA1</i>
HP-6957	<i>E. coli</i>	ERR3393110	Porcine	Czechia	Caecal content	2017	57	64	Yes	IncFIB, IncFIC(FII), IncIγ	—
ECO NRS 187.57	<i>E. coli</i>	ERR3393279	Porcine	Netherlands	Caecal content	2017	88	>64	Yes	IncFIB, IncFIC(FII), IncIγ	—
17-AB00518_0	<i>E. coli</i>	ERR3393142	Bovine	Germany	Caecal content	2017	48	64	Yes	IncIγ , Col(pHAD28)	sul2 , bla_{CTX-M-1}
10047105/3	<i>E. coli</i>	ERR3393194	Porcine	Hungary	Meat	2017	453	>64	Yes	IncFII , IncI γ , IncY	sul2 , <i>bla_{CTX-M-1}</i> , tet(A) , <i>dfrA17</i> , <i>aadA5</i>
HP36996	<i>E. coli</i>	ERR3393109	Porcine	Czechia	Caecal content	2017	165	>64	No	IncFII, IncI γ , Col(MG828)	—
17-AB01235_0	<i>E. coli</i>	ERR3393151	Bovine	Germany	Caecal content	2017	NT ^b	32	No	IncFIA, IncFIB, IncFIC, IncI γ , IncX4	<i>sul2</i> , <i>tet(B)</i> , <i>bla_{CTX-M-1}</i> , <i>fosA7</i>
17-AB02375_0	<i>E. coli</i>	ERR3393160	Porcine	Germany	Caecal content	2017	57	64	No	IncFIA(HI1), IncFIB, IncFIC(FII), IncI γ , IncHI1A, IncHI1B	<i>sul2</i> , <i>sul3</i> , <i>bla_{CTX-M-1}</i> , <i>tet(A)</i> , <i>tet(B)</i> , <i>dfrA12</i> , <i>dfrA1</i> , <i>qnrS1</i> , <i>aadA1</i> , <i>aadA2</i> , <i>aadA5</i> , <i>cmiA</i> , <i>flor</i> , <i>aph(3')-Ib</i> , <i>aph(3')-Ia</i> , <i>aph(6)-Id</i> , <i>bla_{TEM-1B}</i> , <i>ant(3')-1</i> , <i>tet(M)</i>

Accession	Species	Accession	Host	Location	Year	Sample type	Number of reads	Number of contigs	Number of genes	AMR genes	
10849-2GR	<i>E. coli</i>	ERR393337	Porcine	Romania	2017	Caecal content	101	64	No	Col156, ColRNAI, IncFIB, IncFIC(FII), IncIγ, IncFII, Col(pHAD28) IncIα	sul2, bla _{CTX-M-1} , tet(A), aadA5, aadA1, dfrA17, dfrA1, ant(3')-Ia sul2, aadA5, dfrA17, bla _{CTX-M-1}
Ec151	<i>E. coli</i>	ERR11629170	Bovine	Denmark	2019	Caecal content	8863	>64	ND	IncFIB, IncFII	sul1, sul2, bla _{TEM-1B} , aadA1, aac(6')-Ia; tet(A), dfrA1
S308	<i>Salmonella</i> 1,4,[5],12:i:c	ERR11628334	Porcine	Denmark	2017	Clinical	19	>64	No	ColpVC, IncQ1	sul2, aac(6')-Ia; aph(6)-Ia, aph(3')-Ib, fosA7, dfrA17
S514	<i>Salmonella</i> Derby ^d	ERR11628347	Porcine	Denmark	2018	Clinical	40	>16	No		

^aIn bold, plasmid replicons and AMR genes detected in TCs.

^bNot typeable.

^c*S. enterica* subsp. *enterica* serovar 1,4, [5],12:i:-.

^d*S. enterica* subsp. *enterica* serovar Derby.




Figure 1. Sankey plots showing macrolide resistance genes/mutations and azithromycin MICs in the *E. coli* and *Salmonella* isolates in this study. *mph(A)* gene and its combinations with other macrolide resistance genes (a), *erm* genes and their combinations with other macrolide resistance genes (b), other macrolide resistance genes (c). The *rplDV* non-synonymous mutations detected in seven susceptible isolates were not included in the graph (Table S4). It is important to note that in resistant isolates harbouring *mph(A)* in combination with *mph(B)* or *mef(B)*, the full *mph(A)* operon was present, and the promoter region was complete in all cases (Table S4). The 10 resistant isolates (MIC > 16 mg/L) without known azithromycin resistance mechanisms are not included in the graph. This figure appears in colour in the online version of *JAC* and in black and white in the print version of *JAC*.

between azithromycin-susceptible and -resistant isolates. In general, resistant isolates ($n = 136$; 73%) had an intact *mph(A)* operon (Figure 3a), while *mph(A)*-harbouring susceptible isolates ($n = 50$; 27%) had altered operon structure (Figure 3b). Most commonly, susceptible isolates lacked the *mphR(A)* repressor gene and part of the *mrx* gene, disrupted by an *ISEcp1-bla_{CTX-M-1}* transposition unit. Additionally, the susceptible isolates ($n = 50$) had either a 24 bp ($n = 23$) or a 66 bp ($n = 14$) deletion spanning the *mphR(A)* binding site and/or the transcription start site and the ribosomal binding site (RBS). Because it has been previously suggested that bacterial cells sense the presence of macrolides through the MphR(A) repressor by forming a macrolide-repressor complex,²⁹ our interpretation of these results is that *mph(A)* is not expressed in susceptible isolates due to the lack of the macrolide-repressor complex.

Among the resistant *mph(A)*-harbouring isolates ($n = 136$), 117 followed the common *mph(A)* structure for resistant isolates [intact regulatory region and full *mph(A)* operon] (Figure 3a). One isolate (16037780A201X5, MIC = 32 mg/L) had a truncated regulatory region (Table S4) and three isolates had disruptions into the *mph(A)* operon due to Tn insertions (Figure 3c, 1–3). Additionally, four isolates had too short contigs to assess the completeness of the regulatory regions (Table S4). In *S. Blockley* 5601, the complete *mph(A)* operon was chromosomally integrated (Figure 3c, 5) as previously reported for *S. Blockley* strain 159383.³⁰

Exceptions to the *mph(A)* operon structure that is typical in the susceptible isolates [truncated regulatory region and lack of *mphR(A)*] were also observed. The regulatory region was intact in 13 isolates, and in 5 of them *mphR(A)* was absent, whereas in 7 of them *mphR(A)* was present but *mrx* had a deletion mutation at nucleotide position 576 leading to a premature stop codon (PMSC) and truncated Mrx. However, previous cloning experiments

in minicells carrying *mrx* with a non-sense mutation showed that Mph(A) production was still enhanced in presence of erythromycin as in minicells carrying non-mutated (WT) *mrx* gene.²⁹ Therefore, the deletion mutation in *mrx* in the seven isolates in our study might not be the explanation for the observed susceptible phenotype. Additionally, in isolate NRS_2017_ESBL_19.14 (MIC = 8 mg/L), *mphR(A)* was present and the regulatory region intact, but *mrx* was located in two contigs. In isolate U16-0318X (MIC = 8 mg/L) *mph(A)* had a deletion (Figure 3c, 4).

Despite exceptions to each of the two most common *mph(A)* operon structures, the size of the dataset analysed and the high concordance between *mph(A)* operon structures and observed phenotypes suggest that in addition to the *mph(A)* gene, the *mphR(A)* repressor gene is also required for azithromycin resistance. Nevertheless, *in vitro* studies are needed to elucidate the association between the absence of *mphR(A)*, truncation in the regulatory region and susceptibility to azithromycin.

mef(C)-mph(G) genes

The *mef(C)-mph(G)* tandem genes, conferring high-level azithromycin resistance, were first identified in *Photobacterium damsela* subsp. *damsela*³¹ on an MDR plasmid pAQU1 and subsequently found in other marine and enteric bacteria from fish intestines.^{31,32} Since their first report in Enterobacteriales, in a Shiga toxin-producing *E. coli* (STEC) isolate in 2021,⁹ *mef(C)-mph(G)* have been detected in *E. coli* from various origins and geographical locations.^{33–36}

We first detected *mef(C)-mph(G)* in *E. coli* and *Salmonella* isolates included in the 2018 EU harmonized monitoring of AMR and the Danish Programme for surveillance of antimicrobial consumption and resistance (results included in this study). Here, screening





Figure 2. Distribution of macrolide resistance genes in *E. coli* and *Salmonella* isolates in this study according to their origin. Twenty-one percent ($n = 51$) of the isolates carried more than one macrolide resistance gene and are included under each of the genes they carry. The total number of genes in each species or from each source is given in parentheses. Two *mph(A)*- and two *mph(B)*-harbouring *E. coli* isolates for which isolation sources are not available are not included. This figure appears in colour in the online version of JAC and in black and white in the print version of JAC.

of all *E. coli* and *Salmonella* isolates collected between 2014 and 2018 identified *mef(C)-mph(G)* in 11 *E. coli* and two *Salmonella* isolates from bovine and porcine origin. In all isolates, *mef(C)-mph(G)* were associated with an azithromycin-resistant phenotype (MIC > 16 mg/L). No other known macrolide resistance mechanisms were detected in these isolates, except *mph(B)* in isolate ZTA15:00420EB1 (Tables 1 and S4). However, other AMR genes were found in the *mef(C)-mph(G)*-harbouring isolates, i.e. *tet(A)*, *sul2*, *bla_{TEM-52}* and *bla_{CTX-M-1}*, in all cases located on the same contig as *mef(C)-mph(G)* (Figure 4). The *E. coli* isolates carrying *mef(C)-mph(G)*


belonged to eight different STs, and the two *Salmonella* isolates, *S. Derby* and *S. 1,4, [5],12:i:-*, belonged to ST40 and ST19, both of which have frequently been associated with the pork production chain (Table 1).^{37,38}

Since *mef(C)-mph(G)* genes were previously described to be located on plasmids and associated with various mobile genetic elements, including plasmids of various sizes and integrative conjugative elements,^{9,32} we carried out conjugation experiments to assess their transferability. The conjugation performed for 10 *E. coli* and two *Salmonella* isolates showed that *mef(C)-mph(G)* could be located on transferrable plasmids, as in 5 of the *E. coli*

(a) Resistant isolates (MIC > 16 mg/L) - full *mph(A)* operon and intact regulatory region



(b) Susceptible isolates (MIC ≤ 16 mg/L) - disrupted *mphA(R)*, partial *mxr*, and truncated regulatory region



(c) Isolates with other structures due to Tn and IS elements insertions




Figure 3. *mph(A)* operon structure in azithromycin-resistant isolates (MIC > 16 mg/L) (a), azithromycin-susceptible isolates (MIC ≤ 16 mg/L) (b), and in isolates with Tn and IS insertions (c). *S. Blockley 5601* (c5) carries the *mph(A)* operon as part of an MDR region, consisting of the streptomycin resistance cluster *aph(3')-Ib-aph(6)-Id-aph(3')-Ia* and the *tet(A)* gene. NCBI blastn revealed 99.24%–100% identity and 100% coverage between the whole streptomycin and azithromycin resistance genomic region of *S. Blockley* isolate 5601 and *S. Blockley* strain 159383 (GenBank accession number CP043662.1: 4326383–4352831). The MDR region was reconstructed using Bandage (<http://github.com/rwwick/Bandage>). *istA*—IS21-like element IS21 family transposase *IstA*, *istB*—IS21-like element IS21 family ATPase *IstB*. In (a), the difference between the two sequences is T instead of C in the –10 region. Wavy lines at the end(s) of diagrams show where only part of the contig is included. This figure appears in colour in the online version of *JAC* and in black and white in the print version of *JAC*.




Figure 4. Comparative analyses of contigs harbouring *mef(C)*-*mph(G)* in *E. coli* and *Salmonella*. D, donor (included for isolates that did not yield TCs), ND, conjugation not performed. Wavy lines at the end(s) of diagrams show where only part of the contig is included. This figure appears in colour in the online version of *JAC* and in black and white in the print version of *JAC*.

isolates the genes were successfully transferred to the *E. coli* J62-2 recipient (Table 1). Unequivocal association between *mef(C)*-*mph(G)* and a plasmid replicon gene could not be made using short-read sequencing data. However, in all TCs, except TC_NRS-2015 ESB-08-69, only one plasmid replicon was detected, which allowed us to associate *mef(C)*-*mph(G)* with IncFII in isolate TC_10047105-3 and IncI γ in isolates TC_17-AB00518-0, TC_HP-6957 and TC_ECO-NRS-187-57 (Table 1). Analyses of the genetic contexts of *mef(C)*-*mph(G)* in the donor strains and TCs revealed that the genes were associated with truncated TnAs1 consisting of *tnpR* and *tnpA* genes in IncFII plasmids and ISCR2 in IncI γ plasmids, with both elements located downstream of *mef(C)*-*mph(G)* (Figure 4). The ISCR2 is known to be associated with dissemination of multiple resistance genes in various species.³⁹

Other azithromycin resistance genes

The relevance of the plasmid-borne *erm* genes in mediating high-level azithromycin resistance was confirmed in this study as previously described.^{7,8,35,40} The presence of *erm(C)*, *erm(B)* and *erm(42)*, alone or in combination with other macrolide resistance genes, was associated with a resistant phenotype (MIC \geq 64 mg/L) in all cases. Association of *erm* genes with high azithromycin MIC levels has been reported in *E. coli* even in the presence of efflux pump inhibitors.^{6,41,42}

The *mph(B)* gene, the second most detected gene in our collection, encodes macrolide 2'-phosphotransferase II, which phosphorylates 14-membered (e.g. erythromycin) and 16-membered (e.g. tylosin) macrolides.^{43,44} Nevertheless, 6 out of 26 isolates harbouring *mph(B)* as the sole macrolide resistance gene in our collection had azithromycin MIC > 16 mg/L (Table S4). The only *mph(B)*-harbouring isolate in the study of Gomes *et al.*⁷ exhibited an MIC of 16 mg/L in the absence of the efflux pump inhibitor Pa β N. Previous cloning and expression analysis of *mph(B)* in *E. coli* also showed that *mph(B)* did not confer resistance to azithromycin.⁴⁴

In addition, we detected *mef(B)*, *msr(E)*-*mph(E)* or *ere(A)* as the only macrolide resistance gene in isolates with azithromycin MIC > 16 mg/L. While azithromycin-resistant *ere(A)*-carrying *E. coli*, even in the presence of Pa β N, has been observed,⁷ there is no description of *msr(E)*-*mph(E)* or *mef(B)* conferring azithromycin resistance in *E. coli* or *Salmonella*. Generally, the decreased azithromycin susceptibility in isolates with MIC = 32 mg/L without a known mechanism of azithromycin resistance could be explained by the one 2-fold dilution MIC variation from the cut-off for resistance. However, the occurrences of high-level azithromycin resistance (MIC \geq 64 mg/L) that cannot be explained by genes detected in this study suggest that unknown mechanism(s) could be involved in the reduced azithromycin susceptibility. For instance, a recent study using hidden Markov models identified and experimentally validated five novel macrolide resistance

genes that increased the azithromycin resistance when cloned into *E. coli*.⁴⁵

Study limitations

A limitation in our study was that we could not unequivocally determine whether the detected macrolide resistance genes were located on the chromosome or on plasmids due to the nature of the Illumina short-read sequencing technology. Furthermore, the association of AMR genes with IS or Tn elements is not entirely certain, due to the possibility of mis-assemblies, although studies suggest that these tend to be rare for high-coverage data. Both limitations could be overcome by long-read sequencing technologies.

Conclusions

We investigated azithromycin resistance determinants and phenotype-genotype correlation in a collection of 1276 *E. coli* and *Salmonella* isolates from food-producing animals and meat in Europe. Eleven different genes previously associated with azithromycin resistance were detected. Our study highlights that the interpretation of *mph(A)*-mediated phenotypes in Enterobacterales requires consideration of the entire *mph(A)* operon and its regulatory region. Generally, azithromycin-susceptible isolates lacked the *mphR(A)* repressor gene and had a truncated *mphR(A)* binding site, while azithromycin-resistant isolates had an intact *mph(A)* operon and complete *mphR(A)* binding site. We observed elevated azithromycin MIC values (≥ 64 mg/L) in all cases when an *erm* gene was found alone or associated with another macrolide resistance determinant. Moreover, our study provides additional insight into the genetic context and transferability of *mef(C)*-*mph(G)* genes, which are emerging in *E. coli* and *Salmonella*. These genes were found to be associated with various transposable elements, often harboured on conjugative plasmids and containing additional AMR genes, thus highlighting concerns for co-selection and spread.

Acknowledgements

We gratefully acknowledge Birthe S. Rosenqvist Lund, Inge Marianne Hansen, Christina Aaby Svendsen, Jacob Dyring Jensen and Gunhild Larsen for performing antimicrobial susceptibility testing and WGS. The Department of Animal Health and Antimicrobial Strategies (Uppsala, Sweden) is acknowledged for sharing isolates for this study. We thank Veterinary Services (Nicosia, Cyprus), Istituto Zooprofilattico Sperimentale del Lazio e della Toscana 'M. Aleandri' (Rome, Italy), Laboratoire de Médecine Vétérinaire de l'État (LMVE, Dudelange, Luxembourg) and National Veterinary Research Institute (Pulawy, Poland) for the WGS of selected isolates for this study. We would also like to acknowledge the anonymous reviewers for their expertise to improve the manuscript.

Funding

This work was supported by general funding to the European Union Reference Laboratory for Antimicrobial Resistance (EURL-AR) (<http://www.eurl-ar.eu/>).

Transparency declarations

B.G. and P.A.B. are currently employed by the European Food Safety Authority (EFSA) in its BIOHAW Unit that provides scientific and

administrative support to EFSA's scientific activities in the area of microbial risk assessment. The positions and opinions presented in this article are those of the authors alone and are not intended to represent the views or scientific works of EFSA. All other authors: none to declare.

Author contributions

M.I. and V.B. designed the experiments. M.I., A.O. and H.M. carried out the conjugation experiments. M.I., A.O., A.M.S., R.S.H., V.B. and J.S.K. analysed the data at the EURL-AR. B.G. and P.A.B. analysed the data and coordinated the isolate selection at EFSA. M.I. performed the bioinformatic analyses, data visualizations, wrote the draft manuscript and addressed reviewers comments during revisions. V.B., J.S.K. and A.M.V.B. revised the draft manuscript. J.S.K. and A.M.V.B. helped with addressing reviewers' comments. R.S.H., A.M.V.B. and G.H.L. supervised the project. All additional authors contributed to sample preparation, data analysis in the originating countries and revision of the draft manuscript.

Supplementary data

Tables S1–S6 are available as [Supplementary data](#) at JAC Online.

References

- WHO. The WHO list of Critically Important Antimicrobials. 2019. <https://www.who.int/groups/advisory-group-on-the-who-list-of-critically-important-antimicrobials>.
- Hooda Y, Sajib MS, Rahman H et al. Molecular mechanism of azithromycin resistance among typhoidal *Salmonella* strains in Bangladesh identified through passive pediatric surveillance. *PLoS Negl Trop Dis* 2019; **13**: e0007868. <https://doi.org/10.1371/journal.pntd.0007868>
- Lübbert C. Antimicrobial therapy of acute diarrhoea: a clinical review. *Expert Rev Anti Infect Ther* 2016; **14**: 193–206. <https://doi.org/10.1586/14787210.2016.1128824>
- Darton TC, Tuyen HT, The HC et al. Azithromycin resistance in *Shigella* spp. in Southeast Asia. *Antimicrob Agents Chemother* 2018; **62**: e01748-17. <https://doi.org/10.1128/AAC.01748-17>
- Timmermans M, Latour S, Ceysens P-J et al. The AMR-ARRAY: a modular bead array detecting β -lactam, (fluoro) quinolone, colistin, aminoglycoside and macrolide resistance determinants in Gram-negative bacteria. *J Microbiol Methods* 2022; **196**: 106472. <https://doi.org/10.1016/j.mimet.2022.106472>
- Gomes C, Martínez-Puchol S, Palma N et al. Macrolide resistance mechanisms in Enterobacteriaceae: focus on azithromycin. *Crit Rev Microbiol* 2017; **43**: 1–30. <https://doi.org/10.3109/1040841X.2015.1136261>
- Gomes C, Ruiz-Roldán L, Mateu J et al. Azithromycin resistance levels and mechanisms in *Escherichia coli*. *Sci Rep* 2019; **9**: 6089. <https://doi.org/10.1038/s41598-019-42423-3>
- Nguyen MCP, Woerther PL, Bouvet M et al. *Escherichia coli* as reservoir for macrolide resistance genes. *Emerg Infect Dis* 2009; **15**: 1648–50. <https://doi.org/10.3201/eid1510.090696>
- Bizot E, Cointe A, Bidet P et al. Azithromycin resistance in Shiga-toxin producing *Escherichia coli* in France between 2004 and 2020 and detection of *mef(C)*-*mph(G)* genes. *Antimicrob Agents Chemother* 2021; **66**: e0194921. <https://doi.org/10.1128/AAC.01949-21>
- European Commission. Commission implementing decision of 12 November 2013 on the monitoring and reporting of antimicrobial resistance in zoonotic and commensal bacteria (2013/652/EU). https://eur-lex.europa.eu/eli/dec_impl/2013/652/oj.
- European Commission. Commission implementing decision (EU) 2020/1729 of 17 November 2020 on the monitoring and reporting of antimicrobial resistance in zoonotic and commensal bacteria and

- repealing implementing decision 2013/652/EU. (2020/1729/EU). https://eur-lex.europa.eu/eli/dec_impl/2020/1729/oj.
- 12** European Food Safety Authority, ECDC. The European Union Summary Report on Antimicrobial Resistance in zoonotic and indicator bacteria from humans, animals and food in 2020/2021. *EFSA J* 2023; **21**: e07867. <https://doi.org/10.2903/j.efsa.2023.7867>
- 13** European Food Safety Authority, Aerts M, Battisti A *et al.* Technical specifications on harmonised monitoring of antimicrobial resistance in zoonotic and indicator bacteria from food-producing animals and food. *EFSA J* 2019; **17**: 5709. <https://doi.org/10.2903/j.efsa.2019.5709>
- 14** Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 2014; **30**: 2114–20. <https://doi.org/10.1093/bioinformatics/btu170>
- 15** Nurk S, Meleshko D, Korobeynikov A *et al.* metaSPAdes: a new versatile metagenomic assembler. *Genome Res* 2017; **27**: 824–34. <https://doi.org/10.1101/gr.213959.116>
- 16** Gurevich A, Saveliev V, Vyahhi N *et al.* QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 2013; **29**: 1072–5. <https://doi.org/10.1093/bioinformatics/btt086>
- 17** Bortolaia V, Kaas RS, Ruppe E *et al.* ResFinder 4.0 for predictions of phenotypes from genotypes. *J Antimicrob Chemother* 2020; **75**: 3491–500. <https://doi.org/10.1093/jac/dkaa345>
- 18** Gunell M, Kotilainen P, Jalava J *et al.* *In vitro* activity of azithromycin against nontyphoidal *Salmonella enterica*. *Antimicrob Agents Chemother* 2010; **54**: 3498–501. <https://doi.org/10.1128/AAC.01678-09>
- 19** Zhang S, den Bakker HC, Li S *et al.* SeqSero2: rapid and improved *Salmonella* serotype determination using whole-genome sequencing data. *Appl Environ Microbiol* 2019; **85**: e01746–19. <https://doi.org/10.1128/AEM.01746-19>
- 20** Carattoli A, Zankari E, Garcia-Fernández A *et al.* *In silico* detection and typing of plasmids using PlasmidFinder and plasmid multilocus sequence typing. *Antimicrob Agents Chemother* 2014; **58**: 3895–903. <https://doi.org/10.1128/AAC.02412-14>
- 21** Siguier P. ISfinder: the reference centre for bacterial insertion sequences. *Nucleic Acids Res* 2006; **34**: D32–6. <https://doi.org/10.1093/nar/gkj014>
- 22** Wickham H. *Ggplot2: Elegant Graphics for Data Analysis*. 2nd edn. Springer International Publishing, 2016.
- 23** Allaire JJ, Gandrud C, Russell K *et al.* NetworkD3: D3 JavaScript Network Graphs from R. 2017. <https://CRAN.R-project.org/package=networkD3>.
- 24** Hansen KH, Bortolaia V, Nielsen CA *et al.* Host-specific patterns of genetic diversity among IncI1- γ and IncK plasmids encoding CMY-2 β -lactamase in *Escherichia coli* isolates from humans, poultry meat, poultry, and dogs in Denmark. *Appl Environ Microbiol* 2016; **82**: 4705–14. <https://doi.org/10.1128/AEM.00495-16>
- 25** Andrews S. FastQC: A quality control tool for high throughput sequence data. 2010. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>.
- 26** Bushnell B, Rood J, Singer E. BBMerge—accurate paired shotgun read merging via overlap. *PLoS One* 2017; **12**: e0185056. <https://doi.org/10.1371/journal.pone.0185056>
- 27** Bankevich A, Nurk S, Antipov D *et al.* SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 2012; **19**: 455–77. <https://doi.org/10.1089/cmb.2012.0021>
- 28** Xiang Y, Wu F, Chai Y *et al.* A new plasmid carrying *mphA* causes prevalence of azithromycin resistance in enterotoxigenic *Escherichia coli* serogroup O6. *BMC Microbiol* 2020; **20**: 247. <https://doi.org/10.1186/s12866-020-01927-z>
- 29** Noguchi N, Takada K, Katayama J *et al.* Regulation of transcription of the *mph(A)* gene for macrolide 2-phosphotransferase I in *Escherichia coli*: characterization of the regulatory gene *mphR(A)*. *J Bacteriol* 2000; **182**: 5052–8. <https://doi.org/10.1128/JB.182.18.5052-5058.2000>
- 30** Nair S, Ashton P, Doumith M *et al.* WGS for surveillance of antimicrobial resistance: a pilot study to detect the prevalence and mechanism of resistance to azithromycin in a UK population of non-typhoidal *Salmonella*. *J Antimicrob Chemother* 2016; **71**: 3400–8. <https://doi.org/10.1093/jac/dkw318>
- 31** Nonaka L, Maruyama F, Suzuki S *et al.* Novel macrolide-resistance genes, *mef(C)* and *mph(G)*, carried by plasmids from *Vibrio* and *Photobacterium* isolated from sediment and seawater of a coastal aquaculture site. *Lett Appl Microbiol* 2015; **61**: 1–6. <https://doi.org/10.1111/lam.12414>
- 32** Sugimoto Y, Suzuki S, Nonaka L *et al.* The novel *mef(C)*–*mph(G)* macrolide resistance genes are conveyed in the environment on various vectors. *J Glob Antimicrob Resist* 2017; **10**: 47–53. <https://doi.org/10.1016/j.jgar.2017.03.015>
- 33** González-Santamarina B, Weber M, Menge C *et al.* Comparative genomic analysis of antimicrobial-resistant *Escherichia coli* from South American camelids in Central Germany. *Microorganisms* 2022; **10**: 1697. <https://doi.org/10.3390/microorganisms10091697>
- 34** Irrgang A, Tenhagen B-A, Pauly N *et al.* Characterization of VIM-1-producing *E. coli* isolated from a German fattening pig farm by an improved isolation procedure. *Front Microbiol* 2019; **10**: 2256. <https://doi.org/10.3389/fmicb.2019.02256>
- 35** Ma Y, Pirollo M, Subramani P *et al.* Macrolide resistance and *in vitro* potentiation by peptidomimetics in porcine clinical *Escherichia coli*. *mSphere* 2022; **7**: e00402–22. <https://doi.org/10.1128/msphere.00402-22>
- 36** Suzuki S, Kadoya A, Masuda N *et al.* Macrolide resistance genes and mobile genetic elements in waterways from pig farms to the sea in Taiwan. *J Glob Antimicrob Resist* 2022; **29**: 360–70. <https://doi.org/10.1016/j.jgar.2022.04.024>
- 37** Cai Y, Tao J, Jiao Y *et al.* Phenotypic characteristics and genotypic correlation between *Salmonella* isolates from a slaughterhouse and retail markets in Yangzhou, China. *Int J Food Microbiol* 2016; **222**: 56–64. <https://doi.org/10.1016/j.ijfoodmicro.2016.01.020>
- 38** Zhou Z, Li J, Zheng H *et al.* Diversity of *Salmonella* isolates and their distribution in a pig slaughterhouse in Huaian, China. *Food Control* 2017; **78**: 238–46. <https://doi.org/10.1016/j.foodcont.2017.02.064>
- 39** Toleman MA, Bennett PM, Walsh TR. ISCR elements: novel gene-capturing systems of the 21st century? *Microbiol Mol Biol* 2006; **70**: 296–316. <https://doi.org/10.1128/MMBR.00048-05>
- 40** Liu X, Yang X, Ye L *et al.* Genetic characterization of a conjugative plasmid that encodes azithromycin resistance in Enterobacteriaceae. *Microbiol Spectr* 2022; **10**: e00788–22. <https://doi.org/10.1128/spectrum.00788-22>
- 41** Harmer CJ, Holt KE, Hall RM. A type 2 A/C2 plasmid carrying the *aacC4* apramycin resistance gene and the *erm(42)* erythromycin resistance gene recovered from two *Salmonella enterica* serovars. *J Antimicrob Chemother* 2015; **70**: 1021–5. <https://doi.org/10.1093/jac/dku489>
- 42** Chiou C-S, Hong Y-P, Wang Y-W *et al.* Antimicrobial resistance and mechanisms of azithromycin resistance in nontyphoidal *Salmonella* isolates in Taiwan, 2017 to 2018. *Microbiol Spectr* 2023; **11**: e03364–22. <https://doi.org/10.1128/spectrum.03364-22>
- 43** Kono M. Purification and characterization of macrolide 2'-phosphotransferase type II from a strain of *Escherichia coli* highly resistant to macrolide antibiotics. *FEMS Microbiol Lett* 1992; **97**: 89–94. [https://doi.org/10.1016/0378-1097\(92\)90369-y](https://doi.org/10.1016/0378-1097(92)90369-y)
- 44** Chesneau O, Tsvetkova K, Courvalin P. Resistance phenotypes conferred by macrolide phosphotransferases. *FEMS Microbiol Lett* 2007; **269**: 317–22. <https://doi.org/10.1111/j.1574-6968.2007.00643.x>
- 45** Lund D, Kieffer N, Parras-Moltó M *et al.* Large-scale characterization of the macrolide resistome reveals high diversity and several new pathogen-associated genes. *Microb Genomics* 2022; **8**: 000770. <https://doi.org/10.1099/mgen.0.000770>