

Ongoing mpox outbreak in Kamituga, South Kivu province, associated with monkeypox virus of a novel Clade I sub-lineage, Democratic Republic of the Congo, 2024

Leandre Murhula Masirika^{1,2}, Jean Claude Udahemuka^{3,4}, Leonard Schuele⁵, Pacifique Ndishimye^{4,6,7}, Saria Otani⁸, Justin Bengehya Mbiribindi⁹, Jean M. Marekani¹⁰, Léandre Mutimbwa Mambo¹¹, Nadine Malyamungu Bubala¹², Marjan Boter⁵, David F. Nieuwenhuijse⁵, Trudie Lang¹³, Ernest Balyahamwabo Kalalizi², Jean Pierre Musabyimana^{4,14}, Frank M. Aarestrup⁸, Marion Koopmans⁵, Bas B. Oude Munnink^{5,*}, Freddy Belesi Siangoli^{9,*}

1. Centre de Recherche en Sciences Naturelles de Lwiro, DS Bukavu, South Kivu, Bukavu, Democratic Republic of the Congo
2. SaBio Instituto de Investigación en Recursos Cinegéticos IREC (Universidad de Castilla-La Mancha & CSIC), Ciudad Real, Spain
3. Department of Veterinary Medicine, University of Rwanda, Nyagatare, Rwanda
4. Stansile Research Organization, Kigali, Rwanda
5. Department of Viroscience, Erasmus University Medical Center, Rotterdam, The Netherlands
6. Department of Microbiology and Immunology, Dalhousie University, Halifax, Nova Scotia, Canada
7. Research and Innovation Centre, African Institute for Mathematical Sciences, Kigali, Rwanda
8. Research Group for Genomic Epidemiology, National Food Institute, Technical University of Denmark, Kongens Lyngby, Denmark
9. Division Provinciale de la Santé, South Kivu, Bukavu, Democratic Republic of the Congo
10. Unit of Animal Production and Health, Nature Conservation and Development, Department of Biology, Faculty of Science, University of Kinshasa, Kinshasa, Democratic Republic of the Congo
11. Zone de Santé de Kamituga, Kamituga, South Kivu, Bukavu, Democratic Republic of the Congo
12. Hôpital Général de Référence de Kamituga, South Kivu, Bukavu, Democratic Republic of the Congo
13. The Global Health Network, Oxford University, Oxford, United Kingdom
14. Research, innovation and data science division, Rwanda Biomedical Center, Kigali, Rwanda

* These authors contributed equally

Correspondence: Leandre Murhula (leandremurhula@gmail.com)

Citation style for this article:

Masirika Leandre Murhula, Udahemuka Jean Claude, Schuele Leonard, Ndishimye Pacifique, Otani Saria, Mbiribindi Justin Bengehya, Marekani Jean M., Mambo Léandre Mutimbwa, Bubala Nadine Malyamungu, Boter Marjan, Nieuwenhuijse David F., Lang Trudie, Kalalizi Ernest Balyahamwabo, Musabyimana Jean Pierre, Aarestrup Frank M., Koopmans Marion, Oude Munnink Bas B., Siangoli Freddy Belesi. Ongoing mpox outbreak in Kamituga, South Kivu province, associated with monkeypox virus of a novel Clade I sub-lineage, Democratic Republic of the Congo, 2024. *Euro Surveill.* 2024;29(11):pii=2400106. <https://doi.org/10.2807/1560-7917.ES.2024.29.11.2400106>

Article submitted on 14 Feb 2024 / accepted on 13 Mar 2024 / published on 14 Mar 2024

Since the beginning of 2023, the number of people with suspected monkeypox virus (MPXV) infection have sharply increased in the Democratic Republic of the Congo (DRC). We report near-to-complete MPXV genome sequences derived from six cases from the South Kivu province. Phylogenetic analyses reveal that the MPXV affecting the cases belongs to a novel Clade I sub-lineage. The outbreak strain genome lacks the target sequence of the probe and primers of a commonly used Clade I-specific real-time PCR.

In the Democratic Republic of the Congo (DRC), the numbers of people with suspected infection with monkeypox virus (MPXV), the virus that causes mpox, have increased since the start of 2023. A total of 12,569 suspected mpox cases have been reported up to 12 November, the highest number of annual cases ever recorded [1]. The case fatality rate has been estimated at 4.6% [1]; moreover, new cases have occurred in geographical areas of the country where the disease was previously not observed, such as Kinshasa and South


Kivu province [1,2]. Despite this concerning situation, there is only limited genomic information available on the circulating viruses, which suggests that they belong to Clade I [3]. To gain more insight into the characteristics of the strains causing the epidemic, as well as assurance that current and commonly used molecular assays to diagnose MPXV infections can detect these strains, we sequenced monkeypox viral genomes from recently diagnosed cases in South Kivu, DRC.

Case definitions and patient characteristics

A case was listed as ‘suspect’ if presenting with an acute illness with fever, intense headache, myalgia, and back pain, followed by 1 to 3 days of a progressively developing rash often starting on the face and spreading on the body. A confirmed mpox case had a monkeypox virus infection which was laboratory-confirmed by PCR. A case was listed as ‘probable’ when satisfying the clinical definition of a suspected case and having an epidemiological link to a confirmed or

FIGURE 1

Phylogenetic analysis of all near-to-complete genome mpox sequences from Africa available on GISAID and the NCBI including the 2024 sequences from Kamituga, Democratic Republic of the Congo (n = 121 sequences)



DRC: Democratic Republic of the Congo; NCBI: National Center for Biotechnology Information.

The scale under the tree represents the number of substitutions per site. Sequences from Kamituga are presented in magenta, previous sequences from DRC in red and sequences from the 2022 outbreak in blue [14]. Arrows indicate sequences which have been shown in a prior publication to cluster with the previous sequences from a 2023 MPXV outbreak from Kwilu Province near Kinshasa [3]. Lineages are indicated as proposed by Nakazawa et al. [19] and Berthet et al. [20].

We gratefully acknowledge all data contributors, i.e., the authors and their originating laboratories responsible for obtaining the specimens, and their submitting laboratories for generating the genetic sequence and metadata and sharing via the GISAID Initiative, on which this research is based.

probable case; a probable case was not laboratory-confirmed [4].

The study involved patients from South Kivu province in the territory of Mwenga, who were hospitalised in the Kamituga hospital, which is in the Kamituga health zone. The first mpox cases in this area were reported from September 2023 onwards.

A total of 10 patients were included in the study. All were young adults in their late teenage up to the age of mid-20 years and five were male and five females. Regarding professions comprised, the majority of the concerned individuals were sex workers. For these patients, admission to the Kamituga Hospital had been based on clinical diagnosis of mpox by hospital staff. According to routines, skin lesion and oropharyngeal swabs collected from the patients had been sent to the national medical research institute of the DRC (Institut National de la Recherche Biomédicale; INRB) in Goma for MPXV infection confirmation by PCR, and all patients had tested positive for the virus.

DNA library preparation, sequencing and phylogenetic analysis

The samples from the 10 patients had been collected throughout the month of January 2024, stored in virus transport medium and frozen at -20°C . For sequencing, DNA was extracted with the Blood and Tissue Kit (Qiagen) according to the manufacturer's recommendations. Viral transport medium was used as a negative control. MPXV DNA was amplified using a mpox amplification scheme specifically designed for Clade IIb MPXV to generate 2,500bp amplicons and quantified using the Qubit dsHS DNA assay (ThermoFisher) [5]. Sequencing libraries were prepared using the Native Barcoding Kit 24 V14 (SQK-NBD114.24, Oxford Nanopore Technologies; ONT) using a R10 flow cell.

Sequencing reads were basecalled using dorado v7.2.13 with the super-accurate base calling model (ONT). Primer sequences were removed using cutadapt v4.6 [6] and aligned against a MPXV reference (GenBank accession: OQ729808) in MinKNOW v23.11.5 (ONT). The consensus was created from the resulting bam file using samtools v1.19.2 as described [7,8]. NextClade v3.1.0 was used for clade assignment and quality checks [9].

Sequences were aligned using MAFFT v7.520 [10], and the alignment manually curated. Phylogenetic analysis was performed using IQ-TREE v2.2.6 [11], visualised with FigTree v1.4.4 [12], and for additional confirmation of the results, phylogenetic analysis was also performed using NextStrain CLI v8.2.0 and visualised in auspic.

Identification of monkeypox viruses belonging to novel Clade I sub-lineage

At the time of writing, no near-complete sequences of MPXV circulating in DRC from the 2023 outbreak were available on (public) online databases. The targeted amplicon sequencing conducted to characterise MPXV sequences in the 10 hospitalised study patients allowed to generate near-complete MPXV sequences for six of them; all were classified as Clade I. The genome coverage ranged between 93.5%–100% (average of 95.2%) as described in Supplementary Tables 1 and 2.

Phylogenetic analysis was done including 113 near-to-complete genome reference mpox sequences from Africa available from GISAID [13], which include all sequences available on the National Center for Biotechnological Information (NCBI) on 8 February 2024 and two European sequences from the recent 2022 global mpox outbreak [14]. The six sequences from the current study grouped with published Clade I sequences, but in a distinct sub-lineage from all other Clade I sequences, suggesting that the ongoing outbreak in South Kivu results from a separate introduction (Figure 1 and Supplementary Figure 1 and 2). The six sequences have several single nucleotide polymorphism (SNP) differences between them, which suggests ongoing circulation of this outbreak strain for some time already.

The monkeypox virus outbreak strain in South Kivu lacks the target sequence used for identifying Clade I viruses

To check if the strains obtained in the current study could be detected by commonly used molecular assays to diagnose MPXV infections, their sequences were aligned to the closely related Clade I sequence EPI_ISL_13056243. This sequence matches primer and probe sequences recommended by the United States (US) Centers for Disease Control and Prevention (CDC) to diagnose MPXV [15]. The alignment was assessed using an in-house Primer Check Tool (<https://viro-science-emc.shinyapps.io/primer-check/>). While the generic primers and probe still seem to be functional with only one mutation in the reverse primer, the specific Clade I virus real-time PCR target, recommended by the US CDC, is absent in the genomes of the novel MPXV strains (Figure 2 and Supplementary Figure 3). The observed deletion is 1,114nt in size and results in the complete deletion of the OPG032 gene. The coverage of this region ranged between 76× and 941× sequence reads depending on the sample. Due to the deletion, the rapid US CDC method to identify Clade I in newly diagnosed mpox cases is most likely not reliable for detection of the novel sub-lineage identified in the current study.

Discussion

MPXV is an emerging zoonotic virus belonging to the *Poxviridae* family and the genus *Orthopoxvirus*. In the past, MPXV has been primarily detected in West and Central Africa, however, in 2022 the World Health

FIGURE 2

Sequence alignments highlighting genetic sequence features of the MPXV detected in Kamituga resulting in (A) a single nt mismatch with US CDC recommended generic reverse primer (G2R_G reverse primer) and (B) absence of the US CDC recommended clade I specific forward, reverse, and probe target location (C3L), Democratic Republic of the Congo, 2024

A. Single nt mismatch with CDC-recommended generic reverse primer

GCTATCACATAATCTGGAAGCGTA		G2R_G_Reverse_Primer	
194834A.....	194857	mpox_2L
194834A.....	194857	mpox_9L
194834A.....	194857	mpox_1L
194834A.....	194857	mpox_3L
194834A.....	194857	mpox_70
194834A.....	194857	mpox_4L

B. Absence of target sequence for CDC-recommended Clade-I-specific real-time PCR

TTGCTTTCTGTATCCAGGTAGACA		C3L_Forward_Primer	
19782	-----	19805	mpox_2L
19782	-----	19805	mpox_9L
19782	-----	19805	mpox_1L
19782	-----	19805	mpox_3L
19782	-----	19805	mpox_70
19782	-----	19805	mpox_4L

ATCAATGTATTAAACGGAGATGCC		C3L_Reverse_Primer	
19706	-----	19729	mpox_2L
19706	-----	19729	mpox_9L
19706	-----	19729	mpox_1L
19706	-----	19729	mpox_3L
19706	-----	19729	mpox_70
19706	-----	19729	mpox_4L

TCCGGTACCGGTACATTTAGCATATATGGG		C3L_Probe	
19743	-----	19772	mpox_2L
19743	-----	19772	mpox_9L
19743	-----	19772	mpox_1L
19743	-----	19772	mpox_3L
19743	-----	19772	mpox_70
19743	-----	19772	mpox_4L

US CDC: United States Centers for Disease Control and Prevention; MPXV: monkeypox virus.

The sequence matching the US CDC primers or probe annealing sites is on the top line, where each nt is highlighted in a different colour. The lines underneath the top line each represent sequences of the MPXV identified in the current study, which are aligned to the sequence on the top line. At each position of a study sequence, identity to the top sequence is represented by a dot. When a difference is a substitution, the nt present in each study sequences is indicated, when a difference is a deletion, this is represented by a dash highlighted in purple.

Visualised by primer-check tool: <https://viroscience-emc.shinyapps.io/primer-check/>.

Organization (WHO) declared a multi-country global outbreak of MPXV of Clade IIb [14,16]. This outbreak was predominantly described as affecting communities of men having sex with men (MSM), although infections were also reported in a limited number of people not engaging in sexual contact with infected individuals.

In DRC, where human infection with MPXV was first ever recognised in 1970, sporadic mpox outbreaks have been reported since, with increasing frequency over time [17]. The particular rise in the number of suspected MPXV cases recorded in the country in 2023, and their occurrence in unusual geographic areas [2] suggests a possible change in the characteristics of the virus and the epidemiology of the disease. Therefore, obtaining sequence information on the circulating strain(s) is considered crucial and MPXV whole genome Nanopore sequencing training has been provided recently to local scientists from Rwanda and DRC to perform both wet-laboratory sequencing and dry-laboratory sequence analysis.

From the mpox outbreak in Kamituga, South Kivu, six near-to-complete MPXV sequences derived from local patients hospitalised with mpox were obtained. Phylogenetic analyses of these sequences together with those available for other Clade I and II viruses, placed them in a new sub-lineage near the root of Clade I, which suggests that the outbreak in this region results from a new introduction, most likely from a zoonotic reservoir. Although sequences from a small 2023 Kinshasa outbreak are not publicly available, the placement of those sequences in a published phylogenetic tree [3] suggests that the Kamituga outbreak is not related to the outbreak in Kinshasa. Our findings therefore suggest that there are at least two independent outbreaks ongoing in DRC.

Remarkably, a large stretch of sequence in the genomes belonging to the novel MPXV sub-lineage was absent compared to other Clade I genomes, which would lead to failure of the Clade I-specific real-time PCR recommended by the US CDC [15]. A deletion in the same region is also observed in Clade II MPXV, and this was the basis for clade assignment using the CDC PCR. Therefore, if the viruses from the new lineage were to spread internationally, this molecular surveillance tool can no longer be used to rapidly identify these Clade I virus infections while the global Clade IIb outbreak is ongoing.

Multiple amplicon-based assays targeting Clade IIb MPXV have been developed and applied since the global mpox outbreak in 2022 [5,18]. In the current work, the application of a large amplicon assay which was originally designed to target Clade IIb resulted in an average genome coverage of 95.2% in the new Clade I genome, showing that this method can also be used to sequence Clade I viruses, although minor adjustments of primer pairs might result in more complete consensus sequences. Also, the consensus was

generated using a reference-based alignment using Nanopore data. Metagenomic Nanopore or Illumina sequencing might be needed in combination with de novo assembly approaches for refined consensus calling, however, this was not possible here, making this a limitation of the current work. Further studies are needed/ongoing to assess transmissibility and clinical severity associated with the new lineage.

Conclusion

Altogether, the findings of this study strongly suggest that whole genome sequencing of a larger subset of MPXV currently causing mpox cases in DRC, as well as public data sharing, are essential to understand the ongoing epidemic. Further studies, sequencing and analyses are ongoing, but in accordance with the above statement we believe that rapid public sharing of all available information is essential to help to better understand and contain the current mpox emergence.

†Authors' correction

Upon publication of the article, some affiliation numbers in the authors' list were incorrect for the following authors: Justin Bengheya Mbiribindi, Jean Pierre Musabyimana and Freddy Belesi Siangoli. Affiliation numbers were corrected at the request of the authors on 18 March 2024.

Ethical statement

Ethical clearance to conduct this study was obtained from the Ethical Review Committee of the Catholic University of Bukavu (Number UCB/CIES/NC/022/2023).

Data availability

Consensus sequences were uploaded on GISAID with the accession numbers: EPI_ISL_18886301, EPI_ISL_18886588, EPI_ISL_18886467, EPI_ISL_18886467, EPI_ISL_18886588 and EPI_ISL_18886301. Raw sequence data was uploaded to the ENA under the study ID: ERP157439.

Acknowledgements

This work was funded by the EU Horizon 2020 grants VEO (874735) and the Global Health EDCTP3 Joint Undertaking (Global Health EDCTP3) programme under grant agreement No. 101103059 (GREATLIFE). We greatly thank the University of Dalhousie, Canada for supporting the field works, and the Institut National de Recherche Biomedicale (INRB) Goma for PCR diagnostics.

We would like to thank the Provincial Division of Health (DPS) of South-Kivu and Kamituga Health Zone (KHZ) for their collaboration during the study. We greatly thank Wildlife Conservation Network (WCN) and Conservation Action Research Network (CARN) for the scholarship and research supports they awarded to the first author.

We gratefully acknowledge all data contributors, i.e., the Authors and their Originating laboratories responsible for obtaining the specimens, and their Submitting laboratories for generating the genetic sequence and metadata and sharing via the GISAID Initiative, on which this research is based.

Conflict of interest

None declared.

Authors' contributions

LMM, JCU, PN, MK, BBOM, FMA, FBS, JPM conceptualised and designed the study, LMM, MB, JCU, PN, LS, DFN, SO, FMA, BBOM, FBS contributed to data acquisition and interpretation, drafted, cross-reviewed the manuscript. JBM, JMM, LMM, NMB, TL, EBK, were involved in sample collection and investigation. All authors approved the final version.

References

1. World Health Organization (WHO). Mpox (monkeypox) - Democratic Republic of the Congo. Geneva: WHO; 23 Nov 2023. Available from: <https://www.who.int/emergencies/disease-outbreak-news/item/2023-DON493>
2. Multi-country outbreak of mpox. External situation report#30 – 25 November 2023. Geneva: World Health Organization. [Accessed 14 Feb 2024]. Available from: <https://www.who.int/publications/m/item/multi-country-outbreak-of-mpox--external-situation-report-30--25-november-2023>
3. Kibungu EM, Vakaniaki EH, Kinganda-Lusamaki E, Kalonji-Mukendi T, Pukuta E, Hoff NA, et al. , International Mpox Research Consortium. Clade I-Associated Mpox Cases Associated with Sexual Contact, the Democratic Republic of the Congo. *Emerg Infect Dis.* 2024;30(1):172-6. <https://doi.org/10.3201/eid3001.231164> PMID: 38019211
4. Murhula LM, Udahemuka JC, Ndishimye P, Sganzerla Martinez G, Kelvin P, Bubala Nadine M, et al. Epidemiology, clinical characteristics, and transmission patterns of a novel Mpox (Monkeypox) outbreak in eastern Democratic Republic of the Congo (DRC): an observational, cross-sectional cohort study. *medRxiv.* 2024.03.05.24303395. Preprint. <https://doi.org/10.1038/s41598-021-92315-8> PMID: 34158533
5. Welkers M, Jonges M, Van Den Ouden A. Monkeypox virus whole genome sequencing using combination of NextGenPCR and Oxford Nanopore. *protocols.io.* 12 Jul 2022. Available from: <https://protocols.io/view/monkeypox-virus-whole-genome-sequencing-using-comb-ccc7sszn>
6. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J.* 2011;17(1):10. <https://doi.org/10.14806/ej.17.1.200>
7. Schuele L, Boter M, Nieuwenhuijse DF, Götz H, Fanoy E, de Vries H, et al. Circulation, viral diversity and genomic rearrangement in mpox virus in the Netherlands during the 2022 outbreak and beyond. *J Med Virol.* 2024;96(1):e29397. <https://doi.org/10.1002/jmv.29397> PMID: 38235923
8. Nieuwenhuijse DF, van der Linden A, Kohl RHG, Sikkema RS, Koopmans MPG, Oude Munnink BB. Towards reliable whole genome sequencing for outbreak preparedness and response. *BMC Genomics.* 2022;23(1):569. <https://doi.org/10.1186/s12864-022-08749-5> PMID: 35945497
9. Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C, et al. Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics.* 2018;34(23):4121-3. <https://doi.org/10.1093/bioinformatics/bty407> PMID: 29790939
10. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* 2002;30(14):3059-66. <https://doi.org/10.1093/nar/gkf436> PMID: 12136088
11. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Mol Biol Evol.* 2020;37(5):1530-4. <https://doi.org/10.1093/molbev/msaa015> PMID: 32011700
12. Rambaut A. FigTree. 2016. Available from: <http://tree.bio.ed.ac.uk/software/figtree/>
13. Elbe S, Buckland-Merrett G. Data, disease and diplomacy: GISAID's innovative contribution to global health. *Glob Chall.* 2017;1(1):33-46. <https://doi.org/10.1002/gch2.1018> PMID: 31565258
14. World Health Organization (WHO). Multi-country monkeypox outbreak in non-endemic countries. Geneva: WHO; 21 May 2022. Available from: <https://www.who.int/emergencies/disease-outbreak-news/item/2022-DON385>
15. Li Y, Zhao H, Wilkins K, Hughes C, Damon IK. Real-time PCR assays for the specific detection of monkeypox virus West African and Congo Basin strain DNA. *J Virol Methods.* 2010;169(1):223-7. <https://doi.org/10.1016/j.jviromet.2010.07.012> PMID: 20643162
16. World Health Organization (WHO). Monkeypox: experts give virus variants new names. Geneva: WHO; 12 August 2022. Available from: <https://www.who.int/news/item/12-08-2022-monkeypox--experts-give-virus-variants-new-names>
17. Ladnyj ID, Ziegler P, Kima E. A human infection caused by monkeypox virus in Basankusu Territory, Democratic Republic of the Congo. *Bull World Health Organ.* 1972;46(5):593-7. PMID: 4340218
18. Brinkmann A, Pape K, Uddin S, Woelk N, Förster S, Jessen H, et al. Genome sequencing of the mpox virus 2022 outbreak with amplicon-based Oxford Nanopore MinION sequencing. *J Virol Methods.* 2024;325:114888. <https://doi.org/10.1016/j.jviromet.2024.114888> PMID: 38246565
19. Nakazawa Y, Mauldin MR, Emerson GL, Reynolds MG, Lash RR, Gao J, et al. A phylogeographic investigation of African monkeypox. *Viruses.* 2015;7(4):2168-84. <https://doi.org/10.3390/v7042168> PMID: 25912718
20. Berthet N, Descorps-Declère S, Besombes C, Curaudeau M, Nkili Meyong AA, Selekon B, et al. Genomic history of human monkey pox infections in the Central African Republic between 2001 and 2018. *Sci Rep.* 2021;11(1):13085. <https://doi.org/10.1038/s41598-021-92315-8> PMID: 34158533

License, supplementary material and copyright

This is an open-access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0) Licence. You may share and adapt the material, but must give appropriate credit to the source, provide a link to the licence and indicate if changes were made.

Any supplementary material referenced in the article can be found in the online version.

This article is copyright of the authors or their affiliated institutions, 2024.