



Citation for published version:

Aldiss, P, Betts, J, Sale, C, Pope, M, Budge, H & Symonds, ME 2018, 'Exercise-induced 'browning' of adipose tissues', *Metabolism*, vol. 81, no. 4, pp. 63-70. <https://doi.org/10.1016/j.metabol.2017.11.009>

DOI:

[10.1016/j.metabol.2017.11.009](https://doi.org/10.1016/j.metabol.2017.11.009)

Publication date:

2018

Document Version

Peer reviewed version

[Link to publication](#)

Publisher Rights

CC BY-NC-ND

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Exercise-induced 'browning' of adipose tissues

Peter Aldiss¹, James Betts², Craig Sale³, Mark Pope¹, Helen Budge¹ and Michael E Symonds^{1,4}

¹The Early Life Research Unit, Division of Child Health, Obstetrics and Gynaecology, and ⁴Nottingham Digestive Disease Centre and Biomedical Research Centre School of Medicine, University Hospital, University of Nottingham, Nottingham, UK, NG7 2UH

²Department for Health, University of Bath, Bath, BA2 7AY, UK

³Musculoskeletal Physiology Research Group, Sport, Health and Performance Enhancement Research Centre, School of Science and Technology, Nottingham Trent University, Nottingham, UK

Abbreviated title: Exercise and 'browning' of adipose tissue

Key words: Exercise, Brown adipose tissue, Browning

Corresponding author and to whom reprints should be requested:

Professor Michael E. Symonds
Academic Division of Child Health Obstetrics & Gynaecology,
School of Medicine, Queen's Medical Centre,
The University of Nottingham,
Nottingham NG7 2UH,
United Kingdom

Phone: +44 1158230625 Fax: +44 1158230626.

Email: michael.symonds@nottingham.ac.uk

Abstract

Global rates of obesity continue to rise and are necessarily the consequence of a long-term imbalance between energy intake and energy expenditure. This is the result of an expansion of adipose tissue due to both the hypertrophy of existing adipocytes and hyperplasia of adipocyte pre-cursors. Exercise elicits numerous physiological benefits on adipose tissue, which are likely to contribute to the associated cardiometabolic benefits. More recently it has been demonstrated that exercise, through a range of mechanisms, induces a phenotypic switch in adipose tissue from energy storing white adipocytes to thermogenic beige adipocytes. This has generated the hypothesis that the process of adipocyte 'browning' may partially underlie the improved cardiometabolic health in physically active populations. Interestingly, 'browning' also occurs in response to various stressors and could represent an adaptive response. In the context of exercise, it is not clear whether the appearance of beige adipocytes is metabolically beneficial or whether they occur as a transient adaptive process to exercise-induced stresses. The present review discusses the various mechanisms (*e.g.* fatty acid oxidation during exercise, decreased thermal insulation, stressors and angiogenesis) by which the exercise-induced 'browning' process may occur.

Keywords: Exercise, brown adipose tissue, browning, white adipose tissue

1. White, brown and 'beige' adipose tissue

Adipose tissue typically accounts for ~20-28% of total body mass in lean humans, with variance largely due to biological sex, and in the obese state can account for ~80% of body weight¹. It is well understood that adipose tissues vary in their function and are dependent on both the type and anatomical location of the depot. Subcutaneous adipose tissue (ScAT), located under the skin, accounts for the majority of total white adipose tissue (WAT) in humans². Visceral adipose tissue (VAT), located around the kidneys (perirenal), intestines (mesenteric and omental), vasculature (perivascular) and heart (epicardial/paracardial) normally accounts for much less but poses a far greater cardiometabolic risk following expansion³. Far from being "merely" energy storage depots, adipose tissues are potentially the largest endocrine organ in the body and, by the secretion of numerous factors, can regulate a range of physiological functions including metabolic homeostasis, appetite, angiogenesis, immunity and the cardiovascular system⁴.

Conversely, brown adipose tissue (BAT) is located primarily in the supraclavicular region but also in smaller amounts around the kidneys, vasculature and heart⁵. BAT is responsible for adaptive thermogenesis, which preserves homeostasis in response to a thermal stimulus (e.g. temperature and/or energy balance); for example by uncoupling oxidative metabolism from ATP production in favour of heat production via uncoupling protein 1 (UCP1)⁶. The re-discovery of this tissue in adult humans⁷ and its association with multiple parameters of metabolic health⁸⁻¹⁰ and cardiovascular events¹¹ has led to huge interest in the potential to activate this tissue to combat obesity-associated cardiometabolic disease. More recently it has been shown that, following a stimulus (e.g. exercise), brown-like UCP1+ cells termed 'beige' adipocytes appear interspersed in what were previously classical white depots^{12,13}. It has therefore been suggested that these beige adipocytes: a) are distinct in origin, deriving from a lineage not shared with brown or white adipocytes; b) appear due to the trans-differentiation of pre-existing white adipocytes; or c) arise from a combination of the above¹⁴. It is important to note, however, that the molecular signature of both brown and beige adipocytes

differs between small animals and humans¹⁵. In rodents, brown and beige adipocytes are anatomically distinct and clearly distinguishable whereas in humans current evidence suggests BAT is a heterogeneous depot expressing markers of both brown and beige adipocytes^{16,17}. As such, for the purpose of this review we will use the term 'browning' to describe the appearance of UCP1+ adipocytes. Species differences in molecular signature aside, the 'browning' of white adipose tissues has become an attractive therapeutic target due to mounting evidence suggesting that these 'beige' cells are metabolically active, contributing to both thermogenesis and metabolic health^{17,18}.

Exercise and physical activity play a key role in modulating many parameters of cardiometabolic health, eliciting several benefits on adipose tissues¹⁹. A reduction in adipocyte size can be seen with exercise training²⁰, although this effect seems to be sex-specific as females undergoing the same training regime as men exhibited no changes in body mass or adipocyte size²¹. Furthermore, exercise training elicits improvements in adipose tissue inflammation²², vascularity²³ and mitochondrial biogenesis²⁴, increasing the supply of oxygen and nutrients to the tissue and improving their oxidative capacity. More recently it has emerged that factors produced during exercise by skeletal muscle, adipose tissue and potentially the liver act to induce the 'browning' of WAT in an endocrine and/or paracrine manner. Of these, irisin, a PPAR γ coactivator-1 α (PGC1- α) dependent myokine²⁵, has generated the most interest as a browning agent, although there is major controversy surrounding the results and validity of this purported myokine²⁶. Numerous other factors that promote 'browning' are also altered during or following exercise, such as interleukin-6²⁷, B-aminoisobutyric acid²⁸, meteorin-like²⁹, fibroblast growth factor-21³⁰, natriuretic peptides (NP's)³¹⁻³³ and lactate³⁴, leading many to suggest that the long-lasting metabolic effects of exercise may be due to the 'browning' of white adipose tissues. However, many of these indices have only been validated in-vitro or in animal models and as such their role in 'browning' human WAT are currently unknown.

2. Regulation of adipose tissue metabolism during exercise

Despite the therapeutic potential of thermogenic 'beige' cells, it is not entirely clear why they appear during exercise, a time of heat production³⁵. It has been suggested, however, that following the lipolytic response to exercise and subsequent increase in circulating non-esterified fatty acids (NEFA) there is a need for other areas of oxidation and that the browning of white adipocytes provides these sites in order to maintain NEFA flux³⁶. Adipose tissue metabolism during exercise is regulated by multiple factors released both centrally and peripherally to modulate the rate of NEFA uptake and release. The major endocrine mechanism is increased plasma epinephrine which, acting through cyclic AMP (cAMP) and protein kinase A (PKA), phosphorylates adipose triglyceride lipase (ATGL) to stimulate lipolysis and maintain NEFA supply during exercise, a response that is abolished following the blockade of B-adrenoreceptors with propanolol^{37,38}. Other lipolytic factors include norepinephrine (NE), glucagon, cortisol and NP's³⁷. NE only contributes marginally to exercise-induced lipolysis³⁸, whilst glucagon and cortisol concentrations increase later in exercise and bind to stimulatory GTP-binding protein to activate ATGL through cAMP and PKA³⁷. NP's are increased during exercise of moderate intensity and stimulate lipolysis through the activation of cGMP-dependant protein kinase I (cGKI) and subsequent phosphorylation of perilipin 1 and hormone-sensitive lipase (HSL)³⁹. Altered concentrations of adenosine and insulin act against these signals to regulate lipolysis and prevent excess release of NEFA, which is apparent when the typical exercise-induced reduction in insulin is absent⁴⁰. Post-exercise, lipolysis stabilises but remains higher compared to rest for up to 24 hours, thus even a single bout of exercise can influence energy expenditure/balance over the next day⁴¹ and modulate insulin sensitivity for up to 48 hours⁴². Whilst exercise elicits substantial alterations in adipose tissue metabolism, whether the 'browning' that occurs following exercise training occurs during an acute bout of exercise to facilitate the oxidation of excess NEFA as postulated by Virtanen et al.³⁶ remains to be determined. Intriguingly, two recent groups have demonstrated that BAT lipolysis is not essential for cold-induced thermogenesis whereas WAT lipolysis is crucial, fuelling thermogenesis during fasting^{43,44}. It would be interesting to know, in the context of the hypothesis by Virtanen et al.³⁶, whether WAT lipolysis is also essential for the 'browning' seen following exercise.

3. Is exercise induced 'browning' a consequence of reduced adiposity?

Regular exercise can improve body composition, including reduced adiposity and/or increased muscle mass. A particularly interesting recent finding was that the phenotypic switch towards thermogenic brown cells shown following exercise in rats primarily occurred in subcutaneous depots, which was concomitant with alterations in BAT morphology (*i.e.* whitening) and a decreased thermogenic capacity⁴⁵. Exercise training reduced adiposity but the appearance of multilocular adipocytes, thermogenic genes and increase in oxidative capacity were only evident in the inguinal depot (visceral depots being resistant), suggesting depot-specificity. These inter-depot differences, also highlighted by De Jong et al.⁴⁶, are of particular interest as exercise does not induce browning in human WAT biopsies⁴⁷ and suggests a single biopsy is not an appropriate method to detect browning in humans (at least until we better understand how this process is regulated across both a single depot and between depots). Whilst challenging, future human studies should be designed to examine the response to exercise both at multiple sites in the same adipose depot and also across different depots.

The 'whitening' of BAT and reduced thermogenic capacity following exercise training was suggested to be a counter-regulatory mechanism whereby the ability of BAT to expend energy is reduced during times of increased energy expenditure⁴⁸⁻⁵⁰. However, the discovery that exercise induces browning of subcutaneous depots whilst simultaneously increasing lipid content in BAT has led to suggestions for other roles of exercise induced 'browning'. It is now postulated, in rodents, that a reduction in BAT occurs in response to regular increases in core body temperature, whilst the browning of subcutaneous WAT occurs due to a reduction in total mass and thus the degree of insulation conferred by this depot^{45,51}. Whilst this theory is consistent with multiple findings that high-fat feeding and resultant excess subcutaneous adiposity decreases UCP1 content⁵², recent evidence in mice suggests that adipose tissue does not confer any insulatory properties⁵³. However, given that rodents are typically cold-exposed throughout life⁵⁴, the 'browning' seen in subcutaneous adipose tissues following exercise could be an adaptive mechanism to increase the thermal potential of reduced

adiposity, thus offsetting any increased sensation of peripheral cold. Interestingly, recent work by Pepler and colleagues demonstrated that removal of IWAT by lipectomy did not attenuate the metabolic benefits of wheel-running in mice and that a compensatory browning was not shown in epididymal WAT despite the reduction in adiposity⁵⁵. However, compensatory browning in the face of reduced adiposity may be limited to ScAT. Had other ScAT depots, or in fact dermal AT been collected an induction of thermogenic genes may have been present. Whilst the authors note the limited clinical relevance of their model it suggests the importance of exercise-induced browning of IWAT is negligible and is of particular relevance given that humans have not been shown to exhibit 'browning' following exercise but still exhibit numerous metabolic benefits⁵⁶⁻⁵⁸. Whilst human WAT biopsies may not be the most effective way to detect 'browning' due to the limited amount of tissue relative to depot mass, exercise does reduce BAT mass and activity in athletic humans^{47,59}. This is consistent with rodent data and the notion that the thermogenic capacity of BAT may be reduced in the presence of regular increases in core body temperature. Given the inverse relationship between BAT mass and lean mass²³, it is possible that the recruitment of a large muscle mass during exercise training negates the need for BAT, given the potentially superior capacity for muscle to generate heat⁶⁰.

4. 'Browning' as an adaptive mechanism to exercise-induced stress?

There is a growing consensus that the 'browning' of adipose tissues occurs as an adaptive mechanism to certain stresses, aside from the atypical stress of a reduced ambient temperature. This stress-induced 'browning' has been shown in numerous clinical populations including those suffering burn injuries and pheochromocytoma^{13,61}. The hypermetabolic state induced by burn injury is derived from a chronic increase in circulating catecholamines and IL-6 which induce the remodelling of subcutaneous adipose tissue^{13,62}. This chronic stress increases mitochondrial mass in ScAT, expression of thermogenic markers such as UCP1 and PGC1a in addition to whole body energy expenditure⁶³. These factors that are thought to play a major role in the hypermetabolism evident in cancer patients, contributing to the cachexic state^{64,65}. Regular physical exercise induces multiple physiological

stressors such as regular chronic sympathetic activation leading to increases in catecholamine secretion and IL-6 after only 6 minutes of exercise⁶⁶, whilst 25-fold increases in IL-6 are seen during longer running protocols in humans and are maintained 2.5 hours post-exercise⁶⁷. Whether the effects of IL-6 on adipose tissue are due to its pro- or anti-inflammatory properties or due to the synergistic immune response following exercise which includes alterations in IL-1ra and IL-10⁶⁸ for instance is unknown.

Rodent studies suggest that the metabolic benefits of both BAT and the exercise-induced browning of ScAT are dependent on the presence of IL-6, with no effect in IL-6k/o mice; again, whether this is simply an effect of IL-6 or due to the effect of its absence on other factors is unclear^{69,70}. Accumulating evidence suggests that 'browning' is associated with reduced adipose tissue inflammation, so the 'browning' seen with exercise could be an adaptive response to increased inflammatory markers such as IL-6²². There are other factors indicating a role for browning as an adaptive mechanism to exercise-induced stresses. In particular, the recent finding that lactate administration induces browning, thus maintaining intracellular redox state³⁴. Whilst lactate was not reported in response to exercise in that study, both the treatment of murine and human adipocytes with lactate and its exogenous administration in-vivo induced UCP1 in a redox state dependent manner³⁴. Physical exercise is a source of free radicals, occurring primarily during strenuous or prolonged exercise and whilst these free radicals can, depending on the state of antioxidant defence, cause cellular damage at high levels they are also important signalling molecules⁷¹. In the context of exercise, the increase in free radicals could modulate changes in the intracellular redox state of WAT, with lactate being an evolutionary mechanism to modulate redox state by inducing the brown phenotype. BAT has an abundance of antioxidant enzymes, including UCP1 (which itself plays a key role in regulating ROS production)^{72,73} although, as discussed above, chronic exposure to a higher core temperature during exercise training may downregulate its activity. As such, 'browning' in other depots may be a compensatory mechanism for a reduction in BAT mitochondrial function.

Another mechanism by which “browning” could occur is through the increase in NP secreted from cardiac tissues in response to the increased stress and demands placed on the cardiac system during exercise³¹. Whilst NP induce browning in subcutaneous adipose tissues³¹, it is feasible that their prime target is the local adipose tissues around heart and vasculature that modulate both myocardial⁷⁴ and vascular⁷⁵ redox state, in order to attenuate excess oxidative stress within the heart and vasculature.

5. Does exercise-induced angiogenesis modulate the expression of beige cells?

An increase in angiogenesis and vascularisation of tissues and organs, with raised oxygenation, blood flow and nutrient delivery is one of the mechanisms through which exercise improves cardiometabolic health⁷⁶. Unlike classical BAT which shares origins with skeletal muscle, beige adipocytes were originally shown to originate from vascular smooth muscle cells and subsequently it was demonstrated that most, if not all beige adipocytes are derived from the vascular niche^{77,78}. This is particularly interesting given that browning occurs in a distinct region of the inguinal depot along the main vasculature and suggests that angiogenesis and/or the proliferation of vascular smooth muscle cells into beige adipocytes may be key to this regional browning effect⁷⁹. Further evidence to support a role of exercise-induced angiogenic mechanisms in the induction of the beige phenotypes comes from data demonstrating that adipose specific over-expression of vascular endothelial growth factor (VEGF; a key regulator of angiogenesis) reproduces the browning that occurs in an enriched environment, effects which were negated with adipose specific VEGF k/o and VEGF blockade⁸⁰. This suggests that VEGF and related angiogenic mechanisms are essential for the browning effect induced by exercise.

More recently it has been shown that expansion of the adipose vasculature by another member of the VEGF family, VEGFB (plus its receptor VEGFR1), prevents the occurrence of several aspects of obesity-induced metabolic dysfunction concomitant with increased vascularisation and thermogenic markers in WAT⁸¹. In the obese state the dysfunction of adipose tissues and the ‘whitening’ of BAT stems from the adipocytes becoming hypertrophic and outstripping the vasculature to become hypoxic^{82,83}. Thus

it is likely that the restoration of the vascular supply to these tissues can induce or restore the brown phenotype and, in the case of exercise, the induction of angiogenic genes will promote the vascular supply in WAT to induce 'browning'. Whether 'browning' is an essential end-product of this process or a simple by-product of an increased vascularisation and differentiation of pre-cursor cells from the vascular niche remains to be determined. Furthermore, it is worth noting the presence, location and relevance of the lymph node in relation to the regionalisation of 'browning' demonstrated by Barreau et al.⁷⁹.

Exercise training triggers multiple complex changes in immune function⁸⁴ and emerging evidence suggests that innate immune cells control brown and beige adipogenesis⁸⁵. Meteorin-like is produced following resistance training via the induction of the PGC1 α isoform PGC1 α 4 and induces the browning of adipose tissues by stimulating IL-4 and IL-13 secretion from eosinophils and the activation of alternate M2 macrophages²⁹. Whilst the role of catecholamine production from macrophages has recently been questioned, the use of bone-marrow derived macrophages and not tissue resident macrophages leaves such questions unanswered, particularly as the latter may play a local role in immune-adipose crosstalk. Just as UCP2 regulates immune cell metabolism, differentiation and survival, it is feasible UCP1 may play a similar role in adipose tissue^{86,87}. Finally, immune cells produced post-exercise from the lymph node may have a local effect on neighbouring adipocytes leading to the region specific 'browning' seen by others⁷⁹.

6. Summary

Exercise, through various secreted factors and mechanisms induces a 'browning' of WAT. However, the physiological explanation for this process is unclear. From an evolutionary point of view the preservation of energy stores is vital. Why then would exercise cause the appearance of thermogenic adipocytes that can increase energy expenditure? It is plausible that our lean and active ancestors naturally possessed large quantities of brown or beige adipocytes. Our subsequent transition to a modern, largely sedentary population has meant these adipocytes have undergone a 'whitening' and

the 'browning' that occurs in response to exercise is actually a transition back to their natural state. Despite a lack of understanding as to why exercise should elicit these effects on adipose tissue, it is worth noting Orgel's second law: "Evolution is cleverer than you are". Based on this rule there is likely a valid physiological explanation(s) for the 'browning' process that is yet to be discovered.

Whilst exercise induces a brown phenotype in rodent ScAT, it is not yet clear whether exercise elicits similar effects in humans though the available data suggests that, similar to rodents, exercise downregulates both the mass and activity of human BAT. Whether this is compensated for by browning of WAT or is due to the significant increases in muscle mass and therefore greater capacity for shivering thermogenesis remains to be established.

At present both rodent and human evidence is based on either sedentary or exercise-trained populations, with little attention given to the role of physical activity *per se*. Whether regular physical activity rather than exercise training potentiates any thermogenic effects on adipose tissues is unknown but, considering the effects of regular physical activity on adiposity and cardiometabolic health, is an area that merits investigation. Studies investigating regular physical activity would also not be confounded by increases in core body temperature and reductions in adiposity, so would rule out these hypothetical mechanisms.

Acknowledgements: P. Aldiss is funded by the British Heart Foundation (FS/15/4/31184)

Disclosure Statement: The authors report no relationships that could be construed as a conflict of interest.

Summary figure: Mechanisms of exercise-induced browning. Discussed in detail in-text.

References

- 1 Thompson, D., Karpe, F., Lafontan, M. & Frayn, K. Physical activity and exercise in the regulation of human adipose tissue physiology. *Physiol Rev* **92**, 157-191, doi:10.1152/physrev.00012.2011 (2012).
- 2 Thomas, E. L. *et al.* Magnetic resonance imaging of total body fat. *J Appl Physiol (1985)* **85**, 1778-1785 (1998).
- 3 Despres, J. P. Is visceral obesity the cause of the metabolic syndrome? *Ann Med* **38**, 52-63, doi:10.1080/07853890500383895 (2006).
- 4 Coelho, M., Oliveira, T. & Fernandes, R. Biochemistry of adipose tissue: an endocrine organ. *Arch Med Sci* **9**, 191-200, doi:10.5114/aoms.2013.33181 (2013).
- 5 Sacks, H. & Symonds, M. E. Anatomical locations of human brown adipose tissue: functional relevance and implications in obesity and type 2 diabetes. *Diabetes* **62**, 1783-1790, doi:10.2337/db12-1430 (2013).
- 6 Cannon, B. & Nedergaard, J. Brown adipose tissue: function and physiological significance. *Physiol Rev* **84**, 277-359, doi:10.1152/physrev.00015.2003 (2004).
- 7 Cypess, A. M. *et al.* Identification and importance of brown adipose tissue in adult humans. *N Engl J Med* **360**, 1509-1517, doi:10.1056/NEJMoa0810780 (2009).
- 8 van Marken Lichtenbelt, W. D. *et al.* Cold-activated brown adipose tissue in healthy men. *N Engl J Med* **360**, 1500-1508, doi:10.1056/NEJMoa0808718 (2009).
- 9 Matsushita, M. *et al.* Impact of brown adipose tissue on body fatness and glucose metabolism in healthy humans. *Int J Obes (Lond)* **38**, 812-817, doi:10.1038/ijo.2013.206 (2014).
- 10 Yoneshiro, T. *et al.* Age-related decrease in cold-activated brown adipose tissue and accumulation of body fat in healthy humans. *Obesity (Silver Spring)* **19**, 1755-1760, doi:10.1038/oby.2011.125 (2011).
- 11 Takx, R. *et al.* Supraclavicular Brown adipose tissue FDG uptake and cardiovascular disease. *J Nucl Med*, doi:10.2967/jnumed.115.166025 (2016).
- 12 De Matteis, R. *et al.* Exercise as a new physiological stimulus for brown adipose tissue activity. *Nutr Metab Cardiovasc Dis* **23**, 582-590, doi:10.1016/j.numecd.2012.01.013 (2013).
- 13 Sidossis, L. S. *et al.* Browning of Subcutaneous White Adipose Tissue in Humans after Severe Adrenergic Stress. *Cell Metab* **22**, 219-227, doi:10.1016/j.cmet.2015.06.022 (2015).
- 14 Sanchez-Gurmaches, J. & Guertin, D. A. Adipocyte lineages: tracing back the origins of fat. *Biochim Biophys Acta* **1842**, 340-351, doi:10.1016/j.bbadis.2013.05.027 (2014).
- 15 Liu, X., Cervantes, C. & Liu, F. Common and distinct regulation of human and mouse brown and beige adipose tissues: a promising therapeutic target for obesity. *Protein Cell* **8**, 446-454, doi:10.1007/s13238-017-0378-6 (2017).
- 16 Jespersen, N. Z. *et al.* A classical brown adipose tissue mRNA signature partly overlaps with brite in the supraclavicular region of adult humans. *Cell Metab* **17**, 798-805, doi:10.1016/j.cmet.2013.04.011 (2013).
- 17 Wu, J. *et al.* Beige adipocytes are a distinct type of thermogenic fat cell in mouse and human. *Cell* **150**, 366-376, doi:10.1016/j.cell.2012.05.016 (2012).
- 18 Shabalina, I. G. *et al.* UCP1 in brite/beige adipose tissue mitochondria is functionally thermogenic. *Cell Rep* **5**, 1196-1203, doi:10.1016/j.celrep.2013.10.044 (2013).
- 19 Joyner, M. J. & Green, D. J. Exercise protects the cardiovascular system: effects beyond traditional risk factors. *J Physiol* **587**, 5551-5558, doi:10.1113/jphysiol.2009.179432 (2009).
- 20 Despres, J. P., Bouchard, C., Savard, R., Tremblay, A. & Allard, C. Lack of relationship between changes in adiposity and plasma lipids following endurance training. *Atherosclerosis* **54**, 135-143 (1985).
- 21 Despres, J. P. *et al.* The effect of a 20-week endurance training program on adipose-tissue morphology and lipolysis in men and women. *Metabolism* **33**, 235-239 (1984).

- 22 Haczeyni, F. *et al.* Exercise improves adipose function and inflammation and ameliorates fatty liver disease in obese diabetic mice. *Obesity (Silver Spring)* **23**, 1845-1855, doi:10.1002/oby.21170 (2015).
- 23 Disanzo, B. L. & You, T. Effects of exercise training on indicators of adipose tissue angiogenesis and hypoxia in obese rats. *Metabolism* **63**, 452-455, doi:10.1016/j.metabol.2013.12.004 (2014).
- 24 Vieira, V. J. & Valentine, R. J. Mitochondrial biogenesis in adipose tissue: can exercise make fat cells 'fit'? *J Physiol* **587**, 3427-3428, doi:10.1113/jphysiol.2009.175307 (2009).
- 25 Bostrom, P. *et al.* A PGC1-alpha-dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature* **481**, 463-468, doi:10.1038/nature10777 (2012).
- 26 Albrecht, E. *et al.* Irisin - a myth rather than an exercise-inducible myokine. *Sci Rep* **5**, 8889, doi:10.1038/srep08889 (2015).
- 27 Ma, Y., Gao, M., Sun, H. & Liu, D. Interleukin-6 gene transfer reverses body weight gain and fatty liver in obese mice. *Biochim Biophys Acta* **1852**, 1001-1011, doi:10.1016/j.bbadis.2015.01.017 (2015).
- 28 Kammoun, H. L. & Febbraio, M. A. Come on BAIBA light my fire. *Cell Metab* **19**, 1-2, doi:10.1016/j.cmet.2013.12.007 (2014).
- 29 Rao, R. R. *et al.* Meteorin-like is a hormone that regulates immune-adipose interactions to increase beige fat thermogenesis. *Cell* **157**, 1279-1291, doi:10.1016/j.cell.2014.03.065 (2014).
- 30 Giralt, M., Gavalda-Navarro, A. & Villarroya, F. Fibroblast growth factor-21, energy balance and obesity. *Mol Cell Endocrinol* **418 Pt 1**, 66-73, doi:10.1016/j.mce.2015.09.018 (2015).
- 31 Palmer, B. F. & Clegg, D. J. An Emerging Role of Natriuretic Peptides: Igniting the Fat Furnace to Fuel and Warm the Heart. *Mayo Clin Proc* **90**, 1666-1678, doi:10.1016/j.mayocp.2015.08.006 (2015).
- 32 Bordbar, S., Bigi, M. A., Aslani, A., Rahimi, E. & Ahmadi, N. Effect of endurance and strength exercise on release of brain natriuretic peptide. *J Cardiovasc Dis Res* **3**, 22-25, doi:10.4103/0975-3583.91599 (2012).
- 33 Follenius, M. & Brandenberger, G. Increase in atrial natriuretic peptide in response to physical exercise. *Eur J Appl Physiol Occup Physiol* **57**, 159-162 (1988).
- 34 Carriere, A. *et al.* Browning of white adipose cells by intermediate metabolites: an adaptive mechanism to alleviate redox pressure. *Diabetes* **63**, 3253-3265, doi:10.2337/db13-1885 (2014).
- 35 Gagnon, D., Jay, O., Lemire, B. & Kenny, G. P. Sex-related differences in evaporative heat loss: the importance of metabolic heat production. *Eur J Appl Physiol* **104**, 821-829, doi:10.1007/s00421-008-0837-0 (2008).
- 36 Virtanen, K. A. BAT thermogenesis: Linking shivering to exercise. *Cell Metab* **19**, 352-354, doi:10.1016/j.cmet.2014.02.013 (2014).
- 37 Tsiloulis, T. & Watt, M. J. Exercise and the Regulation of Adipose Tissue Metabolism. *Prog Mol Biol Transl Sci* **135**, 175-201, doi:10.1016/bs.pmbts.2015.06.016 (2015).
- 38 Arner, P., Kriegholm, E., Engfeldt, P. & Bolinder, J. Adrenergic regulation of lipolysis in situ at rest and during exercise. *J Clin Invest* **85**, 893-898, doi:10.1172/JCI114516 (1990).
- 39 Sengenès, C. *et al.* Involvement of a cGMP-dependent pathway in the natriuretic peptide-mediated hormone-sensitive lipase phosphorylation in human adipocytes. *J Biol Chem* **278**, 48617-48626, doi:10.1074/jbc.M303713200 (2003).
- 40 Marker, J. C. *et al.* Catecholamines in prevention of hypoglycemia during exercise in humans. *Am J Physiol* **260**, E705-712 (1991).
- 41 Magkos, F., Mohammed, B. S., Patterson, B. W. & Mittendorfer, B. Free fatty acid kinetics in the late phase of postexercise recovery: importance of resting fatty acid metabolism and exercise-induced energy deficit. *Metabolism* **58**, 1248-1255, doi:10.1016/j.metabol.2009.03.023 (2009).

- 42 Bogardus, C. *et al.* Effect of muscle glycogen depletion on in vivo insulin action in man. *J Clin Invest* **72**, 1605-1610, doi:10.1172/JCI111119 (1983).
- 43 Schreiber, R. *et al.* Cold-Induced Thermogenesis Depends on ATGL-Mediated Lipolysis in Cardiac Muscle, but Not Brown Adipose Tissue. *Cell Metab*, doi:10.1016/j.cmet.2017.09.004 (2017).
- 44 Shin, H. *et al.* Lipolysis in Brown Adipocytes Is Not Essential for Cold-Induced Thermogenesis in Mice. *Cell Metab*, doi:10.1016/j.cmet.2017.09.002 (2017).
- 45 Wu, M. V., Bikopoulos, G., Hung, S. & Cедdia, R. B. Thermogenic capacity is antagonistically regulated in classical brown and white subcutaneous fat depots by high fat diet and endurance training in rats: impact on whole-body energy expenditure. *J Biol Chem* **289**, 34129-34140, doi:10.1074/jbc.M114.591008 (2014).
- 46 de Jong, J. M., Larsson, O., Cannon, B. & Nedergaard, J. A stringent validation of mouse adipose tissue identity markers. *Am J Physiol Endocrinol Metab* **308**, E1085-1105, doi:10.1152/ajpendo.00023.2015 (2015).
- 47 Vosselman, M. J. *et al.* Low brown adipose tissue activity in endurance-trained compared with lean sedentary men. *Int J Obes (Lond)*, doi:10.1038/ijo.2015.130 (2015).
- 48 Nozu, T., Kikuchi, K., Ogawa, K. & Kuroshima, A. Effects of running training on in vitro brown adipose tissue thermogenesis in rats. *Int J Biometeorol* **36**, 88-92 (1992).
- 49 Larue-Achagiotis, C., Rieth, N., Gubern, M., Laury, M. C. & Louis-Sylvestre, J. Exercise-training reduces BAT thermogenesis in rats. *Physiol Behav* **57**, 1013-1017 (1995).
- 50 Yamashita, H. *et al.* Effect of running training on uncoupling protein mRNA expression in rat brown adipose tissue. *Int J Biometeorol* **37**, 61-64 (1993).
- 51 Sepa-Kishi, D. M. & Cедdia, R. B. Exercise-Mediated Effects on White and Brown Adipose Tissue Plasticity and Metabolism. *Exerc Sport Sci Rev* **44**, 37-44, doi:10.1249/JES.0000000000000068 (2016).
- 52 Fromme, T. & Klingenspor, M. Uncoupling protein 1 expression and high-fat diets. *Am J Physiol Regul Integr Comp Physiol* **300**, R1-8, doi:10.1152/ajpregu.00411.2010 (2011).
- 53 Fischer, A. W., Csikasz, R., von Essen, G., Cannon, B. & Nedergaard, J. No insulating effect of obesity. *Am J Physiol Endocrinol Metab*, ajpendo 00093 02016, doi:10.1152/ajpendo.00093.2016 (2016).
- 54 Maloney, S. K., Fuller, A., Mitchell, D., Gordon, C. & Overton, J. M. Translating animal model research: does it matter that our rodents are cold? *Physiology (Bethesda)* **29**, 413-420, doi:10.1152/physiol.00029.2014 (2014).
- 55 Peppler, W. T., Townsend, L. K., Knuth, C. M., Foster, M. T. & Wright, D. C. Subcutaneous inguinal white adipose tissue is responsive to, but dispensable for, the metabolic health benefits of exercise. *Am J Physiol Endocrinol Metab*, ajpendo 00226 02017, doi:10.1152/ajpendo.00226.2017 (2017).
- 56 Nakhuda, A. *et al.* Biomarkers of browning of white adipose tissue and their regulation during exercise- and diet-induced weight loss. *Am J Clin Nutr* **104**, 557-565, doi:10.3945/ajcn.116.132563 (2016).
- 57 Ronn, T. *et al.* Extensive changes in the transcriptional profile of human adipose tissue including genes involved in oxidative phosphorylation after a 6-month exercise intervention. *Acta Physiol (Oxf)* **211**, 188-200, doi:10.1111/apha.12247 (2014).
- 58 Scheele, C. Adipose adaptation to exercise training -increased metabolic rate but no signs of browning. *Acta Physiol (Oxf)* **211**, 11-12, doi:10.1111/apha.12280 (2014).
- 59 Singhal, V. *et al.* Effect of Chronic Athletic Activity on Brown Fat in Young Women. *PLoS One* **11**, e0156353, doi:10.1371/journal.pone.0156353 (2016).
- 60 M, U. D. *et al.* Human brown adipose tissue [O]O PET imaging in the presence and absence of cold stimulus. *Eur J Nucl Med Mol Imaging*, doi:10.1007/s00259-016-3364-y (2016).

- 61 Vergnes, L. *et al.* Adipocyte Browning and Higher Mitochondrial Function in Periadrenal But Not SC Fat in Pheochromocytoma. *J Clin Endocrinol Metab* **101**, 4440-4448, doi:10.1210/jc.2016-2670 (2016).
- 62 Patsouris, D. *et al.* Burn Induces Browning of the Subcutaneous White Adipose Tissue in Mice and Humans. *Cell Rep* **13**, 1538-1544, doi:10.1016/j.celrep.2015.10.028 (2015).
- 63 Razzoli, M. *et al.* Stress-induced activation of brown adipose tissue prevents obesity in conditions of low adaptive thermogenesis. *Mol Metab* **5**, 19-33, doi:10.1016/j.molmet.2015.10.005 (2016).
- 64 Hetzler, K. L. *et al.* Sex differences in the relationship of IL-6 signaling to cancer cachexia progression. *Biochim Biophys Acta* **1852**, 816-825, doi:10.1016/j.bbadis.2014.12.015 (2015).
- 65 Tsoli, M., Swarbrick, M. M. & Robertson, G. R. Lipolytic and thermogenic depletion of adipose tissue in cancer cachexia. *Semin Cell Dev Biol*, doi:10.1016/j.semcdb.2015.10.039 (2015).
- 66 Nielsen, H. B., Secher, N. H., Christensen, N. J. & Pedersen, B. K. Lymphocytes and NK cell activity during repeated bouts of maximal exercise. *Am J Physiol* **271**, R222-227 (1996).
- 67 Ostrowski, K. *et al.* A trauma-like elevation of plasma cytokines in humans in response to treadmill running. *J Physiol* **513 (Pt 3)**, 889-894 (1998).
- 68 Scott, J. P. *et al.* Effect of exercise intensity on the cytokine response to an acute bout of running. *Med Sci Sports Exerc* **43**, 2297-2306, doi:10.1249/MSS.0b013e31822113a9 (2011).
- 69 Knudsen, J. G. *et al.* Role of IL-6 in exercise training- and cold-induced UCP1 expression in subcutaneous white adipose tissue. *PLoS One* **9**, e84910, doi:10.1371/journal.pone.0084910 (2014).
- 70 Stanford, K. I. *et al.* Brown adipose tissue regulates glucose homeostasis and insulin sensitivity. *J Clin Invest* **123**, 215-223, doi:10.1172/JCI62308 (2013).
- 71 Powers, S. K. & Jackson, M. J. Exercise-induced oxidative stress: cellular mechanisms and impact on muscle force production. *Physiol Rev* **88**, 1243-1276, doi:10.1152/physrev.00031.2007 (2008).
- 72 Oelkrug, R., Goetze, N., Meyer, C. W. & Jastroch, M. Antioxidant properties of UCP1 are evolutionarily conserved in mammals and buffer mitochondrial reactive oxygen species. *Free Radic Biol Med* **77**, 210-216, doi:10.1016/j.freeradbiomed.2014.09.004 (2014).
- 73 Kazak, L. *et al.* UCP1 deficiency causes brown fat respiratory chain depletion and sensitizes mitochondria to calcium overload-induced dysfunction. *Proc Natl Acad Sci U S A*, doi:10.1073/pnas.1705406114 (2017).
- 74 Antonopoulos, A. S. *et al.* Mutual Regulation of Epicardial Adipose Tissue and Myocardial Redox State by PPAR-gamma/Adiponectin Signalling. *Circ Res* **118**, 842-855, doi:10.1161/CIRCRESAHA.115.307856 (2016).
- 75 Margaritis, M. *et al.* Interactions between Vascular Wall and Perivascular Adipose Tissue Reveal Novel Roles for Adiponectin in the Regulation of eNOS Function in Human Vessels. *Circulation*, doi:10.1161/circulationaha.112.001133 (2013).
- 76 Bloor, C. M. Angiogenesis during exercise and training. *Angiogenesis* **8**, 263-271, doi:10.1007/s10456-005-9013-x (2005).
- 77 Berry, D. C., Jiang, Y. & Graff, J. M. Mouse strains to study cold-inducible beige progenitors and beige adipocyte formation and function. *Nat Commun* **7**, 10184, doi:10.1038/ncomms10184 (2016).
- 78 Long, J. Z. *et al.* A smooth muscle-like origin for beige adipocytes. *Cell Metab* **19**, 810-820, doi:10.1016/j.cmet.2014.03.025 (2014).
- 79 Barreau, C. *et al.* Regionalization of browning revealed by whole subcutaneous adipose tissue imaging. *Obesity (Silver Spring)* **24**, 1081-1089, doi:10.1002/oby.21455 (2016).
- 80 During, M. J. *et al.* Adipose VEGF Links the White-to-Brown Fat Switch With Environmental, Genetic, and Pharmacological Stimuli in Male Mice. *Endocrinology* **156**, 2059-2073, doi:10.1210/en.2014-1905 (2015).

- 81 Robciuc, M. R. *et al.* VEGFB/VEGFR1-Induced Expansion of Adipose Vasculature Counteracts Obesity and Related Metabolic Complications. *Cell Metab* **23**, 712-724, doi:10.1016/j.cmet.2016.03.004 (2016).
- 82 Shimizu, I. *et al.* Vascular rarefaction mediates whitening of brown fat in obesity. *J Clin Invest* **124**, 2099-2112, doi:10.1172/JCI71643 (2014).
- 83 Trayhurn, P. & Alomar, S. Y. Oxygen deprivation and the cellular response to hypoxia in adipocytes - perspectives on white and brown adipose tissues in obesity. *Front Endocrinol (Lausanne)* **6**, 19, doi:10.3389/fendo.2015.00019 (2015).
- 84 Woods, J. A., Vieira, V. J. & Keylock, K. T. Exercise, inflammation, and innate immunity. *Immunol Allergy Clin North Am* **29**, 381-393, doi:10.1016/j.iac.2009.02.011 (2009).
- 85 DiSpirito, J. R. & Mathis, D. Immunological contributions to adipose tissue homeostasis. *Semin Immunol*, doi:10.1016/j.smim.2015.10.005 (2015).
- 86 Chaudhuri, L., Srivastava, R. K., Kos, F. & Shrikant, P. A. Uncoupling protein 2 regulates metabolic reprogramming and fate of antigen-stimulated CD8+ T cells. *Cancer Immunol Immunother* **65**, 869-874, doi:10.1007/s00262-016-1851-4 (2016).
- 87 Emre, Y. & Nubel, T. Uncoupling protein UCP2: when mitochondrial activity meets immunity. *FEBS Lett* **584**, 1437-1442, doi:10.1016/j.febslet.2010.03.014 (2010).

