



Citation for published version:

Hill, M, Wilson, A, Dinoi, C, Mahon, M, Maron, L & Richards, E 2020, 'Calcium Hydride Reduction of Polycyclic Aromatic Hydrocarbons', *Angewandte Chemie International Edition*, vol. 59, no. 3, pp. 1232-1237.
<https://doi.org/10.1002/anie.201913895>

DOI:

[10.1002/anie.201913895](https://doi.org/10.1002/anie.201913895)

Publication date:

2020

Document Version

Peer reviewed version

[Link to publication](#)

This is the peer reviewed version of the following article: A. S. S. Wilson, C. Dinoi, M. S. Hill, M. F. Mahon, L. Maron, E. Richards, *Angew. Chem. Int. Ed.* 2020, 59, 1232. , which has been published in final form at <https://doi.org/10.1002/anie.201913895>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Calcium Hydride Reduction of Polycyclic Aromatic Hydrocarbons

Andrew S. S. Wilson,^[a] Chiara Dinoi,^[b] Michael S. Hill,^{[a]*} Mary F. Mahon,^[a] Laurent Maron^{[b]*} and Emma Richards^[c]

Hydride derivatives have provided a focal point for many recent advances in molecular alkaline earth (Ae = Mg, Ca, Sr and Ba) chemistry.^[1-3] Instigated by Harder and co-workers' report of the calcium hydride, [(^{Dipp}BDI)Ca(THF)H]₂ (**1**, ^{Dipp}BDI = HC((Me)CN-2,6-*i*-Pr₂C₆H₃)₂),^[4,2] β -diketiminate derivatives have been especially prominent. The synthesis of a solvent-free analogue of **1** (**2**, see Scheme 1) highlights how a reduction in coordinative saturation at the calcium center can impact significantly on the observed chemistry.^[5] Compound **2** reacts with a wide range of unactivated terminal alkenes,^[4,6] providing *n*-alkyl derivatives, which are sufficiently reactive to effect the heterolytic cleavage of H₂ or even the direct nucleophilic displacement of hydride from benzene.^[5,6] Subsequent use of a bulkier β -diketiminate ligand in the synthesis of [(^{DIPeP}BDI)SrH]₂ (**3**; ^{DIPeP}BDI = HC((Me)CN-2,6-(Et₂CH)₂C₆H₃)₂) has shown that analogous reactivity may be achieved at strontium.^[3]

Computational analysis using density functional theory (DFT) indicated reactions of **2** with C=C multiple bonds are highly polarized and proceed in a stepwise fashion with no necessary rupture of either the dinuclear hydride or the dinuclear hydrido-alkyl intermediates formed during the reactions. The dimeric structure of **2** is particularly significant during its reaction with 1,3,5,7-cyclooctatetraene (COT) to yield the inverse sandwich dicalcium cyclooctatetraenyl [COT]²⁻ derivative (**4**, Scheme 1).^[7] Although the two electron reduction of COT is typically achieved by its stepwise reaction with a suitable alkali metal,^[8] monitoring of the formation of **4** by EPR spectroscopy did not provide any evidence of single electron transfer or radicaloid intermediates. Rather, DFT analysis indicated that reduction proceeds via a pathway involving consecutive polarized Ca-H/C=C insertion and intramolecular deprotonation, which is imposed by the retention of dicalcium species throughout the reaction.



Scheme 1: Sequential Ca-H/C=C insertion and deprotonation pathway to compound **4**.

[a] Dr Andrew S. S. Wilson, Dr Mary F. Mahon and Prof. Michael S. Hill
Department of Chemistry
University of Bath
Bath, BA2 7AY, UK.
E-mail: msh27@bath.ac.uk

[b] Prof. Laurent Maron, Dr Chiara Dinoi
Université de Toulouse et CNRS, INSA, UPS, UMR 5215, LPCNO,
135 Avenue de Rangueil,
F-31077 Toulouse, France
E-mail: laurent.maron@irsamc.ups-tlse.fr

[c] Dr Emma Richards School of Chemistry, Cardiff University, Park
Place, Cardiff CF10 3AT, UK.


This reactivity raises questions about the broader reductive potential of soluble molecular compounds such as **2** and **3**. Although the elemental Ae metals have been employed as electron sources (E^0 = Mg, -2.36; Ca, -2.76; Sr, -2.90; Ba, -2.91 V vs SCE), more apposite examples of *molecular* group 2 reductants are provided by Jones and Stasch's dinuclear Mg(I) species such as [(^{Dipp}BDI)Mg]₂ (**5**), and [(^{Mes}BDI)Mg]₂ (**6**) (Scheme 2), which have been posited as 'quasi-universal' reducing agents.^[9] In contrast to the heterogeneous nature of any 'dissolving metal' reduction, electron transfer from the hydrocarbon-soluble compounds **5** and **6** is facilitated by the relatively diffuse nature of the Mg-Mg-centered HOMO.

The stepwise reduction of polycyclic aromatic hydrocarbons (PAHs) such as naphthalene and anthracene has been studied since the mid-19th century.^[10] Unlike the 10 π electron aromatization of COT to [COT]²⁻, reduction involves disruption of the aromatic system and population of the relevant antibonding π^* LUMO. The total free energy of the system may be estimated from the PAH reduction potentials which increase, that is become less negative, with the number of constituent C₆ carbocycles. With regard to group 2, the reaction of anthracene (E^0 = -1.98 and -2.44 V vs SCE) and magnesium is particularly well studied and, like the corresponding reactions with calcium, strontium and barium, can access the products of both single and two-electron reduction.^[11,12] Reflecting the electrochemical data, however, only lithium among the elements of group 1 and 2 can effect twofold reduction of naphthalene (E^0 = ca. -3.1 V) to the [C₁₀H₈]²⁻ dianion,^[13] whereas liquid ammonia solutions of naphthalene and Ae metals yield only single electron reduction to produce the naphthalenide radical anion.^[12,14]

Two electron reduction of COT (E^0 = -1.83 and -1.99 V vs SCE) by **5**, occurs readily to deliver the magnesium analogue of **4**, compound **7** (Scheme 2). Treatment of anthracene by compound

6 also provides two-electron reduction and the formation of compound **8**. While the electrochemical reduction potentials of **5** and **6** have never been determined, a very recent report of the six electron reduction of C₆₀ by **6** indicates a reducing capability which is at least in the

order of E^0 = -2.9 V (vs. SCE).^[15] Consistent with this upper limit, however, such Mg(I) species are indefinitely stable in benzene (E^0 = ca. -3.4 V) and no evidence of reactivity with naphthalene (E^0 = ca. -3.1 V) has been reported. Although any operant mechanisms are likely to be very different from the more direct electron transfer implicated in the formation of compounds **7** and **8** (*vide infra*), these observations highlight that the reactivity of **2** with a range of PAHs should provide a semi-quantitative means with which to assess its reducing potential.



Scheme 2: Two electron reduction of COT and anthracene by β -diketiminato Mg(I) derivatives.

Our study commenced with a consideration of the reactivity of **2** with the parent monocyclic arene, benzene. Reminiscent of the behaviour of the group 3 and 4f hydride derivatives, $[\text{Cp}^*_2\text{MH}]_n$ ($n = 1$, $M = \text{Sc}$; $n = 2$, $M = \text{Y}$, La , Sm , Lu),^[16] Harder has reported that moderate heating (60 °C) of a C_6D_6 solution of the molecular strontium hydride (**3**) provides complete H–D exchange with the deuterated solvent to provide $[(^{\text{D}}\text{ippPBDI})\text{SrD}]_2$ and $\text{C}_6\text{D}_5\text{H}$ within 12 hours.^[3] Analogous thermal treatment of a C_6D_6 solution of compound **2** resulted, primarily, in Schlenk-type equilibration to insoluble CaH_2 and unreactive $[(^{\text{D}}\text{ippPBDI})_2\text{Ca}]$.^[17] Monitoring at room temperature by ^1H NMR spectroscopy, however, provided evidence for similar H–D exchange. After three weeks, albeit accompanied by redistribution to ca. 20% $[(^{\text{D}}\text{ippPBDI})_2\text{Ca}]$, the initial 1:1 ratio of intensities of the $^{\text{D}}\text{ippPBDI}$ γ -methine signal and Ca–H signals of **2** at δ 4.83 and δ 4.27 ppm, respectively, changed to approximately 2:1 (Figure S16). During this time period, the singlet hydride signal was transformed into an apparent unresolved 1:2:1 triplet, an observation that we attribute to sequential deuterium ($I = 1$) incorporation and the formation of the previously identified dicalcium hydride-deuteride, **2-d**.^[5] Although mechanistic and kinetic studies of this process were precluded by the aforementioned solution lability, an assessment of a benzene solution of **2** by EPR spectroscopy provided no

evidence of radical formation, even at more elevated temperatures. DFT calculations were, therefore, carried out to provide insight into the mechanism of the H–D exchange reaction of benzene. A reaction enthalpy profile (Figure 1) was determined at room temperature using the B3PW91 functional that has proven its ability to describe group 2 reactivity.^[5,6] Consistent with the slow transformation observed, this process requires the assembly of a number of higher energy intermediates. As in previous studies,^[3,5] the reaction is initiated by nucleophilic attack of a hydride of the dimeric complex **2** at the benzene ring. The resultant four-coordinate carbon center disrupts the π electron delocalization in the benzene ring ensuring that the associated barrier is rather high (35.5 kcal mol⁻¹). The system is able to evolve from this transition state, however, to a dicalcium benzenyl intermediate, **C(benzene)**, that retains a single Ca– μ_2 -H–Ca linkage. While **C(benzene)** is also high in energy (+20.6 kcal mol⁻¹), it is sufficiently stabilized by an interaction between the remaining π density and the calcium centers to allow a facile 90° rotation of the benzenyl ligand to yield a further intermediate, **D(benzene)**. Although this process incurs the loss of the benzenyl π -interaction, its replacement by a C–H interaction with one Ca center facilitates delivery of a benzene hydrogen atom back to calcium. Although the barrier associated with this latter process, **TS-DE(benzene)**, is only 7.8 kcal mol⁻¹ and the overall H/D exchange is effectively athermic, the high energies associated with **C(benzene)** and **D(benzene)** indicate that such processes are unlikely to exert any significant impact during reactions of **2** with more reactive or readily reducible PAH substrates.




Figure 1: Computed enthalpy profile at room temperature for the H/D exchange reaction catalyzed by compound **2**. Gibbs Free energies are given in parenthesis.




Figure 2: ORTEP representation (30% probability ellipsoids) of compound **9**. Hydrogen atoms, except for H1, and *iso*-propyl groups have been removed for clarity. Selected bond lengths (Å) and angles (°): Ca1-N1 2.3327(18), Ca1-N2 2.3735(16), Ca1-C88 2.8526(19), Ca1-C93 2.769(2), Ca1-C94 2.681(2), Ca1-C95 2.701(2), Ca1-C96 2.792(2), Ca1-C97 2.887(2), Ca2-N3 2.3859(16), Ca2-N4 2.4057(18), Ca2-C89 2.962(2), Ca2-C90 2.734(2), Ca2-C91 2.718(2), Ca2-C92 2.937(2), Ca3-N5 2.3744(17), Ca3-N6 2.4176(16), Ca3-C94 2.981(2), Ca3-C95 2.770(2), Ca3-C96 2.770(2), Ca3-C97 2.947(2), N1-Ca1-N2 75.09(6), N3-Ca2-N4 79.33(6), N5-Ca3-N6 77.54(6).

A reaction performed at room temperature between compound **2** and naphthalene in C_6D_6 provided the slow development of a deep blue solution over the course of 60 days. Appraisal by 1H NMR spectroscopy throughout this time period indicated consumption of *ca.* 50% of compound **2** and the production of a new compound (**9**). This latter species comprised two differentiated ^{Dipp}BDI ligand environments, which were most clearly characterized by a pair of γ -methine resonances at δ 5.46 and 5.10 ppm and which emerged in a strict 1:2 ratio of intensities throughout the course of the reaction. Compound **9** also displayed a characteristic (1H) singlet signal at δ 9.05 ppm, which was subsequently assigned as a $[Ca-\mu_2-H-Ca]$ bridging hydride environment (*vide infra*). While further experiments demonstrated that the reaction time under these conditions could be reduced to one week through the introduction of a ten-fold excess of naphthalene, the greatest improvements were obtained by performance of the reaction in the absence of any solvent. Simply heating compound **2** and ten equivalents of naphthalene for 1 hour at 80 °C, after removal of excess naphthalene by sublimation, yielded a blue powder, which was observed to comprise predominantly of compound **9** by 1H NMR spectroscopy. Crystallization of compound **9** from a saturated toluene/hexane

solution provided blue single crystals suitable for X-ray diffraction analysis, the results of which are shown in Figure 2.

The solid-state structure of compound **9** contains three unique ^{Dipp}BDI -ligated calcium environments. Ca1 interacts via a η^6 -interaction with one C_6 ring of a planar naphthalenide dianion. The resultant Ca1...C contacts [range 2.681(2) - 2.887(2) Å] are intermediate between the very short distances observed in Westerhausen's unique Ca(I) species, $[(THF)_3Ca(\mu-C_6H_3-1,3,5-Ph_3)Ca(THF)_3]$ [2.586(3), 2.598(3) Å],^[18] and those of the recently reported charge-separated cation in, $[(^{Dipp}BDI)Ca(C_6H_6)]^+$ $[Al\{OC(CF_3)_3\}_4]^-$ [2.932(5), 2.935(5) Å], wherein a molecule of benzene is bound in a η^6 fashion to the calcium center but without any necessary level of reduction. The remaining two $[(^{Dipp}BDI)Ca]$ units bind to the obverse face of the arene dianion through what are best described as η^4 -interactions with Ca-C distances in the range 2.770(2) – 2.981(2) Å to each of the two fused C_6 carbocycles, while overall charge balance is maintained by a hydride anion which bridges Ca2 and Ca3.

Reflecting their more accessible reduction potentials, reactions of the tri- and tetracyclic PAHs, anthracene and tetracene with compound **2** proceeded more readily in toluene to provide compounds **10** and **11**, respectively. Reminiscent of the data provided by **9**, the 1H NMR spectrum of compound **10** was consistent with a structure comprising a 1:2 ratio of two dissimilar ^{Dipp}BDI environments along with a single residual Ca-H proton environment, which manifested as a 1H singlet resonating at δ 5.12 ppm. In contrast, the NMR data arising from compound **11** were consistent with the presence of two identical $\{(^{Dipp}BDI)Ca\}$ units, symmetrically disposed about a reduced tetracene dianion. The veracity of these deductions was confirmed by the isolation of purple (**10**) and blue (**11**) single crystals of both compounds suitable for X-ray diffraction analysis from toluene and benzene, respectively (Figure 3).

The gross features of compound **10** (Figure 3a), a trimetallic species featuring formally monocationic $[(^{Dipp}BDI)Ca]^+$ and $[(^{Dipp}BDI)Ca-\mu-H-Ca(^{Dipp}BDI)]^+$ units residing on opposite faces of the $[C_{14}H_{10}]^{2-}$ dianion, are reminiscent of the structure of **9**. In the case of **10**, however, the Ca1-containing monometallic moiety interacts with the central C_6 unit whereas both of the hydride-bridged calcium centers are bound with an approximate η^6 disposition to each of the peripheral anthracenyl rings. In common with previously reported alkali metal derivatives of the $[C_{14}H_{10}]^{2-}$ dianion,^[19] the formally antiaromatic nature of the reduced PAH is clearly apparent from a pronounced 'hinging' of the tricyclic structure at the [C88-C89-C94-C95-C96-C101] ring, such that the mean planes defined by the peripheral C_6 -rings intersect to subtend a dihedral angle of 26.5°.

Compound **11** (Figure 3b) presents a contrasting and simpler centrosymmetric structure, whereby two identical $[(^{Dipp}BDI)Ca]$ units interact with the opposing faces of the interior (C30-C34- and C30'-C34'-containing) C_6 -rings of the tetracyclic $[C_{18}H_{12}]^{2-}$ dianion.




Figure 3: ORTEP representations (30% probability ellipsoids) of (a) compound **10** and (b) compound **11**. Hydrogen atoms, except H1 of compound **10**, *iso*-propyl groups and occluded solvent are removed for clarity. Selected bond lengths (Å) and angles (°): (**10**) Ca1-N1 2.3204(17), Ca1-N2 2.3411(17), Ca1-C88 2.651(2), Ca1-C89 2.923(2), Ca1-C94 2.838(2), Ca1-C95 2.624(2), Ca1-C96 3.019(2), Ca2-N3 2.3492(17), Ca2-N4 2.4179(18), Ca2-C96 3.071(2), Ca2-C97 2.981(2), Ca2-C98 2.858(2), Ca2-C99 2.748(2), Ca2-C100 2.712(2), Ca2-C101 2.868(2), Ca3-N5 2.3902(18), Ca3-N6 2.4405(19), Ca3-C89 3.138(2), Ca3-C90 2.995(2), Ca3-C91 2.875(2), Ca3-C92 2.804(2), Ca3-C93 2.801(2), Ca3-C94 2.964(2), N1-Ca1-N2 78.05(6), N3-Ca2-N4 77.51(6), N5-Ca3-N6 78.73(6); (**11**) Ca1-N1 2.3290(12), Ca1-N2 2.3412(12), Ca1-C30 2.7127(15), Ca1-C31 2.8973(15), Ca1-C32 2.7097(14), Ca1-C33 2.7711(14), Ca1-C34 2.7660(15), N1-Ca1-N2 76.92(4). Symmetry operations to generate equivalent atoms 1-x, 1-y, 1-z.

The formation of compounds **9** – **11**, particularly the generation of the naphthalenide dianion in **9**, indicates a reducing potential for compound **2** that is at least equivalent to or surpasses that of elemental calcium metal and molecular Mg(I) compounds such as **5** and **6**. Evans and co-workers have previously made similar observations through comparison of the reactivity of Cp*₂Sm (**12**) and Cp*₂SmH (**13**), which contain formally reducing Sm²⁺ and fully oxidised Sm³⁺ centers, respectively (E^0 Sm³⁺/Sm²⁺ –2.304 V).^[20,21] Although neither samarium compound was reactive toward naphthalene, compound **12** effects two electron reduction of both anthracene and tetracene. Treatment of anthracene with compound **13** was also observed, albeit with less discrimination, to provide the identical reduction product, [(Cp*₂Sm)₂(μ-η²:η²-C₁₄H₁₀)], along with the formation of H₂.^[21] Although no mechanistic discussion was included in these earlier reports, examination of *in situ* ¹H NMR spectra recorded during reactions to provide compounds **9** – **11** confirmed the similar production of dihydrogen, observed as a resonance at δ 4.46 ppm, prompting a more detailed assessment of the mode of reaction of compound **2** with PAHs.

Reactions of **2** with naphthalene, anthracene and tetracene performed at 350 K within the cavity of a CW EPR spectrometer provided no evidence for the production of short lived hydrogen radicals. In each case, however, signals arising from the PAH radical anions, [C₁₀H₈]^{•-}, [C₁₄H₁₀]^{•-} and [C₁₈H₁₂]^{•-}, were readily observed and confirmed by comparison to simulated spectra of the fully delocalized products of single electron reduction (Figure S17). While these data indicate that the formation of the fully reduced PAH dianions is likely to occur via sequential single

electron reduction, they provide no compelling evidence for the mode of electron transfer or the production of the dihydrogen by-product. To further assess the nature of these processes, therefore, DFT calculations were performed on the complete system resulting from the reaction between compound **2** and tetracene. The formation of the resultant bimetallic compound (**11**) was reasoned to provide the necessary insight whilst limiting the computational expense that would have been incurred by a similar study of the formation of either trimetallic derivative, **9** or **10**. Possible reaction profiles yielding the formation of **11** were computed at the DFT level (B3PW91), varying the methine C-H unit involved in the reaction on the arene molecule (denoted as positions **f**, **g** and **h** in Figure 4). As described for the H/D exchange of benzene, reaction is initiated by the nucleophilic addition of a hydride of compound **2** to the tetracene molecule (**TS-BC**). The activation barrier (*ca.* 28-30 kcal mol⁻¹) associated with this attack varies with the position of hydride addition and may be compared to the analogous process computed for COT (21.8 kcal mol⁻¹).^[7] Following the intrinsic reaction coordinate in all cases leads to the formation of dicalcium tetracenyli intermediates (**Cf/Ch/Cg**) that retain a Ca-μ₂-H-Ca linkage similar to that identified during the reaction with benzene. While there is little to discriminate the kinetic facility of the nucleophilic attack, the formation of intermediates **Cf-h** is endothermic due to the disruption of the tetracene π system and their relative stabilities depend on the position of hydride addition (**Cf**, 16.1; **Cg** 13.7; **Ch** 6.2 kcal mol⁻¹). Irrespective of the consequent low productivity of the hydride addition step, however, **Cf-h** can evolve the further intermediates, **Df-h**, that comprise the experimentally observed




Figure 4: Computed enthalpy profile at room temperature for the reaction of **2** with tetracene. The Gibbs Free energies are given in parenthesis. The calculated SOMO of species **D** is shown on the far right hand side.

tetracene radical anion upon loss of hydrogen radical. In contrast to the appreciably exothermic formation of **Df** and **Dg** from their respective parent species, the transformation of **Ch** to **Dh** was computed to be effectively isoenergetic. While production of any of **Df**, **Dg** and **Dh** enables the formation of **11** through a second cascade of nucleophilic hydride addition and hydrogen radical release (see Figure S18), the comparable energies of **Ch** and **Dh** allow the identification of a further kinetically favored pathway to **11**. This alternative process (Figure S19) involves direct H_2 formation through hydride addition and intramolecular hydride deprotonation in a similar manner to that computed for the aromatization of COT by **2** (Scheme 1). Although the production of the tetraceny radical anion was unambiguously observed during the formation of **11**, the simultaneous operation of both one- and two-electron based mechanisms should not be discounted. We are comfortable to suggest, therefore, that the more complex trimetallic structures of compounds **9** and **10** may be a reflection of competing kinetic pathways, both inter- and intramolecular, leading to the two-electron reduction process.

In conclusion, the ability of compound **2** to effect the two electron reduction of naphthalene implies that its reducing potency is at least comparable to now well established dinuclear Mg(I) reagents. The emerging reactivity of heavier group 2 hydrides and related reagents, particularly for the nucleophilic activation of electron rich arenes, however, indicates that they will continue to provide a complementary source of unprecedented reactivity. We and others are continuing to investigate these possibilities.

Acknowledgements

We thank the EPSRC (UK) for the support of a DTP studentship (ASSW).

Conflict of interest

The authors declare no conflict of interest.

Keywords: calcium • reduction • hydride • main group chemistry • density functional theory


- [1] (a) S. Harder, *Chem. Commun.* **2012**, *48*, 11165-11177; (b) D. Mukherjee, J. Okuda, *Angew. Chem. Int. Ed.* **2018**, *57*, 1458-1473; D. Mukherjee, D. Schuhknecht, J. Okuda, *Angew. Chem. Int. Ed.* **2018**, *57*, 9590-9602; (c) J. Spielmann, S. Harder, *Chem. Eur. J.* **2007**, *13*, 8928-8938; (d) A. Causero, G. Ballmann, J. Pahl, C. Farber, J. Intemann, S. Harder, *Dalton Trans.* **2017**, *46*, 1822-1831; (e) B. Maitland, M. Wiesinger, J. Langer, G. Ballmann, J. Pahl, H. Elsen, C. Farber, S. Harder, *Angew. Chem. Int. Ed.* **2017**, *56*, 11880-11884; (f) M. Wiesinger, B. Maitland, C. Farber, G. Ballmann, C. Fischer, H. Elsen, S. Harder, *Angew. Chem. Int. Ed.* **2017**, *56*, 16654-16659; (g) P. Jochmann, J. P. Davin, T. P. Spaniol, L. Maron, J. Okuda, *Angew. Chem. Int. Ed.* **2012**, *51*, 4452-4455; (h) V. Leich, T. P. Spaniol, J. Okuda, *Inorg. Chem.* **2015**, *54*, 4927-4933; (i) C. N. de Bruin-Dickason, T. Sutcliffe, C. A. Lamsfus, G. B. Deacon, L. Maron, C. Jones, *Chem. Commun.* **2018**, *54*, 786-789; (j) X. H. Shi, C. P. Hou, C. L. Zhou, Y. Y. Song, J. H. Cheng, *Angew. Chem. Int. Ed.* **2017**, *56*, 16650-16653.
- [2] S. Harder, J. Brettar, *Angew. Chem. Int. Ed.* **2006**, *45*, 3474-3478.
- [3] S. Harder, B. Rösch, T. Xaver Gentner, H. Elsen, J. Langer, M. Wiesinger, *Angew. Chem. Int. Ed.* **2019**, doi.org/10.1002/anie.201901548, Accepted article.
- [4] A. S. S. Wilson, M. S. Hill, M. F. Mahon, *Organometallics* **2019**, *38*, 351-360.
- [5] A. S. S. Wilson, M. S. Hill, M. F. Mahon, C. Dinoi, L. Maron, *Science* **2017**, *358*, 1168-1171.
- [6] A. S. S. Wilson, C. Dinoi, M. S. Hill, M. F. Mahon, L. Maron, *Angew. Chem. Int. Ed.* **2018**, *57*, 15500-15504.
- [7] M. S. Hill, M. F. Mahon, A. S. S. Wilson, C. Dinoi, L. Maron, E. Richards, *Chem. Commun.* **2019**, *55*, 5732-5735.
- [8] (a) F. J. Smentowski, G. R. Stevenson, *J. Am. Chem. Soc.* **1967**, *89*, 5120-5121; (b) F. J. Smentowski, Stevenson, *J. Am. Chem. Soc.* **1969**, *91*, 7401-7402; (c) F. J. Smentowski, G. R. Stevenson, *J. Phys. Chem.* **1969**, *73*, 340-341.
- [9] (a) A. Stasch, C. Jones, *Dalton Trans.* **2011**, *40*, 5659-5672; (b) C. Jones, *Nature Rev. Chem.* **2017**, *1*, 0059.
- [10] (a) A. V. Zabula, M. A. Petrukhina, *Adv. Organometal. Chem.*, **2013**, *61*, 375-462; (b) J. E. Ellis, *Dalton Trans.* **2019**, *48*, 9538-9563.
- [11] (a) B. Bogdanovic, *Acc. Chem. Res.* **1988**, *21*, 261-267; (b) E. Bartmann, B. Bogdanovic, N. Janke, S. J. Liao, K. Schlichte, B. Spliethoff, J. Treber, U. Westeppe, U. Wilczok, *Chem. Ber. Rec.* **1990**, *123*, 1517-1528; (c) H. Bonnemann, B. Bogdanovic, R. Brinkmann, N. Egeler, R. Benn, I. Topalovic, K. Seevogel, *Main Group Metal Chem.* **1990**, *13*, 6; (d) A. Uresk, N. J. O'Neil, Z. Zhou, Z. Wei, M. A. Petrukhina, *J. Organometal. Chem.* **2019**, *897*, 57-63.
- [12] N. G. Connelly, W. E. Geiger, *Chem. Rev.* **1996**, *96*, 877-910.

- [13] (a) J. J. Brooks, W. Rhine, G. D. Stucky, *J. Am. Chem. Soc.* **1972**, *94*, 7346-7347; (b) C. Melero, A. Guijarro, M. Yus, *Dalton Trans.* **2009**, 1286-1289.
- [14] S. Krieck, L. Yu, M. Reiher, M. Westerhausen, *Eur. J. Inorg. Chem.* **2010**, 197-216.
- [15] S. R. Lawrence, C. A. Ohlin, D. B. Cordes, A. M. Z. Slawin, A. Stasch, *Chem. Sci.* **2019**, Advance Article, DOI: 10.1039/C9DT03976G.
- [16] (a) M. E. Thompson, S. M. Baxter, A. R. Bulls, B. J. Burger, M. C. Nolan, B. D. Santarsiero, W. P. Schaefer, J. E. Bercaw, *J. Am. Chem. Soc.* **1987**, *109*, 203-219; (b) M. Booi, B. J. Deelman, R. Duchateau, D. S. Postma, A. Meetsma, J. H. Teuben, *Organometallics* **1993**, *12*, 3531-3540; (c) G. Jeske, H. Lauke, H. Mauermann, H. Schumann, T. J. Marks, *J. Am. Chem. Soc.* **1985**, *107*, 8111-8118.
- [17] (a) S. Harder, *Organometallics* **2002**, *21*, 3782-3787; (b) A. G. Avent, M. R. Crimmin, M. S. Hill, P. B. Hitchcock, *Dalton Trans.* **2005**, 278-284.
- [18] S. Krieck, H. Görls, L. Yu, M. Reiher, M. Westerhausen, *J. Am. Chem. Soc.* **2009**, *131*, 2977-2985.
- [19] (a) W. E. Rhine, J. Davis, G. Stucky, *J. Am. Chem. Soc.* **1975**, *97*, 2079-2085; (b) W. E. Rhine, J. H. Davis, G. Stucky, *J. Organometal. Chem.* **1977**, *134*, 139-149; (c) H. Bock, K. Gharagozloo-Hubmann, M. Sievert, T. Prisner, Z. Havlas, *Nature* **2000**, *404*, 267-269.
- [20] W. J. Evans, S. L. Gonzales, J. W. Ziller, *J. Am. Chem. Soc.* **1994**, *116*, 2600-2608.
- [21] W. J. Evans, B. M. Schmiede, S. E. Lorenz, K. A. Miller, T. M. Champagne, J. W. Ziller, A. G. DiPasquale, A. L. Rheingold, *J. Am. Chem. Soc.* **2008**, *130*, 8555-8563.

COMMUNICATION

Entry for the Table of Contents (Please choose one layout)

COMMUNICATION



*Andrew S. S. Wilson, Chiara Dinoi,
Michael S. Hill, Mary F. Mahon, Laurent
Maron and Emma Richards*

Page No. – Page No.

**Calcium Hydride Reduction of
Polycyclic Aromatic Hydrocarbons**

A Fragrant Reduction: A molecular calcium hydride effects the two electron reduction of polyaromatic hydrocarbons, including naphthalene ($E^0 = -3.1$ V).