



*Citation for published version:*

Mulhearn, B, Li, D, McMorrow, F, Lu, H, McHugh, NJ & Tansley, SL 2022, 'A Commercial Anti-TIF1 $\gamma$  ELISA Is Superior to Line and Dot Blot and Should Be Considered as Part of Routine Myositis-Specific Antibody Testing', *Frontiers in Immunology*, vol. 13, 804037. <https://doi.org/10.3389/fimmu.2022.804037>

*DOI:*

[10.3389/fimmu.2022.804037](https://doi.org/10.3389/fimmu.2022.804037)

*Publication date:*

2022

*Document Version*

Peer reviewed version

[Link to publication](#)

**University of Bath**

**Alternative formats**

If you require this document in an alternative format, please contact:  
[openaccess@bath.ac.uk](mailto:openaccess@bath.ac.uk)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1

2

3 Brief Report

4 Title

5 A commercial anti-TIF1 $\gamma$  ELISA is superior to line and dot blot and  
6 should be considered as part of routine myositis-specific antibody  
7 testing.

8 Ben Mulhearn<sup>1,2</sup>, Danyang Li<sup>1</sup>, Fionnuala McMorrow<sup>1</sup>, Hui Lu<sup>1</sup>, Neil McHugh<sup>1</sup> and Sarah  
9 Tansley<sup>1,2</sup>

10 <sup>1</sup>Department of Pharmacy and Pharmacology, University of Bath, Claverton Down, Bath,  
11 UK, BA2 7AY

12 <sup>2</sup>Royal National Hospital for Rheumatic Diseases, Royal United Hospitals, Combe Park, Bath,  
13 UK, BA1 3NG

14 Corresponding email address: [bm765@bath.ac.uk](mailto:bm765@bath.ac.uk)

15

16

## 17 1) Abstract

18 Objectives: Anti-TIF1 $\gamma$  is an important autoantibody in the diagnosis of cancer-associated  
19 dermatomyositis and the most common autoantibody in juvenile onset dermatomyositis. Its  
20 reliable detection is important to instigate further investigations into underlying malignancy  
21 in adults. We previously showed that commercial assays using line and dot blots do not  
22 reliably detect anti-TIF1 $\gamma$ . We aimed to test a new commercial ELISA and compare with  
23 previously obtained protein immunoprecipitation.

24 Methods: Radio-labelled immunoprecipitation had previously been used to determine the  
25 autoantibody status of patients with immune-mediated inflammatory myopathies and  
26 several healthy controls. ELISA was undertaken on healthy control and anti-TIF1 $\gamma$  sera and  
27 compared to previous immunoprecipitation data.

28 Results: A total of 110 serum samples were analysed: 42 myositis patients with anti- TIF1 $\gamma$   
29 and 68 autoantibody negative healthy control sera. Anti-TIF1 $\gamma$  was detected by ELISA in 41  
30 out of 42 of the anti-TIF1 $\gamma$ -positive samples by immunoprecipitation, and in none of the  
31 healthy controls, giving a sensitivity of 97.6% and specificity of 100%. The false negative rate  
32 was 2%.

33 Conclusion: ELISA is an affordable and time-efficient method which is accurate in detecting  
34 anti-TIF1 $\gamma$ .

35

## 36 Keywords

37 IIMs, DM, TIF1 $\gamma$ , ELISA, cancer, autoantibodies, myositis

38

39

## 40 2) Introduction

41 The ability to detect myositis -specific and -associated antibodies (MSAs and MAAs), which  
42 can be found in the sera of 60 – 70% patients with immune-mediated inflammatory  
43 myopathies (IIMs) (1), has greatly improved the diagnosis and phenotyping of these rare  
44 diseases. Not only do they aid diagnosis, but they also guide further investigation and  
45 management (2). For instance, it is well-known that IIMs, and dermatomyositis (DM) in  
46 particular, are strongly linked with cancer, with estimates varying between 7 and 32 % (3).

47 Anti-transcription intermediary factor 1 $\gamma$  (TIF1 $\gamma$ ) autoantibodies are found in both juvenile  
48 dermatomyositis (JDM) and adult IIMs. They are present in 7% of European adults with DM  
49 and 20 – 30% of children affected by JDM (2). Strikingly, 38 – 84% of patients adult DM  
50 patients  $\geq$  39 years of age who are TIF1 $\gamma$ -positive in both European and Japanese cohorts  
51 develop cancer in the 3 years before and after DM diagnosis (4–6). Anti-TIF1 $\gamma$  detection in  
52 patients with a new diagnosis of DM  $\geq$  39 years of age may therefore prompt a thorough  
53 investigation for the detection of cancer and reduce cancer mortality rates, making the  
54 accurate detection of anti-TIF1 $\gamma$  a research priority.

55 Currently the reference standard in the detection of MSAs is immunoprecipitation (IP) due  
56 to its ability to detect well-described and novel autoantibodies. However, this technique is  
57 impractical for use in clinical practice owing to its expense and the length of time it takes to  
58 reach a result which usually takes a minimum of 2 – 3 weeks. For this reason, several  
59 commercially available immunoassays have become available which are low cost, easy to use,  
60 and are reported to detect an array of MSAs. However, these immunoassays are subject to  
61 both false positives and false negatives. A number of them have recently been tested by our  
62 group and others (7,8). In particular, anti-TIF1 $\gamma$  was found to be particularly problematic with  
63 false negatives found in 40% samples analysed by line blot and 76% by dot blot (7). Espinosa-  
64 Ortega *et al.* (8) also found low concordance between anti-TIF1 $\gamma$  detected by line/dot blot  
65 and immunoprecipitation, with a Cohen's kappa of 0.56. This is likely because anti-TIF1 $\gamma$   
66 frequently target a conformational epitope, meaning the tertiary antigen structure is required  
67 to remain intact to be recognised by the autoantibody (9). Whereas line and dot blot  
68 immunoassays utilise denatured antigen, enzyme-linked immunosorbent assays (ELISAs)

69 maintain the tertiary structure of the protein. Fujimoto *et al.* (10) recently tested a newly-  
70 developed ELISA in a Japanese cohort of patients with a spectrum of IIMs, and found this  
71 approach to be highly effective with 100% sensitivity and specificity which was a result  
72 comparable to immunoprecipitation.

73 In this study, we aimed to test the same commercial ELISA kit (Medical & Biological  
74 Laboratories Co. Ltd., Nagoya, Aichi, Japan) for the detection of TIF1 $\gamma$  autoantibodies in a  
75 European cohort of adult IIM patients and compared results with samples previously analysed  
76 using immunoprecipitation .

77

### 78 3) Methods

#### 79 Sample selection

80 Myositis serum samples used in this study were chosen as previously described (7) from a  
81 biobank of more than 3000 samples collected for research or diagnostic purposes (2,11). All  
82 serum samples had previously been analysed by immunoprecipitation locally and contain at  
83 least one MSAA. Twenty-five anti-TIF1 $\gamma$  samples had also been previously analysed by line  
84 and dot blot (7). Briefly, sera were stored at -20°C prior to analysis in a facility at the University  
85 of Bath. The study had ethical approval through the host Institute (University of Bath EIRA  
86 reference number 17-01211). All samples from research cohorts had existing ethics in place.

#### 87 ELISA

88 ELISA was performed on 5 $\mu$ L of diluted serum sample as per the manufacturer's instructions  
89 (Medical & Biological Laboratories Co. Ltd., Nagoya, Aichi, Japan). All samples were run in  
90 duplicate. Briefly, samples were thawed and diluted to a 1:101 concentration and incubated  
91 on a microwell plate for 30 minutes. Wells were then incubated with a horseradish  
92 peroxidase-conjugated goat anti-human IgG antibody conjugate for 30 minutes followed by a  
93 TMB/peroxide substrate for 15 minutes. The reaction was terminated by 0.25 mol/L sulfuric  
94 acid. All incubations took place at room temperature with 4 wash cycles between steps. The

95 absorbance of each well was read on a FLUOstar Omega microplate reader (BMG Labtech  
96 Ltd., Aylesbury, Buckinghamshire, Great Britain) at 450 nm wavelength. Positive and negative  
97 cut off values were calculated according to previous work described by Fujimoto *et al.* (10)  
98 and expressed in arbitrary units (au).

## 99 Immunoprecipitation

100 Radio-immunoprecipitation had been previously undertaken as described by Tansley *et al.*  
101 (7). Briefly, sera were mixed with protein-A-Sepharose beads and a 35(S)methionine  
102 radiolabelled K562 cell extract, followed by fractionation by SDS-PAGE and analysis by  
103 autoradiography. A characteristic doublet band at 155/140 was read as being positive for  
104 TIF1 $\gamma$  (12).

## 105 Data analysis

106 Statistical analysis was undertaken using Prism 9 version 9.2.0 for macOS (GraphPad  
107 Software, LLC., San Diego, CA, USA). Confidence intervals (CI) are expressed at 95%.

108

## 109 4) Results

110 A total of 110 serum samples were analysed, of which 42 were known to have anti-TIF1 $\gamma$  and  
111 68 were healthy control sera. Immunoprecipitation data was held for all samples. Diagnoses  
112 included DM (n=27), clinically amyopathic DM (n=4), JDM (n=5), polymyositis (n=4), and  
113 overlap syndrome (n=2). All HC samples tested were autoantibody negative by  
114 immunoprecipitation.


## 115 Commercial TIF1 $\gamma$ ELISA performed as well as immunoprecipitation

116 Forty-one patient samples with anti-TIF1 $\gamma$  tested positive by ELISA as defined by a cut-off  
117 point of 32 au. None of the HC samples tested positive using this cut-off point. The remaining  
118 anti-TIF1 $\gamma$  positive sample was just under the cut-off for positivity (30.2 au). This gives an area  
119 under the ROC curve (AUC) of 0.988 (CI 0.961 – 1.000, P < 0.0001) which is equivalent to

120 sensitivity of 97.6% (CI 87.7% - 99.9%) and a specificity of 100% (CI 94.65% - 100%). In this  
121 case, Cohen's Kappa would give a value of 1.

122 Quantitative results for the ELISA values are shown in figure 1. Briefly, the median ELISA assay  
123 result for HC samples was 5.99 au. (median CI 4.74 – 7.87) and for the TIF1 $\gamma$  samples was  
124 128.5 au. (median CI 110.4 – 135.4).

**Figure 1. TIF1 $\gamma$  ELISA values**



125

126 **Figure 1. TIF1 $\gamma$  ELISA values for 68 healthy controls and 42 TIF1 $\gamma$  serum samples**


127 Graph showing the relative ELISA titres for healthy control and TIF1 $\gamma$  samples expressed in  
128 arbitrary units for each individual serum sample (circles). Dashed line represents the positive  
129 cut-off point as previous described (10). All 68 healthy control (HC) samples were underneath  
130 the cut-off and all but one of the 42 TIF1 $\gamma$  samples were above the cut-off. The TIF1 $\gamma$  sample  
131 below the cut-off had a weak band in the 140/155 kDa region.

132 **Low anti-TIF1 $\gamma$  ELISA titres are associated with false negative line blot results**

133 Given that our group previously tested 25 anti-TIF1 $\gamma$  samples by line blot, we were able to  
134 compare ELISA titres in this study with this data to try and understand which samples might  
135 test negative by line blot. The results are shown in figure 2. All anti-TIF1 $\gamma$  positive samples  
136 by ELISA with low titres (between 30 – 100 au.) tested negative by line blot. However, 3 out  
137 of the 9 samples testing negative by line blot had high anti- TIF1 $\gamma$  titres (> 100 au.). The  
138 difference in ELISA titres between those testing negative and positive by line blot was  
139 statistically significant (P = 0.0041, two-tailed Mann-Whitney test), suggesting that lower

140 anti-TIF1 $\gamma$  antibody titres lead to false negative line blot results. Similarly, dot blot samples  
141 returned only 7/24 (29%) true positives out of the anti-TIF1 $\gamma$  samples that tested positive by  
142 ELISA and immunoprecipitation.

**Figure 2. Comparison of anti-TIF1 $\gamma$  ELISA titre and line blot result**



143

**Figure 2. Comparison of anti-TIF1 $\gamma$  ELISA titre and line blot result**

144 Graph showing a comparison between anti-TIF1 $\gamma$  ELISA titre and line blot result, as previously  
145 tested by our group (7). ELISA titres are expressed in arbitrary units and calculated as per the  
146 manufacturer's instructions. Lines and error bars represent median values with 95%  
147 confidence intervals. The line blot results are expressed as negative (-), low positive (+),  
148 moderately positive (++), and high positive (+++). The median ELISA values for negative, low  
149 positive, moderately positive, and high positive results were 63.5 au., 123.5 au., 145.8 au.,  
150 and 151.2 au., respectively. A two-tailed Mann Whitney test comparing ELISA titres between  
151 negative (-) and positive (+, ++, +++) line blots found a statistical difference between the two  
152 groups (P = 0.0041).  
153

154

## 155 5) Discussion

156 This data has shown that accurate detection of anti-TIF1 $\gamma$  can be achieved by ELISA and  
157 confirms the findings made by Fujimoto *et al.* (10). The accuracy of detection is high and  
158 would be acceptable for use in clinical practice. Compared to other cost- and time- effective  
159 methods such as line and dot blot which have false negative rate of 40% - 70% (7), this data



160 found that ELISA has a false negative rate of 1/42 (2%). This data has also shown that anti-  
161 TIF1 $\gamma$  titre correlates with a positive line blot result. This result is not unexpected given that  
162 the line blot is a semi-quantitative method of detecting autoantibodies. Importantly, where  
163 ELISA was able to detect samples with low titres of anti-TIF1 $\gamma$  (between 30 – 100 au.), line  
164 blot was unable to do so. Line blot also failed to detect some samples with high anti-TIF1 $\gamma$   
165 titres (> 100 au.). Taken together, anti-TIF1 $\gamma$  ELISA performs better than line blot in detecting  
166 this clinically important autoantibody.

167 Anti-TIF1 $\gamma$  status by immunoprecipitation was determined by recognition of 155/140kDa  
168 bands alongside an anti-TIF1 $\gamma$  standard control. It remains possible that the sample negative  
169 by ELISA has an unknown autoantibody with an identical band pattern although this would  
170 seem unlikely. Furthermore, the sample produced an ELISA result just below the positive  
171 threshold and may simply be a low-titre positive. The ELISA threshold could be adjusted to  
172 reduce the likelihood of this occurring, but this is likely to lead to some false positives. The  
173 most appropriate cut-off threshold may depend on the clinical context, for example, a low  
174 false positive rate may be tolerable in patients with confirmed dermatomyositis to inform the  
175 intensity of malignancy screening.

176 The current study was not designed to investigate the relationship between anti-TIF1 $\gamma$  titres  
177 and cancer detection rates. It would, however, be useful to investigate how anti-TIF1 $\gamma$  titre  
178 using ELISA correlates with malignancy. Recent work by Fiorentino *et al.* (13) found anti-TIF1 $\gamma$   
179 titre positively correlated with cancer detection rate in DM, ranging from 8% detection for  
180 low titres to 36% detection for high titres. Furthermore, some of our healthy control samples  
181 had low anti-TIF1 $\gamma$  titres just below the positive cut-off and it would be of interest to  
182 investigate if these healthy subjects had a higher malignancy rate compared to a negative  
183 anti-TIF1 $\gamma$  control population.

184 The detection of anti-TIF1 $\gamma$  in adult DM patients should be considered a red flag for  
185 malignancy (4–6). Accurate and timely detection of anti-TIF1 $\gamma$  autoantibodies is therefore  
186 vital for these patients to ensure underlying malignancy is diagnosed and treated promptly.  
187 We suggest that, when investigating IIMs, anti-TIF1 $\gamma$  ELISA is undertaken alongside, ANA  
188 testing and a multiplex immunoblot assay to ensure accurate detection of this important  
189 autoantibody.

190 Funding: This work was supported by funding from CureJM and the Bath Institute of  
191 Rheumatic Diseases. ELISA kits were provided by MBL.

192 Disclosure statement: The authors have declared no conflicts of interest.

### 193 Key messages

- 194 1. Anti-TIF1 $\gamma$  is a key autoantibody in the diagnosis of cancer-associated  
195 dermatomyositis and juvenile dermatomyositis
- 196 2. ELISA is a quick and easy method in accurately detecting anti-TIF1 $\gamma$  autoantibodies
- 197 3. Diagnosis of IIMs should include ANA immunofluorescence, line or dot blot, and anti-  
198 TIF1 $\gamma$  ELISA

199    **References**

200    1.    McHugh NJ, Tansley SL. Autoantibodies in myositis. *Nat Rev Rheumatol* (2018)  
201        **14**:290–302. doi:10.1038/nrrheum.2018.56

202    2.    Betteridge Z, Tansley S, Shaddick G, Chinoy H, Cooper RG, New RP, Lilleker JB,  
203        Vencovsky J, Chazarain L, Danko K, et al. Frequency, mutual exclusivity and clinical  
204        associations of myositis autoantibodies in a combined European cohort of idiopathic  
205        inflammatory myopathy patients. *J Autoimmun* (2019) **101**:  
206        doi:10.1016/j.jaut.2019.04.001

207    3.    Aussy A, Boyer O, Cordel N. Dermatomyositis and immune-Mediated Necrotizing  
208        Myopathies: A window on Autoimmunity and Cancer. *Front Immunol* (2017) **8**:  
209        doi:10.3389/fimmu.2017.00992

210    4.    Oldroyd A, Sergeant JC, New P, McHugh NJ, Betteridge Z, Lamb JA, Ollier WE, Cooper  
211        RG, Chinoy H. The temporal relationship between cancer and adult onset anti-  
212        transcriptional intermediary factor 1 antibody-positive dermatomyositis. *Rheumatol*  
213        (*United Kingdom*) (2019) **58**: doi:10.1093/rheumatology/key357

214    5.    Hida A, Yamashita T, Hosono Y, Inoue M, Kaida K, Kadoya M, Miwa Y, Yajima N,  
215        Maezawa R, Arai S, et al. Anti-TIF1- $\gamma$  antibody and cancer-associated myositis: A  
216        clinicohistopathologic study. *Neurology* (2016) **87**:  
217        doi:10.1212/WNL.0000000000002863

218    6.    Fiorentino DF, Chung LS, Christopher-Stine L, Zaba L, Li S, Mammen AL, Rosen A,  
219        Casciola-Rosen L. Most patients with cancer-associated dermatomyositis have  
220        antibodies to nuclear matrix protein NXP-2 or transcription intermediary factor 1 $\gamma$ .  
221        *Arthritis Rheum* (2013) **65**: doi:10.1002/art.38093

222    7.    Tansley SL, Li D, Betteridge ZE, McHugh NJ. The reliability of immunoassays to detect  
223        autoantibodies in patients with myositis is dependent on autoantibody specificity.  
224        *Rheumatol (United Kingdom)* (2020) **59**: doi:10.1093/rheumatology/keaa021

225    8.    Espinosa-Ortega F, Holmqvist M, Alexanderson H, Storfors H, Mimori T, Lundberg IE,  
226        Rönnelid J. Comparison of autoantibody specificities tested by a line blot assay and  
227        immunoprecipitation-based algorithm in patients with idiopathic inflammatory  
228        myopathies. *Ann Rheum Dis* (2019) **78**:858–860. doi:10.1136/ANNRHEUMDIS-2018-  
229        214690

230    9.    Targoff IN, Mamyrova G, Trieu EP, Perurena O, Koneru B, O’Hanlon TP, Miller FW,  
231        Rider LG. A novel autoantibody to a 155-kd protein is associated with  
232        dermatomyositis. *Arthritis Rheum* (2006) **54**: doi:10.1002/art.22164

233    10.    Fujimoto M, Murakami A, Kurei S, Okiyama N, Kawakami A, Mishima M, Sato S,  
234        Seishima M, Suda T, Mimori T, et al. Enzyme-linked immunosorbent assays for  
235        detection of anti-transcriptional intermediary factor-1 gamma and anti-Mi-2  
236        autoantibodies in dermatomyositis. *J Dermatol Sci* (2016) **84**:  
237        doi:10.1016/j.jdermsci.2016.09.013

238    11.    Tansley SL, Simou S, Shaddick G, Betteridge ZE, Almeida B, Gunawardena H, Thomson  
239        W, Beresford MW, Midgley A, Muntoni F, et al. Autoantibodies in juvenile-onset  
240        myositis: Their diagnostic value and associated clinical phenotype in a large UK

241 cohort. *J Autoimmun* (2017) **84**: doi:10.1016/j.jaut.2017.06.007

242 12. Fujimoto M, Hamaguchi Y, Kaji K, Matsushita T, Ichimura Y, Kodera M, Ishiguro N,  
243 Ueda-Hayakawa I, Asano Y, Ogawa F, et al. Myositis-specific anti-155/140  
244 autoantibodies target transcription intermediary factor 1 family proteins. *Arthritis*  
245 *Rheum* (2012) **64**:513–522. doi:10.1002/art.33403

246 13. Fiorentino DF, Gutierrez-Alamillo L, Hines D, Yang Q, Casciola-Rosen L. Distinct  
247 dermatomyositis populations are detected with different autoantibody assay  
248 platforms. *Clin Exp Rheumatol* (2019) **37**:1048–1051. Available at:  
249 <http://www.ncbi.nlm.nih.gov/pubmed/31376258>

250