

## Notulae to the Italian flora of algae, bryophytes, fungi and lichens: 15

Sonia Ravera<sup>1,2</sup>, Marta Puglisi<sup>3</sup>, Alfredo Vizzini<sup>4</sup>, Cecilia Totti<sup>5</sup>, Mattia M. Azzella<sup>6</sup>, Giulia Bacilliere<sup>3</sup>, Rossano Bolpagni<sup>7</sup>, Othmar Breuss<sup>8</sup>, Annalena Cogoni<sup>9</sup>, Antonio B. De Giuseppe<sup>10</sup>, Zuzana Fačková<sup>11</sup>, Felix Faltner<sup>12</sup>, Gabriele Gheza<sup>13</sup>, Paolo Giordani<sup>14</sup>, Deborah Isocrono<sup>15</sup>, Petra Mair<sup>16</sup>, Jiří Malíček<sup>17</sup>, Pasquale Marino<sup>18</sup>, Helmut Mayrhofer<sup>19</sup>, Juri Nascimbene<sup>13</sup>, Silvia Ongaro<sup>15</sup>, Luca Paoli<sup>20</sup>, Nicodemo G. Passalacqua<sup>10,21</sup>, Silvia Poponessi<sup>9</sup>, Domenico Puntillo<sup>10</sup>, Francesco Maria Raimondo<sup>22</sup>, Giovanni Sicoli<sup>21</sup>, Wilhelm Tratter<sup>23</sup>

**1** *Dipartimento di Scienze e Tecnologie Biologiche Chimiche e Farmaceutiche (STEBICEF), Università di Palermo, Via Archirafi 38, 90123, Palermo, Italy* **2** *NBFC, National Biodiversity Future Center, Palermo 90133, Italy* **3** *Dipartimento di Scienze Biologiche, Geologiche e Ambientali, Sezione di Biologia vegetale, Università di Catania, Via A. Longo 19, 95125, Catania, Italy* **4** *Institute for Sustainable Plant Protection (IPSP) – CNR, Viale P.A. Mattioli 25, 10125, Torino, Italy* **5** *Dipartimento di Scienze della Vita e dell'Ambiente, Università Politecnica delle Marche, via Breccie Bianche, 60131, Ancona, Italy* **6** *Dipartimento di Pianificazione, Design, Tecnologia dell'Architettura – La Sapienza Università di Roma, Via Emanuele Gianturco 2, 00196, Roma, Italy* **7** *Dipartimento di Scienze Chimiche, della Vita e della Sostenibilità Ambientale dell'Università degli Studi di Parma, Parco Area delle Scienze 33/A, 43124, Parma, Italy* **8** *Department of Botany and Biodiversity Research, University of Vienna, Rennweg 14, 1030, Vienna, Austria* **9** *Dipartimento di Scienze della Vita e dell'Ambiente, Università di Cagliari, Viale S. Ignazio, 09123, Cagliari, Italy* **10** *Museo di Storia Naturale della Calabria ed Orto Botanico, Università della Calabria, 87036 Arcavacata di Rende (Cosenza), Italy* **11** *Plant Science and Biodiversity Centre, Slovak Academy of Sciences, Dúbravská cesta 9, SK-84523, Bratislava, Slovakia* **12** *Reinthalstraße 4a, 39031, Bruneck/Reischach, Italy* **13** *BIOME Lab, Dipartimento di Scienze Biologiche, Geologiche e Ambientali, Alma Mater Studiorum, Università di Bologna, Via Irnerio 42, 40126, Bologna, Italy* **14** *Dipartimento di Farmacia (DIFAR), Università di Genova, viale Cembrano 4, 16148, Genova, Italy* **15** *Dipartimento di Scienze Agrarie, Forestali e Alimentari, Università degli Studi di Torino, Largo Paolo Braccini, 10095, Grugliasco (Torino) Italy* **16** *Museo di Scienze Naturali dell'Alto Adige, via dei Bottai 1, 39100, Bolzano, Italy* **17** *Institute of Botany, The Czech Academy of Sciences, Zámek 1, CZ-252 43, Průhonice, Czech Republic* **18** *Bona Furtuna LLC, P.O. Box 277 95031 Los Gatos, CA, USA* **19** *Division of Plant Sciences, Institute of Biology, NAWI Graz, University of Graz, Holteigasse 6, 8010, Graz, Austria* **20** *Dipartimento di Biologia, Università di Pisa, Via Luca Ghini 13, 56126, Pisa, Italy* **21** *Dipartimento di Biologia, Ecologia e Scienze della Terra, Università della Calabria, 87036, Arcavacata di Rende (Cosenza), Italy* **22** *PLANTA/Centro autonomo di Ricerca, Documentazione e Formazione, Via Serraglio Vecchio 28, 90123, Palermo, Italy* **23** *Alpreid 77, 39010, St. Pankraz (Bolzano), Italy*

Corresponding author: Sonia Ravera ([sonia.ravera@unipa.it](mailto:sonia.ravera@unipa.it))

---

Academic editor: Lorenzo Peruzzi | Received 23 March 2023 | Accepted 11 April 2023 | Published 19 April 2023

---

**Citation:** Ravera S, Puglisi M, Vizzini A, Totti C, Azzella MM, Bacilliere G, Bolpagni R, Breuss O, Cogoni A, De Giuseppe AB, Fačková Z, Faltner F, Gheza G, Giordani P, Isocrono D, Mair P, Malíček J, Marino P, Mayrhofer H, Nascimbene J, Ongaro S, Paoli L, Passalacqua NG, Poponessi S, Puntillo D, Raimondo FM, Sicoli G, Tratter W (2023) Notulae to the Italian flora of algae, bryophytes, fungi and lichens: 15. Italian Botanist 15: 35–47. <https://doi.org/10.3897/italianbotanist.15.103781>

---

## Abstract

In this contribution, new data concerning algae, bryophytes, fungi and lichens of the Italian flora are presented. It includes new records and confirmations for the algal genus *Nitella*, for the bryophyte genera *Anthoceros*, *Dicranodontium*, *Fontinalis*, and *Riccia*, the fungal genera *Inocybe* and *Xerophorus*, and the lichen genera *Bagliettoa*, *Biatora*, *Calicium*, *Cladonia*, *Coniocarpon*, *Lecanora*, *Opegrapha*, *Placynthium*, *Rhizocarpon*, *Scytinium*, *Solenopsis*, *Stereocaulon*, and *Verrucaria*.

## Keywords

Ascomycota, Basidiomycota, Biodiversity, Bryidae, Charophyceae

## How to contribute

The text of the records should be submitted electronically to: Cecilia Totti (c.totti@univpm.it) for algae, Marta Puglisi (mpuglisi@unict.it) for bryophytes, Alfredo Vizzini (alfredo.vizzini@unito.it) for fungi, Sonia Ravera (sonia.ravera@unipa.it) for lichens. Each text should be within 1,000 characters (spaces included).

## Floristic records

### Algae

#### *Nitella capillaris* (Krocker) J.Groves & Bullock-Webster (Characeae)

+ **LAZ:** Ceprano (Frosinone), tiny temporary pool on the southern side of the via Sfratti woodland (WGS84: 33T 371658.4600583), 130 m, 25 February 2023, leg. R. Bolpagni, det. M.M. Azzella (RO). – Species confirmed for the flora of Lazio.

Both female and male individuals were found growing sparsely in a small (not more than 1 m<sup>2</sup>) temporary pool not very far from the highway crossing (via Carl Vepu). In Italy, *Nitella capillaris* has been recently confirmed for Sicilia (Romanov et al. 2019), while the reports for peninsular Italy date back to more than 50 years ago (Bazzichelli and Abdelahad 2009). This record is a confirmation of the species for the flora of Lazio 60 years after the last reports (Corillion and Guerlesquin 1963).

M.M. Azzella, R. Bolpagni

## Bryophytes

### *Anthoceros agrestis* Paton (Anthocerotaceae)

+ **TAA**: Walchhorn, E of Reischach, Brunico, Val Pusteria (Bolzano), on the edge of a stubble cornfield, near hedges on siliceous soil (UTM WGS84: 32T 726187.5184209), 1038 m, 24 September 2022, *F. Faltner*, conf. *P. Mair* (Herb. Faltner; BOZ BRYO 8353). – Species new for the flora of Trentino-Alto Adige.

*Anthoceros agrestis* is quite similar to *A. punctatus* L., from which it can be distinguished essentially for the size of the mature antheridia, mostly about 80 µm long. It is mainly distributed in the temperate zone of central Europe but is rather rare in the Mediterranean-Atlantic parts (Hodgetts and Lockart 2020). In Italy, *A. agrestis* is signaled with old records for Piemonte, Toscana, Campania, and Sardegna, while more recent data (after 1968) are available for Lombardia (Brusa 2019), Lazio, Vatican City State, Calabria, and Sicilia (Aleffi et al. 2020). In the new locality it was found with *Riccia* sp. pl. and *Tortula truncata* (Hedw.) Mitt. This species is assessed as Near Threatened in Europe (Hodgetts et al. 2019).

F. Faltner, P. Mair

### *Dicranodontium asperulum* (Mitt.) Broth. (Leucobryaceae)

+ **EMR**: Passo della Scalucchia, Alpe di Succiso (Reggio Emilia) (UTM WGS84 32T 5979945.4912109), 1346 m, 21 July 2018, *S. Poponessi* (Bryo/AR002). – Specie new for the flora of Emilia-Romagna.

*Dicranodontium asperulum* is a Boreal-montane species with main distribution in North America, SE Asia (China, Japan Nepal, India), and central Europe, while it is very rare in the Mediterranean countries (Hodgetts and Lockart 2020). In Italy it occurs only in Piemonte, Veneto, Umbria, and was recorded before 1968 in Lombardia and Trentino-Alto Adige (Aleffi et al. 2020). The species prefers sheltered moist sites where it grows on acidic soils in montane heath, on ledges or crevices among acidic rocks. It is easily recognized by the wide-spreading leaves, standing out from the stem at 45° or more when wet, the serrate to serrulate shoulders of the leaf bases and the long arista which is serrate all around. In the new locality, *D. asperulum* was found on acidic soil at the base of *Fagus sylvatica* L., together with *Plagiotecium denticulatum* (Hedw.) Schimp. var. *denticulatum*, *Radula complanata* (L.) Dumort., *Brachytheciastrum velutinum* (Hedw.) Ignatov & Huttunen, and *Pterygynandrum filiforme* Hedw.

S. Poponessi, G. Bacilliere

### *Fontinalis hypnoides* var. *hypnoides* C.Hartm. (Fontinalaceae)

+ **SAR**: Rio Santa Lucia, Assemini (Cagliari) (UTM WGS84 32T 50226.4338222), on riverbed rocks of the river Rio Santa Lucia, 82 m, 24 September 2020, *S. Poponessi*, *A. De Agostini*, *A. Cogoni* (CAG SA2.2.3). – Variety confirmed for the flora of Sardegna.

*Fontinalis hypnoides* var. *hypnoides* is an aquatic moss growing submerged in standing or slow-moving water, including slightly eutrophic lakes, slow flowing rivers, or in reed-beds, normally attached to stones and wood. It is quite common in Europe (Hodgetts and Lockart 2020). According to Aleffi et al. (2020), *F. hypnoides* var. *hypnoides* is known in Italy from Veneto, Abruzzo, Puglia, Calabria and, with old records, Lombardia, Lazio and Sardegna. In Sardegna it was previously signalled more than a century ago from a few sites located in the eastern part of the island (Cardot 1882; Barbey 1884; Fleischer 1893). *Fontinalis hypnoides* var. *hypnoides* is distinguished from other species in the genus by its medium size and monomorphic leaves consistently plane at the apices.

S. Poponessi, A. Cogoni

***Riccia breidleri* Jur. ex Steph. (Ricciaceae)**

+ **TAA:** Oberer Hintergratsee, Sulden, Stilfser Joch/Stelvio National Park, Ortler Alps (Bozen/Bolzano), on damp, muddy, fine sandy shores (calcareous morainic material on phyllitic bedrock) of the small remnant of this alpine lake, partly drying out in summer (UTM WGS84: 32T 621420.5150986), 2650 m, 7 July 2017, *W. Tratter* (BOZ BRYO 8351); *ibidem*, 12 September 2022, *leg. P. Mair*, *conf. H. Köckinger* (BOZ BRYO 8352); East side of the Unterer Hintergratsee, Sulden, Stilfser Joch/Stelvio National Park, Ortler Alps (Bozen/Bolzano), on damp and dry, muddy shore edges, (UTM WGS84: 32T 621585.5150646), 2596 m, 12 September 2022, *P. Mair*. – Species new for the flora of Trentino-Alto Adige.

*Riccia breidleri* is endemic to Alps (Hodgetts et al. 2020), growing above 2000 m of elevations. This species grows on acidic soil in temporary pools that fill with melting snow water in the spring. In Italy it is only known for Piemonte, Valle d'Aosta, and Lombardia (Aleffi et al. 2020). *Riccia breidleri* is included in Annex II of the Habitats Directive of the EU and in the Appendix I of the Bern Convention, and considered Vulnerable in the European red list of bryophytes (Hodgetts et al. 2019).

P. Mair, W. Tratter

**Fungi**

***Inocybe langei* R.Heim (Inocybaceae)**

+ **CAL:** Botanical Garden, University of Calabria, Rende (Cosenza), on the ground under the crown of downy oak (*Quercus pubescens* Willd.) trees (UTM WGS84: 33S 605921.435728), 210 m, 5 September 2022, *G. Sicoli, A.B. De Giuseppe, N.G. Pissalacqua* (CLU F323). – Species new for the flora of Calabria.

A group of solitary, but gregarious, basidiomata belonging to *Inocybe langei* was observed on the ground among the litter of *Quercus pubescens* trees. The pilei were broadly conical and slightly umbonate when young, then almost applanate with age, less than 3 cm wide, straw-coloured, but a bit darker ochre in the centre, and fibrillose.

The stipe was pale, rather short and robust, with a slightly bulbous base, covered by crested caulocystidia especially along the upper half. The gills were beige-gray, the edge showing ventricose-lageniform and crested cheilocystidia. Spores were amygdaliform and  $6.5\text{--}8.5 \times 3.5\text{--}5 \mu\text{m}$  in size, thus shorter than those belonging to the closely-related species *I. hirtella* Bres., which is also characterised by the smell of almonds not recorded in our samples (Printz 1992; Courtecuisse and Duhem 1995). So far, *Inocybe langei* has been reported only in northern Italy (Onofri et al. 2013).

G. Sicoli, A.B. De Giuseppe, N.G. Passalacqua

### ***Xerophorus donadinii* (Bon) Vizzini, Consiglio & M. Marchetti (Callistosporiaceae)**

+ **CAL:** Botanical Garden, University of Calabria, Rende (Cosenza) (UTM WGS84: 33S 605857. 4357281), on the ground under the crown of a pedunculate oak tree (*Quercus robur* L.), 200 m, 21 October 2022, A.B. De Giuseppe, N.G. Passalacqua, G. Sicoli (CLU F325). – Species new for the flora of Calabria.

A group of six small, gregarious and collybioid basidiomata was observed on the ground on the litter under a planted tree of *Quercus robur*. The youngest pilei were hemisphaeric convex and with an involute edge; the most mature ones were more appanate, matt, smooth, dark brown with a reddish tone in the centre, paler at the edge, and 1.0–2.5 cm wide. The lower side of the pileus showed distant, emarginated and pale yellowish gills supported by a lemon-yellow, slender, flexuose and fibrillose stipe with white rhizomorphs attached at the base. Spores were amygdaliform, smooth, hyaline, and  $7\text{--}9 \times 4\text{--}5 \mu\text{m}$  in size. Cheilocystidia were scattered and cylindrical, no pleurocystidia were observed. Smell and taste resulted indistinct. Based on the above characters and ecology, this fungus was identified as *Xerophorus donadinii*. This species, was first described as *Callistosporium olivascens* var. *donadinii* Bon, ecologically distinct from the proper *C. olivascens* (Boud.) Bon, currently *X. olivascens* (Bon) Vizzini, Consiglio & M. Marchetti, which prefers association with cedar trees. Further differences are also morphological: *X. donadinii* has pilei with smaller diameters, without an evident umbo, with a more reddish shade in the centre, and a paler and more involute edge; the stipe is also slenderer than in *X. olivascens* (Courtecuisse and Duhem 1995; Vizzini et al. 2020). In Italy, *X. donadinii* has so far been reported always on broadleaved trees, only in Emilia-Romagna, Marche, and Puglia (Vizzini et al. 2020).

A.B. De Giuseppe, N.G. Passalacqua, G. Sicoli

## **Lichens**

### ***Bagliettoa baldensis* (A. Massal.) Vězda (Verrucariaceae)**

+ **CAL:** Campotenese, E Monte Pollino (Cosenza) (UTM WGS84 33S 592440.441491), on limestone, c. 1000 m, 30 May 1979, H. Mayrhofer (no. 22024 GZU). – Species confirmed for the flora of Calabria.

This species is widespread in Italy on compact calcareous rocks with optimum in the submediterranean belt (Nimis 2016; Nimis and Martellos 2023). The only previous record in Calabria was from Valle di Fiume Argentino (Halda 2003: 35, sub *Verrucaria baldensis*) but it is currently not included in the Information System on Italian Lichens (Nimis and Martellos 2023).

H. Mayrhofer, O. Breuss

### ***Biatora mendax* Anzi (Ramalinaceae)**

+ **LOM**: Bagni di Masino, Valmasino (Sondrio), on bark of *Fagus sylvatica* L. in an old-growth beech forest (UTM WGS84: 32T 546327.5121063), 1137 m, 23 August 2019, leg. G. Gheza, det. H. Mayrhofer (PAV). – Species confirmed for the flora of Lombardia.

*Biatora mendax* was previously known from Lombardia only from Val di Tartano, Val del Bitto and Val Malenco (Anzi 1862; Anzi Lich. Rar. Lang. Exs. n.168: Printzen 1995); its distribution is now extended to Val Masino. It was collected in a moist, old-growth beech stand (Natura 2000 Habitat 9110 “Luzulo-Fagetum beech forests”), which is coherent with it having its optimum in humid beech forests (Nimis and Martellos 2023). It is listed in the Red List of Italian epiphytic lichens as “endangered” (Nascimbene et al. 2013).

G. Gheza, H. Mayrhofer

### ***Calicium denigratum* (Vain.) Tibell (Caliciaceae)**

+ **ITA (TAA)**: Fiè allo Sciliar (Bolzano) (UTM WS84: 32T 694257.5153572), on bark of *Pinus sylvestris* L., 1165 m, 4 May 2012, J. Nascimbene, conf. J. Hafellner (CLU No. 18230). – Species new for the flora of Trentino-Alto Adige.

*Calicium denigratum* is recognizable for its tall, thin, and shiny black ascomata with stalk of sclerotized, dark, irregularly interwoven hyphae and distinct bell-shaped capitulum lacking pruina. Spores have a coarsely cracked surface (Tibell 1999). The species could be confused with *C. glaucellum* Ach. and *C. abietinum* Pers. but the former has an obovoid to lenticular capitulum with white pruina and coarsely cracked spores with ridges and the latter has a black lenticular capitulum with minutely warted spores. This species occurs in open canopy woodlands in Europe and Siberia (Tibell 1999). It is reported also for North America (McMullin et al. 2012). This is the first Italian collection but the species is likely widespread throughout the Italian Alps.

J. Nascimbene, D. Puntillo

### ***Cladonia trassii* Ahti (Cladoniaceae)**

+ **LOM**: Valle delle Messi, Ponte di Legno (Brescia), on soil above siliceous rock in an open larch stand (UTM WGS84: 32T 614582.5129878), 1686 m, 19 August 2020, G. Gheza (BOLO). – Species confirmed for the flora of Lombardia.

*Cladonia trassii* is an arctic-alpine species with squamulose verticillate podetia with very narrow scyphi (Gheza and Nimis 2021). It was previously recorded in Lombardia only from the high elevation coniferous forests of Mount Confinale, in Valtellina (Anzi 1868, as “*Cladonia stricta*”, cited by Rivellini and Valcuvia Passadore 1996). Our specimen contains fumarprotocetraric acid and atranorin. It was collected together with *Cladonia arbuscula* (Wallr.) Flot., *C. chlorophaea* (Sommerf.) Spreng., *C. furcata* (Huds.) Schrad., *C. macroceras* (Delise) Hav., *C. pleurota* (Flörke) Schaer., and *C. rangiferina* (L.) F.H.Wigg. (Gheza et al. 2022).

G. Gheza, H. Mayrhofer

### ***Coniocarpon fallax* (Ach.) Grube (Arthoniaceae)**

+ **LOM**: road between Dezzo di Scalve and Passo della Presolana, Colere (Bergamo) (UTM WGS84: 32T 585785.5090970), on bark of *Alnus incana* (L.) Moench in a sheltered and moist broadleaved stand within a mixed forest, 810 m, 4 January 2023, G. Gheza (BOLO). – Species confirmed for the flora of Lombardia.

*Coniocarpon fallax* is an oceanic crustose lichen harbouring a trentepohlioid photobiont characterized by the orange-rusty red pruina along the margin of confluent elongated or stellate lirellae (Frisch et al. 2020). The only previous record from Lombardia was from the surroundings of Como by Anzi (1860); a collection by the latter author (*Lich. Min. Rar. Ital.* n. 317) was also reported in Stizenberger (1882) and Sundin and Tehler (1998).

G. Gheza

### ***Lecanora subcarpineae* Szatala (Lecanoraceae)**

+ **SAR**: along a road in pasture, Gavoi (Nuoro) (UTM 32T 515790.4448723), on bark of *Prunus* sp., 850 m, 1 May 2012, J. Malíček (PRA). – Species new for the flora of Sardegna.

*Lecanora subcarpineae* is a crustose epiphytic lichen belonging to the *L. carpineae*-*L. leptyroides* complex, found on smooth and base-rich bark of isolated trees, which occurs also in well-lit or sparse forests, from the mesomediterranean to the subalpine belt. According to Nimis and Martellos (2023), it appears that this species has not always been correctly identified, and the Italian material collected so far would need revision. This sample, collected in the typical habitat of *L. subcarpineae*, is characterized by strongly pruinose apothecia and a distinct Pd+ bright yellow reaction of the apothecial margin, which is coherent with the taxon description.

J. Malíček, S. Ravera

### ***Opegrapha vermicellifera* (Kunze) J.R.Laundon (Opegraphaceae)**

+ **TAA**: Toblino lake, path in the holm oak forest above the lake, eastern slope (Trento) (UTM WGS84 32T: 651601.5102080), on *Quercus ilex* L., 200 m, 25 May 2005, P. Giordani, F. Cristofolini (GE). – Species new for the flora of Trentino-Alto Adige.

*Opegrapha vermicellifera* is a crustose lichen with *Trentepohlia* as photobiont, characterized by a pale grey to whitish thallus. Thalli without apothecia, such as the one found in Toblino, have numerous and prominent pycnidia. In Italy, the species usually colonises large trees in humid areas, especially along large rivers and lakes, both in the Padanian and subalpine regions (Nimis and Martellos 2023) and in the Mediterranean (Nascimbene et al. 2021).

P. Giordani

***Placynthium nigrum* (Huds.) Gray (Placynthiaceae)**

+ **VDA:** Val Veny, Courmayeur (Aosta) (UTM WGS84: 32T 341129.5075178), on north-east facing walls, 1353 m, 8 November 2021, *S. Ongaro, D. Isocrono* (ORO). – Species confirmed for the flora of Valle d’Aosta.

*Placynthium nigrum* is a saxicolous species distinguished by its dark thallus with a conspicuous bluish-black prothallus. Despite being a widespread species in the Alps (Nimis et al. 2018), *P. nigrum* is rarely collected in Valle d’Aosta (Pievittori and Isocrono 1999), where it is uncommon due to the regional climatic and geological characteristics. The only two previous records for this region were from a coniferous forest in Valpelline (Henry 1910), and from broad-leaved forest in Arnad (Henry 1914).

D. Isocrono, S. Ongaro

***Rhizocarpon norvegicum* Räsänen (Rhizocarpaceae)**

+ **LOM:** ridge between Monte Pagano and Monte Pianaccio, Monno (Brescia), on *Pleopsidium flavum* (Trevis.) Körb. on a schist rock outcrop (UTM WGS84: 32T 604483.5123088), 2170 m, 12 August 2022, *G. Gheza* (BOLO); ridge between Monte Tonale Occidentale and Cima Cadi, Ponte di Legno (Brescia), on *P. flavum* on a schist rock outcrop (UTM WGS84: 32T 620459.5126438), 2610 m, 4 June 2022, *G. Gheza* (Herb. Gheza). – Species new for the flora of Lombardia.

*Rhizocarpon norvegicum* is a yellow-greenish crustose chlorolichen, which often starts its life cycle on thalli of Acarosporaceae on schistaceous rocks (Nimis and Martellos 2023). In Italy, it was reported so far only from Friuli Venezia Giulia and Trentino-Alto Adige (Nimis and Martellos 2023), but it is likely more widespread in the Italian Alps.

G. Gheza, J. Nascimbene

***Scytinium fragrans* (Sm.) Otálora, P.M.Jørg. & Wedin (Collemataceae)**

+ **SIC:** Corleone (Palermo), on *Ulmus minor* subsp. *canescens* Bartolucci & Galasso (UTM WGS84: 33S 350022.4180983), 625 m, 28 June 2022, leg. *P. Marino, F.M. Raimondo, S. Ravera*, det. *S. Ravera* (PAL). – Species confirmed for the flora of Sicilia.

*Scytinium fragrans* is a small-foliose to subsquamulose epiphytic cyanolichen occurring in natural or semi-natural habitats, extremely rare or extinct in northern Italy



(Nimis and Martellos 2023), whose last report in Sicilia dates back to the beginning of the 1900s (Jatta 1909–1911).

S. Ravera, P. Marino, F.M. Raimondo

### ***Solenopsora liparina* (Nyl.) Zahlbr. (Leprocaulaceae)**

+ **LOM:** IlGropo, Brallo di Pregola (Pavia), on serpentine outcrops close to the road Casone-Pregola (UTM WGS84: 32T 521498.4955086), 947 m, 17 September 2022, Z. Fačková, L. Paoli (Herb. SAV) – Species new for the flora of Lombardia.

*Solenopsora liparina* has a crustose-placodioid thallus, forming orbicular (up to 2.5 cm) rosettes, with the central parts sometimes falling off and leaving semicircular arcs of olivaceous grey or grey-green lobes, with rounded white-pruinose margins. It is a Mediterranean species colonizing inclined surfaces of ultrabasic rocks such as serpentines, often preferring fissures, in shaded situations also on vertical faces, in habitats with low eutrophication. So far in Italy it was recorded only from Liguria and Toscana (Guttová and Nimis 2021; Nimis and Martellos 2023). The distribution of *S. liparina* along the Apennine Peninsula seems strongly associated to a combination of geology (i.e. presence of serpentine and other ultramafic outcrops) and climate (preference for localities with temperate climate and warm winters) and recently developed habitat suitability maps highlighted higher probabilities of occurrence along the Tyrrhenian side (Guttová et al. 2019).

L. Paoli, Z. Fačková

### ***Stereocaulon incrustatum* Flörke (Stereocaulaceae)**

+ **PIE:** Sesia riverbed near Greggio (Vercelli) (UTM WGS84: 32T 452951.5032689), on siliceous sandy-pebbly soil in a dry grassland colonizing the riverbed, 157 m, 23 December 2020, G. Gheza (PAV). – Species confirmed for the flora of Piemonte.

*Stereocaulon incrustatum* typically grows on mineral oligotrophic soils, often near rivers (Nimis and Martellos 2023). It was reported from “the surroundings of Vercelli” with specimens collected in 1858 (Lamb 1977; Watson 2014; Oset 2015; TSB-25735; E.C.I. n. 194). This record confirms its occurrence in the area near Vercelli after more than 150 years since the latest record, and after more than a century since the last record from the same administrative region (Martel 1911). The analysed specimen contained only atranorin, which is reported as the most widespread chemotype by Oset (2015).

G. Gheza

### ***Verrucaria endocarpoides* Servít (Verrucariaceae)**

+ **MAR:** Badia di S. Pietro, Monte Conero, Sirolo (Ancona) (UTM WGS84: 33T 388234.4822510), in a *Quercus ilex* L. wood on limestone, c. 470 m, 5 June 1979, H. Mayrhofer (GZU no. 22022). – Species new for the flora of Marche.

+ **TOS**: along the lane from Carrara to Campocecina, E of the hill La Pizza, surroundings of Carrara (Massa-Carrara) (UTM WGS84: 32T 588905.4884717), E-facing limestone cliff, c. 800 m, 1 June 1978, *H. Mayrhofer* (GZU no. 22023). – Species new for the flora of Toscana.

*Verrucaria endocarpoides* is one out of many species with a thick, brown, areolate thallus. It is characterized by immersed perithecia with an involucrellum reaching down to exciple-base level, medium-sized ascospores, and stout periphyses. The species is apparently widespread but poorly known. In Italy it was previously reported from Friuli (Breuss 2008a) and from its type locality in Liguria (Breuss 2008b, Servít 1952).

H. Mayrhofer, O. Breuss

***Verrucaria lecideoides* (A.Massal.) Trevis. (Verrucariaceae)**

+ **CAL**: Campotenese, E Monte Pollino (Cosenza) (UTM WGS84 33S 592440.441491), on limestone, c. 1000 m, 30 May 1979, *H. Mayrhofer* (no. 22021 GZU). – Species new for the flora of Calabria.

*Verrucaria lecideoides* belongs to a small group of species with a distinctly areolate, epilithic, autonomous thallus and with perithecia mostly at the margins of, or between the areoles; its perithecia are encircled by an involucrellum that extends down to exciple-base level and incurves beneath. It does not fit within the (core group of) genus *Verruculopsis* that is morphologically different, lacks an involucrellum and is parasitic on Teloschistacean hosts (Navarro-Rosinés et al. 2007). The species is widespread in Italy on exposed calciferous rocks (Nimis 2016; Nimis and Martellos 2023).

O. Breuss, H. Mayrhofer

## Acknowledgements

Sonia Ravera acknowledges the support of NBFC to University of Palermo, funded by the Italian Ministry of University and Research, PNRR, Missione 4 Componente 2, “Dalla ricerca all’impresa”, Investimento 1.4, Project CN00000033.

## References

- Aleffi M, Tacchi R, Poponessi S (2020) New checklist of the bryophytes of Italy. *Cryptogamie, Bryologie* 41: 147–195. <https://doi.org/10.5252/cryptogamie-bryologie2020v41a13>
- Anzi M (1860) *Catalogus Lichenum Quos in Provincia Sondriensi et Circa Novum-Comum Collegit et in Ordinem Systematicum Digessit Presbyter Martinus Anzi*. Tipografia C. Franchi, Como, 126 pp.
- Anzi M (1862) *Manipulus lichenum rariorum vel novorum, quos in Longobardia et Etruria collegit et enumeravit Presb. Martinus Anzi*. *Commentari della Società Crittogamologica Italiana* 1: 130–166.

- Anzi M (1868) *Analecta lichenum rariorum vel novorum Italiae superioris*. Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano 11: 156–181.
- Barbey W (1884) *Florae Sardoae Compendium*. Catalogue raisonné des végétaux observés dans l'île de Sardaigne. George Bridel Editeur, Losanna.
- Bazzichelli G, Abdelahad N (2009) *Algha d'acqua dolce d'Italia*. Flora analitica delle Caroficce. Università degli Studi di Roma La Sapienza – Ministero dell'Ambiente e della Tutela del Territorio e del Mare, Roma, 73 pp.
- Bruss O (2008a) Neue Funde pyrenocarper Flechten aus den Julischen Alpen (Slowenien und Italien). *Herzogia* 21: 85–92.
- Bruss O (2008b) Bemerkungen zu einigen Arten der Flechtengattung *Verrucaria*. *Sauteria* 15: 121–138.
- Brusa G (2019) L'epatica *Phaeoceros carolinianus* (Anthocerotophyta) nuova per l'Italia. Segnalazioni e brevi note. *Pianura* 37: 113–116.
- Cardot J (1892) Monographie des Fontinalacées. Mémoires de la Société des Sciences Naturelles et Mathématiques de Cherbourg 28: 1–152.
- Corillion R, Guerlesquin M (1963) Contribution à l'étude des végétations de Charophycées d'Italie péninsulaire (étage méditerranéen). *Bulletin de la Société scientifique de Bretagne* 38: 193–211.
- Courtecuisse R, Duhem B (1995) *Mushrooms and Toadstools of Britain and Europe*. Harper-Collins Publishers, Ramsbury, Wiltshire, UK.
- Fleischer M (1893) Contribuzioni alla Briologia della Sardegna. *Malpighia* 7: 313–344.
- Frisch A, Moen VS, Grube M, Bendiksby M (2020) Integrative taxonomy confirms three species of *Coniocarpon* (Arthoniaceae) in Norway. *MycKeys* 62: 27–51. <https://doi.org/10.3897/mycokeys.62.48480>
- Gheza G, Nimis PL (2021) Keys to the lichens of Italy – 61) Cladoniaceae (*Cladonia*, *Pilophorus* and *Pycnothelia*). [https://italic.units.it/flora/index.php?procedure=ext\\_key\\_home&key\\_id=3975](https://italic.units.it/flora/index.php?procedure=ext_key_home&key_id=3975)
- Gheza G, Vallese C, Nascimbene J (2022) Enhancing lichen inventories in Italy: new records of *Cladonia*, *Nephroma* and *Peltigera* from the mountains of Lombardia. *Borziana* 3: 5–17. <https://doi.org/10.7320/Borz.003.005>
- Guttová A, Nimis PL (2021) The genus *Solenopsora* (Lichenized Ascomycetes, Leprocaulaceae) in Italy. *Flora Mediterranea* 31: 55–65. <https://doi.org/10.7320/FlMedit31SI.055>
- Guttová A, Fačkovcová Z, Martellos S, Paoli L, Munzi S, Pittao E, Ongaro S (2019) Ecological specialization of lichen congeners with a strong link to Mediterranean-type climate: a case study of the genus *Solenopsora* in the Apennine Peninsula. *Lichenologist* 51: 75–88. <https://doi.org/10.1017/S0024282918000543>
- Halda J (2003) A taxonomic study calcicolous endolithic species of the genus *Verrucaria* (Ascomycotina, Verrucariales) with the lid-like and radiately opening involucrellum. *Acta Musei Richnoviensis Sect. Natur* 10: 1–148.
- Henry JM (1910) Contribution à la lichenologie valdôtaine. *Bulletin de la Société de la Flore Valdôtaine* 6: 8–17.
- Henry JM (1914) Campagne lichénique 1912. *Bulletin de la Société de la Flore Valdôtaine* 9: 29–32.

- Hodgetts N, Lockhart N (2020) Checklist and country status of European bryophytes – update 2020. Irish Wildlife Manuals, No. 123. National Parks and Wildlife Service, Department of Culture, Heritage and the Gaeltacht, Ireland, 214 pp.
- Hodgetts N, Cáliz M, Englefield E, Fettes N, García Criado M, Patin L, Nieto A, Bergamini A, Bisang I, Baisheva E, Campisi P, Cogoni A, Hallingbäck T, Konstantinova N, Lockhart N, Sabovljevic M, Schnyder N, Schröck C, Sérgio C, Sim Sim M, Vrba J, Ferreira CC, Afonina O, Blockeel T, Blom H, Caspari S, Gabriel R, Garcia C, Garilleti R, González Mancebo J, Goldberg I, Hedenäs L, Holyoak D, Hugonnot V, Huttunen S, Ignatov M, Ignatova E, Infante M, Juutinen R, Kiebacher T, Köckinger H, Kučera J, Lönnell N, Lüth M, Martins A, Maslovsky O, Papp B, Porley R, Rothero G, Söderström L, Ștefănuț S, Syrjänen K, Untereiner A, Váňa J, Vanderpoorten A, Vellak K, Aleffi M, Bates J, Bell N, Brugués M, Cronberg N, Denyer J, Duckett J, During HJ, Enroth J, Fedosov V, Flatberg K-I, Ganeva A, Gorski P, Gunnarsson U, Hassel K, Hespánhol H, Hill M, Hodd R, Hylander K, Ingerpuu N, Laaka-Lindberg S, Lara F, Mazimpaka V, Mežaka A, Müller F, Orgaz JD, Patiño J, Pilkington S, Puche F, Ros RM, Rumsey F, Segarra-Moragues JG, Seneca A, Stebel A, Virtanen R, Weibull H, Wilbraham J, Żarnowiec J (2019) A miniature world in decline: European Red List of Mosses, Liverworts and Hornworts. IUCN, Brussels. <https://doi.org/10.2305/IUCN.CH.2019.ERL.2.en>
- Jatta A (1909–1911) Flora italica Cryptogama. Pars III. Lichenes. Tip. Cappelli, Rocca di S. Casciano, 958 pp.
- Lamb IM (1977) A conspectus of the lichen genus *Stereocaulon* (Schreb.) Hoffm. Journal of the Hattori Botanical Laboratory 43: 191–355.
- Martel E (1911) Contribuzione alla lichenologia del Piemonte. Memorie della Reale Accademia delle Scienze di Torino (2) 61: 135–176.
- McMullin RT, Selva SB, Maloles JR, Newmaster SC (2012) *Calicium denigratum* (Vain.) Tibell, a new lichen record for North America. North America Fungi 7: 1–5. <https://doi.org/10.2509/naf2012.007.011>
- Nascimbene J, Gheza G, Hafellner J, Mayrhofer H, Muggia L, Obermayer W, Thor G, Nimis PL (2021) Refining the picture: new records to the lichen biota of Italy. MycoKeys 82: 97–137. <https://doi.org/10.3897/mycokeys.82.69027>
- Nascimbene J, Nimis PL, Ravera S (2013) Evaluating the conservation status of epiphytic lichens of Italy: a red list. Plant Biosystems 147: 898–904. <https://doi.org/10.1080/11263504.2012.748101>
- Navarro-Rosinés P, Roux C, Gueidan C (2007) La genroj *Verrucula* kaj *Verruculopsis* (Verrucariaceae, Verrucariales). Bulletin de la Société Linnéenne de Provence 58: 133–180.
- Nimis PL (2016) The lichens of Italy – A second annotated catalogue. EUT Edizioni Università di Trieste, Trieste, 740 pp.
- Nimis PL, Martellos S (2023) ITALIC – The Information System on Italian Lichens. Version 7.0. University of Trieste, Dept. of Biology. [www.italic.units.it](http://www.italic.units.it) [accessed 9.2.2023]
- Nimis PL, Hafellner J, Roux C, Clerc P, Mayrhofer H, Martellos S, Bilovitz PO (2018) The lichens of the Alps – an annotated checklist. MycoKeys 31: 1–634. <https://doi.org/10.3897/mycokeys.31.23568>

- Onofri S, Bernicchia A, Filipello Marchisio V, Padovan F, Perini C, Ripa C, Savino E, Venturella G, Vizzini A, Zotti M, Zucconi L (2013) Checklist of the macrobasidiomycetes of Italy. <http://dryades.units.it/macrobasidiomiceti/index.php> [accessed 2.3.2022]
- Oset M (2015) The lichen genus *Stereocaulon* (Schreb.) Hoffm. in Poland. A taxonomic and ecological study. *Monographiae Botanicae* 104: 1–81. <https://doi.org/10.5586/mb.2014.001>
- Piervittori R, Isocrono D (1999) I licheni della Valle d'Aosta. I. Museo Regionale di Scienze Naturali di Saint-Pierre, Monografie I. – Valle d'Aosta, 264 pp.
- Printz P (1992) *Inocybe* (Fr.) Fr. In: Hansen L, Knudsen H (Eds) *Nordic Macromycetes* (Vol. 2). Nordsvamp, Copenhagen, Denmark.
- Printzen C (1995) Die Flechtengattung *Biatora* in Europa. *Bibliotheca Lichenologica* 60: 1–275.
- Rivellini G, Valcuvia Passadore M (1996) I licheni appartenenti ai generi *Cladonia* e *Stereocaulon* in provincia di Sondrio (Lombardia, Italia settentrionale). *Il Naturalista Valtellinese. Atti del Museo Civico di Storia Naturale di Morbegno* 7: 3–32.
- Romanov R, Napolitano T, van de Weyer K, Troia A (2019) New records and observations to the Characean flora of Sicily (Italy). *Webbia* 74: 111–119. <https://doi.org/10.1080/00837792.2019.1609258>
- Servít M (1952) Neue und weniger bekannte Arten der Familie Verrucariaceae und Dermocarpaceae. *Preslia* 24: 347–390.
- Stizenberger E (1882) *Lichenes Helvetici eorumque stationes et distributio*. Jahresbericht der St. Gallischen Naturwissenschaftlichen Gesellschaft 22: 225–522.
- Sundin R, Tehler A (1998) Phylogenetic studies of the genus *Arthonia*. *The Lichenologist* 30: 381–413. <https://doi.org/10.1006/lich.1998.0155>
- Tibell L (1999) *Caliciales Nordic Lichen Flora* 1. Bohuslän, Uddevalla.
- Vizzini A, Consiglio G, Marchetti M, Alvarado P (2020) Insights into the *Tricholomatineae* (Agaricales, Agaricomycetes): a new arrangement of *Biannulariaceae* and *Callistosporium*, *Callistosporiaceae* fam. nov., *Xerophorus* stat. nov., and *Pleurocollybia* incorporated into *Callistosporium*. *Fungal Diversity* 101: 211–259. <https://doi.org/10.1007/s13225-020-00441-x>
- Watson P (2014) *Birmingham Botany Collections – Lichens*. Birmingham Museums, Birmingham, 186 pp.