

INVITED REVIEW

Mediterranean springs: Keystone ecosystems and biodiversity refugia threatened by global change

M. Fernández-Martínez^{1,2,3}  | J. Barquín⁴  | N. Bonada^{5,6}  | M. Cantonati⁷  |
 C. Churro^{8,9}  | J. Corbera²  | C. Delgado¹⁰  | M. Dulsat-Masvidal¹¹  | G. Garcia¹²  |
 O. Margalef^{1,13}  | R. Pascual¹²  | J. Peñuelas^{1,14}  | C. Preece¹⁵  | F. Sabater^{1,2,3}  |
 H. Seiler¹⁶ | J. M. Zamora-Marín¹⁷  | E. Romero^{1,3} 

Correspondence

Marcos Fernández-Martínez and Estela Romero, CREAL, Campus de Bellaterra (UAB), Cerdanyola del Vallès, Spain.
 Email: m.fernandez@creaf.uab.cat and estela.romero@creaf.uab.cat

Funding information

European Research Council, Grant/Award Number: ERC-StG-2022-101076740; Generalitat de Catalunya, Grant/Award Number: SGR2021-01333; Ministerio de Ciencia e Innovación, Grant/Award Number: PID2021-128778OA-I00 and RYC2021-031511-I; Spanish Research Agency, Grant/Award Number: CEX2018-000828-S; Spanish Ministry of Science and Innovation; European Union NextGeneration, Grant/Award Number: FJC2021-046923-I

Abstract

Mediterranean spring ecosystems are unique habitats at the interface between surface water and groundwater. These ecosystems support a remarkable array of biodiversity and provide important ecological functions and ecosystem services. Spring ecosystems are influenced by abiotic, biotic, and anthropogenic factors such as the lithology of their draining aquifers, their climate, and the land use of their recharge area, all of which affect the water chemistry of the aquifer and the spring discharges. One of the most relevant characteristics of spring ecosystems is the temporal stability of environmental conditions, including physicochemical features of the spring water, across seasons and years. This stability allows a wide range of species to benefit from these ecosystems (particularly during dry periods), fostering an unusually high number of endemic species. However, global change poses important threats to these freshwater ecosystems. Changes in temperature, evapotranspiration, and precipitation patterns can alter the water balance and chemistry of spring water. Eutrophication due to agricultural practices and emergent pollutants, such as pharmaceuticals, personal care products, and pesticides, is also a growing concern for the preservation of spring biodiversity. Here, we provide a synthesis of the main characteristics and functioning of Mediterranean spring ecosystems. We then describe their ecological value and biodiversity patterns and highlight the main risks these ecosystems face. Moreover, we identify existing knowledge gaps to guide future research in order to fully uncover the hidden biodiversity within these habitats and understand the main drivers that govern them. Finally, we provide a brief summary of recommended actions that should be taken to effectively manage and preserve Mediterranean spring ecosystems for future generations. Even though studies on Mediterranean spring ecosystems are still scarce, our review shows there are sufficient data to conclude that their future viability as functional ecosystems is under severe threat.

For Affiliation refer page on 15

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Global Change Biology* published by John Wiley & Sons Ltd.

KEYWORDS

biodiversity conservation, freshwater ecosystems, groundwater, semiarid regions, sources

1 | INTRODUCTION

The Mediterranean climate is characterized by warm, dry summers, mild winters, and a high degree of temporal variability, both within (i.e., seasonally) and between years (Rundel et al., 2016). Worldwide, Mediterranean climate regions are among the ecosystems most threatened by global change due to the expected decreases in water availability and intensive human impacts (Cramer et al., 2018; Mekonnen & Hoekstra, 2016). These regions often align with arid or semiarid conditions and can be found around the Mediterranean basin, California, central Chile, the Western Cape Region (South Africa), and southwestern Australia (Blondel & Aronson, 1999; Rundel et al., 2016). The characteristics of these regions make their springs, and spring ecosystems, of paramount importance for human development and biodiversity conservation. Spring ecosystems from these regions (hereafter, Mediterranean spring ecosystems) have been described as refugia for a wide array of organisms, including a large number of endemic species found only in one single spring (Fensham, Ponder, et al., 2023; Freyhof et al., 2017). Spring ecosystems have been reported to support high values of both freshwater (Bes et al., 2018; Cantonati, Füreder, et al., 2012; Fernández-Martínez et al., 2021; Sabatino et al., 2003) and terrestrial biodiversity (Votto et al., 2020; Zamora-Marín, Zamora-López, et al., 2021).

Mediterranean springs can be natural (lacking any human intervention) or seminatural (i.e., human modified), the latter consisting of a small construction, usually of local rock and concrete, with a spout that drains the water from the aquifer to a sink or cistern, maintaining a pool of water before it runs off. These ecosystems are usually small in size (sometimes even less than 1 m²) but nonetheless complex, representing ecotones linking surface and underground water bodies (Cantonati & Ortler, 1998). Spring ecosystems are characterized by a mosaic of numerous microhabitats (Illies & Botosaneanu, 1963), and such features could explain the high number of taxa coexisting in just a few square meters (Lencioni et al., 2018; Mezquita et al., 2000). This spatial heterogeneity also influences the distribution of plant and animal species, the interactions between species, and the trophic structure of biological communities. Even though spring ecosystems and their biodiversity are threatened worldwide, those from Mediterranean-type regions suffer the consequences of global change more strongly because their intrinsic water limitation exacerbates other potential problems, such as eutrophication (Fernández-Martínez et al., 2020; Martín et al., 2024). Furthermore, it is in these regions where spring ecosystems play an important role in sustaining biodiversity because they represent isolated habitats with permanent water availability surrounded by habitats where water is not typically available (Fernández-Martínez, Berloso, et al., 2019; Zamora-Marín, Ilg, et al., 2021).

Spring ecosystems are also of great interest from the viewpoint of freshwater quality monitoring because their particular biodiversity and limited size make them extremely sensitive to any disturbance

(Cantonati et al., 2006; Lencioni et al., 2018; Williams, 1991), thus allowing the effects of different environmental stressors (e.g., water pollution or groundwater depletion) to be assessed within river basins. Pollution from diffuse sources is difficult to track—particularly in aquifers—but this is essential for assessing nutrient legacy effects and safeguarding freshwater resources (Goyette et al., 2018; Van Meter et al., 2018).

Currently, the main threats to the conservation of spring ecosystems (Stevens et al., 2021) can be summarized as: (i) habitat modification for anthropogenic uses, including the overexploitation of water from the source or its related aquifer (Barquín & Scarsbrook, 2008); (ii) eutrophication, mainly related to the disruption of regional and global N and P biogeochemical cycles due to intensive livestock farming and agriculture, as well as leakages from urbanized areas (Fernández-Martínez et al., 2020; Martín et al., 2024); and (iii) climate change, through several processes: (a) Increases in temperature lead to increases in evapotranspiration (ETR) by the vegetation, which has largely expanded in many regions of the planet, reducing water in aquifers (Gallart & Llorens, 2003; García-Ruiz et al., 2011); (b) an increase in torrentiality also reduces infiltration and groundwater recharge; and (c) if total precipitation also decreases (−5% to −30% is expected for the Mediterranean basin, Cramer et al., 2018), this will also reduce the water available for aquifers and springs. In fact, water demand for human and industrial uses is expected to increase in the near future in Mediterranean climate regions as a consequence of global warming (Mekonnen & Hoekstra, 2016), exacerbating the effects of all these threats on spring ecosystems.


Hence, this review aims to first provide a synthesis of the functioning of Mediterranean spring ecosystems, focusing on their main features, geomorphology, and the determinants of their water chemistry. Second, we synthesize the current knowledge regarding general biodiversity patterns in Mediterranean spring ecosystems. Third, current and future threats to these vulnerable ecosystems are discussed. Finally, we provide a roadmap for enhancing spring conservation and management efforts, offering practical recommendations to safeguard spring ecosystem integrity.

2 | FUNCTIONING AND GEOLOGICAL FEATURES OF SPRING ECOSYSTEMS

2.1 | Hydrogeological features

Springs constitute discrete points of groundwater discharge in the landscape (van der Kamp, 1995), and their characteristics are determined by geomorphological (relief) and geological (lithology, tectonics) attributes (Figure 1). For example, high reliefs favor the presence of springs because water discharge can occur when the topography intersects the water table (depression springs). Faults also play an

FIGURE 1 Spring hydrogeological characteristics, processes, and physical and chemical features of groundwater. On the left, in blue, are the processes dominating the water chemistry of the aquifer. OC, organic carbon.



important role in the occurrence of springs: They displace materials and generate conduits that allow groundwater to flow. Contact or lateral springs, which occur because of the lateral contact between the aquifer and a poorly permeable unit, are also often related to faults, as are most thermal springs (i.e., those that discharge water at a temperature higher than the average air temperature). Geological characteristics also determine the difference between the elevation or pressure of the hydraulic head in the aquifer and the discharge, resulting in either gravity (descending) or artesian (ascending) springs.

Springs located in karst systems, or in extrusive volcanic rocks, show larger drainage flows compared to other lithologies (e.g., granites). In particular, karstified rocks rapidly transmit a large percentage of precipitation events as newly infiltrated water. Karst aquifers have primary porosity, common rock discontinuities such as fractures or secondary porosity, and enlarged voids, such as channels and waterways, developed from initial discontinuities (Kresic, 2010). The drainage or catchment areas of karst springs often extend beyond topographic divides. This is why karst aquifers give rise to the largest springs in the world. Given their strong dependence on hydrogeological features, the density of springs can vary widely across geographical areas (see [Supporting Information](#), Section S1; [Table S1](#)).

2.2 | Water chemistry

The chemistry of spring water is mainly determined by the natural characteristics of the parental aquifer and the organic and inorganic processes that occur along the aquifer (Figure 1). The concentration of ions in water depends on the solubilization and re-precipitation of bedrock minerals and is thus related to the composition of the rocks. For example, it has been shown that the Ca/Mg ratio in spring water is indicative of the type of carbonate rock (dolomite vs. limestone) that forms the aquifer (Cantonati et al., 2007). The hydrogeological properties of the parent material (porosity, hydraulic conductivity, transmissivity) determine the length of flow paths, the volume

of aquifer recharge and the residence time of the water, and these properties influence ion exchange and biogeochemical transformations (Welch & Allen, 2014). Longer residence time in the aquifer, often accompanied by increased weathering, leads to an increase in the ion content of the water and determines factors such as alkalinity, and sulfate and chloride ratios (Morell, 1992; Roca, 1990), all of them very influential for spring flora and fauna. In deeper aquifer springs, the water chemistry is highly dependent on temperature and pressure conditions. Water from such springs is usually rich in CO₂, Fe, and H₂S (Lachassagne et al., 2021), leading to naturally occurring sparkling water springs.

Climate characteristics also influence spring water chemistry. Temperature affects weathering rates: the higher the temperature, the higher the reactivity and the greater the weathering. Precipitation may also facilitate weathering by accelerating chemical reactions when water is flowing through dry soils, but it eventually tends to dilute the concentration of ions in the water. Proximity to the coast mainly increases the concentration of Na⁺ and Cl⁻ due to continuous deposition of sea spray (Fernández-Martínez, Margalef, et al., 2019). On top of lithology and climate, vegetation and soil properties can also affect the water chemistry of the aquifer. Vegetated and well-structured soils favor water infiltration, and as it flows through the soil, the water is enriched in organic matter. Additionally, soil properties can also buffer the pH of the infiltrated water. Connectivity between shallow and deep soil layers due to biological factors such as root structure and root exudation, or soil fauna and the microbiome may also interact with the chemistry of the infiltrated water (Neary et al., 2009; Scanlon et al., 2005). The importance of these factors affecting the chemistry of spring water, however, remains unknown.

2.3 | Hydrogeomorphology

Two zones are usually distinguished in springs: the eucrenon (i.e., the source area) and the hypocrenon (i.e., spring brooks running down

from the outlet) (Illies & Botosaneanu, 1963), but the threshold between both is hard to delimit in Mediterranean springs because it varies widely throughout the different periods (i.e., wet and dry seasons). Especially during the dry season, the extension of the spring outlets (eucrenon) is vague because the running flow is significantly reduced. Temperature differences and bryophyte coverage have been some criteria suggested to establish the extension limits of the eucrenon (Crema et al., 1996; Erman & Erman, 1995; Illies, 1961; von Fumetti et al., 2007), but there is still no general consensus. Alternatively, the hydrogeomorphological approach is particularly suitable for defining the limits of Mediterranean springs. It has been long recognized that springs are composed of a particular mosaic of microhabitats, each one characterized by size, water dynamics and type of substratum (Gerecke et al., 1998; Myers & Resh, 2002; Stoch et al., 2011; White & Pickett, 1985). Consequently, we could define the spring limits as the border of the area filled by these eucrenon microhabitats. The latter can be identified by the influence of the spring water on the community of hygrophytes (i.e., aquatic or semi-aquatic organisms). This approach was proposed by Stevens et al. (2011) and is easily applicable to Mediterranean environments because of the typical lack of hygrophytes outside of the spring domain, unless the spring is located on a riverbank.

Typically, Mediterranean springs from arid and semiarid areas (annual precipitation below 500 mm per year) have low-discharge flow, their influence area (eucrenic or crenic ecosystem) is relatively small (usually in the metric or decametric range), and the hypocrenon is poorly developed or nonexistent. Consequently, in these springs, the most frequent situation is the lack of connection between springs and the river network, except for short periods of heavy rainfall. In these semiarid regions, springs are typically isolated freshwater habitats that predominantly act as biodiversity refuges during drought periods, when the fluvial network is dry. In contrast, during hyper-humid periods (which may occur only a few days a year), they can aid in the dispersion of organisms that live in the spring, thus contributing to maintaining biological communities in adjacent rivers and freshwater habitats.

2.4 | Spring classifications

Springs are usually classified according to their morphology, size, or hydrological regime (Stevens et al., 2021). The first spring classification systems (Steinmann et al., 1915; Thienemann, 1926) distinguished three main types according to hydromorphological criteria: (1) *rheocrenes* or flowing springs, in which water flow feeds small streams; (2) *limnocrenes* or pool springs, where the flow is low and creates lentic habitats; and (3) *helocrenes* or seepages that generate shallow or soggy wet zones. This basic classification remains accepted today, but modern approaches also distinguish them according to their geology, hydrology, water chemistry, water temperature, ecology, and human use (Glazier, 2014). For instance, Stevens et al. (2021) used site-specific source geomorphology to define 13 types of springs. Cantonati (2022), instead, used a multicriteria

approach to distinguish seven macrotypes of springs. The high spring diversity resulting from the great variety of geological and morphological features in regions with Mediterranean climate implies that almost all types of both Stevens' and Cantonati's classification are present (except for geysers and desert springs).

However, most Mediterranean springs have been *domesticated*, so their current status is seminatural. In fact, a great proportion of small springs, the most widespread representatives (Cantonati, 2022), which had been originally helocrenes or rheo-helocrenes, have been transformed to rheocrenes by digging a hole in the ground to reach the subterranean aquifer, or through water mines or wells to concentrate the flow through a spout, a pipe, or a channel. In addition, other artificial elements are sometimes present, such as cisterns or small ditches to transport the water. Nevertheless, unless water abstraction is very high, these artificial elements do not generate strong negative impacts on the biodiversity of springs and they can still support relevant populations of threatened species (e.g., when a sink is created, it can be used by amphibians for reproduction) (Buono et al., 2019; Egea-Serrano, Oliva-Paterna, Tejado, et al., 2006; Egea-Serrano, Oliva-Paterna, & Torralva, 2006).

3 | BIODIVERSITY OF MEDITERRANEAN SPRING ECOSYSTEMS

3.1 | General patterns of spring biodiversity

The characteristics of spring ecosystems are determinant in providing refugia or structuring freshwater biotic assemblages in the landscape. When the springs are stable and isolated, the environmental conditions can favor the development of a highly specialized flora and fauna. These specialized spring-dwelling organisms are often called *crenobionts* and correspond to species that have necessarily adapted to these spring habitats to survive and complete their life cycles. Some authors also consider springs as potential refuges for relict species (Botosaneanu, 1995; Cantonati, Füreder, et al., 2012; Taxböck et al., 2017). In contrast, in densely populated landscapes with heavily exploited aquifers, most of these species have already disappeared.


The biodiversity of spring ecosystems is strongly linked to the ecohydrogeological characteristics of the aquifer (Cantonati, Stevens, et al., 2020). The climate, hydrostructure, and lithology of the aquifers determine whether a spring has permanent or intermittent flow (Cantonati, Segadelli, et al., 2020, see also Section 2). In some regions, springs fed by large karst systems become even more vital as stable hydrological refuges for biodiversity as climate change advances (Cartwright et al., 2020). For example, larger and more permanent springs might have a higher contribution as freshwater refuges in arid and semiarid regions (Davis et al., 2013; Work, 2023), while the role of smaller and more temporary springs might be critical as stepping stones in metapopulation and metacommunity dynamics (Collier & Smith, 2006), with an influence on alpha, beta, and gamma diversity. Data from a survey of springs in the NE Iberian Peninsula

using bryophytes (Fernández-Martínez et al., 2021) were used here to calculate diversity measures through the R packages *vegan* and *betapart* (Baselga, 2010; Oksanen et al., 2018; R Core Team, 2023). The results confirm that alpha and gamma diversity are higher in springs under wetter climates compared to those from drier climates (Figure 2a). Beta diversity, instead, is higher among springs located in drier regions, since they are generally more isolated environments as compared to springs from more humid regions. Hence, the relative contribution of species turnover to their community composition is also higher compared to that in wetter areas (Figure 2b,c). This highlights the high degree of uniqueness of spring community composition in drier areas. In this regard, the degree of connectivity or isolation of a spring habitat to other surface waters could be determinant to prevent the colonization of invasive species or to conserve their unique biota.

Despite most Mediterranean springs being small in size, the structure of microhabitats can always be recognized. Biological richness (or biodiversity) in springs has been shown to increase with the diversity of hydrogeomorphological microhabitats (Reiss

& Chiffard, 2015; Taxböck et al., 2020). Distinct organisms occupy different microhabitats of the crenic environment as depicted in Figure 3. Those organisms living in the actual source of spring water are referred to as crenobionts (e.g., springsnails), while those being associated with the spring sink during a given phase of their life cycle are called crenophiles (e.g., amphibians or emerging insects). Several plant and animal species (e.g., ferns or terrestrial vertebrates) that do not inhabit spring ecosystems (crenoxens) may also routinely benefit from many ecological functions provided by springs, such as food, habitat, or water supply (Zamora-Marín et al., 2023). With regard to the microhabitat selected within the springscape (i.e., the natural spatial unit of springs encompassing abiotic and biotic factors), species may be classified as stygobionts when they inhabit the groundwater ecosystem, stygophiles when they mostly inhabit the spring sink, and rheophiles when they use spring downstream waters. This water continuum from aquifer to spring and to streams may be severed in periods of intense drought. Therefore, the degree of endemism is more likely to occur among stygophile and crenophile taxa, often mites and mollusks (see Section 3.3), than among

FIGURE 2 Alpha, beta, and gamma diversity of bryophytes in 58 and 59 springs (~1 m²) for the Littoral mountain range and the Catalan Pyrenees regions, respectively. Temperature and water availability differ between these two regions, with the Littoral mountain range being warmer and drier than the Pyrenees (respectively [mean ± standard deviation], mean annual temperature: 14.0 ± 0.8, 11.7 ± 2.0°C; annual precipitation: 726 ± 67, 1060 ± 64 mm year⁻¹). Panel (a) shows alpha diversity, with the inset stating the gamma diversity in the two regions. Springs from more arid regions are more important as biodiversity refugia. Panel (b) shows total beta diversity and panel (c) shows the different components of beta diversity. B_{SOR} stands for Sorensen's beta diversity (total beta diversity), B_{SIM} stands for Simpsons' beta diversity (species turnover), and B_{NES} stands for nestedness ($B_{SOR} - B_{SIM}$). These analyses were performed using freely available data published in Fernández-Martínez et al. (2021) and the experimental design can be found therein. Calculations of diversity were performed using packages *vegan* and *betapart* in R (Baselga, 2010; Oksanen et al., 2018; R Core Team, 2023). See Data Availability Statement to access data and code to generate these results.






FIGURE 3 Conceptual models representing the geomorphological structure of a typical spring ecosystem (a) and the distribution of different spring-associated biotic groups across the springscape (b). Panel (a) shows the main geomorphological elements composing a Mediterranean spring ecosystem, and terms for spring-associated biodiversity are also depicted. Panel (b) displays the distribution of typical spring-associated biotic groups in relation to their spring dependence degree and the expected endemism ratio.

rheophile taxa which are much more connected to downstream waters. Overall, the list of taxa that has been described to inhabit spring ecosystems is very large (Figure 4; Table S2; Sections 3.2 and 3.3), even though they have received far less attention compared to many other freshwater ecosystems.

3.2 | Plants, algae, and microorganisms

Autotrophs form the basis of the food webs in spring ecosystems as primary producers (Cantonati et al., 2015; Cantonati, Rott, et al., 2012). Vascular plants, bryophytes, algae, and microbial mats can all be found in Mediterranean spring ecosystems and provide important functions, such as stabilizing soils, retaining water, and offering food and shelter to aquatic invertebrates (Cantonati, Füreder, et al., 2012; Žutinić et al., 2018). Spring vegetation contributes significantly to habitat structure and plays a central role in various physicochemical processes (e.g., limestone precipitation). Generally, European springs are known to harbor high plant diversity (e.g., see Cantonati & Ortler, 1998; Seiler et al., 2021), including numerous

specialized and rare plant species, especially bryophytes (Cantonati et al., 2006; Geissler, 1976). Spring bryophyte communities can also sustain large communities of associated microflora (e.g., diatoms, green algae) and fauna (e.g., nematodes, tardigrades, and arthropods, Cantonati, 2022; see Section 3.3). However, in contrast to other regions (e.g., the Alps), there are very few studies on the vegetation of Mediterranean spring ecosystems. For example, Pascual et al. (2020) found similar plant species richness in springs in the mountains of Mallorca and Catalan pre-littoral mountains compared to springs in the Alps.

Further studies with bryophytes carried out in Catalonia (NW Mediterranean) indicate that these habitats play an important role in semiarid regions as shelters where water is constantly available, representing important biodiversity refuges for these species (Bes et al., 2018; Fernández-Martínez, Berloso, et al., 2019). Some typical plant species associated with Mediterranean spring ecosystems are currently considered near-threatened or endangered in Europe (e.g., Christenhusz et al., 2017; de Foucault, 2015), as well as many other narrow-ranging or endemic plant and animal species (Deleuil, 1974; Jiménez-Alfaro et al., 2013; Jiménez-Mejías &

FIGURE 4 Simplified representation of the Mediterranean spring biodiversity, showing most of the taxonomic groups related to this ecosystem and their microhabitat preferences in a seminatural spring. 1, Odonata, *Ischnura graellsii*; 2, Diptera, *Culex*; 3, Rotifera, *Cephalodella forficula*; 4, Ostracoda, *Notodromas*; 5, Acari, Hydrachnidia; 6, Tardigrada; 7, Bacillariophyceae, *Navicula*; 8, Cyanobacteria, *Phormidium*; 9, Nematoda; 10, Aves, *Turdus merula*; 11, Magnoliophyta, *Carex pendula*; 12, Mammalia, *Capreolus*; 13, Bryophyta, *Apopellia endiviifolia*; 14, Bryophyta, *Eucladium verticillatum*; 15, Fungi, *Scutellinia* sp.; 16, Pteridophyta, *Adiantum capillus-veneris*; 17, Amphibia, *Salamandra*; 18, Copepoda, *Cyclops*; 19, Cladocera, *Daphnia*; 20, Amphipoda, *Echinogammarus*; 21, Pteridophyta, *Equisetum telmateia*; 22, Amphibia, *Alytes obstetricans*; 23 Amphibia (larva), *Alytes obstetricans*; 24, Gastropoda, *Physa*; 25, Odonata (nymph), *Chalcolestes viridis*; 26, Lepidoptera, *Celastrina argiolus*; 27, Magnoliophyta, *Carex remota*.



Luceño, 2009; Rivas-Martínez et al., 2011). Although only a few lichen species are known from freshwater habitats, they may offer valuable information about the quality and integrity of spring ecosystems (Cantonati, 2022). There are indications that Mediterranean freshwater habitats contain lichen communities that differ in their taxonomic and functional composition from other European regions and likely host yet-undiscovered species (Nascimbene et al., 2023), which deserves further research.

Diatoms and cyanobacteria (accompanied by bacteria and fungi) are the dominant autotrophs found in microbial mats (Cantonati et al., 2015; Kolda et al., 2019). Diatom diversity has been shown to increase from 0 to 25°C and starts to decrease at temperatures above 25 or 30°C, mainly in thermal springs (Dallas, 2008; Delgado et al., 2020; Kishi et al., 2005; Patrick et al., 1969). Hence, climate warming will likely change diatom communities of springs. Springs are indeed important habitats for diatoms with unique environmental requirements, and this may explain why new species are regularly discovered in spring ecosystems (Cantonati, Angeli, et al., 2016;

Cantonati & Lange-Bertalot, 2006; Delgado et al., 2013; Sabater & Roca, 1990). From the 22 taxonomic orders of Cyanobacteria, only four (Oscillatoriales, Chroococcales, Leptolyngbyales and Gomontiellales) are mainly present in freshwater springs. Many of the identified taxa, however, comprise a plethora of poorly reported and uncommon species (Cantonati, 2008; Cantonati, Rott, et al., 2012; Nowicka-Krawczyk & Żelazna-Wieczorek, 2013, 2017). Unlike hot springs where cyanobacteria diversity is well characterized, little is known about species composition in ambient springs (Cantonati et al., 2015; Jasser et al., 2022).

3.3 | Animals: Aquatic and terrestrial fauna

Mediterranean spring ecosystems show an extremely high environmental heterogeneity, promoting the uniqueness of individual spring assemblages (Cantonati, Segadelli, et al., 2020) and leading to increased values of faunistic dissimilarity (i.e., higher beta

diversity mediated by greater contribution of species turnover as compared to more humid environments; see Figure 2) and ultimately enabling higher levels of gamma diversity (Pascual et al., 2020; Pešić et al., 2017; Płóciennik et al., 2016). Mediterranean spring ecosystems support outstanding ratios of endemic and rare freshwater fauna as a result of their role as biodiversity paleoreugia, but also provide key ecological functions to associated terrestrial species and adjacent ecosystems. Even though previous studies on animal communities in Mediterranean spring ecosystems are notably scarce, the available information points toward disproportionately high values of richness for several animal groups (Fensham et al., 2011; Rossini et al., 2018), including spring-dwelling and spring-associated taxa.

Several genera of spring-dwelling invertebrates evolved strong ecological links to spring habitats, such as in the case of water mites (Hydrachnidia), rotifers and springsnails (Cantonati, Segadelli, et al., 2020; Miracle et al., 1995), all being considered representative components of spring ecosystems and showing high levels of spring-specific endemism. For instance, Italian spring ecosystems host up to 18 endemic species of water mites and provide habitat for a total of 163 water mite species, which represent about 43% of the total water mite diversity recorded in Italian inland waters (Sabatino et al., 2003). Likewise, Mediterranean spring ecosystems placed in the eastern Iberian Peninsula support a non-negligible proportion (32%) of the total diversity of non-marine ostracods inhabiting Iberian inland waters (Mezquita et al., 1999). High values of endemism have also been found in outcrop and discharge springs from the Eastern Lake Eyre Basin (Australia), which support at least 18 endemic mollusk species (Fensham et al., 2011), most of them belonging to a single genus (*Jardinella*). In turn, these spring ecosystems support a higher number of endemic and narrow-ranging mollusk species as compared to other freshwater habitats within the same region. In this context, three new genera of gastropods have also been recently described as endemic from Tunisian spring ecosystems (Khaloufi et al., 2020), and two new species of ostracods have recently been found in Italian spring ecosystems (Rossetti et al., 2022). Also among crustaceans, the amphipod genus *Echinogammarus* is a paradigmatic case. It has more than 20 species inhabiting springs and upper zones of small rivers in the Mediterranean Basin. The distribution of most of these species is very restricted, and a few of them are only found in a single location (Pinkster, 1993). However, the highest values of spring-specific species endemism worldwide are found in Australian springs from the Great Artesian Basin, with at least 51 mollusk and 24 crustacean species known as being endemic from this spring biodiversity hotspot (Rossini et al., 2018).

Although the vast majority of mollusks inhabiting spring ecosystems correspond to snails (Gastropoda), some clam species (e.g., taxa from the Sphaeriidae or Lasaeidae families) may also occur in these singular environments and even show clear spring-habitat preferences (Rassam et al., 2021; Rossini et al., 2018). In addition to passively dispersing organisms (e.g., crustaceans, mollusks, and water mites), Mediterranean spring ecosystems also host a high diversity of actively dispersing invertebrates and even comprise the main habitat for several threatened taxa, such as some conservation concern and

spring-dependent Odonata species (Assandri et al., 2020; Vilenica et al., 2021). In fact, about 60% of the total Odonata species occurring in Montenegro have been recorded at spring ecosystems (Pešić et al., 2017). In the same country, spring surveys targeting water bugs (Heteroptera) found 13 new species previously unrecorded at national scale (Gligorović et al., 2016). In line with the above studies, spring ecosystems from the Rif region (North of Morocco) have been identified as priority conservation habitats to ensure the effective protection of threatened water beetle species (Bennas et al., 2009).

Mediterranean spring ecosystems, and those from more arid regions, also support vertebrate species including endemic or conservation concern species. For instance, outcrop and discharge springs from the Eastern Lake Eyre Basin (Australia) host three fish species which are endemic from a single site or highly concentrated to a local area (Fensham et al., 2011). Likewise, the Australian Artesian springs harbor up to eight fish species endemic to single spring sites or confined to extremely restricted areas (Rossini et al., 2018). Spring ecosystems from some Mediterranean countries also support endemic or narrow-ranging fish species, mostly killifishes (i.e., genus *Aphanius*). For instance, *Aphanius sirhani* is endemic to the Azraq Oasis in Jordan, whereas *A. kruppi*, *A. dispar*, and *A. richardsoni* are highly concentrated to spring ecosystems in Oman, Sinai, and Israel, respectively (Freyhof et al., 2017). Mediterranean spring ecosystems also comprise critical habitats for amphibians and several aquatic reptiles, as populations of many species appear highly concentrated to springscapes or are even spring dependent. For instance, threatened populations of the Betic midwife toad (*Alytes dickhilleni*) and the fire salamander (*Salamandra salamandra morenica*) from the south of the Iberian Peninsula mostly rely on spring-fed drinking troughs and artificial pools as breeding habitats (Egea-Serrano, Oliva-Paterna, Tejedo, et al., 2006; Egea-Serrano, Oliva-Paterna, & Torralva, 2006). Springs also deliver important ecological services to terrestrial wildlife and may largely contribute to ecosystem functioning in adjacent landscapes. In this context, Mediterranean spring-fed small water bodies act as key landscape elements for about 70% of the breeding bird species inhabiting adjacent ecosystems (Zamora-Marín et al., 2022), performing pivotal cross-system functions such as the provision of drinking, bathing, and foraging sites (Zamora-Marín et al., 2024). In fact, spring ecosystems may play a key role in nutrient exchange with the adjacent terrestrial ecosystems through the provision of emerging insects (e.g., in the form of chironomids or mayflies) to both aerial (e.g., bats and swifts) and terrestrial (e.g., spiders and lizards) consumers, as reported for other small water bodies (Lewis-Phillips et al., 2020). However, quantitative assessments have not yet been conducted in spring ecosystems.

3.4 | Temporal dynamics of biodiversity in Mediterranean springs

As highlighted in the previous sections, the importance of perennial springs as biodiversity refuges for Mediterranean flora and fauna is largely due to the fact that they are shelters where water




FIGURE 5 Hypothetical temporal dynamics (i.e., seasonal and interannual) of biodiversity in perennial and temporary spring ecosystems under the effects of global change (e.g., drought). Spring ecosystems from more mesic climates (i.e., temperate) are more likely to be better connected than those from arid regions, which are more isolated and may represent an evolutionary refuge for endemic species. The increase/drop in alpha diversity is only shown in Mediterranean climates to highlight that seasonal and interannual hydrological differences are more pronounced in these regions. This potential fluctuation in alpha diversity is also related to differences in connectivity, which in Mediterranean regions lies between that of mesic and arid regions. In mesic climates, even if the springs are temporary, the dry phase is usually short and there is high connectivity in the hydrological network, so we would expect variations in alpha diversity to be low. In arid climates, weather is predominantly dry, perennial springs have a high degree of isolation, and the hydrological disconnection is even greater in temporary springs, but again, we would expect low seasonal and interannual differences.

is constantly available in the midst of often dry environments. However, temporary springs are also common in the Mediterranean as a result of the high seasonality and interannual variability imposed by the climate (Bonada & Resh, 2013). The effect of global change, as described below (Section 4), could displace the environmental conditions of mesic (i.e., temperate) springs (and their surrounding regions) into Mediterranean-like or arid conditions (Figure 5). Similarly, perennial springs under any type of climate may become temporary due to the effects of global change. We predict that those changes could have profound impacts on spring biodiversity. As previously shown (Section 3.1; Figure 2), spring ecosystems from wetter regions tend to have higher diversity per spring (alpha diversity) and a higher pool of species (gamma diversity) because springs have a higher connectivity locally (e.g., downstream reaches) and regionally (e.g., a larger set of potential habitats and larger colonizable areas for crenophillic biodiversity). Hence, we would expect alpha and gamma diversity to decrease as we move from mesic, to Mediterranean and arid climates, as the latter ones are much more isolated (i.e., drying results in a decrease of richness and a loss of connectivity) (Soria et al., 2017).

In mesic springs, changes in gamma diversity should not be evident when perennial springs become temporary, as other surrounding habitats with available water should still preserve the overall pool of species. Nonetheless, given the high number of

spring endemisms, the disappearance of species cannot be disregarded. Alpha diversity (at the spring scale), however, should be more sensitive to shifts in the environmental conditions. While mesic spring ecosystems should keep their alpha diversity more or less steady with time, Mediterranean springs should experience variations in their alpha diversity due to the occurrence of drought periods, either seasonal (i.e., summer droughts) or meteorological (i.e., prolonged periods of unusual drought conditions). In perennial Mediterranean springs, drought periods can actually increase alpha diversity. This is because organisms that typically do not rely on spring ecosystems, both spring-dwelling (e.g., those depending on the springs, such as invertebrates and amphibians) and spring-subsidized species (e.g., those that visit them, such as mammals and birds), may be drawn to these springs in order to survive. Conversely, in temporary Mediterranean springs, drought periods are likely to coincide with the dry phase when the springs have the potential to run dry. This is expected to reduce alpha diversity because several species that rely on these springs may temporarily disappear. In mesic climates, dry periods may not be long enough to significantly impact the alpha diversity of temporary springs in most cases. Even if the spring does not flow, ponds and high humidity may persist, providing sustenance for all species. However, in arid climates, the dry conditions may be the norm. In such environments, alpha diversity will likely only decrease when a spring

becomes temporary. The extended dry phases in arid climates require all present species to be adapted to tolerate the lack of water, preventing a significant decrease in alpha diversity. In arid regions, springs function as oases and are highly isolated habitats regionally, with low alpha and gamma diversity, and a high number of endemic species (Souza et al., 2006). Therefore, the importance of springs as refuges increases as we move away from mesic conditions, where they act as ecological refuges, to more arid conditions where they act as evolutionary refuges (hence the large number of endemisms, see sections above).

4 | CURRENT AND FUTURE RISKS OF SPRING ECOSYSTEMS

4.1 | Reduction of water availability

In Mediterranean regions, climate change is expected to severely affect freshwater availability (Ali et al., 2022; Cramer et al., 2018), due to more frequent, longer and intense droughts and heat waves (Hartmann et al., 2014), leading to drier conditions in the coming years (Dubrovský et al., 2014). Heat waves and droughts will reduce water availability in aquifers and springs through various mechanisms: (i) lack of precipitation and therefore direct declines in soil water content and groundwater (Barros et al., 2014); (ii) higher temperatures, with higher ETR rates by vegetation; (iii) higher water demand for agriculture and livestock, as well as for the population, and thus increased groundwater abstraction (Sivelle et al., 2021). Together with longer drought and the general decrease in precipitation, future scenarios indicate a trend toward extreme precipitation events and flash floods (Dubrovský et al., 2014; Llasat et al., 2016), which would further decrease infiltration rates and limit groundwater recharge (Hartmann et al., 2014; Nerantzaki & Nikolaidis, 2020).

In addition, climate change interacts with other environmental issues that can exacerbate water scarcity in spring ecosystems, such as increased urbanization and land use changes. Increased urbanization entails an increase in impermeable surfaces and reduced infiltration rates. The increase in forests following cropland abandonment has increased ETR and reduced water availability, with numerous examples from European Mediterranean countries (Gallart & Llorens, 2003; García-Ruiz et al., 2011; Teuling et al., 2019) and other arid and semiarid regions (Farley et al., 2005; Filoso et al., 2017). Degraded and compacted soils, along with desertification, can also lead to lower infiltration rates and a decrease in the water recharge of the aquifers (Ferreira et al., 2022).

A clear example of how global change may affect spring water quantity can be found in a recent regional study (Figure 6). In January–February 2013, we measured water runoff from a total of 60 springs across the Littoral mountain range (NE of the Iberian Peninsula). Ten years later, in January–February 2023, we resampled 31 of them (selected randomly) and found an overall 92% reduction in water runoff, with 45% of the springs being completely dry. These




FIGURE 6 Water flow differences (10-year period) recorded at 31 springs located in the Littoral mountain range, in Catalonia. More than 45% of the springs surveyed in 2013 had dried up by 2023, and a reduction in 92% of total flow was observed.

changes are fully consistent with changes in piezometric levels from four wells located around the same area, recording a reduction of 1.1 ± 0.4 m in the water table from January 2013 to January 2023 ($p = .001$; Figure S1). The most likely factors affecting the decrease in water runoff of these springs are: (1) lack of management, (2) a 2-year persistent drought in the region, accounting for around 30% less precipitation during the 2021–2022 period, compared to the 2011–2012 period, and (3) an increase in mean temperature of 0.6°C during the latter period compared to the first one. In this area, monthly and annual real ETR derived from MODIS (product MOD16A2GF, Running et al., 2019) is positively related to precipitation and temperature (Figure 7). We found that for every increase of 1°C , ETR increases from $3.8\% \pm 0.4\%$ during the spring season, to $15.1\% \pm 0.9\%$ in winter, with an annual average of $7.2\% \pm 1.6\%$ increase per $^\circ\text{C}$. Since 2003 (when data were available), there have been considerable increases in both mean annual temperature ($0.49 \pm 0.11^\circ\text{C}$ per decade, $p < .001$) and ETR ($6.8 \pm 1.0\%$ per decade, $p < .001$). In this case study, an increase in water abstraction from these springs seems very unlikely because they are mainly located in mountainous regions with very little human influence. The data presented here clearly show that Mediterranean spring ecosystems




FIGURE 7 Partial residual plots showing effect of temperature and precipitation on real evapotranspiration (ETR) of the studied region (Littoral mountain range, Catalonia). Panel (a) shows a positive but saturating relationship between monthly ETR and precipitation (MAP, here precipitation of the last 12 months). Panel (b) shows how the effect of monthly temperature on monthly ETR depends on the season. Panels (c, d) show, respectively, the effect of mean annual temperature, and precipitation, on annual ETR. Finally, panel (e) shows the temporal trend of ETR from 2003 to 2022. During this same period, MAT increased by $0.49 \pm 0.07^\circ\text{C}$ per decade ($p < .001$, weather data from the town of Mataró $41^\circ33'23''\text{N}$ to $2^\circ26'10''\text{E}$: www.meteomataro.com). We used ETR data from MODIS [product MOD16A2GF (Running et al., 2019)] for central coordinates for the two main massifs of the study region (Cèllec and Montnegre, respective coordinates: 41.67831N , 2.57941E ; and 41.56768N , 2.36892E) covering an area of $20.5 \times 20.5\text{ km}$. Results from panels (a, b) were obtained using a mixed effects model in which the response variable was monthly ETR (ln-transformed) and the fixed predictors were mean monthly temperature, season, their interaction, and monthly precipitation of the last 12 months (ln-transformed). We included the massif as the random factor (Cèllec or Montnegre) and an AR1 (auto-regressive function for lag 1) temporal autocorrelation structure. Results in panels (c, d) followed the same approach for annual values, but excluding the season. Results in panel (e) followed the same procedure but including only year as a fixed predictor. Shaded areas represent 95% confidence intervals. See the data availability statement to access data and code to reproduce these results.

are under threat due to global change. If the climate continues to become warmer and drier, as expected, many of these ecosystems will likely be lost in the near future.

Moreover, Mediterranean springs will become increasingly disconnected from downstream areas, which will shift from perennial to intermittent conditions. This change will lead to

substantial alterations in biological communities. While this shift will impact spring biodiversity and its temporal dynamics (as depicted in Figure 5), it will also serve to reduce the risk of downstream colonization by non-native species (Costello et al., 2011). Additionally, a decrease in water quantity will affect the water quality of aquifers and springs. This reduction diminishes the potential for pollutant dilution, as discussed in Section 4.2 (e.g., Mas-Pla & Menció, 2019; Van Vliet et al., 2017). The higher concentrations of pollutants in spring water are likely to result in a reduction in spring biodiversity (Fernández-Martínez et al., 2020).

4.2 | Water quality

4.2.1 | Eutrophication

Intensive agricultural practices have relied for decades on the over-fertilization of crops with synthetic fertilizers and manure, and much of the excess nutrients applied are carried by runoff into surface watercourses and through infiltrated water into aquifers (Galloway et al., 2008; Sutton et al., 2011). Intensive livestock farming is also responsible for much of the N and P leaching into freshwater systems (Bouwman et al., 2013). Once produced, the removal of nutrient pollution (mainly nitrates) in groundwater is difficult to achieve (Mas-Pla & Menció, 2019), and given that the residence time in aquifers can be long, sometimes the pollution detected in springs lags years behind the time when pollution occurred (Alley et al., 2002).

Intensive agricultural practices have already compromised groundwater quality in Mediterranean areas of southern Europe and North African countries (Morocco, Algeria, Tunisia). In Spain, Italy, France, Greece, Croatia and Malta, both nitrogen surpluses and high abstraction pressures have been identified (European Environment Agency, 2022; Psomas et al., 2021), and many of their aquifers are in nitrate vulnerable zones. High concentrations of nutrients in the groundwater that feed springs can lead to eutrophication and drastic changes in the flora and fauna of the spring ecosystems (Fernández-Martínez et al., 2020; Jacoby et al., 2008; Martín et al., 2024). Some studies have reported plant community shifts from adapted submerged aquatic plants to systems dominated by benthic and attached filamentous algae, while some others link elevated nitrate concentrations with alterations in the reproductive functioning of amphibians and fish (Jacoby et al., 2008).

4.2.2 | Wastewater

Human activities, such as intensive livestock farming and wastewater discharges, have also contributed to the degradation of the microbiological quality of groundwater, aquifers, and springs (Savio et al., 2018; Stevens et al., 2022; Takuissu et al., 2022). Fecal contamination infiltrates soils and reaches groundwater and aquifers, even at great depths. The extent of contamination depends on several factors, including soil type, vegetation, terrain and bedrock fracture,

precipitation, and land use (Aqso et al., 2014; Davidson et al., 2016; Murphy et al., 2017; Savio et al., 2018). Recent studies have indicated a general underestimation of aquifer contamination through enlarged cracks and fissures (Hartmann et al., 2021). Pathogenic bacteria, such as *Escherichia coli* and other coliforms, and enteric viruses like *Norovirus*, *Rotavirus*, hepatitis A and E viruses, as well as protozoa like *Cryptosporidium* and *Giardia*, are increasing, compromising springs as secure freshwater resources (An & Breindenbach, 2005; Aqso et al., 2014; Blanco et al., 2017; Davidson et al., 2016; Murphy et al., 2017; Ryu et al., 2019; Takuissu et al., 2022).

4.2.3 | Emergent pollutants: Synthetic chemicals, drugs, and microplastics

Soils act as a sink of organic pollutants; however, organic contaminants can also leach from soils to groundwater (Burri et al., 2019), especially in karst systems (Chen et al., 2022). Diffuse pollution from agriculture is the most common pressure to the quality of groundwater, affecting 19% of the total groundwater body area in the EU (European Environment Agency, 2022). Pesticides can migrate through soil or surface water to streams and groundwater, where they can have unintended ecological impacts such as accumulation in aquatic organisms and loss of ecosystem biodiversity (McGinley et al., 2023). Among pesticides, triazine herbicides have been highlighted as being one of the most widespread pesticides in European groundwaters (Loos et al., 2010), and the most frequently detected herbicides in water resources in Spain (Herrero-Hernández et al., 2017; Llamas et al., 2022). The impact of pesticides on groundwater can last for long periods due to the slow renewal rate of groundwater and the slow release of retained pesticides and their metabolites in the soil (Baran et al., 2022).

Pharmaceuticals may enter the aqueous environment through anthropogenic activities such as sewage discharge, livestock, fertilizing and landfill leachate, resulting in their presence in groundwater at high concentration levels of ngL^{-1} to μgL^{-1} (Sui et al., 2015). Recently, awareness has grown about the presence of antibiotic residues in groundwaters related to the presence of antibiotic-resistant bacteria (ARB) (Andrade et al., 2020; Kaiser et al., 2022; Kampouris et al., 2022; Singh et al., 2022). The incorrect usage and overuse of antibiotics in human and veterinary health treatments has created an anthropogenic selective pressure on environmental bacteria, and the emergence of multidrug-resistant “superbugs” is now a reality (Andrade et al., 2020; Huang et al., 2023). A recent review by Andrade et al. (2020) states that groundwater represents a major global reservoir for ARB. Several recent studies have reported high amounts of ARB, antibiotic-resistant genes, and mobile genetic elements in groundwater and springs (Gros et al., 2023; Huang et al., 2023; Kaiser et al., 2022; Kampouris et al., 2022).

Nonsteroidal anti-inflammatory drugs (NSAIDs) are other pharmaceuticals frequently detected in water resources due to their widespread use. The main source of contamination is excretion

through urine and feces in wastewater, which contain unmetabolized or conjugated and transformed forms (Singh et al., 2014). Jurado et al. (2021) studied 13 groundwater samples in an urban aquifer in Barcelona and concluded that individual NSAIDs and their mixtures posed minimal risk to human health. However, the risk posed by the sum of different pharmaceuticals that can simultaneously be detected in groundwater remains elusive. Additionally, polycyclic aromatic hydrocarbons (PAHs) are concerning compounds for wildlife and human health due to their carcinogenic, mutagenic, and endocrine disruptor potential (Wallace et al., 2020). Forest fires are a well-known natural source of PAHs that play an important role in their distribution in both terrestrial and aquatic compartments (Campos & Abrantes, 2021). Mansilha et al. (2019) studied the impact of wildfires and posterior precipitation to water from springs and streams from Caramulo Mountain (Portugal) indicating that burnt areas presented from 1.2 to 4 times higher concentrations of PAHs compared to unburned areas.

Among emerging pollutants, microplastics (MPs, plastic particles smaller than 5 mm) are becoming important due to their persistence in aquatic environments (Viaroli et al., 2022). The penetration of MPs through pores and fractures, as well as interaction with colloidal aggregates, can partially affect the dynamics of MPs in the subsoil, making their detection difficult in groundwater systems and springs. MPs in groundwater may have several sources, including the atmosphere, the interaction with surface water bodies, urban infrastructures, or agricultural soils (Viaroli et al., 2022). The characterization and quantification of MPs in groundwater are challenging due to the wide variety of sizes and compositions (Alfonso et al., 2021). Some studies focused on aquifer and spring water MP pollution have been performed in the USA, Europe, and South Africa (Minténig et al., 2018; Panno et al., 2019). Despite the low number of studies, most of them have found the presence of MPs in groundwater (Boyle & Örmeci, 2020), highlighting the fact that MP contamination of groundwater is a problem that deserves further attention.

4.3 | Biological invasions

The introduction of alien species is an ongoing and widespread conservation issue for spring ecosystems, especially because of the constant conditions of these freshwater habitats (Fensham et al., 2011; Kodric-Brown & Brown, 2007). Although the magnitude of this threat has not yet been quantified for these freshwater environments, several alien plant and animal species have been widely introduced in Mediterranean springs, where they became established and now maintain self-sustaining populations (e.g., Aksu et al., 2021; Emiroğlu et al., 2016; Saber et al., 2022).

For instance, invasive diatoms have been introduced into an oasis mineral spring in Egypt, with potential impacts on native diatom assemblages (Saber et al., 2022). The occurrence of alien plant species in springs has been associated with increased anthropogenic disturbance, ultimately being correlated with decreased richness of native flora (Nielson et al., 2019). Cases of animal introductions

in Mediterranean spring ecosystems are comparatively much more abundant, particularly concerning exotic fish and crayfish (Kodric-Brown & Brown, 2007). Mediterranean spring ecosystems are particularly prone to biological invasions due to the scarce representation of native spring-dwelling predators (e.g., fish) and the high popularity of springs among local people, which acts as an intentional (e.g., releases for aesthetic or recreational purposes) or unintentional vector for alien species introduction. In fact, the establishment of alien species in spring ecosystems likely involves greater impacts on native spring-dwelling biodiversity as compared to other larger freshwater ecosystems, because their limited size and delicate ecological balance magnify the ecological impacts of introduced alien species. Conversely, the high degree of isolation that characterizes Mediterranean spring ecosystems could reduce the propagule pressure of alien species and the invasion risk through natural spread mechanisms (Stevens et al., 2021).

4.4 | Changes in local culture and traditions: Spring abandonment

In Mediterranean regions, the presence of natural springs has been a key part of the culture since the earliest civilizations. The location of the Acropolis in Athens was likely selected not only for its defensive capacity but also because an aquifer and a spring were located there, and the aqueduct system of Ancient Rome was predominantly supplied with water taken from springs (Mays et al., 2007). The influence of springs in Mediterranean culture is also evident in numerous ancient literary works, often highlighting their spiritual and biodiversity importance as shown in the book “Llibre de meravelles” by Ramón Llull (1287–1289 AD), in a fragment that translates as: “The philosopher was sitting under a tree, full of leaves and flowers. The tree was watered by a spring, in which there were lots of sweetly singing birds. The philosopher was contemplating the greatness and goodness of God in the arrangement of the tree, the spring and the birds” (see original text in Section S2 of Supporting Information). This short text exemplifies how deeply ingrained these ecosystems are within Mediterranean culture.

Large technological advances and important lifestyle changes occurred during the 20th century in many countries, leading to the abandonment of rural areas and increasing human population density in cities. This has consequently led to the abandonment of traditional practices as well, even though they may have been crucial for human livelihoods just a few decades before. Some of these activities were strongly related to spring ecosystems, particularly in Mediterranean and arid regions. The management of human-modified springs for providing both humans and livestock with drinking water, but also for supporting irrigated crops and other uses, has sharply decreased in just a few decades. The abandonment of traditional croplands or farmhouses has been directly related to the loss of human-modified springs and their related ecosystems. For instance, the decline of livestock transhumance and other regimes of extensive pastoralism across the Mediterranean basin is leading

to the abandonment and consequent collapse of spring-fed drinking troughs (Manenti et al., 2017), particularly in the Iberian southeast (Zamora-Marin et al., 2022).

In addition to land abandonment, changes in lifestyle and beliefs have further affected the conservation of springs. Religious celebrations associated with springs went from being very popular and common to becoming almost completely extinct. Since the second half of the 19th century, spring-mediated tourism to drink naturally occurring sparkling water, which was believed to have curative properties, was also very popular among wealthy people. Again, social changes have almost completely put an end to both activities and left spring-based health resorts and their saints as anecdotes from old times, thus causing the progressive loss of springs caused by vegetation encroachment. These cultural manifestations, based on traditions, legends, customs, and habits occurring in the surroundings of springs, have been previously referred to as “spring culture” (Fernández-Martínez et al., 2017). Losing these springs implies not only losing the biodiversity linked to those ecosystems, but sometimes also losing ancient local knowledge and rich cultural legacy often only preserved through oral transmission (Table S3 provides a list of cultural events linked to spring ecosystems in Mediterranean and arid regions around the world).

In spite of the general trend of spring cultural abandonment, new social trends are emerging with an increasing interest in protecting nature, including movements in several Mediterranean regions to study and protect spring ecosystems. In Catalonia, several local groups have engaged different projects to study and protect springs (e.g., *Grup de Fonts d'Argentona*; *Projecte Fonts* by several of the signing authors; *Grup de Defensa del Ter*; *Fonts del Montseny*; or the *CercaFonts*, a mobile app with information on 5000 springs). In Andalusia, the *Asociación Proyecto Conoce Tus Fuentes* manages the first participatory online catalog of springs in Andalusia, with more than 13,000 springs inventoried to date. Likewise, in Mallorca (Balearic Islands), an exhaustive survey of springs was performed across the Tramuntana mountain range. In Castilla-La Mancha, the project *Apadrina una Fuente* aims to find, signpost, and restore all the springs in the municipality of Cifuentes (whose name means “one hundred springs”). Some other initiatives of international relevance have emerged in order to put springs and their conservation into the spotlight, such as the Fellowship of the Spring (Fensham, Adinehvand, et al., 2023), or the Springs Stewardship Institute. Whether the spring culture will remain alive for future generations or not depends mostly on our ability to preserve spring ecosystems.

5 | CONSERVATION AND MANAGEMENT

Most Mediterranean spring ecosystems are managed in order to provide water for human activities, thus mostly ignoring the conservation of their biodiversity and ecological functions. Hence, it should be a top priority for biodiversity conservation to preserve, as intact as possible, those few natural springs that have not yet been disturbed. However, as mentioned above, most Mediterranean

spring ecosystems are now seminatural ecosystems. In these cases, managers should oversee a trade-off between conserving or restoring (fully naturalizing) those ecosystems. In cases where the spring remains in active use, we strongly recommend maintaining its current utilization. This approach is advisable because the existing biota should already be in equilibrium with the spring's current conditions, as well as the social services these ecosystems offer (see Section 4.4). Preserving springs that are still in use can be achieved at a very low cost, and the benefits for biodiversity conservation can be substantial, particularly in arid regions and in springs inhabited by endemic species. However, if the spring is no longer in use or has dried up, certain actions should be taken: (i) To restore the spring to a previous state: This involves rejuvenating the spring to a condition where it can sustain a more or less constant flow of water. Sometimes, issues like collapsed water mines or obstructed pipes due to plant roots can be addressed to revive spring biodiversity. (ii) Fully re-naturalize the spring: In this approach, any artificial construction or modifications are removed to return the spring to its natural state. The choice between these two options depends on the specific circumstances. The first option is more likely to successfully recover the spring ecosystem. However, the latter solution may not be practical for preserving spring ecosystems that have been significantly altered by human intervention, especially those fed through water mines, because restoring them to a state before any human intervention would imply irreversibly losing any potential water flow. Nonetheless, in humid regions, the second option could be a valuable approach in order to increase the number of natural springs. Furthermore, in human-modified springs, conservation actions should also promote environmental heterogeneity at spring scale (i.e., microhabitats), thus boosting diversified biotic communities (Zamora-Marin et al., 2024). For that purpose, management actions could be focused on promoting adjacent terrestrial vegetation, allowing the occurrence of ground-level puddles from water leakages or overflow, and providing structural refuges (e.g., stone walls) for crenophilic wildlife, among others. In this regard, regional authorities should provide management guidelines to promote a more nature-friendly management from stakeholders and spring users, some of which are currently subjected to extensive livestock or farming purposes.

Legislation to protect spring ecosystems, however, is somewhat scarce. The Habitats Directive of the EU, which forms part of the legal basis for the Natura 2000 Network, includes in its list of protected habitats only limestone-precipitating springs, leading to a situation in which many spring habitats remain unprotected (Cantonati, Segadelli, et al., 2016). For example, Spampinato et al. (2023) listed soft water springs and crenel brooks of southern Italy as habitats deserving protection but neglected by the Habitats Directive. Indirect protection of spring ecosystems may be afforded by the protection of, for instance, calcareous spring mires and other wetlands, given that these habitats often include springs within their heterogeneous structure. However, although spring ecosystems are meaningful for regional biodiversity and threatened in many regions by human activity, few nations bear a legal framework explicitly protecting spring

habitats. Comprehensive legislation for the conservation of spring ecosystems worldwide is urgently required (Cantonati et al., 2021).

An important step toward the large-scale protection of the biodiversity of springs is the comprehensive research on their ecological integrity. Stevens et al. (2022) found in their global review that studies of spring ecosystems are sparse, erratic, and widely scattered. Contributing scientists of Mediterranean biogeographical regions generally reported both insufficient data on springs as well as high estimated endangerment levels (e.g., for the Iberian Peninsula, South Africa, Greece, and Morocco). National or regional spring studies such as those underway in Andalusia and Catalonia (Section 4.4) can help to close this gap.

6 | CONCLUSIONS

Spring ecosystems are interface habitats between subterranean and surface freshwater environments, and they harbor unique biodiversity favored by both surface and groundwater organism populations providing important ecological functions. However, they probably constitute one of the rarest, most fragile ecosystems threatened by the effects of global change and the overexploitation of water, which might impair their capability to function as “time capsules of biodiversity” (Beasley-Hall et al., 2023). This is particularly critical for Mediterranean springs where the combination of natural (drying) and anthropogenic disturbances jeopardize their biodiversity and the ecological functions and ecosystem services they provide. Given that future scenarios for Mediterranean climate regions suggest an increase in aridity, it is imperative to prioritize the conservation of Mediterranean springs and develop specific management plans for these ecosystems. Our review demonstrates the urgent need for immediate action to preserve these ecosystems ensuring the retention or restoration of their ecological and cultural value.

AUTHOR CONTRIBUTIONS

Marcos Fernández-Martínez: Conceptualization; data curation; formal analysis; funding acquisition; investigation; methodology; project administration; resources; software; supervision; validation; visualization; writing – original draft; writing – review and editing. **José Barquín:** Conceptualization; investigation; methodology; validation; visualization; writing – original draft; writing – review and editing. **Nuria Bonada:** Data curation; investigation; methodology; validation; visualization; writing – original draft; writing – review and editing. **Marco Cantonati:** Conceptualization; investigation; methodology; validation; visualization; writing – original draft; writing – review and editing. **Catarina Churro:** Conceptualization; investigation; methodology; validation; visualization; writing – original draft; writing – review and editing. **Jordi Corbera:** Conceptualization; data curation; formal analysis; investigation; methodology; resources; validation; visualization; writing – original draft; writing – review and editing. **Cristina Delgado:** Conceptualization; investigation; methodology; validation; visualization; writing – original draft; writing – review and editing. **Maria Dulsat-Masvidal:** Conceptualization;

investigation; methodology; validation; visualization; writing – original draft; writing – review and editing. **Guillermo Garcia:** Conceptualization; investigation; methodology; validation; visualization; writing – original draft; writing – review and editing. **Olga Margalef:** Conceptualization; investigation; methodology; validation; visualization; writing – original draft; writing – review and editing. **Roger Pascual:** Conceptualization; investigation; methodology; validation; visualization; writing – original draft; writing – review and editing. **Josep Peñuelas:** Conceptualization; investigation; methodology; validation; visualization; writing – original draft; writing – review and editing. **Catherine Preece:** Conceptualization; investigation; methodology; resources; validation; visualization; writing – original draft; writing – review and editing. **Francesc Sabater:** Conceptualization; investigation; methodology; resources; validation; visualization; writing – original draft; writing – review and editing. **Hallie Seiler:** Conceptualization; investigation; methodology; validation; visualization; writing – original draft; writing – review and editing. **José Manuel Zamora-Marín:** Conceptualization; investigation; methodology; validation; visualization; writing – original draft; writing – review and editing. **Estela Romero:** Conceptualization; data curation; funding acquisition; investigation; methodology; project administration; resources; supervision; validation; visualization; writing – original draft; writing – review and editing.

AFFILIATIONS

¹CREAF, Campus de Bellaterra (UAB), Cerdanyola del Vallès, Spain

²Delegació de la Serralada Litoral Central – ICHN, Mataró, Spain

³Department of Evolutionary Biology, Ecology and Environmental Sciences (BEECA-UB), University of Barcelona, Barcelona, Spain

⁴Instituto de Hidráulica Ambiental de la Universidad de Cantabria (IHCantabria), Santander, Spain

⁵Freshwater Ecology, Hydrology and Management Research Group (FEHM), Department of Evolutionary Biology, Ecology and Environmental Sciences, University of Barcelona, Barcelona, Spain

⁶Institut de Recerca de la Biodiversitat (IRBio), University of Barcelona, Barcelona, Spain

⁷BIOME Lab, Department of Biological, Geological and Environmental Sciences – BiGeA, Alma Mater Studiorum – University of Bologna, Bologna, Italy

⁸Laboratory of Virology and Molecular Biology and Laboratory of Phytoplankton, Department of the Sea and Marine Resources, Portuguese Institute for the Sea and Atmosphere (IPMA), Lisbon, Portugal

⁹Blue Biotechnology and Ecotoxicology (BBE), CIIMAR – Centro Interdisciplinar de Investigação Marinha e Ambiental, Universidade do Porto, Matosinhos, Portugal

¹⁰Departamento de Ecología e Biología Animal, Facultade de Ciencias, Universidade de Vigo, Vigo, Spain

¹¹IDAEA-CSIC, Institute of Environmental Assessment and Water Research, Barcelona, Spain

¹²BioSciCat, The Catalan Society of Sciences for the Conservation of Biodiversity, Tarragona, Spain

¹³Departament de Dinàmica de la Terra i de l'Oceà, GRC RISKINAT, UB-Geomodels, Facultat de Ciències de la Terra, University of Barcelona, Barcelona, Spain

¹⁴CSIC, Global Ecology Unit, CREAM-CSIC-UAB, Barcelona, Spain

¹⁵Institute of Agrifood Research and Technology (IRTA), Sustainability in Biosystems Programme, Barcelona, Spain

¹⁶Vegetation Ecology, Institute of Natural Resource Sciences (IUNR), Zurich University of Applied Sciences (ZHAW), Wädenswil, Switzerland

¹⁷Department of Applied Biology, Centro de Investigación e Innovación Agroalimentaria (CIAGRO-UMH), Miguel Hernández University of Elche, Elche, Spain

ACKNOWLEDGEMENTS

This review was written thanks to the funding provided by the CREAM's Severo Ochoa CEX2018-000828-S Synthesis actions program. This research was supported by the Catalan government project SGR2021-01333, the European Research Council project ERC-StG-2022-101076740 STOIKOS and the Spanish MCIN project KALORET (PID2021-128778OA-I00). M.F.-M. was supported by a Ramón y Cajal fellowship (RYC2021-031511-I) funded by the Spanish Ministry of Science and Innovation, the NextGenerationEU program of the European Union, the Spanish plan of recovery, transformation and resilience, and the Spanish Research Agency. E.R. was supported by the Severo Ochoa Excellence Program (CEX2018-000828-S) of the Spanish Research Agency. J.M.Z.-M. was supported by a postdoctoral grant funded by the Spanish Ministry of Science and Innovation and the European Union NextGeneration EU/PRTR (FJC2021-046923-I). We acknowledge the Institució Catalana d'Història Natural (ICHN) and the Secció de Ciències Biològiques de l'Institut d'Estudis Catalans (IEC) for additional funding for studying Mediterranean springs. We thank Jordi Gomis (www.meteomataro.com) for providing publicly available climate data for the region of study. We finally thank the Direcció de Cultura, Ajuntament de Mataró for providing the venue to hold the workshop that promoted this review. We also thank Zeynep Ersoy, Ricardo Figueroa, Verónica Ferreira, Roderick J. Fensham (The Fellowship of the Spring), Manuel Graça, Vincent H. Resh, Boudjéma Samraoui, Farrah Samraoui, and Nikos Skoulidakis for providing invaluable information regarding cultural activities and traditions related to springs.

CONFLICT OF INTEREST STATEMENT

All signing authors declare that they have no conflicts of interest.

DATA AVAILABILITY STATEMENT

Data and code to perform analyses presented in this study can be openly found at Figshare: <https://doi.org/10.6084/m9.figshare.23220998.v1>, reference number 23220998.

ORCID

M. Fernández-Martínez  <https://orcid.org/0000-0002-5661-3610>
 J. Barquín  <https://orcid.org/0000-0003-1897-2636>
 N. Bonada  <https://orcid.org/0000-0002-2983-3335>
 M. Cantonati  <https://orcid.org/0000-0003-0179-3842>
 C. Churro  <https://orcid.org/0000-0003-2890-4338>
 J. Corbera  <https://orcid.org/0000-0003-3583-3929>
 C. Delgado  <https://orcid.org/0000-0003-3954-0146>
 M. Dulsat-Masvidal  <https://orcid.org/0000-0002-7677-1333>
 O. Margalef  <https://orcid.org/0000-0002-3036-3182>
 R. Pascual  <https://orcid.org/0000-0001-9937-9644>
 J. Peñuelas  <https://orcid.org/0000-0002-7215-0150>
 C. Preece  <https://orcid.org/0000-0001-6584-3541>
 F. Sabater  <https://orcid.org/0000-0001-6767-231X>
 J. M. Zamora-Marín  <https://orcid.org/0000-0002-7021-267X>
 E. Romero  <https://orcid.org/0000-0003-3115-7572>

REFERENCES

- Aksu, S., Başkurt, S., Emiroğlu, Ö., & Tarkan, A. S. (2021). Establishment and range expansion of non-native fish species facilitated by hot springs: The case study from the Upper Sakarya Basin (NW, Turkey). *Oceanological and Hydrobiological Studies*, 50(3), 247–258.
- Alfonso, M. B., Arias, A. H., Ronda, A. C., & Piccolo, M. C. (2021). Continental microplastics: Presence, features, and environmental transport pathways. *Science of the Total Environment*, 799, 149447. <https://doi.org/10.1016/j.scitotenv.2021.149447>
- Ali, E., Cramer, W., Carnicer, J., Georgopoulou, E., Hilmi, N. J. M., Le Cozannet, G., & Lionello, P. (2022). *Cross-chapter. Paper 4: Mediterranean region. U: Climate change 2022: Impacts, adaptation and vulnerability*. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change (H.-O. Pörtner, D. C. Roberts, M. Tignor, E. S. Poloczanska, K. Mintenbeck, A. Alegria, M. Craig, S. Langsdorf, S. Lösckhe, V. Möller, A. Okem, & B. Rama, Eds.). IPCC.
- Alley, W. M., Healy, R. W., LaBaugh, J. W., & Reilly, T. E. (2002). Flow and storage in groundwater systems. *Science*, 296(5575), 1985–1990. <https://doi.org/10.1126/science.1067123>
- An, Y. J., & Breindenbach, G. P. (2005). Monitoring *E. coli* and total coliforms in natural spring water as related to recreational mountain areas. *Environmental Monitoring and Assessment*, 102(1), 131–137. <https://doi.org/10.1007/s10661-005-4691-9>
- Andrade, L., Kelly, M., Hynds, P., Weatherill, J., Majury, A., & O'Dwyer, J. (2020). Groundwater resources as a global reservoir for antimicrobial-resistant bacteria. *Water Research*, 170, 115360. <https://doi.org/10.1016/j.watres.2019.115360>
- Aqso, A., Özkul, K., & Karakaya, H. (2014). An investigation on the bacterial contents of natural springs in a rural area of the middle Black Sea region in Turkey. *Applied Ecology and Environmental Sciences*, 2(5), 123–129. <https://doi.org/10.12691/aees-2-5-3>
- Assandri, G., Bazzi, G., Maggioni, D., Galimberti, A., & Kunz, B. (2020). Distribution, autecology, genetic characterization, and conservation of the Western Mediterranean endemic dragonfly *Orthetrum nitidinerve* (Selys, 1841): Insights from Italy. *International Journal of Odonatology*, 23(4), 405–422. <https://doi.org/10.1080/13887890.2020.1828194>
- Baran, N., Rosenbom, A. E., Kozel, R., & Lapworth, D. (2022). Pesticides and their metabolites in European groundwater: Comparing regulations and approaches to monitoring in France, Denmark, England and Switzerland. *Science of the Total Environment*, 842, 156696. <https://doi.org/10.1016/j.scitotenv.2022.156696>
- Barquín, J., & Scarsbrook, M. (2008). Management and conservation strategies for coldwater springs. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 18(5), 580–591. <https://doi.org/10.1002/aqc.884>
- Barros, V. R., Field, C. B., Dokken, D. J., Mastrandrea, M. D., Mach, K. J., Bilir, T. E., Chatterjee, M., Ebi, K. L., Estrada, Y. O., & Genova, R. C. (2014). *Climate change 2014: Impacts, adaptation, and vulnerability. Part A: Global and sectoral aspects* (pp. 35–94). Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press.
- Baselga, A. (2010). Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography*, 19(1), 134–143. <https://doi.org/10.1111/j.1466-8238.2009.00490.x>
- Beasley-Hall, P. G., Murphy, N. P., King, R. A., White, N. E., Hedges, B. A., Cooper, S. J. B., Austin, A. D., & Guzik, M. T. (2023). Time capsules of biodiversity: Future research directions for groundwater-dependent ecosystems of the Great Artesian Basin. *Frontiers in Environmental Science*, 10, 1021987. <https://doi.org/10.3389/fenvs.2022.1021987>
- Bennas, N., Sánchez-Fernández, D., Abellán, P., & Millán, A. (2009). Analyse de la vulnérabilité des coléoptères aquatiques dans la rive

- sud méditerranéenne: Cas du Rif Marocain. *Annales de La Société Entomologique de France*, 45(3), 309–320. <https://doi.org/10.1080/00379271.2009.10697616>
- Bes, M., Corbera, J., Sayol, F., Bagaria-Morató, G., Jover, M., Preece, C., Viza, A., Sabater i Comas, F., & Fernández-Martínez, M. (2018). On the influence of water conductivity, pH and climate on bryophyte assemblages in Catalan semi-natural springs. *Journal of Bryology*, 40, 149–158. <https://doi.org/10.1080/03736687.2018.1446484>
- Blanco, A., Guix, S., Fuster, N., Fuentes, C., Bartolomé, R., Cornejo, T., Pintó, R. M., & Bosch, A. (2017). Norovirus in bottled water associated with gastroenteritis outbreak, Spain, 2016. *Emerging Infectious Diseases*, 23(9), 1531–1534. <https://doi.org/10.3201/eid2309.161489>
- Blondel, J., & Aronson, J. (1999). *Biology and wildlife of the Mediterranean Region*. Oxford University Press.
- Bonada, N., & Resh, V. H. (2013). Mediterranean-climate streams and rivers: Geographically separated but ecologically comparable freshwater systems. *Hydrobiologia*, 719(1), 1–29. <https://doi.org/10.1007/s10750-013-1634-2>
- Botosaneanu, L. (1995). Springs as refugia for geographic relicts. *Crunoecia*, 4, 4–9.
- Bouwman, A. F., Bierkens, M. F. P., Griffioen, J., Hefting, M. M., Middelburg, J. J., Middelkoop, H., & Slomp, C. P. (2013). Nutrient dynamics, transfer and retention along the aquatic continuum from land to ocean: Towards integration of ecological and biogeochemical models. *Biogeosciences*, 10(1), 1–22. <https://doi.org/10.5194/bg-10-1-2013>
- Boyle, K., & Örmeci, B. (2020). Microplastics and nanoplastics in the freshwater and terrestrial environment: A review. *Water*, 12(9), 2633. <https://doi.org/10.3390/w12092633>
- Buono, V., Bissattini, A. M., & Vignoli, L. (2019). Can a cow save a newt? The role of cattle drinking troughs in amphibian conservation. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 29(6), 964–975. <https://doi.org/10.1002/aqc.3126>
- Burri, N. M., Weatherl, R., Moeck, C., & Schirmer, M. (2019). A review of threats to groundwater quality in the Anthropocene. *Science of the Total Environment*, 684, 136–154. <https://doi.org/10.1016/j.scitotenv.2019.05.236>
- Campos, I., & Abrantes, N. (2021). Forest fires as drivers of contamination of polycyclic aromatic hydrocarbons to the terrestrial and aquatic ecosystems. *Current Opinion in Environmental Science & Health*, 24, 100293. <https://doi.org/10.1016/j.coesh.2021.100293>
- Cantonati, M. (2008). Cyanoprocaryotes and algae other than diatoms in springs and streams of the Dolomiti Bellunesi National Park (Northern Italy). *Algological Studies*, 126, 113–136. <https://doi.org/10.1127/1864-1318/2008/0126-0113>
- Cantonati, M. (2022). Springs—Groundwater-borne ecotones—A typology, with an overview on the diversity of photoautotrophs in springs. In T. Mehner & K. Tockner (Eds.), *Encyclopedia of inland waters* (Vol. 3, 2nd ed., pp. 488–509). Elsevier Science.
- Cantonati, M., Angeli, N., Spitale, D., & Lange-Bertalot, H. (2016). A new *Navicula* (Bacillariophyta) species from low-elevation carbonate springs affected by anthropogenic disturbance. *Fottea*, 16(2), 255–265. <https://doi.org/10.5507/fot.2016.013>
- Cantonati, M., Decet, F., Corradini, F., & Bertuzzi, E. (2007). The significance of chemical and physical factors influencing the ecology of springs, and a case study in the south-eastern Alps (Dolomiti Bellunesi National Park). In M. Cantonati, E. Bertuzzi, & D. Spitale (Eds.), *The spring habitat: Biota and sampling methods* (Vol. 4, pp. 45–76). Museo Tridentino di Scienze Naturali.
- Cantonati, M., Fensham, R. J., Stevens, L. E., Gerecke, R., Glazier, D. S., Goldscheider, N., Knight, R. L., Richardson, J. S., Springer, A. E., & Tockner, K. (2021). Urgent plea for global protection of springs. *Conservation Biology*, 35(1), 378–382. <https://doi.org/10.1111/cobi.13576>
- Cantonati, M., Füreder, L., Gerecke, R., Jüttner, I., & Cox, E. J. (2012). Crenic habitats, hotspots for freshwater biodiversity conservation: Toward an understanding of their ecology. *Freshwater Science*, 31(2), 463–480. <https://doi.org/10.1899/11-111.1>
- Cantonati, M., Gerecke, R., & Bertuzzi, E. (2006). Springs of the Alps – Sensitive ecosystems to environmental change: From biodiversity assessments to long-term studies. *Hydrobiologia*, 562(1), 59–96. <https://doi.org/10.1007/s10750-005-1806-9>
- Cantonati, M., Komárek, J., & Montejano, G. (2015). Cyanobacteria in ambient springs. *Biodiversity and Conservation*, 24(4), 865–888. <https://doi.org/10.1007/s10531-015-0884-x>
- Cantonati, M., & Lange-Bertalot, H. (2006). *Achnanthydium dolomiticum* sp. nov. (Bacillariophyta) from oligotrophic mountain springs and lakes fed by dolomite aquifers. *Journal of Phycology*, 42(6), 1184–1188. <https://doi.org/10.1111/j.1529-8817.2006.00281.x>
- Cantonati, M., & Ortler, K. (1998). Using spring biota of pristine mountain areas for long-term monitoring. *Proceedings of the Headwaters '98 Conference* (Vol. 248, pp. 379–385). Meran, Italy. IAHS-AISH Publication.
- Cantonati, M., Rott, E., Spitale, D., Angeli, N., & Komárek, J. (2012). Are benthic algae related to spring types? *Freshwater Science*, 31(2), 481–498. <https://doi.org/10.1899/11-048.1>
- Cantonati, M., Segadelli, S., Ogata, K., Tran, H., Sanders, D., Gerecke, R., Rott, E., Filippini, M., Gargini, A., & Celico, F. (2016). A global review on ambient limestone-precipitating springs (LPS): Hydrogeological setting, ecology, and conservation. *Science of the Total Environment*, 568, 624–637. <https://doi.org/10.1016/j.scitotenv.2016.02.105>
- Cantonati, M., Segadelli, S., Spitale, D., Gabrieli, J., Gerecke, R., Angeli, N., De Nardo, M. T., Ogata, K., & Wehr, J. D. (2020). Geological and hydrochemical prerequisites of unexpectedly high biodiversity in spring ecosystems at the landscape level. *Science of the Total Environment*, 740, 140157. <https://doi.org/10.1016/j.scitotenv.2020.140157>
- Cantonati, M., Stevens, L. E., Segadelli, S., Springer, A. E., Goldscheider, N., Celico, F., Filippini, M., Ogata, K., & Gargini, A. (2020). Ecohydrogeology: The interdisciplinary convergence needed to improve the study and stewardship of springs and other groundwater-dependent habitats, biota, and ecosystems. *Ecological Indicators*, 110, 105803. <https://doi.org/10.1016/j.ecolind.2019.105803>
- Cartwright, J. M., Dwire, K. A., Freed, Z., Hammer, S. J., McLaughlin, B., Misztal, L. W., Schenk, E. R., Spence, J. R., Springer, A. E., & Stevens, L. E. (2020). Oases of the future? Springs as potential hydrologic refugia in drying climates. *Frontiers in Ecology and the Environment*, 18(5), 245–253. <https://doi.org/10.1002/fee.2191>
- Chen, W., Zhang, Z., Zhu, Y., Wang, X., Wang, L., Xiong, J., Qian, Z., Xiong, S., Zhao, R., Liu, W., Su, Q., Zhou, J., Zhou, H., Qi, S., & Jones, K. C. (2022). Distribution, sources and transport of polycyclic aromatic hydrocarbons (PAHs) in karst spring systems from Western Hubei, Central China. *Chemosphere*, 300, 134502. <https://doi.org/10.1016/j.chemosphere.2022.134502>
- Christenhusz, M., Bento-Elias, R., Dyer, R., Ivanenko, Y., Rouhan, G., Rumsey, F., & Väre, H. (2017). IUCN Red List of Threatened Species: *Isoetes malinverniana*. IUCN Red List of Threatened Species 2017, e.T162124A85431493. <https://doi.org/10.2305/IUCN.UK.2017-2.RLTS.T162124A85431493.en>
- Collier, K. J., & Smith, B. J. (2006). Distinctive invertebrate assemblages in rockface seepages enhance lotic biodiversity in northern New Zealand. *Biodiversity and Conservation*, 15(11), 3591–3616. <https://doi.org/10.1007/s10531-005-5395-8>
- Costello, D. M., Tiegs, S. D., & Lamberti, G. A. (2011). Do non-native earthworms in Southeast Alaska use streams as invasional corridors in watersheds harvested for timber? *Biological Invasions*, 13(1), 177–187. <https://doi.org/10.1007/s10530-010-9800-1>
- Cramer, W., Guiot, J., Fader, M., Garrabou, J., Gattuso, J.-P., Iglesias, A., Lange, M. A., Lionello, P., Llasat, M. C., Paz, S., Peñuelas, J., Snoussi, M., Toreti, A., Tsimplis, M. N., & Xoplaki, E. (2018). Climate

- change and interconnected risks to sustainable development in the Mediterranean. *Nature Climate Change*, 8(11), 972–980. <https://doi.org/10.1038/s41558-018-0299-2>
- Crema, S., Ferrarese, U., Golo, D., Modena, P., Sambugar, B., & Gerecke, R. (1996). Ricerche sulla fauna bentonica ed interstiziale di ambienti sorgentizi in area alpina e prealpina. *Report Del Centro Di Ecologia Alpina*, 8, 1–104.
- Dallas, H. (2008). Water temperature and riverine ecosystems: An overview of knowledge and approaches for assessing biotic responses, with special reference to South Africa. *Water*, 34(3), 393–404. <https://doi.org/10.4314/wsa.v34i3>
- Davidson, P. C., Kuhlenschmidt, T. B., Bhattarai, R., Kalita, P. K., & Kuhlenschmidt, M. S. (2016). Overland transport of rotavirus and the effect of soil type and vegetation. *Water*, 8(3), 78. <https://doi.org/10.3390/w8030078>
- Davis, J., Pavlova, A., Thompson, R., & Sunnucks, P. (2013). Evolutionary refugia and ecological refuges: Key concepts for conserving Australian arid zone freshwater biodiversity under climate change. *Global Change Biology*, 19(7), 1970–1984. <https://doi.org/10.1111/gcb.12203>
- de Foucault, B. (2015). Contribution au prodrome des végétations de France: Les *Adiantetea capilli-veneris* Braun-Blanq. ex Braun-Blanq., Roussine & Nègre 1952. *Acta Botanica Gallica*, 162(4), 375–403. <https://doi.org/10.1080/12538078.2015.1108868>
- Deleuil, G. (1974). Introduction phytogéographique générale. *Bulletin de La Société Botanique de France*, 121(1), 11–26. <https://doi.org/10.1080/00378941.1974.10835571>
- Delgado, C., Ector, L., Novais, M. H., Blanco, S., Hoffmann, L., & Pardo, I. (2013). Epilithic diatoms of springs and spring-fed streams in Majorca Island (Spain) with the description of a new diatom species *Cymbopleura margalefii* sp. nov. *Fottea*, 13(2), 87–104. <https://doi.org/10.5507/fot.2013.009>
- Delgado, C., Feio, M. J., Pardo, I., & Almeida, S. F. P. (2020). Effects of water temperature over benthic diatom communities: Insights from thermal springs. *Plant Ecology & Diversity*, 13(3–4), 325–337. <https://doi.org/10.1080/17550874.2020.1762133>
- Dubrovský, M., Hayes, M., Duce, P., Trnka, M., Svoboda, M., & Zara, P. (2014). Multi-GCM projections of future drought and climate variability indicators for the Mediterranean region. *Regional Environmental Change*, 14(5), 1907–1919. <https://doi.org/10.1007/s10113-013-0562-z>
- Egea-Serrano, A., Oliva-Paterna, F. J., Tejado, M., & Torralva, M. (2006). Breeding habitat selection of an endangered species in an arid zone: The case of *Alytes dickhilleni* Arntzen & García-Paris, 1995. *Acta Herpetologica*, 1(2), 81–94. https://doi.org/10.13128/Acta_Herpetol-1290
- Egea-Serrano, A., Oliva-Paterna, F. J., & Torralva, M. (2006). Breeding habitat selection of *Salamandra salamandra* (Linnaeus, 1758) in the most arid zone of its European distribution range: Application to conservation management. *Hydrobiologia*, 560(1), 363–371. <https://doi.org/10.1007/s10750-005-1589-z>
- Emiroğlu, Ö., Ekmekçi, F., Aksu, S., Başkurt, S., Atalay, M., & Tarkan, A. (2016). Introduction and establishment of tropical ornamental fish, *Pterygoplichthys* spp. (Actinopterygii: Siluriformes: Loricariidae) in hot springs: Aquarium trade as a potential risk for biodiversity in Turkey. *Acta Ichthyologica et Piscatoria*, 46(4), 351–356. <https://doi.org/10.3750/AIP2016.46.4.07>
- Erman, N. A., & Erman, D. C. (1995). Spring permanence, *Trichoptera* species richness, and the role of drought. *Journal of the Kansas Entomological Society*, 68(2), 50–64.
- European Environment Agency. (2022). *Europe's groundwater: A key resource under pressure*. Publications Office of the European Union. <https://doi.org/10.2800/50592>
- Farley, K. A., Jobbágy, E. G., & Jackson, R. B. (2005). Effects of afforestation on water yield: A global synthesis with implications for policy. *Global Change Biology*, 11(10), 1565–1576. <https://doi.org/10.1111/j.1365-2486.2005.01011.x>
- Fensham, R., Ponder, W., Souza, V., & Lawrence, E. S. (2023). Extraordinary concentrations of local endemism associated with arid-land springs. *Frontiers in Environmental Science*, 11, 3378. <https://doi.org/10.3389/fenvs.2023.1143378>
- Fensham, R. J., Adinehvand, R., Babidge, S., Cantonati, M., Currell, M., Daniele, L., & Villholth, K. G. (2023). Fellowship of the spring: An initiative to document and protect the world's oases. *Science of the Total Environment*, 887, 163936.
- Fensham, R. J., Silcock, J. L., Kerezsy, A., & Ponder, W. (2011). Four desert waters: Setting arid zone wetland conservation priorities through understanding patterns of endemism. *Biological Conservation*, 144(10), 2459–2467. <https://doi.org/10.1016/j.biocon.2011.06.024>
- Fernández-Martínez, M., Berloso, F., Corbera, J., García-Porta, J., Sayol, F., Preece, C., & Sabater, F. (2019). Towards a moss sclerophylly continuum: Evolutionary history, water chemistry and climate control traits of hygrophytic mosses. *Functional Ecology*, 33(12), 2273–2289. <https://doi.org/10.1111/1365-2435.13443>
- Fernández-Martínez, M., Corbera, J., Domene, X., Sayol, F., Sabater, F., & Preece, C. (2020). Nitrate pollution reduces bryophyte diversity in Mediterranean springs. *Science of the Total Environment*, 705, 135823. <https://doi.org/10.1016/j.scitotenv.2019.135823>
- Fernández-Martínez, M., Corbera, J., Torner, G., Bagaria, G., & Sayol, F. (2017). Les fonts del Montseny: Entre amenaces i oportunitats. *La Sitja Del Llop*, 42, 9–11.
- Fernández-Martínez, M., Margalef, O., Sayol, F., Asensio, D., Bagaria, G., Corbera, J., Sabater, F., Domene, X., & Preece, C. (2019). Sea spray influences water chemical composition of Mediterranean semi-natural springs. *Catena*, 173, 414–423. <https://doi.org/10.1016/j.catena.2018.10.035>
- Fernández-Martínez, M., Preece, C., Corbera, J., Cano, O., García-Porta, J., Sardans, J., Janssens, I. A., Sabater, F., & Peñuelas, J. (2021). Bryophyte C:N:P stoichiometry, biogeochemical niches and elementome plasticity driven by environment and coexistence. *Ecology Letters*, 24(7), 1375–1386. <https://doi.org/10.1111/ele.13752>
- Ferreira, C. S. S., Seifollahi-Aghmiuni, S., Destouni, G., Ghajarnia, N., & Kalantari, Z. (2022). Soil degradation in the European Mediterranean region: Processes, status and consequences. *Science of the Total Environment*, 805, 150106. <https://doi.org/10.1016/j.scitotenv.2021.150106>
- Filoso, S., Bezerra, M. O., Weiss, K. C. B., & Palmer, M. A. (2017). Impacts of forest restoration on water yield: A systematic review. *PLoS One*, 12(8), e0183210. <https://doi.org/10.1371/journal.pone.0183210>
- Freyhof, J., Weissenbacher, A., & Geiger, M. (2017). *Aphanius kruppi*, a new killifish from Oman with comments on the *Aphanius dispar* species group (Cyprinodontiformes: Aphaniidae). *Zootaxa*, 4338(3), 557–573. <https://doi.org/10.11646/zootaxa.4338.3.10>
- Gallart, F., & Llorens, P. (2003). Catchment management under environmental change: Impact of land cover change on water resources. *Water International*, 28(3), 334–340. <https://doi.org/10.1080/02508060308691707>
- Galloway, J. N., Townsend, A. R., Erisman, J. W., Bekunda, M., Cai, Z., Freney, J. R., Martinelli, L. A., Seitzinger, S. P., & Sutton, M. A. (2008). Transformation of the nitrogen cycle: Recent trends, questions, and potential solutions. *Science*, 320(5878), 889–892. <https://doi.org/10.1126/science.1136674>
- García-Ruiz, J. M., López-Moreno, J. I., Vicente-Serrano, S. M., Lasanta-Martínez, T., & Beguería, S. (2011). Mediterranean water resources in a global change scenario. *Earth-Science Reviews*, 105(3), 121–139. <https://doi.org/10.1016/j.earscirev.2011.01.006>
- Geissler, P. (1976). Zur Vegetation alpiner Fließgewässer: Pflanzensoziologisch-ökologische Untersuchungen hygrophiler Moosgesellschaften in den östlichen Schweizer Alpen. *Beiträge Zur Kryptogamenflora Der Schweiz*, 14, 1–55.

- Gerecke, R., Meisch, C., Stoch, F., Aciri, F., & Franz, H. (1998). Eucrenon-hypocrenon ecotone and spring typology in the Alps of Berchtesgaden (Upper Bavaria, Germany). A study of microcrustacea (Crustacea: Copepoda, Ostracoda) and water mites (Acari: Halacaridae, Hydrachnellae). *Studies in Crenobiology*. In L. Botosaneanu (Ed.), *The biology of springs and Springbrooks* (pp. 167–182). Backhuys Publishers.
- Glazier, D. S. (2014). Springs. In S. A. Elias (Ed.), *Reference module in Earth systems and environmental sciences* (pp. 1–78). Elsevier.
- Gligorović, B., Savić, A., Protić, L., & Pešić, V. (2016). Ecological patterns of water bug (Hemiptera: Heteroptera) assemblages in karst springs: A case study from central Montenegro. *Oceanological and Hydrobiological Studies*, 45(4), 554–563. <https://doi.org/10.1515/ohs-2016-0046>
- Goyette, J. O., Bennett, E. M., & Maranger, R. (2018). Low buffering capacity and slow recovery of anthropogenic phosphorus pollution in watersheds. *Nature Geoscience*, 11(12), 921–925. <https://doi.org/10.1038/s41561-018-0238-x>
- Gros, M., Mas-Pla, J., Sánchez-Melsió, A., Čelić, M., Castaño, M., Rodríguez-Mozaz, S., Borrego, C. M., Balcázar, J. L., & Petrović, M. (2023). Antibiotics, antibiotic resistance and associated risk in natural springs from an agroecosystem environment. *Science of the Total Environment*, 857, 159202. <https://doi.org/10.1016/j.scitotenv.2022.159202>
- Hartmann, A., Jasechko, S., Gleeson, T., Wada, Y., Andreo, B., Barberá, J. A., Brielmann, H., Bouchaou, L., Charlier, J.-B., Darling, W. G., Filippini, M., Garvelmann, J., Goldscheider, N., Kralik, M., Kunstmann, H., Ladouche, B., Lange, J., Lucianetti, G., Martín, J. F., ... Wagener, T. (2021). Risk of groundwater contamination widely underestimated because of fast flow into aquifers. *Proceedings of the National Academy of Sciences of the United States of America*, 118(20), e2024492118. <https://doi.org/10.1073/pnas.2024492118>
- Hartmann, A., Mudarra, M., Andreo, B., Marín, A., Wagener, T., & Lange, J. (2014). Modeling spatiotemporal impacts of hydroclimatic extremes on groundwater recharge at a Mediterranean karst aquifer. *Water Resources Research*, 50(8), 6507–6521. <https://doi.org/10.1002/2014WR015685>
- Herrero-Hernández, E., Rodríguez-Cruz, M. S., Pose-Juan, E., Sánchez-González, S., Andrades, M. S., & Sánchez-Martín, M. J. (2017). Seasonal distribution of herbicide and insecticide residues in the water resources of the vineyard region of La Rioja (Spain). *Science of the Total Environment*, 609, 161–171. <https://doi.org/10.1016/j.scitotenv.2017.07.113>
- Huang, F.-Y., Zhao, Y., Neilson, R., Zhou, X.-Y., Li, H., Ding, L., Zhou, S.-Y.-D., & Su, J.-Q. (2023). Antibiotic resistome in groundwater and its association with mountain springs and river. *Ecotoxicology and Environmental Safety*, 252, 114603. <https://doi.org/10.1016/j.ecoenv.2023.114603>
- Illies, J. (1961). Versuch einer allgemeinen biozönotischen Gliederung der Fließgewässer. *Internationale Revue der Gesamten Hydrobiologie Und Hydrographie*, 46(2), 205–213. <https://doi.org/10.1002/iroh.19610460205>
- Illies, J., & Botosaneanu, L. (1963). Problèmes et méthodes de la classification et de la zonation écologique des eaux courantes, considérées surtout du point de vue faunistique. *Internationale Vereinigung Für Theoretische Und Angewandte Limnologie: Mitteilungen*, 12(1), 1–57. <https://doi.org/10.1080/05384680.1963.11903811>
- Jacoby, C. A., Frazer, T. K., & Philips, E. J. (2008). Nutrient effects on spring flora and fauna. In M. T. Brown, K. C. Reiss, M. J. Cohen, J. M. Evans, K. R. Reddy, P. W. Inglett, K. S. Inglett, T. K. Frazer, C. A. Jacoby, E. J. Philips, R. L. Knight, S. K. Notestein, & K. A. McKee (Eds.), *Summary and synthesis of the available literature on the effects of nutrients on spring organisms and systems* (pp. 179–230). University of Florida Water Institute Report.
- Jasser, I., Panou, M., Khomutovska, N., Sandzewicz, M., Panteris, E., Niyatbekov, T., Łach, Ł., Kwiatowski, J., Kokociński, M., & Gkelis, S. (2022). Cyanobacteria in hot pursuit: Characterization of cyanobacteria strains, including novel taxa, isolated from geothermal habitats from different ecoregions of the world. *Molecular Phylogenetics and Evolution*, 170, 107454. <https://doi.org/10.1016/j.ympev.2022.107454>
- Jiménez-Alfaro, B., Fernández-Menéndez, S., Bueno, Á., & Fernández-Prieto, J. A. (2013). Vegetation and hydrogeology along the distribution range of *Centaureum somedanum*, an endemic plant of mountain calcareous springs. *Alpine Botany*, 123(1), 31–39. <https://doi.org/10.1007/s00035-013-0114-7>
- Jiménez-Mejías, P., & Luceño, M. (2009). *Carex castroviejoi* Luceño & Jiménez Mejías (Cyperaceae), a new species from North Greek mountains. *Acta Botanica Malacitana*, 34, 231–233. <https://doi.org/10.24310/abm.v34i0.6911>
- Jurado, A., Vázquez-Suñé, E., & Pujades, E. (2021). Urban groundwater contamination by non-steroidal anti-inflammatory drugs. *Water*, 13(5), 720. <https://doi.org/10.3390/w13050720>
- Kaiser, R. A., Polk, J. S., Datta, T., Parekh, R. R., & Agga, G. E. (2022). Occurrence of antibiotic resistant bacteria in urban karst groundwater systems. *Water*, 14(6), 960. <https://doi.org/10.3390/w14060960>
- Kampouris, I. D., Alygizakis, N., Klümper, U., Agrawal, S., Lackner, S., Cacace, D., Kunze, S., Thomaidis, N. S., Slobodnik, J., & Berendonk, T. U. (2022). Elevated levels of antibiotic resistance in groundwater during treated wastewater irrigation associated with infiltration and accumulation of antibiotic residues. *Journal of Hazardous Materials*, 423, 127155. <https://doi.org/10.1016/j.jhazmat.2021.127155>
- Khalloufi, N., Béjaoui, M., & Delicado, D. (2020). Two new genera and three new subterranean species of Hydrobiidae (Caenogastropoda: Truncatelloidea) from Tunisia. *European Journal of Taxonomy*, 648. <https://doi.org/10.5852/ejt.2020.648>
- Kishi, D., Murakami, M., Nakano, S., & Maekawa, K. (2005). Water temperature determines strength of top-down control in a stream food web. *Freshwater Biology*, 50(8), 1315–1322. <https://doi.org/10.1111/j.1365-2427.2005.01404.x>
- Kodric-Brown, A., & Brown, J. H. (2007). Native fishes, exotic mammals, and the conservation of desert springs. *Frontiers in Ecology and the Environment*, 5(10), 549–553. <https://doi.org/10.1890/070002>
- Kolda, A., Petrić, I., Mucko, M., Gottstein, S., Žutinić, P., Goretta, G., Ternje, I., Rubinić, J., Radišić, M., & Udovič, M. G. (2019). How environment selects: Resilience and survival of microbial mat community within intermittent karst spring Krčić (Croatia). *Ecohydrology*, 12(2), e2063. <https://doi.org/10.1002/eco.2063>
- Kresic, N. (2010). Chapter 2—Types and classifications of springs. In N. Kresic & Z. Stevanovic (Eds.), *Groundwater hydrology of springs* (pp. 31–85). Butterworth-Heinemann. <https://doi.org/10.1016/C2009-0-19145-6>
- Lachassagne, P., Dewandel, B., & Wyns, R. (2021). Review: Hydrogeology of weathered crystalline/hard-rock aquifers—Guidelines for the operational survey and management of their groundwater resources. *Hydrogeology Journal*, 29(8), 2561–2594. <https://doi.org/10.1007/s10040-021-02339-7>
- Lencioni, V., Mezzanotte, E., Spagnol, C., & Latella, L. (2018). Effects of human impacts on diversity and distribution of chironomids (Diptera: Chironomidae) in prealpine springs. *Journal of Limnology*, 77(1), 203–212. <https://doi.org/10.4081/jlimnol.2018.1804>
- Lewis-Phillips, J., Brooks, S. J., Sayer, C. D., Patmore, I. R., Hilton, G. M., Harrison, A., Robson, H., & Axmacher, J. C. (2020). Ponds as insect chimneys: Restoring overgrown farmland ponds benefits birds through elevated productivity of emerging aquatic insects. *Biological Conservation*, 241, 108253. <https://doi.org/10.1016/j.biocon.2019.108253>
- Llamas, M. I., Jiménez-Gavilán, P., Luque-Espinar, J. A., Benavente-Herrera, J., Candela, L., Sanmiguel-Martí, M., Rambla-Nebot, J.,

- Aranda-Mares, J. L., & Vadillo-Pérez, I. (2022). Hydrogeological, hydrodynamic and anthropogenic factors affecting the spread of pharmaceuticals and pesticides in water resources of the Granada plain (Spain). *Journal of Hydrology*, 610, 127791. <https://doi.org/10.1016/j.jhydrol.2022.127791>
- Llasat, M. C., Marcos, R., Turco, M., Gilabert, J., & Llasat-Botija, M. (2016). Trends in flash flood events versus convective precipitation in the Mediterranean region: The case of Catalonia. *Journal of Hydrology*, 541, 24–37. <https://doi.org/10.1016/j.jhydrol.2016.05.040>
- Loos, R., Locoro, G., Comero, S., Contini, S., Schwesig, D., Werres, F., Balsaa, P., Gans, O., Weiss, S., Blaha, L., Bolchi, M., & Gawlik, B. M. (2010). Pan-European survey on the occurrence of selected polar organic persistent pollutants in ground water. *Water Research*, 44(14), 4115–4126. <https://doi.org/10.1016/j.watres.2010.05.032>
- Manenti, R., Zanetti, N., Pennati, R., & Scari, G. (2017). Factors driving semi-aquatic predator occurrence in traditional cattle drinking pools: Conservation issues. *Journal of Limnology*, 76(1), 1447. <https://doi.org/10.4081/jlimnol.2016.1447>
- Mansilha, C., Duarte, C. G., Melo, A., Ribeiro, J., Flores, D., & Marques, J. E. (2019). Impact of wildfire on water quality in Caramulo Mountain ridge (Central Portugal). *Sustainable Water Resources Management*, 5(1), 319–331. <https://doi.org/10.1007/s40899-017-0171-y>
- Martín, A., Corbera, J., Cano, O., Preece, C., Peñuelas, J., Sabater, F., & Fernández-Martínez, M. (2024). The influence of nitrate pollution on elemental and isotopic composition of aquatic and semi-aquatic bryophytes. *Aquatic Botany*, 190, 103710. <https://doi.org/10.1016/j.aquabot.2023.103710>
- Mas-Pla, J., & Menció, A. (2019). Groundwater nitrate pollution and climate change: Learnings from a water balance-based analysis of several aquifers in a western Mediterranean region (Catalonia). *Environmental Science and Pollution Research*, 26(3), 2184–2202. <https://doi.org/10.1007/s11356-018-1859-8>
- Mays, L. W., Koutsoyiannis, D., & Angelakis, A. N. (2007). A brief history of urban water supply in antiquity. *Water Supply*, 7(1), 1–12. <https://doi.org/10.2166/ws.2007.001>
- McKinley, J., Healy, M. G., Ryan, P. C., O'Driscoll, H., Mellander, P.-E., Morrison, L., & Siggins, A. (2023). Impact of historical legacy pesticides on achieving legislative goals in Europe. *Science of the Total Environment*, 873, 162312. <https://doi.org/10.1016/j.scitotenv.2023.162312>
- Mekonnen, M. M., & Hoekstra, A. Y. (2016). Four billion people facing severe water scarcity. *Science Advances*, 2(2), e1500323. <https://doi.org/10.1126/sciadv.1500323>
- Mezquita, F., Sanz-Brau, A., & Wansard, G. (2000). Habitat preferences and population dynamics of Ostracoda in a helocrene spring system. *Canadian Journal of Zoology*, 78(5), 840–847.
- Mezquita, F., Tapia, G., & Roca, J. R. (1999). Ostracoda from springs on the eastern Iberian Peninsula: Ecology, biogeography and palaeolimnological implications. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 148(1), 65–85. [https://doi.org/10.1016/S0031-0182\(98\)00176-X](https://doi.org/10.1016/S0031-0182(98)00176-X)
- Minteni, S. M., Bäuerlein, P. S., Koelmans, A. A., Dekker, S. C., & van Wezel, A. P. (2018). Closing the gap between small and smaller: Towards a framework to analyse nano- and microplastics in aqueous environmental samples. *Environmental Science: Nano*, 5(7), 1640–1649. <https://doi.org/10.1039/C8EN00186C>
- Miracle, M. R., Alfonso, M. T., Vicente, E., & Koste, W. (1995). Rotifers of spring pools in the coastal marshland of Albufera of Valencia Natural Park. *Limnetica*, 11, 39–47.
- Morell, I. (1992). *Los manantiales de la provincia de Castellón*. Diputación de Castellón.
- Murphy, H. M., Prioleau, M. D., Borchardt, M. A., & Hynds, P. D. (2017). Review: Epidemiological evidence of groundwater contribution to global enteric disease, 1948–2015. *Hydrogeology Journal*, 25(4), 981–1001. <https://doi.org/10.1007/s10040-017-1543-y>
- Myers, M., & Resh, V. (2002). Trichoptera and other macroinvertebrates in springs of the Great Basin: Species composition, richness, and distribution. *Western North American Naturalist*, 62(1), 1–13.
- Nascimbene, J., Nimis, P. L., Klüßendorf, J., & Thüs, H. (2023). Freshwater lichens, including new species in the Genera *Verrucaria*, *Placopyrenium* and *Circinaria*, associated with *Lobothallia hydrocharis* (Poelt & Nimis) Sohrabi & Nimis from watercourses of Sardinia. *Journal of Fungi*, 9(3), 380. <https://doi.org/10.3390/jof9030380>
- Nearly, D. G., Ice, G. G., & Jackson, C. R. (2009). Linkages between forest soils and water quality and quantity. *Forest Ecology and Management*, 258(10), 2269–2281. <https://doi.org/10.1016/j.foreco.2009.05.027>
- Nerantzaki, S. D., & Nikolaidis, N. P. (2020). The response of three Mediterranean karst springs to drought and the impact of climate change. *Journal of Hydrology*, 591, 125296. <https://doi.org/10.1016/j.jhydrol.2020.125296>
- Nielson, K. G., Gill, K. M., Springer, A. E., Ledbetter, J. D., Stevens, L. E., & Rood, S. B. (2019). Springs ecosystems: Vulnerable ecological islands where environmental conditions, life history traits, and human disturbance facilitate non-native plant invasions. *Biological Invasions*, 21(9), 2963–2981. <https://doi.org/10.1007/s10530-019-02025-6>
- Nowicka-Krawczyk, P., & Żelazna-Wieczorek, J. (2017). Dynamics in cyanobacterial communities from a relatively stable environment in an urbanised area (ambient springs in Central Poland). *Science of the Total Environment*, 579, 420–429. <https://doi.org/10.1016/j.scitotenv.2016.11.080>
- Nowicka-Krawczyk, P. B., & Żelazna-Wieczorek, J. (2013). Cyanobacteria microflora in a limestone spring (Troniny spring, Central Poland). *Acta Societatis Botanicae Poloniae*, 82(3), 219–224. <https://doi.org/10.5586/asbp.2013.017>
- Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., & Wagner, H. (2018). *Vegan: Community ecology package*. R package version 2.5-2.
- Panno, S. V., Kelly, W. R., Scott, J., Zheng, W., McNeish, R. E., Holm, N., Hoellein, T. J., & Baranski, E. L. (2019). Microplastic contamination in karst groundwater systems. *Groundwater*, 57(2), 189–196. <https://doi.org/10.1111/gwat.12862>
- Pascual, R., Gomà, J., Pedrocchi, C., Cadiach, O., García, G., & Solé, J. (2020). First data on the biological richness of Mediterranean springs. *Limnetica*, 39(1), 121–139. <https://doi.org/10.23818/limn.39.09>
- Patrick, R., Crum, B., & Coles, J. (1969). Temperature and manganese as determining factors in the presence of diatom or blue-green algal floras in streams. *Proceedings of the National Academy of Sciences of the United States of America*, 64(2), 472–478. <https://doi.org/10.1073/pnas.64.2.472>
- Pešić, V., Gligorović, B., Savić, A., & Buczyński, P. (2017). Ecological patterns of Odonata assemblages in karst springs in central Montenegro. *Knowledge & Management of Aquatic Ecosystems*, 418, 3. <https://doi.org/10.1051/kmae/2016035>
- Pinkster, S. (1993). A revision of the genus *Echinogammarus* Stebbing, 1899 with some notes on related genera (Crustacea, Amphipoda). *Memorie del Museo Civico di Storia Naturale*, 2(10), 9–183.
- Plóciennik, M., Dmitrović, D., Pešić, V., & Gadawski, P. (2016). Ecological patterns of Chironomidae assemblages in Dinaric karst springs. *Knowledge and Management of Aquatic Ecosystems*, 417, 11. <https://doi.org/10.1051/kmae/2015044>
- Psomas, A., Vryzidis, I., Spyridakos, A., & Mimikou, M. (2021). MCDA approach for agricultural water management in the context of water-energy-land-food nexus. *Operational Research*, 21(1), 689–723. <https://doi.org/10.1007/s12351-018-0436-8>
- R Core Team. (2023). *R: A language and environment for statistical computing* (v. 4.2.3). R Foundation for Statistical Computing.
- Rassam, H., Ghamizi, M., Benaissa, H., Clewing, C., & Albrecht, C. (2021). The fingernail clams (Bivalvia: Veneroidea: Sphaeriidae)

- of Morocco: Diversity, distribution and conservation status. *Biodiversity Data Journal*, 9, e73346. <https://doi.org/10.3897/BDJ.9.e73346>
- Reiss, M., & Chiffard, P. (2015). Hydromorphology and biodiversity in headwaters—An eco-faunistic substrate preference assessment in forest springs of the German Subdued Mountains. In Y. H. Lo, J. A. Blanco, & S. Roy (Eds.), *Biodiversity in ecosystems - Linking structure and function* (pp. 223–258). InTech Open. <https://doi.org/10.5772/59072>
- Rivas-Martínez, S., Penas, Á., Díaz-González, T. E., Ladero-Álvarez, M., Asensi-Marfil, A., Díez-Garretas, B., Molero-Mesa, J., Valle-Tendero, F., Cano, E., Costa-Talens, M., López, M. L., Fernández-Prieto, J. A., Llorens, L., del Arco, M., Pérez de Paz, P. L., de la Torre, W., Sánchez-Mata, D., Fernández, F., Masalles-Raurell, R., & Herrero, L. (2011). *Mapa de series, geoserias y geopermaseries de vegetación de España (Memoria del mapa de vegetación potencial de España). Parte II*. <https://digital.csic.es/handle/10261/108186>
- Roca, J. R. (1990). Tipología físico-química de las fuentes de los Pirineos centrales: síntesis regional. *Limnetica*, 6, 57–78.
- Rossetti, G., Stoch, F., & Mazzini, I. (2022). A reassessment of the origin and distribution of the subterranean genus *Pseudolimnocythere* Klie, 1938 (Ostracoda, Loxoconchidae), with description of two new species from Italy. *Subterranean Biology*, 43, 33–60. <https://doi.org/10.3897/subtblol.43.82158>
- Rossini, R. A., Fensham, R. J., Stewart-Koster, B., Gotch, T., & Kennard, M. J. (2018). Biogeographical patterns of endemic diversity and its conservation in Australia's artesian desert springs. *Diversity and Distributions*, 24(9), 1199–1216. <https://doi.org/10.1111/ddi.12757>
- Rundel, P. W., Arroyo, M. T. K., Cowling, R. M., Keeley, J. E., Lamont, B. B., & Vargas, P. (2016). Mediterranean Biomes: Evolution of their vegetation, floras, and climate. *Annual Review of Ecology, Evolution, and Systematics*, 47(1), 383–407. <https://doi.org/10.1146/annurev-ecolsys-121415-032330>
- Running, S., Mu, Q., Zhao, M., & Moreno, A. (2019). MOD16A2GF MODIS/Terra net evapotranspiration gap-filled 8-day L4 global 500 m SIN grid V006. NASA EOSDIS Land Processes DAAC.
- Ryu, S., Won, S. A., Uh, J., & Song, J. Y. (2019). Hepatitis A virus infection from a contaminated tap of ground water facility in a neighborhood park, Republic of Korea. *Infection & Chemotherapy*, 51(1), 62–66. <https://doi.org/10.3947/ic.2019.51.1.62>
- Sabater, S., & Roca, J. R. (1990). Some factors affecting distribution of diatom assemblages in Pyrenean springs. *Freshwater Biology*, 24(3), 493–507.
- Sabatino, A. D., Cicolani, B., & Gerecke, R. (2003). Biodiversity and distribution of water mites (Acari, Hydrachnidia) in spring habitats. *Freshwater Biology*, 48(12), 2163–2173. <https://doi.org/10.1046/j.1365-2427.2003.01151.x>
- Saber, A. A., Borrini, A., Saber, H., El-Sheekh, M., Gontcharov, A. A., & Cantonati, M. (2022). A marine invasive benthic diatom species [*Licmophora normaniana* (Greville) Wahrer, 1985] in an inland oasis mineral spring in Egypt. *BioInvasions Record*, 11(1), 13–22. <https://doi.org/10.3391/bir.2022.11.1.02>
- Savio, D., Stadler, P., Reischer, G. H., Kirschner, A. K. T., Demeter, K., Linke, R., Blaschke, A. P., Sommer, R., Szewzyk, U., Wilhartitz, I. C., Mach, R. L., Stadler, H., & Farnleitner, A. H. (2018). Opening the black box of spring water microbiology from alpine karst aquifers to support proactive drinking water resource management. *WIREs Water*, 5(3), e1282. <https://doi.org/10.1002/wat2.1282>
- Scanlon, B. R., Reedy, R. C., Stonestrom, D. A., Prudic, D. E., & Dennehy, K. F. (2005). Impact of land use and land cover change on groundwater recharge and quality in the southwestern US. *Global Change Biology*, 11(10), 1577–1593. <https://doi.org/10.1111/j.1365-2486.2005.01026.x>
- Seiler, H., Küry, D., Billeter, R., & Dengler, J. (2021). Regional typology of spring vegetation in Parc Ela (Grisons, Switzerland). *Vegetation Classification and Survey*, 2, 257–274. <https://doi.org/10.3897/VCS/2021/69101>
- Singh, A. K., Kaur, R., Verma, S., & Singh, S. (2022). Antimicrobials and antibiotic resistance genes in water bodies: Pollution, risk, and control. *Frontiers in Environmental Science*, 10, 830861. <https://doi.org/10.3389/fenvs.2022.830861>
- Singh, K. P., Rai, P., Singh, A. K., Verma, P., & Gupta, S. (2014). Occurrence of pharmaceuticals in urban wastewater of north Indian cities and risk assessment. *Environmental Monitoring and Assessment*, 186(10), 6663–6682. <https://doi.org/10.1007/s10661-014-3881-8>
- Sivelle, V., Jourde, H., Bittner, D., Mazzilli, N., & Trambly, Y. (2021). Assessment of the relative impacts of climate changes and anthropogenic forcing on spring discharge of a Mediterranean karst system. *Journal of Hydrology*, 598, 126396. <https://doi.org/10.1016/j.jhydrol.2021.126396>
- Soria, M., Leigh, C., Datry, T., Bini, L. M., & Bonada, N. (2017). Biodiversity in perennial and intermittent rivers: A meta-analysis. *Oikos*, 126(8), 1078–1089. <https://doi.org/10.1111/oik.04118>
- Souza, V., Espinosa-Asuar, L., Escalante, A. E., Eguarte, L. E., Farmer, J., Forney, L., Lloret, L., Rodríguez-Martínez, J. M., Soberón, X., Dirzo, R., & Elser, J. J. (2006). An endangered oasis of aquatic microbial biodiversity in the Chihuahuan desert. *Proceedings of the National Academy of Sciences of the United States of America*, 103(17), 6565–6570. <https://doi.org/10.1073/pnas.0601434103>
- Spampinato, G., Tomaselli, V., Forte, L., Strumia, S., Stinca, A., Croce, A., Fascetti, S., Rosati, L., Di Pietro, R., Mantino, F., Laface, V. L. A., & Musarella, C. M. (2023). Relevant but neglected habitat types by the Directive 92/43 EEC in southern Italy. *Rendiconti Lincei. Scienze Fisiche e Naturali*, 34(2), 457–482. <https://doi.org/10.1007/s12210-023-01136-6>
- Steinmann, P., Siegrist, R., & Gams, H. (1915). *Praktikum der Süßwasserbiologie: Teil 1. Die Organismen des fließenden Wassers*. Borntraeger Publishing Services.
- Stevens, L. E., Aly, A. A., Arpin, S. M., Apostolova, I., Ashley, G. M., Barba, P. Q., Barquín, J., Beauger, A., Benaabidate, L., Bhat, S. U., Bouchaou, L., Cantonati, M., Carroll, T. M., Death, R., Dwire, K. A., Felipe, M. F., Fensham, R. J., Fryar, A. E., Garsaball, R. P. I., ... Voldoire, O. (2022). The ecological integrity of spring ecosystems: A global review. In D. A. DellaSala & M. I. Goldstein (Eds.), *Reference module in Earth systems and environmental sciences* (pp. 436–451). Elsevier. <https://doi.org/10.1016/B978-0-12-821139-7.00111-2>
- Stevens, L. E., Schenk, E. R., & Springer, A. E. (2021). Springs ecosystem classification. *Ecological Applications*, 31(1), e2218. <https://doi.org/10.1002/eap.2218>
- Stevens, L. E., Springer, A. E., & Ledbetter, J. D. (2011). *Inventory and monitoring protocols for springs ecosystems*. Spring Stewardship Institute.
- Stoch, F., Gerecke, R., Pieri, V., Rossetti, G., & Sambugar, B. (2011). Exploring species distribution of spring meiofauna (Annelida, Acari, Crustacea) in the south-eastern Alps. *Journal of Limnology*, 70(1), 65–76. <https://doi.org/10.4081/jlimnol.2011.s1.65>
- Sui, Q., Cao, X., Lu, S., Zhao, W., Qiu, Z., & Yu, G. (2015). Occurrence, sources and fate of pharmaceuticals and personal care products in the groundwater: A review. *Emerging Contaminants*, 1(1), 14–24. <https://doi.org/10.1016/j.emcon.2015.07.001>
- Sutton, M. A., Howard, C. M., Erisman, J. W., Billen, G., Bleeker, A., Grennfelt, P., van Grinsven, H., & Grizzetti, B. (2011). *The European Nitrogen Assessment: Sources, effects and policy perspectives*. Cambridge University Press.
- Takuissu, G. R., Kenmoe, S., Ndip, L., Ebogo-Belobo, J. T., Kengne-Ndé, C., Mbagu, D. S., Bowo-Ngandji, A., Oyono, M. G., Kenfack-Momo, R., Tchatchouang, S., Kenfack-Zanguim, J., Lontuo Fogang, R., Zeuko'o Menkem, E., Kame-Ngasse, G. I., Magoudjou-Pekam, J. N., Nkies Esemu, S., Veneri, C., Mancini, P., Bonanno Ferraro, G., ... La Rosa, G. (2022). Hepatitis E virus in water environments: A systematic review and meta-analysis. *Food and Environmental*

- Virology*, 14(3), 223–235. <https://doi.org/10.1007/s12560-022-09530-3>
- Taxböck, L., Karger, D. N., Kessler, M., Spitale, D., & Cantonati, M. (2020). Diatom species richness in Swiss springs increases with habitat complexity and elevation. *Water*, 12(2), 449. <https://doi.org/10.3390/w12020449>
- Taxböck, L., Linder, H. P., & Cantonati, M. (2017). To what extent are swiss springs refugial habitats for sensitive and endangered diatom taxa? *Water*, 9(12), 967. <https://doi.org/10.3390/w9120967>
- Teuling, A. J., de Badts, E. A. G., Jansen, F. A., Fuchs, R., Buitink, J., Hoek van Dijke, A. J., & Sterling, S. M. (2019). Climate change, reforestation/afforestation, and urbanization impacts on evapotranspiration and streamflow in Europe. *Hydrology and Earth System Sciences*, 23(9), 3631–3652. <https://doi.org/10.5194/hess-23-3631-2019>
- Thienemann, A. (1926). Hydrobiologische untersuchungen an Quellen VII. Insekten aus norddeutschen Quellen mit besonderer Berücksichtigung der Dipteren. *Deutsche Entomologische Zeitschrift*, 1926(1), 1–50. <https://doi.org/10.1002/mmnd.192619260102>
- van der Kamp, G. (1995). The hydrogeology of springs in relation to the biodiversity of spring fauna: A review. *Journal of the Kansas Entomological Society*, 68(2), 4–17.
- Van Meter, K. J., Van Cappellen, P., & Basu, N. B. (2018). Legacy nitrogen may prevent achievement of water quality goals in the Gulf of Mexico. *Science*, 360(6387), 427–430. <https://doi.org/10.1126/science.aar4462>
- Van Vliet, M. T. H., Flörke, M., & Wada, Y. (2017). Quality matters for water scarcity. *Nature Geoscience*, 10, 800–802. <https://doi.org/10.1038/ngeo3047>
- Viaroli, S., Lancia, M., & Re, V. (2022). Microplastics contamination of groundwater: Current evidence and future perspectives. A review. *Science of the Total Environment*, 824, 153851. <https://doi.org/10.1016/j.scitotenv.2022.153851>
- Vilenica, M., Kulijer, D., Gligorović, B., Gligorović, A., & Knijf, G. D. (2021). Distribution, habitat requirements, and vulnerability of *Caliaeschna microstigma* at the north-western edge of its range (Odonata: Aeshnidae). *Odonatologica*, 50(3–4), 203–225. <https://doi.org/10.5281/zenodo.5703202>
- von Fumetti, S., Nagel, P., & Baltes, B. (2007). Where a springhead becomes a springbrook a regional zonation of springs. *Fundamental and Applied Limnology*, 169, 37–48. <https://doi.org/10.1127/1863-9135/2007/0169-0037>
- Votto, S. E., Dyer, F. J., Caron, V., & Davis, J. A. (2020). Thermally-driven thresholds in terrestrial avifauna waterhole visitation indicate vulnerability to a warming climate. *Journal of Arid Environments*, 181, 104217. <https://doi.org/10.1016/j.jaridenv.2020.104217>
- Wallace, S. J., de Solla, S. R., Head, J. A., Hodson, P. V., Parrott, J. L., Thomas, P. J., Berthiaume, A., & Langlois, V. S. (2020). Polycyclic aromatic compounds (PACs) in the Canadian environment: Exposure and effects on wildlife. *Environmental Pollution*, 265, 114863. <https://doi.org/10.1016/j.envpol.2020.114863>
- Welch, L. A., & Allen, D. M. (2014). Hydraulic conductivity characteristics in mountains and implications for conceptualizing bedrock groundwater flow. *Hydrogeology Journal*, 22(5), 1003–1026. <https://doi.org/10.1007/s10040-014-1121-5>
- White, P. S., & Pickett, S. T. A. (1985). Chapter 1 – Natural disturbance and patch dynamics: An introduction. In S. T. A. Pickett & P. S. White (Eds.), *The ecology of natural disturbance and patch dynamics* (pp. 3–13). Academic Press Inc.
- Williams, D. D. (1991). The spring as an interface between groundwater and lotic faunas and as a tool in assessing groundwater quality. *Internationale Vereinigung für Theoretische und Angewandte Limnologie: Verhandlungen*, 24, 1621–1624.
- Work, K. (2023). The distribution, magnitude, and endemic species of US springs. *Frontiers in Environmental Science*, 10, 1022424. <https://doi.org/10.3389/fenvs.2022.1022424>
- Zamora-Marín, J. M., Ilg, C., Demierre, E., Bonnet, N., Wezel, A., Robin, J., Vallod, D., Calvo, J. F., Oliva-Paterna, F. J., & Oertli, B. (2021). Contribution of artificial waterbodies to biodiversity: A glass half empty or half full? *Science of the Total Environment*, 753, 141987. <https://doi.org/10.1016/j.scitotenv.2020.141987>
- Zamora-Marín, J. M., Zamora-López, A., Jiménez-Franco, M. V., Calvo, J. F., & Oliva-Paterna, F. J. (2021). Small ponds support high terrestrial bird species richness in a Mediterranean semiarid region. *Hydrobiologia*, 848(7), 1623–1638. <https://doi.org/10.1007/s10750-021-04552-7>
- Zamora-Marín, J. M., Zamora-López, A., Oliva-Paterna, F. J., Torralva, M., Sánchez-Montoya, M. M., & Calvo, J. (2023). From small waterbodies to large multi-service providers: Assessing their ecological multifunctionality for terrestrial birds in Mediterranean agroecosystems. *Agriculture, Ecosystems & Environments*, 359, 108760. <https://doi.org/10.1016/j.agee.2023.108760>
- Zamora-Marín, J. M., Zamora-López, A., Sánchez-Fernández, D., Calvo, J. F., & Oliva-Paterna, F. J. (2022). Traditional small waterbodies as key landscape elements for farmland bird conservation in Mediterranean semiarid agroecosystems. *Global Ecology and Conservation*, 37, e02183. <https://doi.org/10.1016/j.gecco.2022.e02183>
- Žutinić, P., Petrić, I., Gottstein, S., Udovič, M. G., Borojević, K. K., Kamberović, J., Kolda, A., Plenković-Moraj, A., & Ternjeji, I. (2018). Microbial mats as shelter microhabitat for amphipods in an intermittent karstic spring. *Knowledge & Management of Aquatic Ecosystems*, 419, 7. <https://doi.org/10.1051/kmae/2017061>


SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Fernández-Martínez, M., Barquín, J., Bonada, N., Cantonati, M., Churro, C., Corbera, J., Delgado, C., Dulsat-Masvidal, M., García, G., Margalef, O., Pascual, R., Peñuelas, J., Preece, C., Sabater, F., Seiler, H., Zamora-Marín, J. M., & Romero, E. (2023). Mediterranean springs: Keystone ecosystems and biodiversity refugia threatened by global change. *Global Change Biology*, 00, e16997. <https://doi.org/10.1111/gcb.16997>

1 Supplementary Materials

2 **Figure S1:** Piezometric levels from January 2013 and 2023 of four wells located in the same
3 region as the 30 monitored springs (Maresme, Spain). The data were analysed using a mixed
4 effect model in which piezometric level was the response variable, year was a 2-level fixed
5 factor, and the well was the random factor. Piezometric data for the Maresme aquifers was
6 downloaded from the public dataset of the Catalan Water Agency, Generalitat de Catalunya:
7 <https://aplicacions.aca.gencat.cat/sdim21/>. See Data availability statement to access data and
8 code to reproduce these results.



10 **Section 1, supplementary text: Land density of springs**

11 Given their strong dependence on hydrogeomorphological features, the density of springs
12 can vary widely across geographical areas (**Table S1**). Regional and national inventories
13 tend to overlook small springs and seepages, while inventories focused on small areas
14 (natural parks, municipalities, etc.) provide much more complete estimates. This was pointed
15 out by Glazier, (2014) on the global scale who used several spring inventories from the
16 Iberian Peninsula and found a significant inverse potential relationship between the scale of
17 the study site and the observed spring density. Considering such bias, Glazier (2014)
18 estimated a global average spring density of >0.3 springs km^{-2} . Local studies from
19 mountainous areas provide higher estimates: $>1-4$ springs km^{-2} in the Mediterranean ranges
20 of Catalonia (**Table S1**); 3.8 springs km^{-2} in Kalkalpen National Park, Austria (Cantonati,
21 Füreder, et al., 2012); or 2.6 springs km^{-2} in Berchtesgaden Alps (Gerecke et al., 1998). The
22 spring density in sedimentary plains remains largely unknown but is expected to be much
23 lower. For instance, (Gray, 2005) reports densities in the range of 0.11-0.40 springs km^{-2} in
24 alluvial floodplains of different New Zealand braided river catchments.

25

26 **Section 2, supplementary text: Original text from "Llibre de meravelles" by Ramón Llull**
27 **(1287-1289 AD):**

28 *"Seia lo filosof sots un bell arbre carregat de fulles e de flors; una bella fontana regava aquell*
29 *arbre, en la qual havia molts ocells que dolçament cantaven. Segons disposició de l'arbre e*
30 *de la font e dels ocells contemplava lo filosof la granesa e la bonesa de Déu [...]"*

31 **Table S1:** Data regarding the density of springs in different sites, together with the intensity
 32 and extent of the survey.

Study site	Approx. area (km²)	Number of inventoried springs	Spring density (km²)	Reference
Montblanc municipality	90	350	3.89	Pallisé, J. (2019). <i>Las fuentes naturales frente a Escila y Caribdis</i> . In: SICEF'19 1st Iberian Symposium on the Conservation of Spring Ecosystems. Barcelona, 10-12 June 2019. https://fuentes-naturales.org/sicef19
Collserola range	100	332	3.32	<i>Les fonts de Collserola</i> . Associació Fes Fonts Fent Fonting. URL: https://fontscollserola.com/ . [Accessed 04/09/2020]
Montserrat range	175	398	2.27	MN Consultors en Ciències de la Conservació (2012). <i>Determinació de les característiques fisicoquímiques i de l'estat sanitari de les fonts naturals de la serra de Montsant. Fase 1: Localització i inventariat de les fonts i proposta d'estudi sanitari</i> . Parc Natural de Montsant, Generalitat de Catalunya.
Montserrat range	500	850	1.70	Farrerons-Vidal, O. (2019). <i>Una visión multidimensional etnográfica, cultural, histórica y patrimonial de las fuentes de la reserva de la biosfera del Montseny</i> . In: SICEF'19 1st Iberian Symposium on the Conservation of Spring Ecosystems. Barcelona, 10-12 June 2019. URL: https://fuentes-naturales.org/sicef19
Tramuntana range*	1000	1167	1.17	Morell, A. (2020). <i>Fonts de Mallorca</i> . URL: https://sites.google.com/a/fontsdetramuntana.com/fontsa pps/home . [Accessed 04/09/2020]
Comarca de la Sierra de Albarracín	1414	1200	0.85	Ibáñez, R. (2019). <i>Albaqua. Un proyecto integral para el conocimiento de las fuentes de la sierra de albarracín</i> . In: SICEF'19 1st Iberian Symposium on the Conservation of Spring Ecosystems. Barcelona, 10-12 June 2019. https://fuentes-naturales.org/sicef19
La Rioja	5054	1357	0.27	<i>Inventario de Fuentes y Manantiales de La Rioja</i> (2020). Gobierno de la Rioja. URL: https://www.larioja.org/medio-ambiente/es/agua/inventario-fuentes-manantiales-rioja . [Accessed 04/09/2020]
Catalonia	31895	9846	0.31	García, G. (2019). <i>Las fuentes naturales ibéricas: comprender y conservar un "ecosistema difuso"</i> . In: SICEF'19 1st Iberian Symposium on the Conservation of Spring Ecosystems. Barcelona, 10-12 June 2019. https://fuentes-naturales.org/sicef19

Andalucía	87268	12384	0.14	<i>Conoce tus Fuentes. Manantiales y Fuentes de Andalucía</i> (2020). Instituto Universitario de Investigación del Agua de la Universidad de Granada. URL: http://www.conocetusfuentes.com/home.php . [Accessed 04/09/2020]
Peninsular Spain & Balearic Islands	499003	26060	0.05	García, G. (2019). Las fuentes naturales ibéricas: comprender y conservar un "ecosistema difuso". In: SICEF'19 1st Iberian Symposium on the Conservation of Spring Ecosystems. Barcelona, 10-12 June 2019. https://fuentes-naturales.org/sicef19

33 * This region belongs to the Balearic Islands (Mallorca), close to the eastern coast of the Iberian Peninsula

34

35 **Table S2.** Selection of works dealing with biodiversity of spring ecosystems arranged by broad
 36 taxonomic groups. The number of springs sampled, the number of taxa identified, and the
 37 geographic region are included. We also include some citations from nearby countries with the
 38 intention of improving knowledge on spring biodiversity.

39

Taxa	Reference	N of taxa	N of springs	Geographic region	Observations
Cyanobacteria	Cantonati, 2008	65	16	Southern Alps, Italy	
Cyanobacteria	Gesierich & Kofler, 2010	63	27	Eastern Alps, Austria	
Cyanobacteria	Cantonati et al., 2012a	88	70	Southern Alps, Italy	
Cyanobacteria	Nowicka-Krawczyk & Żelazna-Wieczorek, 2013	30	1	Poland	
Cyanobacteria	Pascual et al., 2020	41	20	Catalonia and Balearic Islands, Spain	
Bacillariophyceae	Sabater & Roca, 1992	193	28	Pyrenees, Spain	
Bacillariophyceae	Roca, 1996	3	25	Catalonia, Spain	
Bacillariophyceae	Aboal et al., 1998	147	135	Castelló, Spain	
Bacillariophyceae	Cantonati et al., 2012b	≥58	126	Southern Alps, Italy	
Bacillariophyceae	Guasch et al., 2012	42	1	Catalonia, Spain	Fe spring
Bacillariophyceae	Delgado et al., 2013	111	6	Balearic Islands, Spain	
Bacillariophyceae	Žutinić et al., 2018	25	1	Croatia	
Bacillariophyceae	Lai et al., 2020	162	8	Sardinia, Italy	
Bacillariophyceae	Pascual et al., 2020	154	20	Catalonia and Balearic Islands, Spain	

Chlorophyta	Cantonati et al., 2012a	10	70	Southern Alps, Italy	
Chlorophyta	Pascual et al., 2020	8	20	Catalonia and Balearic Islands, Spain	
Charophyta	Pascual et al., 2020	11	20	Catalonia and Balearic Islands, Spain	
Rhodophyta	Cantonati et al., 2012a	5	70	Southern Alps, Italy	
Rhodophyta	Pascual et al., 2020	2	20	Catalonia and Balearic Islands, Spain	
Bryophyta	Roca & del Castillo, 1993	17	94	Pyrenees, Spain	
Bryophyta	Roca, 1996	9	25	Catalonia, Spain	
Bryophyta	Suanjak, 2007	60	13	Austria	
Bryophyta	Bes et al., 2017	58	198	Catalonia, Spain	
Bryophyta	Pascual et al., 2020	41	20	Catalonia and Balearic Islands, Spain	
Bryophyta	Fernández-Martínez et al., 2020	77	338	Catalonia, Spain	
Bryophyta	Seiler et al., 2021	95	20	Switzerland	
Tracheophytes	Roca & del Castillo, 1993	12	94	Pyrenees, Spain	
Tracheophytes	Suanjak, 2007	97	13	Austria	
Tracheophytes	Seiler et al., 2021	164	20	Switzerland	
Rotifera	Miracle et al., 1995	107	4	Valencia, Spain	
Rotifera	Jekatierynczuk-Rudczyk & Ejsmont-Karabin, 2023	101	47	Poland	
Nematoda	Gadea, 1968	6	?	Spain	

Nematoda	Ocaña, 1991	44	38	Andalucia, Spain	
Nematoda	Roca, 1996	8	25	Catalonia, Spain	
Nematoda	Zullini et al., 2011	103	108	Trentino, Italy	
Platyhelminthes, Turbellaria	Roca et al., 1992	4	149	Pyrenees, Spain	
Annelida	Roca, 1996	3	25	Catalonia, Spain	
Annelida	Stoch et al., 2011	39	110	Trentino, Italy	
Mollusca, Gastropoda	Gasull, 1971	17	~11	Southeastern Spain	
Mollusca, Gastropoda	Roca, 1996	13	25	Catalonia, Spain	
Mollusca, Gastropoda	Khalloufi et al., 2020	3	3	Tunisia	Endemic species
Mollusca, Gastropoda	Pascual et al., 2020	18	20	Catalonia and Balearic Islands, Spain	
Mollusca, Gastropoda	Falniowski et al., 2021	8		Eastern Herzegovina	
Crustacea, Branchiopoda	Sahuquillo & Miracle, 2013	21	8	Comunitat Valenciana, Spain	
Crustacea, Cladocera	Pascual et al., 2020	5	20	Catalonia and Balearic Islands, Spain	
Crustacea, Copepoda	Bottazzi et al., 2008	19	19	Italy	
Crustacea, Copepoda	Galassi et al., 2011	37	39	Central Italy	
Crustacea, Copepoda	Stoch et al., 2011	24	110	Trentino, Italy	
Crustacea, Copepoda	Fiasca et al., 2014	22	1	Central Italy	2000 m2 seepage area
Crustacea, Ostracoda	Roca & Baltanas, 1993	21	149	Pyrenees, Spain	

Crustacea, Ostracoda	Roca, 1996	9	25	Catalonia, Spain	
Crustacea, Ostracoda	Mezquita et al., 1999	38	117	Eastern Spain	
Crustacea, Ostracoda	Külköylüoğlu, 2005	10	2	Turkey	
Crustacea, Ostracoda	Külköylüoğlu & Yilmaz, 2006	6	3	Turkey	
Crustacea, Ostracoda	Bottazzi et al., 2008	13	19	Italy	
Crustacea, Ostracoda	Stoch et al., 2011	18	110	Trentino, Italy	
Crustacea, Ostracoda	Altınşaçlı et al., 2015	14	7	Turkey	
Crustacea, Ostracoda	Rosati et al., 2017	67	360	Southern Europe	
Crustacea, Ostracoda	Pascual et al., 2020	8	20	Catalonia and Balearic Islands, Spain	
Crustacea, Ostracoda	Rossetti et al., 2022	2	2	Northern Apennines, Italy	
Crustacea, Amphipoda	Žutinić et al., 2018	1	1	Croatia	
Crustacea, Amphipoda	Manenti & Barzaghi, 2021	1	4	Northern Italy	
Crustacea, Amphipoda	Manenti et al., 2023	2	15	Northern Italy	
Crustacea, Isopoda, Asellota	Margalef, 1952	1	1	Balearic Islands, Spain	Endemic subspecies
Insecta, Coleoptera	Roca & del Castillo, 1993	9	94	Pyrenees, Spain	
Insecta, Coleoptera	Roca, 1996	12	25	Catalonia, Spain	
Insecta, Coleoptera	Ilmonen & Paasivirta, 2005	15	6	Southern Finland	
Insecta, Coleoptera	Pascual et al., 2020	40	20	Catalonia and Balearic Islands, Spain	

Insecta, Diptera, Chironomidae	Lencioni et al., 2012	81	124	Alps and Prealps, Italy	
Insecta, Diptera	Ilmonen & Paasivirta, 2005	50	6	Southern Finland	
Insecta, Diptera	Pascual et al., 2020	45	20	Catalonia and Balearic Islands, Spain	
Insecta, Ephemeroptera	Puig & Roca, 1987	6	50	Catalonia, Spain	
Insecta, Ephemeroptera	Roca & del Castillo, 1993	6	94	Pyrenees, Spain	
Insecta, Ephemeroptera	Roca, 1996	9	25	Catalonia, Spain	
Insecta, Ephemeroptera	Savić et al. 2017	6	50	Bosnia and Herzegovina	
Insecta, Ephemeroptera	Pascual et al., 2020	6	20	Catalonia and Balearic Islands, Spain	
Insecta, Hemiptera	Pascual et al., 2020	13	20	Catalonia and Balearic Islands, Spain	
Insecta, Odonata	Roca & del Castillo, 1993	3	94	Pyrenees, Spain	
Insecta, Odonata	Roca, 1996	7	25	Catalonia, Spain	
Insecta, Odonata	Pešić et al., 2017	91	44	Montenegro	
Insecta, Odonata	Pascual et al., 2020	18	20	Catalonia and Balearic Islands, Spain	
Insecta, Odonata	Manenti et al., 2023	3	15	Northern Italy	
Insecta, Plecoptera	Puig & Roca, 1987	6	50	Catalonia, Spain	
Insecta, Plecoptera	Roca & del Castillo, 1993	5	94	Pyrenees, Spain	
Insecta, Plecoptera	Roca, 1996	3	25	Catalonia, Spain	
Insecta, Plecoptera	Ilmonen & Paasivirta, 2005	3	6	Southern Finland	

Insecta, Plecoptera	Savić et al. 2017	5	50	Bosnia and Herzegovina	
Insecta, Trichoptera	Roca & del Castillo, 1993	11	94	Pyrenees, Spain	
Insecta, Trichoptera	Roca, 1996	12	25	Catalonia, Spain	
Insecta, Trichoptera	Cianficconi et al., 1998	136	397	Italy	
Insecta, Trichoptera	Savić et al. 2017	17	50	Bosnia and Herzegovina	
Insecta, Trichoptera	Cerjanec et al., 2020	39	3	Central Croatia	
Insecta, Trichoptera	Ilmonen & Paasivirta, 2005	8	6	Southern Finland	
Insecta, Trichoptera	Pascual et al., 2020	18	20	Catalonia and Balearic Islands, Spain	
Acari	Gil & Roca, 1990	23	33	Pyrenees, Spain	
Acari	Roca & Gil, 1992	22	149	Pyrenees, Spain	
Acari	Gil et al., 1994	28	50	Balearic Islands, Spain	
Acari	Roca, 1996	8	25	Catalonia, Spain	Including an endemic species
Acari	Gerecke et al., 2009	178	104	Trentino, Italy	
Acari	Di Sabatino et al., 2003	163	>300	Italy	
Acari	Stoch et al., 2011	57	110	Trentino, Italy	
Acari	Pascual et al., 2020	5	20	Catalonia and Balearic Islands, Spain	
Amphibia	Roca, 1996	2	25	Catalonia, Spain	
Amphibia	Manenti, 2008	6	95	Lombardy, Italy	
Amphibia	Manenti et al., 2009	1	13	Lombardy, Italy	

Amphibia	Pascual et al., 2020	3	20	Catalonia and Balearic Islands, Spain
Amphibia	Manenti et al., 2023	1	15	Northern Italy

40

41

42 **References from Table S2**

43 Aboal, M., Puig, M. A., & Prefasi, M. (1998). Diatom assemblages in springs in Castellón
 44 province, Eastern Spain. *Algological Studies/Archiv Für Hydrobiologie, Supplement Volumes,*
 45 *90.* https://doi.org/10.1127/algol_stud/90/1998/79

46 Altinsaçli, S., Altinsaçli, S., & Perçin Paçal, F. (2015). Diversity, species composition and
 47 habitat preferences of Ostracoda (Crustacea) in seven karstic springs of varying salinity in
 48 Milas (Muğla, Turkey). *International Journal of Fisheries and Aquatic Studies, 3*(2), 374–389.

49 Bes, M., Corbera, J., Sayol, F., Bagaria, G., Jover Benjumea, M., Preece, C., ... Fernández
 50 Martínez, M. (2017). Efecte de les variables ambientals i hidrològiques sobre la riquesa i
 51 distribució dels briòfits fontinals a la Catalunya oriental. *Butlletí de la Institució Catalana*
 52 *d'Història Natural, 81,* 121–134. <https://raco.cat/index.php/ButlletilCHN/article/view/335567>

53 Bottazzi, E., Bruno, M. C., Mazzini, M., Pieri, V., & Rossetti, G. (2008). First report on
 54 Copepoda and Ostracoda (Crustacea) from northern Apenninic springs (N. Italy): A faunal
 55 and biogeographical account. *Journal of Limnology, 67*(1), 56.
 56 <https://doi.org/10.4081/jlimnol.2008.56>

57 Cantonati, M. (2008). Cyanoprokaryotes and algae other than diatoms in springs and
 58 streams of the Dolomiti Bellunesi National Park (Northern Italy). *Algological Studies, 126.*
 59 <https://doi.org/10.1127/1864-1318/2008/0126-0113>

60 Cantonati, M., Angeli, N., Bertuzzi, E., Spitale, D., & Lange-Bertalot, H. (2012). Diatoms in
 61 springs of the Alps: Spring types, environmental determinants, and substratum. *Freshwater*
 62 *Science, 31*(2), 499–524. <https://doi.org/10.1899/11-065.1>

63 Cantonati, M., Füreder, L., Gerecke, R., Jüttner, I., & Cox, E. J. (2012). Crenic habitats,
 64 hotspots for freshwater biodiversity conservation: toward an understanding of their ecology.
 65 *Freshwater Science, 31*(2), 463–480. <https://doi.org/10.1899/11-111.1>

66 Cantonati, M., Rott, E., Spitale, D., Angeli, N., & Komárek, J. (2012). Are benthic algae
 67 related to spring types? *Freshwater Science, 31*(2), 481–498. <https://doi.org/10.1899/11-048.1>
 68

- 69 Cerjanec, D., Kučinić, M., Vilenica, M., Čukušić, A., Čuk, R., Ibrahim, H., ... Ruk, D. (2020).
70 Ecological and faunistic features of caddisflies (Insecta: Trichoptera) in different types of
71 habitats in the Dinaric karst area (Central Croatia). *Ecologica Montenegrina*, 36, 6–39.
72 <https://doi.org/10.37828/em.2020.36.2>
- 73 Cianficconi, F., Corallini, C., & Moretti, G. P. (1998). Trichopteran fauna of the Italian springs.
74 In *Studies in crenobiology* (pp. 125–140). Leiden: Backhuys Publishers.
- 75 Delgado, C., Ector, L., Novais, M. H., Blanco, S., Hoffmann, L., & Pardo, I. (2013). Epilithic
76 diatoms of springs and spring-fed streams in Majorca Island (Spain) with the description of a
77 new diatom species *Cymbopleura margalefii* sp. nov. *Fottea*, 13(2), 87–104.
78 <https://doi.org/10.5507/fot.2013.009>
- 79 Di Sabatino, A., Cicolani, B., & Gerecke, R. (2003). Biodiversity and distribution of water
80 mites (Acari, Hydrachnidia) in spring habitats. *Freshwater Biology*, 48(12), 2163–2173.
81 <https://doi.org/10.1007/s10641-003-0000-0>
- 82 Falniowski, A., Lewarne, B., Rysiewska, A., Osikowski, A., & Hofman, S. (2021). Crenobiont,
83 stygophile and stygobiont molluscs in the hydrographic area of the Trebišnjica River Basin
84 Launched to accelerate biodiversity research. *ZooKeys*, 1047, 61–89.
85 <https://doi.org/10.3897/zookeys.1047.64034>
- 86 Fernández-Martínez, M., Corbera, J., Domene, X., Sayol, F., Sabater, F., & Preece, C.
87 (2020). Nitrate pollution reduces bryophyte diversity in Mediterranean springs. *Science of the*
88 *Total Environment*, 705, 135823. <https://doi.org/10.1016/j.scitotenv.2019.135823>
- 89 Fiasca, B., Stoch, F., Olivier, M. J., Maazouzi, C., Petitta, M., Di Cioccio, A., & Galassi, D. M.
90 P. (2014). The dark side of springs: What drives small-scale spatial patterns of subsurface
91 meiofaunal assemblages? *Journal of Limnology*, 73(1).
92 <https://doi.org/10.4081/jlimnol.2014.848>
- 93 Gadea, E. (1968). La fauna nematòdica de les torberes. *Treballs de la Societat Catalana de*
94 *Biologia*, 23, 30–36.
- 95 Galassi, D. M. P., Fiasca, B., & Del Tosto, D. (2011). Patterns of copepod diversity
96 (Copepoda: Cyclopoida, Harpacticoida) in springs of central Italy: Implications for
97 conservation issues. In D. Defaye, E. Suarez-Morales, & J. C. von Vaupel Klein (Eds.),
98 *Studies on Freshwater Copepoda: a Volume in Honour of Bernard Dussart* (pp. 199–226).
99 Leiden: Brill. <https://doi.org/10.1163/ej.9789004181380.i-566.78>

100 Gasull, L. (1971). Fauna malacológica de las aguas continentales dulces y salobres del
101 sudeste ibérico. *Boletín de la Sociedad de Historia Natural de Baleares*, 16, 23–92.

102 Gerecke, R., Schatz, H., & Wohltmann, A. (2009). The mites (Chelicerata: Acari) of the
103 CRENODAT project: Faunistic records and ecological data from springs in the autonomous
104 province of Trento (Italian Alps). *International Journal of Acarology*, 35(4), 303–333.
105 <https://doi.org/10.1080/01647950903059452>

106 Gil, M. J., & Roca, J. R. (1990). Hidracnelas de fuentes de los Pirineos Centrales. *Scientia
107 Gerundensis*, 16(1), 107–113.

108 Gil, M. J., Llobera, M., & Ferriol, A. (1994). Water mites (Hydrachnellae, Acari) in Majorcan
109 springs. *Internationale Vereinigung Für Theoretische Und Angewandte
110 Limnologie:Verhandlungen*, 25(3), 1449–1451.
111 <https://doi.org/10.1080/03680770.1992.11900412>

112 Guasch, H., Acosta, X. G., Urrea, G., & Bañeras, L. (2012). Changes in the microbial
113 communities along the environmental gradient created by a small Fe spring. *Freshwater
114 Science*, 31(2), 599–609. <https://doi.org/10.1899/10-127.1>

115 Ilmonen, J., & Paasivirta, L. (2005). Benthic macrocrustacean and insect assemblages in
116 relation to spring habitat characteristics: patterns in abundance and diversity. *Hydrobiologia*,
117 533(1–3), 99–113. <https://doi.org/10.1007/s10750-004-2399-4>

118 Jekatierynczuk-Rudczyk, E., & Ejsmont-Karabin, J. (2023). Rotifers of Inter-Forest Springs.
119 *Diversity*, 15(2), 153. <https://doi.org/10.3390/d15020153>

120 Khalloufi, N., Béjaoui, M., & Delicado, D. (2020). Two new genera and three new
121 subterranean species of Hydrobiidae (Caenogastropoda: Truncatelloidea) from Tunisia.
122 *European Journal of Taxonomy*, 648.

123 Kùlköylüođlu, O. (2005). Ecological requirements of freshwater Ostracoda (Crustacea) in two
124 limnocene springs (Bolu, Turkey). *Annales de Limnologie*, 41(4), 237–246.
125 <https://doi.org/10.1051/limn/2005016>

126 Kùlköylüođlu, O., & Yilmaz, F. (2006). Ecological requirements of Ostracoda (Crustacea) in
127 three types of springs in Turkey. *Limnologica*, 36(3), 172–180.
128 <https://doi.org/10.1016/j.limno.2006.03.003>

129 Lai, G. G., Padedda, B. M., Ector, L., Wetzel, C. E., Lugliè, A., & Cantonati, M. (2020).
130 Mediterranean karst springs: diatom biodiversity hotspots under the pressure of hydrological

- 131 fluctuation and nutrient enrichment. *Plant Biosystems - An International Journal Dealing with*
132 *All Aspects of Plant Biology*, 154(5), 673–684.
133 <https://doi.org/10.1080/11263504.2019.1674402>
- 134 Lencioni, V., Marziali, L., & Rossaro, B. (2012). Chironomids as bioindicators of
135 environmental quality in mountain springs. *Freshwater Science*, 31(2), 525–541.
136 <https://doi.org/10.1899/11-038.1>
- 137 Manenti, R. (2008). Amphibiens des sources et galeries drainantes en territoire préalpin,
138 l'exemple du Mont Barro et du Mont de Brianza (Lombardie, Italie). *Bulletin de La Société*
139 *Herpétologique de France*, 128, 25–40.
- 140 Manenti, R., & Barzaghi, B. (2021). Diel activity of *Niphargus* amphipods in spring habitats.
141 *Crustaceana*. <https://doi.org/10.1163/15685403-bja10117>
- 142 Manenti, R., Ficetola, G. F., Bianchi, B., & de Bernardi, F. (2009). Habitat features and
143 distribution of *Salamandra salamandra* in underground springs. *Acta Herpetologica*, 4(2),
144 143–151.
- 145 Manenti, R., Galbiati, M., Lapadula, S., Forlani, M., Barzaghi, B., Melotto, A., & Ficetola, G.
146 F. (2023). Behavioural drivers of ecotone exploitation: activity of groundwater animals in
147 spring. *Behavioral Ecology and Sociobiology*, 77(2), 23. [https://doi.org/10.1007/s00265-023-](https://doi.org/10.1007/s00265-023-03297-8)
148 [03297-8](https://doi.org/10.1007/s00265-023-03297-8)
- 149 Margalef, R. (1952). Une *Jaëra* dans les eaux douces des Baléares, *Jaëra balearica* nov. sp.
150 (Isopoda Asellota). *Hydrobiologia*, 4(1–2), 209–213. <https://doi.org/10.1007/BF00052900>
- 151 Mezquita, F., Tapia, G., & Roca, J. R. (1999). Ostracoda from springs on the eastern Iberian
152 Peninsula: ecology, biogeography and palaeolimnological implications. *Palaeogeography,*
153 *Palaeoclimatology, Palaeoecology*, 148(1–3), 65–85. [https://doi.org/10.1016/S0031-](https://doi.org/10.1016/S0031-0182(98)00176-X)
154 [0182\(98\)00176-X](https://doi.org/10.1016/S0031-0182(98)00176-X)
- 155 Miracle, M. R., Alfonso, M. T., Vicente, E., & Koste, W. (1995). Rotifers of spring pools in the
156 coastal marshland of Albufera of Valencia natural park. *Limnetica*, 11(2), 39–47.
157 <https://doi.org/10.23818/limn.11.12>
- 158 Nowicka-Krawczyk, P. B., & Żelazna-Wieczorek, J. (2013). Cyanobacteria microflora in a
159 limestone spring (Troniny spring, Central Poland). *Acta Societatis Botanicorum Poloniae*,
160 82(3), 219–224. <https://doi.org/10.5586/asbp.2013.017>
- 161 Ocaña, A. (1991). Relationship between nematode species and the physico-chemical
162 characteristics of spring waters. I. Conductivity. *Nematologia Mediterranea*, 19, 19–24.

- 163 Pascual, R., Gomà, J., Pedrocchi, C., Cadiach, O., García, G., & Solé, J. (2020). First data
164 on the biological richness of Mediterranean springs. *Limnetica*, 39(1), 121–139.
165 <https://doi.org/10.23818/limn.39.09>
- 166 Pešić, V., Gligorović, B., Savić, A., & Buczyński, P. (2017). Ecological patterns of Odonata
167 assemblages in karst springs in central Montenegro. *Knowledge & Management of Aquatic*
168 *Ecosystems*, (418), 3. <https://doi.org/10.1051/kmae/2016035>
- 169 Puig, M., & Roca, J. R. (1987). Efemerópteros y plecópteros de las fuentes del Prepirineo
170 Leridano (NE-España). In *Resúmenes del IV Congreso Español de Limnografía*. Sevilla.
- 171 Roca, J. R. (1996). La vida a les fonts del Montsec. In E. Fanlo (Ed.), *El Patrimoni Natural*
172 *del Montsec. Actes de les VI Jornades de Coneixement del Patrimoni Natural del Montsec*
173 (pp. 159–172). Lleida: Institut d'Estudis Ilerdencs.
- 174 Roca, J. R., & Baltanas, A. (1993). Ecology and Distribution of Ostracoda in Pyrenean
175 Springs. *Journal of Crustacean Biology*, 13(1), 165-174. <https://doi.org/10.2307/1549131>
- 176 Roca, J. R., & del Castillo, M. (1993). Typology of the Pyrenean springs based on their
177 biological communities. *Limnetica*, 6(1), 283–289. <https://doi.org/10.23818/actas.06.36>
- 178 Roca, J. R., & Gil, M. J. (1992). Ecological and historical factors affecting the distribution of
179 water-mites (Hydracnellae, Acari) in pyrenean springs. *Archiv Für Hydrobiologie*, 125(2),
180 227–244. <https://doi.org/10.1127/archiv-hydrobiol/125/1992/227>
- 181 Roca, J. R., Ribas, M., & Baguna, J. (1992). Distribution, ecology, mode of reproduction and
182 karyology of freshwater planarians (Platyhelminthes; Turbellaria; Tricladida) in the springs of
183 the central Pyrenees. *Ecography*, 15(4), 373–384. <https://doi.org/10.1111/j.1600-0587.1992.tb00047.x>
- 184 Rosati, M., Rossetti, G., Cantonati, M., Pieri, V., Roca, J. R., &
185 Mesquita-Joanes, F. (2017). Are aquatic assemblages from small water bodies more
186 stochastic in dryer climates? An analysis of ostracod spring metacommunities.
187 *Hydrobiologia*, 793(1), 199–212. <https://doi.org/10.1007/s10750-016-2938-9>
- 188 Rossetti, G., Stoch, F., & Mazzini, I. (2022). A reassessment of the origin and distribution of
189 the subterranean genus *Pseudolimnocythere* Klie, 1938 (Ostracoda, Loxoconchidae), with
190 description of two new species from Italy. *Subterranean Biology*, 43, 33–60.
191 <https://doi.org/10.3897/subtbiol.43.82158>
- 192 Sabater, S., & Roca, J. R. (1992). Ecological and biogeographical aspects of diatom
193 distribution in Pyrenean springs. *British Phycological Journal*, 27(2), 203–213.
194 <https://doi.org/10.1080/00071619200650201>

195 Sahuquillo, M., & Miracle, M. R. (2013). The role of historic and climatic factors in the
196 distribution of crustacean communities in Iberian Mediterranean ponds. *Freshwater Biology*,
197 58(6), 1251–1266. <https://doi.org/10.1111/fwb.12124>

198 Savić, A., Dmitrović, D., & Pešić, V. (2017). Ephemeroptera, plecoptera, and trichoptera
199 assemblages of karst springs in relation to some environmental factors: A case study in
200 central bosnia and herzegovina. *Turkish Journal of Zoology*, 41(1), 119–129.
201 <https://doi.org/10.3906/zoo-1512-31>

202 Seiler, H., Küry, D., Billeter, R., & Dengler, J. (2021). Regional typology of spring vegetation
203 in Parc Ela (Grisons, Switzerland). *Vegetation Classification and Survey*, 2, 257–274.
204 <https://doi.org/10.3897/VCS/2021/69101>

205 Stock, F., Gerecke, R., Pieri, V., Rossetti, G., & Sambugar, B. (2011). Exploring species
206 distribution of spring meiofauna (Annelida, Acari, Crustacea) in the south-eastern Alps.
207 *Journal of Limnology*, 70(Suppl. 1), 65–76. <https://doi.org/10.3274/JL11-70-S1-05>

208 Suanjak, M. (2007). *Die Vegetation moosdominierter Quellfluren im Nationalpark Gesäuse*.
209 Weng im Gesäuse: Nationalpark Gesäuse GmbH.

210 Zullini, A., Gatti, F., & Ambrosini, R. (2011). Microhabitat preferences in springs, as shown by
211 a survey of nematode communities of Trentino (south-eastern Alps, Italy). *Journal of*
212 *Limnology*, 70(1s), 93–105. <https://doi.org/10.4081/jlimnol.2011.s1.93>

213 Žutinić, P., Petrić, I., Gottstein, S., Gligora Udovič, M., Kralj Borojević, K., Kamberović, J., ...
214 Ternjej, I. (2018). Microbial mats as shelter microhabitat for amphipods in an intermittent
215 karstic spring. *Knowledge & Management of Aquatic Ecosystems*, 419, 7.
216 <https://doi.org/10.1051/kmae/2017061>

217

218

219 **Table S3.** List of traditions and cultural manifestations linked to fountains in Mediterranean
 220 regions around the world.
 221

Country	Cultural manifestation and use	Reference
Chile	<p>Springs are still used nowadays in small towns and villages and for agricultural practices, and their use dates back to pre-hispanic times. However, in cities springs have not been respected and most of them have been lost. Springs were intensively used during the earthquake of 1939 to ensure the survival of the population, but were forgotten soon after.</p> <p>Water from the “Pilonos de Antuco” are supposed to provide good health for those that touch it or drink it.</p>	<p>Ricardo Figueroa Muñoz and Rocha, 2020. Webpage #1</p>
Portugal	<p>Springs are used as a source of drinking water. In the past, they were also used for doing the laundry.</p> <p>Big karst springs are also used as swimming pools in summer and speleologists go diving there, and in Roman times, they were part of a complex water supply system for nearby cities.</p>	<p>Manuel Graça Verónica Ferreira</p>
USA	<p>Springs were important landmarks in the landscape for California tribes (e.g., the Mattole in northwestern California), who used them to demarcate tribal territories and mentioned them in myths and legends. They were amongst the landscape features which were interpreted as signs of the activities of ancestors. Springs also became sources of conflict with white settlers who began to exploit them commercially (e.g., Warner Hot Springs, Cupeños people).</p> <p>There are also records from healing springs in southern California.</p> <p>Santa Monica, a city adjoining Los Angeles, hosts a spring system that was of religious importance to the Tongva people, and is currently registered as a California Landmark as well as a Sacred Site by the Native American Heritage Commission. The springs are also supposedly related to how Santa Monica got its name: Franciscan missionaries likened the spring water to the tears shed by Saint Monica</p> <p>Some springs in California have almost a mystical quality. Their water is drunk for healing during ceremonies around</p>	<p>Anderson 2005 Webpage #2 Webpage #3 Vincent H. Resh</p>

	<p>the time of their festivity. There are also scores of commercial springs used for bottled water. Some water bars have 20 or more types of spring water for tasting (at very high prices). Soaking in springs for health and meditation is widespread in Mediterranean areas in California, and many are very successful commercial enterprises. All advertise the healing, mental, physical, and spiritual values of springs.</p> <p>Cattle troughs are directly filled by spring water and their overflow support rich biological communities.</p>	
Greece	<p>In Greek mythology, under the collective name Pigiiai Nymfai, there are well-known personifications of natural forces (beyond any association with springs). They had the form of virgins, who were considered the direct daughters of Zeus. The Spring Nymphs were partially identified in places with the Naiads, the river, "Fountain" and "Lake" Nymphs.</p> <p>The Waters of Styx are the sources of the Krathis River in Helmos, Kalavryta District in Achaia at an altitude of 2,100 metres. The waters of the Styx were associated with theological and philosophical ideas, such as those from the Eleusinian Mysteries and Orphic beliefs about reincarnation. According to mythology, Styx was an Ocean woman who had her palace in Tartarus and was guarded day and night by sleepless dragons. It was believed that the waters of the spring came from there, from Tartarus and the palace of Styx. In the waters of the Styx all the gods, even the Sun, swore an oath: it was the greatest oath a god could make, and the gods served their sentences there when they were punished. They said that any living being that drank from its water died, and any metal dipped in its waters melted. It was there that Nereida Thetis dipped her son Achilles and he became invulnerable, but she held him by the heel and that remained the only vulnerable spot on his body.</p> <p>Legends, traditions and mythical beings, such as the nymphs of antiquity or the soothsayer of folk tradition, accompany the waters of springs and rivers.</p> <p>Fountains were a reason for the establishment of settlements and churches in a certain location, as well as for the creation of roads. They were a main pole of social attraction, especially in villages. The spring was a daily</p>	<p>Nikos Skoulikidis Σουέρεφ, 2000 Webpage #4 Ginouvés et al., 1994 Photos-Jones et al., 2016 Håland 2009</p>

	<p>meeting point for women. They constituted a place for the performance of important folkloric and cultic customs, especially when they had or were thought to have healing, magical or other properties, such as eugenics or eutecnia.</p> <p>The Castalian spring in the sacred precinct of Delphi consists of two monumental fountains that received the water from the emergence, dating back to the <u>Archaic</u> and <u>Roman</u> periods.</p> <p>Sanctuaries and oracles (e.g., Trophonius at Levadeia, Apollo at Delphi), and healing centres (e.g., Asklepios at Epidauros) were always in the proximity of springs, from where the water was used for drinking.</p> <p>In Lemnos there are six holy springs associated with a saint that are built close to a chapel. Recent studies did not find substantial hydrogeological differences between holy and ordinary springs.</p> <p>Still nowadays, Greek people keep water-rituals in springs in caves that relate to pre-Christian traditions. While in ancient times, springs were related to water-nymphs, nowadays springs are dedicated to the Virgin Mary. In Athens, during the first Friday after Easter Sunday, people celebrate the “life-giving spring” festival, on the Southern Slope of the Acropolis, in Panagia’s chapel inside a circular Spring House.</p>	
Palestina	Some springs are believed to be haunted, and in them one can find water demons. Both, springs and demons, eventually became part of the Greek cultural and religious heritage.	Canaan 1992
Italy	<p><i>Fontinalia</i> was a festival of the Roman religion dedicated to the sacred springs whose god was <i>Fontus</i> (or <i>Fons</i> according to Oxford Classical Dictionary). During the <i>Fontinalia</i> garlands of flowers were thrown into the fountains and sacrifices of wine, oil, etc. were offered to the god. The festival took place on October 13th at the Fontinalis gate of the Servian walls.</p> <p>Fonte Aretusa (Spring Arethusa) is known since the 7th century BCE when the Greeks established one of their colonies there. Mythology explains that Arethusa was a nymph who was turned into a fountain by the goddess Artemis to protect her from Alpheus. It is mentioned by a</p>	<p>Webpage #5</p> <p>Luzzini 2015</p> <p>Bouffier 2019</p> <p>Polto 2001</p>

	<p>large number of poets, writers and travelers of all times (e.g., Pindalo, Ovidio, Virgilio, Ciceron amongst others). Kyane was another nymph that also became a spring in Syracuse.</p>	
<p>Maghreb, Algeria and North Africa</p>	<p>Springs have not only quenched the physical thirst of the Maghrebians, but have also nourished their spiritual and cultural identity, creating a harmonious blend of tradition, faith, and natural beauty. This enduring connection between springs and culture is a testament to the resilience of ancient practices in the face of changing times.</p> <p>Springs are part of the cultural fabric of the Maghreb, where their presence is not only felt but celebrated in various ceremonies and festivals. The landscape of the Maghreb is adorned with the names of springs, each with a unique history and meaning. Many towns and villages bear Arabic or Tamazigh names such as Aïn Makhoulf (Makhoulf's Spring), Aïn Beïda (White Spring), Aïn Berda (Cold Spring), and Tala Rana (Spring of the Frogs), which are living testimony to the vital role of springs in this region.</p> <p>In Islam, a cornerstone of Maghrebian identity, water is considered a symbol of purity. Therefore, many pious people or saints find their eternal rest near these life-giving springs, which gives these natural places a sacred aura. The tradition of dedicating festivals to these revered saints and to the springs themselves, although criticized by Islamic purists, has its roots in times long before the arrival of Islam in the region. If one goes to the region around Annaba, one encounters an abundance of such "holy" springs, each with its own history. Kattara, the 'oozing or leaking spring,' and Aïn Bent Soltane, the 'spring of the sultan's daughter,' are just a few examples that illustrate the enduring connection between springs and the spiritual life of the Maghreb. This fusion of faith and nature not only serves practical purposes, but also strengthens the spiritual bond between people and their environment.</p> <p>Moreover, the ancient Roman heritage in the Maghreb is characterized by an extensive network of aqueducts, baths and fountains that harnessed the power of springs. For example, the ancient city of Hippone (now Annaba), where Saint Augustine lived and died, was supplied by Roman aqueducts that brought spring water from Mount Edough.</p>	<p>Boudjéma Samraoui</p> <p>Farrah Samraoui</p>

	<p>Large white marble Gorgon masks decorated the fountains where the water was distributed.</p> <p>Finally, the art of the Foggaras is an essential part of the cultural and economic heritage of the Maghreb, especially in the arid expanses of the Sahara. Foggaras, an elaborate system of underground tunnels and canals, are remarkable water infrastructures that have enabled Maghrebians to harness the precious water hidden beneath the harsh desert landscape (wadis, sand dunes and mountain springs). In folk culture, foggaras are also a symbol of unity and community effort. The construction and maintenance of foggaras require the collective efforts of entire communities and strengthen the bonds between their members by instilling values of cooperation, shared responsibility, and mutual support. The knowledge and skills of the foggaras' water measurers were recognized as Intangible Cultural Heritage by UNESCO in 2018.</p>	
France	<p>The fountain of Lourdes was discovered by Bernadette Soubirous, who claimed that the Virgin Mary told her “come to the spring, drink its water and clean yourself with it”. It is considered to be miraculous water. Around 70 miraculous healings of various diseases thanks to drinking the spring water have been acknowledged by the Catholic Church. Nowadays, the Sanctuary of Lourdes and its spring is a reference site of pilgrimage for Christian people, where they can drink its water freely.</p> <p>In Provence, several of Pagnol's novels (Jean de Florette, Manon de la Source) are based on the power and importance of springs for survival.</p>	<p>Webpage #6</p> <p>Vincent H. Resh</p>
Turkey	<p>There is a legend about a Saint called Munzur (a shepherd) in the Munzur Valley (Eastern Turkey). It is believed that he disappeared here when he was carrying a pail with milk. The milk was spilled and water began flowing out of the earth there, forming the Munzur Spring. The Alevi Kurds believe this place is sacred. They go there to make a vow, light candles, take water from the spring, hang fabrics on the trees, and to make wishes. It is considered a religious visit. Munzur Spring, however, is not the only spring they visit for these purposes, they also visit different ones around the region.</p>	<p>Zeynep Ersoy</p> <p>Webpage #7</p>
Spain	<p>In several regions of Spain, springs are often dedicated to saints and the Virgin Mary. It is traditional to make a celebration in the spring on the day of the saint, often</p>	<p>Signing authors.</p> <p>Farrerons Vidal</p>

	<p>accompanied with music, food and drinking the water of the spring. In Catalonia these celebrations receive the name of “aplecs” and were very popular until a few decades ago.</p> <p>Multiple legends are also attributed to springs. The Springs of Llobregat (where the Llobregat river starts) are supposed to be the tears of girls that were killed by an evil witch who wanted her ugly daughter to become the most beautiful girl in town. In Montseny, the “Spring of the Big Boy” receives that name after a scrawny little boy became big and strong after drinking a gulp of water from that spring every day. In Covadonga, if someone drinks from all the spouts of the “Marriage Spring” or the “Spring of the Seven Spouts” without breathing in between, they will get married within a year.</p> <p>Some springs have also been attributed healing or medicinal properties for all sort of illnesses (called <i>mineromedicinal</i> water). Just to provide a few examples, bicarbonated water is supposed to help with digestion and liver problems, ferruginous and sparkling water have been described to cure anemia, infertility, and even spermatorrhea, while sulphurous springs have been suggested to help with eye and skin problems. These beliefs led to the appearance of multiple health resorts surrounding these springs.</p>	<p>and Prat, 2020</p> <p>Webpage #8</p> <p>Webpage #9</p>
Australia	<p>In very arid regions in South Australia springs were of paramount importance for trading routes and survival of indigenous people. Hence, springs were very present in the culture of the people, through myths, songs and rituals. However, these traditions rapidly decayed only a few decades after the European occupation. People very strongly identified with springs in the Kuyani, Arabana and Wangkangurru traditions.</p>	Harris 2002

222

223 **References from Table S3**

224 Anderson, M.K. (2005). *Tending the Wild: Native American Knowledge and the Management*
 225 *of California’s Natural Resources*. University of California Press, Berkeley. pp. 39, 107.

226 Bouffier, S. (2019). Arethusa and Kyane, Nymphs and Springs in Syracuse: Between Greece
 227 and Sicily. *Ancient Waterlands*, 159-181. <https://doi.org/10.4000/books.pup.40610>

- 228 Canaan, T. (1922). Haunted springs and water demons in Palestine. *Studies in Palestinian*
229 *customs and folklore (vol II)*. The Palestine Oriental Society. Jerusalem. pp. 153 – 170.
- 230 Σουέρεφ, Κ. (2000). Υδάτινες σχέσεις, το νερό ως πηγή ζωής κατά την αρχαιότητα.
231 Θεσσαλονίκη: University Studio Press.
232
- 233 Farrerons Vidal, O., & Prat, F. (2020). Elements químics a l'aigua de les de fonts del
234 Montseny. *L'Atzavara*, 2020(30), 89-96.
- 235 Ginouvès, R., Guimier-Sorbets, A.-M., Jouanna, I and Villard, L., (eds.), 1994, L'eau, la santé
236 et la maladie dans le monde greco Actes du colloque de Paris (25-27 nov. 1992), Bulletin de
237 Correspondance Hellénique, Supplément XXVIII.
- 238 Håland, E. J. (2009). Water sources and the sacred in modern and ancient Greece and
239 beyond. *Water History*, 1(2), 83-108.
- 240 Harris, C. (2002). Culture and geography: South Australia's mound springs as trade and
241 communication routes. *Historic Environment*, 16(2), 8-11.
- 242 Luzzini, F. (2015). Il mistero e la bellezza. La Fonte Aretusa tra mito, storia e scienza. *Acque*
243 *Sotteranee-Italian Journal of Groundwater*, 4(3). <https://doi.org/10.7343/as-127-15-0154>
- 244 Muñoz, I., & Rocha, J. (2020). Vertientes y acequias: explotación de los recursos hídricos y
245 el sistema de riego en el valle de Azapa en tiempos prehispánicos. *Idesia (Arica)*, 38(2), 65-
246 76. <https://dx.doi.org/10.4067/S0718-34292020000200065>
- 247 Photos-Jones, E., Zagana, E., & Roumelioti, P. (2016). The ayiasmata (holy springs) of
248 Lemnos, NE Greece: An archaeo-hydrogeological study. In *Proceedings of the 6th*
249 *Symposium of the Hellenic Society for Archaeometry; Photos-Jones, E., Basiakos, I.,*
250 *Filippaki, E., Hein, A., Karatasios, I., Kilikoglou, V., Kouloumpi, E., Eds* (pp. 221-229).
- 251 Polto, C. (2001). La Fontana Aretusa tra mito e realtà. In C. Masetti (Ed.), *Chiare, fresche e*
252 *dolci acque. Le sorgenti nell'esperienza odeporica e nella storia del territorio* (pp. 11–25).
253 Roma: CISGE.
- 254
- 255 *Webpages (all accessed 19/09/2023)*
- 256 1. [https://www.latribuna.cl/noticias/2021/02/27/la-leyenda-de-las-aguas-que-emanan-de-los-](https://www.latribuna.cl/noticias/2021/02/27/la-leyenda-de-las-aguas-que-emanan-de-los-pilones-de-antuco.html?amp=1)
257 [pilones-de-antuco.html?amp=1](https://www.latribuna.cl/noticias/2021/02/27/la-leyenda-de-las-aguas-que-emanan-de-los-pilones-de-antuco.html?amp=1).

- 258 2. [https://www.latimes.com/california/story/2021-12-28/a-brief-history-of-southern-californias-](https://www.latimes.com/california/story/2021-12-28/a-brief-history-of-southern-californias-hot-springs)
259 [hot-springs](https://www.latimes.com/california/story/2021-12-28/a-brief-history-of-southern-californias-hot-springs)
- 260 3. http://gabrielinosprings.com/wpsite/?page_id=385
- 261 4. <https://en.wikipedia.org/wiki/Delphi>
- 262 5. <https://it.wikipedia.org/wiki/Fontinalia>
- 263 6. https://fr.wikipedia.org/wiki/Eau_de_Lourdes
- 264 7. <http://www.munzurvalley.com/#intro1>
- 265 8. <https://conversesacatalunya.cat/les-fonts-del-llobregat-i-la-seva-llegenda/>
- 266 9. [https://parcs.diba.cat/web/el-patrimoni-cultural-immaterial-del-montseny/inventari/detall/-](https://parcs.diba.cat/web/el-patrimoni-cultural-immaterial-del-montseny/inventari/detall/-/contingut/29193465/llegenda-la-font-del-noi-gran)
267 [/contingut/29193465/llegenda-la-font-del-noi-gran](https://parcs.diba.cat/web/el-patrimoni-cultural-immaterial-del-montseny/inventari/detall/-/contingut/29193465/llegenda-la-font-del-noi-gran)
- 268
- 269