



Connor-Davidson Resilience Scale: a systematic review psychometrics properties using the COSMIN

Hamid Sharif-Nia, PhD^{a,b}, David Sánchez-Teruel, PhD^j, Erika Sivarajan Froelicher, PhD^j, Sima Hejazi, PhD^d, Lida Hosseini, PhD^e, Fatemeh Khoshnavay Fomani, PhD^f, Mozhgan Moshtagh, PhD^g, Fereshteh Mollaei, PhD^h, Amir Hossein Goudarzian, PhD^{c,f,*}, Amir Babaei, BSc^c

Background: Psychometrical evaluation of persons of diverse contexts and different populations, including general or clinical.

Objective: This review study aimed to evaluate the psychometrics quality of resilience scales.

Methods: International and Iranian databases were searched with MESH terms, including “psychometric”, “validity”, “reliability”, “Connor-Davidson resilience scale”, “Resilience scale”, for published articles up to 1 February 2023. For each of the selected studies, the risk of bias was evaluated using the COSMIN Risk of Bias Checklist. Then the COSMIN checklist was used to evaluate the entire text of the article for methodological quality.

Results: Considering the inclusion criteria, 80 documents were evaluated. According to the COSMIN’s criteria for evaluating the risk of bias, the current study findings revealed the included studies’ limitations in assessing the three versions of CD-RISC cross-cultural and content validity as well as their stability (e.g. conducting test re-test), whereas the majority of psychometric studies of CD-RISC-25, and CD-RISC-2 rated as very good or adequate in terms of structural validity. In terms of quality assessment of the included studies, the current study indicated that investigating the structural validity of the CD-RISC was mainly done based on exploratory factor analysis (EFA), and confirmatory factor analysis was absent.

Conclusion: The general result indicates the acceptability of the quality of the studies. However, concerns for measurement properties such as responsiveness and criterion validity as well as the standard error of measurement have been neglected.

Keywords: Connor-Davidson Resilience Scale, COSMIN, psychometrics, resilience, systematic review

Introduction

Resilience is a multidimensional and complex concept that has achieved popularity in positive psychology as a significant human strength due to its impact on psycho-somatic and social health^[1,2]. Facing stress (internal and external) is not avoidable in different stages of the human life course^[3]. Thus, maintaining individual function or bio-psychological balance, and even reaching evolution or thriving after dealing with adversities and disturbances, has resulted in scientists’ interest in resilience in different disciplines^[4]. Based on various definitions, resilience is an adaptive response to stressful or threatening conditions^[5].

Individuals can help themselves through different mechanisms, including enhancing their strength by relying on available resources and capacities or improving their flexibility and stamina by applying coping strategies efficiently^[6]. As mentioned earlier, resilience is a complex concept, several scales, therefore, have been developed; among them, the Connor-Davidson scale of Resilience (CD-RISC) has achieved the highest rating and validation in different contexts^[7,8]. However, psychometric properties or the number of factors and items vary in those studies.

Being cross-cultural and complicated conception, professionals in different disciplines have multi-criteria perspectives on resilience definition and debates regarding its determinants^[6].

^aTraditional and Complementary Medicine Research Center, Addiction Institute Mazandaran University of Medical Sciences, Sari, Iran, ^bDepartment of Nursing, Amol Faculty of Nursing and Midwifery, Mazandaran University of Medical Sciences, Sari, Iran, ^cStudent Research Committee, Mazandaran University of Medical Sciences, Sari, Iran, ^dBojnurd Faculty of Nursing, North Khorasan University of Medical Sciences, Bojnurd, Iran, ^eSchool of Nursing and Midwifery, Iran University of Medical Sciences Tehran, Iran, ^fSchool of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran Iran, ^gSocial Determinants of Health Research Center, Birjand University of Medical Sciences, Birjand Iran, ^hGuilan University of Medical Sciences, Rasht, Iran, ⁱDepartment of Personality, Assessment and Psychological Treatment, University of Granada, Granada, Spain and ^jDepartment of Physiological Nursing, School of Nursing, and Department of Epidemiology & Biostatistics, School of Medicine, University of California San Francisco, San Francisco, CA

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

*Corresponding author. Address: Student Research Committee, Mazandaran University of Medical Sciences, Sari, Iran. Tel.: +98 938 138 1456. E-mail: amir_sari@yahoo.com (A. H. Goudarzian).

Copyright © 2024 The Author(s). Published by Wolters Kluwer Health, Inc. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Annals of Medicine & Surgery (2024) 86:2976–2991

Received 25 January 2024; Accepted 6 March 2024

Published online 27 March 2024

<http://dx.doi.org/10.1097/MS9.0000000000001968>

Some researchers have focused on attitudes (internal capacity), but others believe in behaviours and relations (supportive resources) or a combination of both as aspects of resilience^[9,10]. Since the items mentioned above, examples of adaptations in different countries should also be included, showing the diversity of existing versions of the CD-RISC: original 25 items, 10 items and 2 items^[10].

Based on the evidence, resilience has a significant association with mental and social health as it significantly affects the process and outcome of developmental transitions in the life course^[11–13]. Therefore, a valid and standard measure such as CD-RISC would be necessary for its assessment. Psychometric evaluation among diverse contexts and different populations, including general or clinical, and individuals under stressful situations such as life-threatening conditions, will require the development of psychosocial interventions. This review study aimed to evaluate the psychometrics quality of CD-RISC.

Methods

A systematic review was conducted in February 2023 on all psychometric adaptations of the CD-RISC using the PRISMA, Supplemental Digital Content 1, <http://links.lww.com/MS9/A411> and AMSTAR 2, Supplemental Digital Content 2, <http://links.lww.com/MS9/A412> (Assessing the methodological quality of systematic reviews) guidelines^[14,15].

Search strategy and selection criteria

We used international databases, including PubMed, Scopus, Web of Science, and Iranian databases, including SID, and Magiran and evaluated studies published up to 1 February 2023. MESH terms included “psychometric”, “validity”, “reliability”, “Connor-Davidson resilience scale”, “Resilience scale”, [with the use of ‘OR’ and ‘AND’ operators] and were used for assessment of the identified databases. Appendix 1 provides details about the search strategy, Supplemental Digital Content 3, <http://links.lww.com/MS9/A413>.

First, duplicate studies from the initial search were removed after selected studies were entered into the Endnote software. Two researchers with doctoral degrees (A.H.G. and S.H.) obtained original articles and then carefully evaluated them for inclusion. Studies needed to meet the following criteria for inclusion: (1) use of CD-RISC in study; and (2) Reliability assessment; (3) Validity assessment; (4) Evaluation of the diagnostic accuracy of CD-RISC. Studies that did not provide information regarding validity and/or reliability assessments (e.g. factor structure); (1) had an insufficient sample size (< 50 participants), (2) they used tools other than the CD-RISC to measure resilience, (3) unable to access full-text of the article, (4) only used Item Response Theory (IRT) for data analysis, (5) were written in languages other than English or Farsi were excluded. The phases of article selection were based on PRISMA guidelines are shown in Fig. 1.

Risk of bias assessment

For each of the selected studies, the risk of bias was evaluated using the COSMIN Risk of Bias Checklist (include reference(s) to this procedure here). This tool includes three parts with 10 boxes. Boxes 1 and 2 are in the first part, which deal with content

HIGHLIGHTS

- Connor-Davidson Resilience Scale is a popular scale in psychology.
- This was the first study that assessed the risk of bias and quality of methodological studies about the Connor-Davidson Resilience Scale.
- The majority of methodological studies were rated as very good or adequate in terms of the structural validity of this scale.

validity. This section evaluates how well each item fits the target construct and population, as well as its relevance and comprehensibility. The internal structure is addressed in the box of the second part) (boxes 3, 4, and 5, which also include information on structural validity, internal consistency, and cross-cultural validity/measurement invariance. The third section is broken down into boxes 6, 7, 8, 9, and 10 and deals with the remaining measurement properties, such as responsiveness, criterion validity, reliability, measurement error, and reliability of hypotheses. Instead of emphasizing individual items, the third section concentrates on the overall quality of the (sub) scale^[16]. This and the following procedures were carried out by researchers other than the above (L.H. and S.H.).

Quality assessment

The COSMIN checklist was used (by L.H. and S.H.) to evaluate the entire text of the article for methodological quality. A variety of psychometric properties are evaluated by the COSMIN checklist, including (i) structural validity, (ii) internal consistency, (iii) reliability, (iv) measurement error, (v) hypotheses testing for construct validity, (vi) Cross-cultural validity\measurement invariance, (vii) criterion validity, and (viii) responsiveness. The result of every single study on a measurement property is rated against the updated criteria for good measurement properties as either sufficient (+), insufficient (–), or indeterminate^[17]. The quality criteria of the measured properties were then analyzed using Terwee’s study criteria^[18,19]. In updated criteria for good measurement properties, the assessment of cross-cultural validity involves determining whether a scale remains consistent in its measurement across different cultures or if there are variations in how specific items are responded to. This is evaluated through measurement invariance (MI) and differential item functioning (DIF) analyses. MI and non-DIF investigate whether individuals from diverse groups while accounting for group differences, respond similarly to a given item at the same underlying level of the trait being measured. Regression analyses or confirmatory factor analysis (CFA) are effective methods for adequately evaluating cross-cultural validity using Classical Test Theory (CTT). In the present study, decisions were made to remove the criteria for determining the suitability of a gold standard. Instead, the review team first identifies which outcome measurement instrument can be considered a reasonable gold standard before assessing the methodological quality of the studies. If a study included in the review utilizes this designated gold standard instrument to evaluate validity, it can be categorized as a study on criterion validity. The COSMIN panel has reached a consensus that there are no established gold standards for PROMs, with the




Figure 1. PRISMA flow chart.

exception being when a shortened version of an instrument is compared to its original long version. In such cases, the original long version can be considered the gold standard^[19]. Any disagreements were settled by discussion and arbitration. The Cohen’s Kappa value was used to evaluate the inter-reviewer reliability. Any disagreements were settled by discussion and arbitration.

Data collection and construct validity assessment

For each study, the following details were considered: authors, publication year, country, average age of participants (or range), type of sample, sample size, sampling method, number of factors, total variance explained and variance of each factor, and reliability outcomes. Principal Component Analysis (PCA) and Exploratory Factor Analysis (EFA), were two methods used to extract factors, have differences, but many researchers have used them interchangeably, and PCA is frequently used when EFA would be more appropriate^[20]. As a result, PCA will have a higher percentage of the extracted variance than EFA because it concentrates on the total variation among the variables rather than the shared variance of the variables^[21]. Around 60% of the total extracted variance is typically explained by factor analysis models in the social and behavioural sciences^[22].

Results

Study characteristics

Considering the inclusion criteria, 80 documents were evaluated^[7,8,23–99]. All of the documents were original articles.

Out of these three articles, 3.75% used Item Response Theory methods in addition to the Classic Test Method for validating the tool. Different versions of the CD-RISC were evaluated in the studies. In 10 articles, 12.5% evaluated more than one version of the scale. The 25-item version of the CD-RISC was psychometrically assessed in 44 studies, the 10-item version in 40 studies, and the 2-item version in 7 studies. For the 25-item version, a minimum of one factor and a maximum of five factors were extracted^[42]. For the 10-item version, one and eventually two^[26,62,74,83,100] factors were identified.

The majority of studies took place in the United States (15 studies, 18.75%), followed by China (12 studies, 15.00%). The publication period of the articles included in the current study ranged from 2003 to 2023, with the highest number of articles (13 articles, 16.25%) published in 2016. A diverse spectrum of individuals was examined as the research population, with the majority being the general population (13 studies), students (12 studies), adolescents and youth (12 studies). Patients, adults with autism and psychiatric disorders, employees, firefighters, family caregivers, nurses and healthcare personnel, earthquake survivors, veterans, and older adults were other populations who were assessed by researchers. Further details regarding the characteristics of the studies are provided in Appendix 2, Supplemental Digital Content 3, <http://links.lww.com/MS9/A413>.

Risk of Bias

After selecting the documents based on inclusion criteria, the first step in evaluating the articles was to assess the potential risk of bias using the COSMIN risk of bias checklist^[16]. The results of

the risk of bias assessment for different versions of the CD-RISC are provided below:

CD-RISC-25

The method used in the PROM Development dimension of the 25-item version in 26 studies (59.1%) received a very good score in terms of parameters related to bias risk. Out of 44 studies, 39 studies (88.6%) did not evaluate the content validity of the 25-item version of the scale. Overall, more than two-thirds of the studies received a very good (26 studies, 59.1%) or adequate (9 studies, 20.5%) score in terms of structural validity, which is notable. Based on the COSMIN criterion, three studies were classified as inadequate or doubtful for assessing structural validity. Six studies (13.6%) did not evaluate the structural validity. In terms of internal consistency, 20 studies (45.5%) received a very good score in terms of the risk of bias. Out of the 44 reviewed studies, 32 studies (72.7%) did not evaluate cross-cultural validity or measurement invariance overall, and only one study (2.3%) received a very good score on these criteria. The majority of studies (28 studies, 63.8%) did not assess the reliability of the CD-RISC-25-item version, and only 3 studies (6.8%) received a very good score in terms of the risk of bias rating in the reliability dimension. Out of 44 studies, in 43 studies (97.7%), the measurement error was not assessed, and one study that evaluated this property received an inadequate score based on COSMIN's criteria. According to the definition used in this study for criterion validity, which was based on COSMIN's definition of criterion validity, none of the 44 reviewed studies evaluated the criterion validity of the 25-item version, so there was no possibility of assessing bias risk in this domain. In the hypothesis testing for the construct validity domain, the method and results of the reviewed articles were in an acceptable state, and a total of 29 articles (65.9%) received a very good or adequate score in this aspect. Only one study evaluated the responsiveness of the 25-item version of the CD-RISC, which received a doubtful score according to the COSMIN risk of bias checklist (Table 1).

CD-RISC-10

In the first domain examined, 18 studies (45%) received a very good score for the 10-item version. Similar to the 25-item version, the majority of studies (34 studies, 80%) did not evaluate the content validity of the 10-item version; as a result, there was no possibility to assess the risk of bias in the results of studies in this domain.

In terms of structural validity, the 10-item version was deemed acceptable, with 27 studies (67%) receiving a very good score in this domain. Similarly, in the domain of internal consistency, the studies' status for the 10-item version was acceptable, with 32 studies (80%) receiving a very good score from COSMIN's risk of bias checklist. Out of 40 studies, 21 studies examined the cross-cultural validity or measurement invariance of the 10-item version, and 8 studies received an insufficient score in this domain. Additionally, only 18 studies out of 40 evaluated the reliability of the 10-item version, and out of these, 9 studies received an inadequate score on the risk of bias checklist. Moreover, only five studies out of 40 addressed the measurement error evaluation of the 10-item version, and out of these five studies, three received a doubtful or inadequate score on the checklist. Only three studies out of 40 evaluated the criterion validity of the 10-item version according to COSMIN's definition, and all three studies received

an inadequate score. In the Hypothesis Testing for Construct Validity domain, 23 studies out of 40 evaluated this property, and similar to the 25-item version, the 10-item version's status was acceptable, with 22 studies receiving a very good score. Out of 40 studies, only one study evaluated the responsiveness of the 10-item version and allocated an adequate score to itself on the checklist (Table 2).

CD-RISC-2

Four out of seven studies did not evaluate the parameters related to the PROM Development dimension in the risk of bias Assessment Checklist for the 2-item version. Among the three studies that evaluated these parameters, two studies obtained a very good score, and one study obtained a doubtful score. None of the studies examined the content and structural validity of the 2-item version, so it was not possible to evaluate this aspect using the checklist. Out of the seven studies, five studies examined the internal consistency of the scale, with four studies receiving a very good score and one study receiving an inadequate score. Only one study examined the cross-cultural validity or measurement invariance of the 2-item version, which had a doubtful score. Three studies (49.2%) received an inadequate score in the assessment of reliability, and one study (14.3%) received an adequate score, while the remaining studies (3 studies, 42.9%) did not assess reliability. Only one study examined the measurement error of the 2-item version and received an adequate score, while the other six studies did not measure this parameter, so it was not possible to evaluate them using the checklist. Out of the seven studies, only two studies (28.6%) assigned a very good score to the criterion validity assessment based on COSMIN's definition. Similar to the other two versions, the 2-item version had good status in terms of hypothesis testing for construct validity, with five studies (74.4%) receiving a very good score and two studies (28.6%) receiving an adequate score. Only one study (14.3%) examined the responsiveness of the 2-item version of CD-RISC and received an inadequate score from the risk of bias Checklist (Table 3).

Quality assessment

After evaluating the risk of bias, the quality of the studies was assessed based on criteria for good measurement properties (quality criteria) introduced by COSMIN^{10,11}. The results, categorized by psychometric parameters and different versions of the scale, are as follows. Additional information is provided in Tables 4 to 6.

Structural validity

Only 22 studies (50.00%) out of the 44 reviewed articles provided "sufficient" evidence for the structural validity of the 25-item version. In terms of the quality assessment of the evidence provided regarding the psychometric properties of the 10-item version, 24 studies (60%) out of 40 studies reported sufficient evidence for structural validity. Many studies only conducted Exploratory Factor Analysis (EFA) and, therefore, failed to provide high-quality evidence for structural validity according to the COSMIN criteria. For the 2-item version, no study succeeded in producing high-quality evidence for structural validity. However, since conducting CFA is necessary to obtain a sufficient score according to the COSMIN criteria, it seems that conducting CFA

Table 1
The COSMIN risk of bias checklist (CD-RISC-25-items).

	BOX 1	BOX 2	BOX 3	BOX 4	BOX 5	BOX 6	BOX 7	BOX 8	BOX 9	BOX 10
Author (year)	PROM development	Content validity	Structural validity	Internal consistency	Cross-cultural validity \ measurement invariance	Reliability (stability with test re-test)	Measurement error	Criterion validity	Hypotheses testing for construct validity	Responsiveness
Abdi <i>et al.</i> (2019) ^[23]	Very good	—	Very good	Very good	—	—	—	—	—	—
Tsigkaropoulou <i>et al.</i> (2018) ^[87]	Very good	—	Adequate	Very good	—	Very good	—	—	Very good	—
Bakhshayesh Eghbali <i>et al.</i> (2022) ^[29]	Very good	Adequate	Very good	Very good	—	—	—	—	—	—
Bezjian <i>et al.</i> (2017) ^[30]	Very good	—	—	Inadequate	—	—	—	—	Very good	—
Campbell-Sills and Stein (2007) ^[33]	Very good	—	Very good	Inadequate	—	Inadequate	—	—	Very good	—
Connor and Davidson (2003) ^[4]	Very good	—	Adequate	Doubtful	—	Adequate	—	—	Very good	Doubtful
Solano <i>et al.</i> (2016) ^[84]	Very good	Doubtful	Adequate	Very good	—	Very good	—	—	Very good	—
Derakhshanrad <i>et al.</i> (2014) ^[37]	Very good	—	—	Doubtful	—	—	—	—	—	—
Kuiper <i>et al.</i> (2019) ^[59]	Very good	—	—	Very good	—	Inadequate	—	—	Very good	—
Dominguez-Cancino <i>et al.</i> (2022) ^[38]	Very good	Adequate	Very good	Doubtful	—	—	—	—	—	—
Anjos <i>et al.</i> (2019) ^[27]	Very good	—	Very good	Doubtful	—	—	—	—	Very good	—
Fujikawa <i>et al.</i> (2013) ^[40]	Very good	—	Very good	Doubtful	Doubtful	Inadequate	—	—	Very good	—
Gao <i>et al.</i> (2021) ^[41]	Very good	—	Very good	Doubtful	Doubtful	Inadequate	—	—	—	—
García León <i>et al.</i> (2019) ^[42]	Very good	—	Adequate	Doubtful	Doubtful	—	—	—	Very good	—
Goins <i>et al.</i> (2013) ^[43]	Very good	—	Doubtful	Very good	Inadequate	—	—	—	Very good	—
Gonzalez <i>et al.</i> (2016) ^[44]	Very good	—	Very good	Very good	Doubtful	—	Inadequate	—	Very good	—
Gouda <i>et al.</i> (2022) ^[46]	Very good	—	Inadequate	Inadequate	—	Inadequate	—	—	—	—
Guihard <i>et al.</i> (2018) ^[50]	Very good	Inadequate	Very good	Inadequate	—	Very good	—	—	—	—
Guamizo Guzmán <i>et al.</i> (2019) ^[49]	Very good	—	Very good	Very good	Very good	—	—	—	—	—
Baek <i>et al.</i> (2010) ^[28]	Very good	—	Adequate	Doubtful	—	Doubtful	—	—	Very good	—
Jorgensen and Seedat (2008) ^[53]	Very good	—	Very good	Inadequate	Doubtful	—	—	—	—	—
Jung <i>et al.</i> (2012) ^[54]	Very good	—	Very good	Inadequate	—	Inadequate	—	—	Very good	—
Karairmak (2010) ^[55]	Very good	—	Very good	Very good	—	Inadequate	—	—	Very good	—
Khoshouei (2009) ^[57]	Very good	—	Adequate	Doubtful	—	Doubtful	—	—	—	—
Kidd <i>et al.</i> (2019) ^[58]	Very good	—	Adequate	Very good	—	—	—	—	Very good	—
Wu <i>et al.</i> (2017) ^[93]	—	—	Very good	—	—	—	—	—	Very good	—
Madewell <i>et al.</i> (2016) ^[64]	—	—	Very good	Very good	—	—	—	—	Adequate	—
Manzano Garcia and Ayala Calvo (2013) ^[65]	—	—	Very good	Inadequate	—	—	—	—	—	—
Martinez <i>et al.</i> (2021) ^[66]	—	—	Very good	Very good	Inadequate	—	—	—	Adequate	—
Bizri <i>et al.</i> (2022) ^[31]	—	—	—	Inadequate	—	—	—	—	Adequate	—
McGillivray and Ho (2016) ^[67]	—	—	Adequate	Very good	—	—	—	—	Adequate	—

Table 2
The COSMIN risk of bias checklist (CD-RISC-10-items).

	BOX 1	BOX 2	BOX 3	BOX 4	BOX 5	BOX 6	BOX 7	BOX 8	BOX 9	BOX 10
Author (year)	PROM development	Content validity	Structural validity	Internal consistency	Cross-cultural validity Measurement invariance	Reliability	Measurement error	Criterion validity	Hypotheses testing for construct validity	Responsiveness
Alarcón <i>et al.</i> (2020) ^[24]	Very good	—	Very good	Very good	—	—	—	—	Very good	—
Aloba <i>et al.</i> (2016A) ^[26]	Very good	—	Very good	Very good	—	—	—	—	Very good	—
Aloba <i>et al.</i> (2016B) ^[25]	Very good	—	Adequate	Very good	—	Inadequate	—	—	Very good	—
Blanco <i>et al.</i> (2019) ^[32]	Very good	—	Very good	Very good	—	Inadequate	—	—	Very good	—
Cheng <i>et al.</i> (2020) ^[34]	Very good	—	Very good	Very good	Very good	Inadequate	—	Inadequate	Very good	—
Coates <i>et al.</i> (2013) ^[35]	Very good	—	Very good	Very good	—	—	—	—	Very good	—
Daniel-González <i>et al.</i> (2020) ^[36]	Very good	Inadequate	Very good	Inadequate	Very good	Inadequate	—	—	Very good	—
Kuiper <i>et al.</i> (2019) ^[59]	Very good	—	—	Very good	—	Inadequate	—	—	Very good	—
Duong and Hurst (2016) ^[39]	Very good	Inadequate	Very good	Doubtful	Doubtful	Inadequate	—	—	Very good	—
Goins <i>et al.</i> (2013) ^[43]	Very good	—	Doubtful	Very good	Inadequate	—	—	—	Very good	—
Gonzalez <i>et al.</i> (2016) ^[44]	Very good	—	Very good	Very good	Doubtful	—	Inadequate	—	Very good	—
Gorman <i>et al.</i> (2021) ^[45]	Very good	—	Very good	—	—	Inadequate	—	—	—	—
Gras <i>et al.</i> (2019) ^[47]	Very good	—	Very good	—	Doubtful	—	—	—	—	—
Guarnizo Guzmán <i>et al.</i> (2019) ^[49]	Very good	—	Very good	Very good	Very good	—	—	—	—	—
Kwan <i>et al.</i> (2019) ^[60]	Very good	Adequate	Very good	Doubtful	—	Very good	Very good	—	Very good	—
Hwang <i>et al.</i> (2020) ^[51]	Very good	—	Very good	Inadequate	Inadequate	Inadequate	—	—	Very good	—
Keyhani <i>et al.</i> (2015) ^[56]	Very good	—	Doubtful	Inadequate	—	—	—	—	Very good	—
Lauridsen <i>et al.</i> (2017) ^[62]	—	Inadequate	Doubtful	Very good	—	—	—	—	Adequate	Adequate
Levey <i>et al.</i> (2021) ^[63]	—	—	Adequate	Very good	—	—	—	—	Very good	—
Madewell <i>et al.</i> (2016) ^[64]	—	—	Very good	Very good	—	—	—	Inadequate	Adequate	—
Meng <i>et al.</i> (2019) ^[69]	—	—	Very good	Very good	Very good	Doubtful	—	—	—	—
Miller <i>et al.</i> (2021) ^[70]	—	—	—	Very good	—	—	—	Inadequate	Very good	—
Minh-Uyen and Im (2021) ^[71]	—	—	Very good	Very good	—	—	—	—	—	—
Nartova-Bochaver <i>et al.</i> (2021) ^[73]	Doubtful	—	Very good	Very good	Doubtful	—	—	—	Adequate	—
Nguyen and Dinh (2022) ^[74]	Doubtful	—	Very good	Very good	Inadequate	—	—	—	Very good	—
Notario-Pacheco <i>et al.</i> (2011) ^[76]	—	—	Doubtful	Very good	Inadequate	Doubtful	—	—	Doubtful	-
Pretorius and Padmanabhanunni (2022) ^[77]	—	—	Doubtful	Very good	Inadequate	—	Inadequate	—	Adequate	—
Rezaeipandari <i>et al.</i> (2022) ^[7]	Doubtful	Doubtful	Very good	Doubtful	Inadequate	Doubtful	—	—	—	—
She <i>et al.</i> (2020) ^[80]	—	—	Very good	Very good	very good	—	—	—	Very good	—
Sharma <i>et al.</i> (2018) ^[79]	Doubtful	—	Very good	Very good	—	Adequate	Adequate	—	Very good	—
Shin <i>et al.</i> (2018) ^[81]	—	—	Doubtful	Very good	—	—	—	—	Adequate	—
Smith <i>et al.</i> (2019) ^[83]	—	—	Very good	Very good	Adequate	—	—	—	Adequate	—
Tourunen <i>et al.</i> (2021) ^[86]	—	—	Very good	Very good	Inadequate	Doubtful	—	—	Adequate	—
Vongsirimas <i>et al.</i> (2017) ^[90]	—	—	Doubtful	Very good	—	—	—	—	—	—
Waddimba <i>et al.</i> (2022) ^[91]	—	—	Very good	Very good	Very good	Inadequate	—	—	Adequate	—
Wang <i>et al.</i> (2010) ^[92]	—	—	Doubtful	Very good	—	Doubtful	—	—	9a: Very good 9b: Adequate	—
Ye <i>et al.</i> (2017) ^[95]	—	Very good	Very good	Very good	—	Doubtful	—	—	Adequate	—
Zhang <i>et al.</i> (2021) ^[98]	—	—	Very good	Very good	Doubtful	—	—	—	Very good	—

CD-RISC, Connor-Davidson Resilience Scale.

Table 3
The COSMIN risk of bias checklist (CD-RISC-2-items).

Author (year)	PROM development	BOX 2 Content validity	BOX 3 Structural validity	BOX 4 Internal consistency	BOX 5 Cross-cultural validity \ measurement invariance	BOX 6 Reliability	BOX 7 Measurement error	BOX 8 Criterion validity	BOX 9 Hypotheses testing for construct validity	BOX 10 Responsiveness
Kuiper et al. (2019) ^[69]	Very good	—	—	Very good	—	Inadequate	—	—	Very good	—
Jeong et al. (2015) ^[62]	Very good	—	—	Inadequate	Doubtful	—	—	—	Very good	—
Ni et al. (2016) ^[75]	—	—	—	Very good	—	—	—	Very good	Very good	—
Sharma et al. (2018) ^[79]	Doubtful	—	—	—	—	Adequate	Adequate	Very good	Very good	—
Vaishnavi et al. (2007) ^[88]	—	—	—	—	—	Inadequate	—	—	9a: Adequate 9b: Doubtful	Inadequate
Waddimba et al. (2022) ^[91]	—	—	—	Very good	—	Inadequate	—	Inadequate	Adequate	—
Zhang et al. (2021) ^[90]	—	—	—	Very good	—	—	—	Inadequate	Very good	—

CD-RISC, Connor-Davidson Resilience Scale.

measurement invariance, and one study provided “insufficient” evidence in this domain.

Criterion validity

For the 25-item version of the scale, all studies were in an “indeterminate” state in terms of quality for criterion validity assessment. Similarly, for the 10-item version of the scale, the majority of studies (25 studies, 62.5%) were in an “indeterminate” state. For the 2-item version, three studies (42.9%) successfully provided “sufficient” evidence, three studies (42.9%) were in a state of “indeterminate” evidence, and the evidence produced by one study was “insufficient”.

Responsiveness

In the responsiveness domain, only one study (2.3%) for the 25-item version was able to provide “sufficient” results according to the criteria of quality assessment. For the 10-item version, no study successfully produced evidence of “sufficient” quality for responsiveness, and all 40 studies were in an “indeterminate” state. Similarly, in the 2-item version, the majority of studies (six studies, 85.7%) did not produce evidence in the domain of responsiveness, so they were in an “indeterminate” state.

Discussion

The CD-RISC stands as a robust and widely utilized tool for assessing resilience across various populations and contexts. Through its carefully crafted items, the scale captures individual differences in the ability to bounce back from adversity, adapt to challenging circumstances, and maintain psychological well-being. Its psychometric properties have been extensively studied and validated, demonstrating high levels of reliability and validity across different demographic groups, including diverse cultural backgrounds and clinical populations. Moreover, the CD-RISC has proven to be a valuable instrument in both research and clinical practice, providing insights into individuals’ resilience levels and predicting outcomes related to mental health, functioning, and overall adjustment. Its versatility and effectiveness make it a valuable asset for researchers, clinicians, and practitioners alike, offering a standardized means of assessing resilience that can inform interventions and support strategies tailored to individuals’ needs.

Among the various measures developed to assess psychological resilience, the CD-RISC has been used in a large-scale study population and settings and has exhibited different but mostly good psychometric evaluation results. The current systematic review sought to investigate the psychometric properties of three versions of CD-RISC by summarizing and analyzing the existing literature in this field. The current study revealed that the maximum number of published papers in the field of psychological resilience originated in the United States and China in 2016. According to Scimago Journal & Country Rank (SJR), the United States published more than 707 000 scientific documents about Resilience, and China, with more than 502 000 published documents were the top-rank countries in the world in 2016. Furthermore, the United States was the first-rank country with the highest scientific publication in applied psychology ($n = 3717$ documents) and clinical psychology ($n = 5910$) in 2016^[102]. Although these statistics can support the current study

Table 4
Quality assessment (CD-RISC-25-items).

Author (year)	Criteria for good measurement properties							
	Structural validity	Internal consistency	Reliability	Measurement error	Hypotheses testing for construct validity	Cross-cultural validity\measurement invariance	Criterion validity	Responsiveness
Abdi <i>et al.</i> (2019) ^[23]	+	+	?	?	?	?	?	?
Tsigkaropoulou <i>et al.</i> (2018) ^[87]	?	+	+	?	+	?	?	?
Bakhshayesh Eghbali <i>et al.</i> (2022) ^[29]	+	+	?	?	?	?	?	?
Bezdzian <i>et al.</i> (2017) ^[30]	?	?	?	?	+	?	?	?
Campbell-Sills and Stein (2007) ^[33]	+	?	?	?	+	?	?	?
Connor and Davidson (2003) ^[4]	?	+	+	?	+	?	?	+
Solano <i>et al.</i> (2016) ^[84]	?	+	+	?	+	?	?	?
Derakhshanrad <i>et al.</i> (2014) ^[37]	?	+	?	?	?	?	?	?
Kuiper <i>et al.</i> (2019) ^[59]	?	+	?	?	+	?	?	?
Dominguez-Cancino <i>et al.</i> (2022) ^[38]	+	-	?	?	?	?	?	?
Anjos <i>et al.</i> (2019) ^[27]	+	-	?	?	+	?	?	?
Fujikawa <i>et al.</i> (2013) ^[40]	+	+	?	?	+	+	?	?
Gao <i>et al.</i> (2021) ^[41]	+	+	?	?	?	+	?	?
García León <i>et al.</i> (2019) ^[41]	-	+	?	?	+	+	?	?
Goins <i>et al.</i> (2013) ^[43]	+	+	?	?	+	?	?	?
Gouda <i>et al.</i> (2022) ^[46]	?	-	?	?	?	?	?	?
Green <i>et al.</i> (2014) ^[48]	+	+	?	?	+	?	?	?
Guihard <i>et al.</i> (2018) ^[50]	+	-	+	?	?	?	?	?
Guarnizo Guzmán <i>et al.</i> (2019) ^[49]	+	+	?	?	?	+	?	?
Baek <i>et al.</i> (2010) ^[28]	+	-	?	?	+	?	?	?
Jorgensen and Seedat (2008) ^[53]	+	-	?	?	?	?	?	?
Jung <i>et al.</i> (2012) ^[54]	+	-	?	?	+	?	?	?
Karairmak (2010) ^[55]	+	+	?	?	+	?	?	?
Khoshouei (2009) ^[57]	?	+	?	?	?	?	?	?
Kidd <i>et al.</i> (2019) ^[58]	?	+	?	?	+	?	?	?
Wu <i>et al.</i> (2017) ^[93]	+	-	?	?	+	?	?	?
Madewell <i>et al.</i> (2016) ^[64]	-	+	?	?	+	?	?	?
Manzano García and Ayala Calvo (2013) ^[65]	?	-	?	?	?	?	?	?
Martinez <i>et al.</i> (2021) ^[66]	+	+	?	?	+	+	?	?
Bizri <i>et al.</i> (2022) ^[31]	?	?	?	?	+	?	?	?
Mealer <i>et al.</i> (2016) ^[68]	-	+	?	?	+	?	?	?
Ni <i>et al.</i> (2016) ^[75]	?	?	?	?	+	?	?	?
Miller <i>et al.</i> (2021) ^[70]	?	?	?	?	+	?	?	?
Nagle <i>et al.</i> (2021) ^[72]	+	?	?	?	+	?	?	?
Sexton <i>et al.</i> (2016) ^[78]	?	?	?	?	+	?	?	?
Sharif-Nia <i>et al.</i> , (2021) ^[99]	+	+	?	?	?	?	?	?
Sharif Nia <i>et al.</i> (2023) ^[8]	+	-	?	?	?	+	?	?
Singh and Yu (2010) ^[82]	-	-	?	?	+	?	?	?
Velickovic <i>et al.</i> (2020) ^[89]	-	?	?	?	?	-	?	?
Yu and Zhang (2007) ^[96]	-	-	?	?	+	?	?	?
Yu <i>et al.</i> (2011) ^[97]	+	-	?	?	+	?	?	?
Xie <i>et al.</i> (2016) ^[96]	-	+	+	?	+	?	?	?

CD-RISC, Connor-Davidson Resilience Scale.

Table 5
Quality assessment (CD-RISC-10-items).

Author (year)	Criteria for good measurement properties							
	Structural validity	Internal consistency	Reliability	Measurement error	Hypotheses testing for construct validity	Cross-cultural validity\measurement invariance	Criterion validity	Responsiveness
Alarcón <i>et al.</i> (2020) ^[24]	+	+	?	?	+	?	?	?
Aloba <i>et al.</i> (2016A) ^[26]	+	+	?	?	+	?	?	?
Aloba <i>et al.</i> (2016B) ^[25]	?	+	?	?	+	?	?	?
Blanco <i>et al.</i> (2019) ^[32]	+	+	?	?	+	?	?	?
Cheng <i>et al.</i> (2020) ^[34]	+	+	?	?	+	+	?	?
Coates <i>et al.</i> (2013) ^[35]	+	+	?	?	+	?	?	?
Daniel-González <i>et al.</i> (2020) ^[36]	+	+	?	?	+	+	?	?
Kuiper <i>et al.</i> (2019) ^[59]	?	+	?	?	+	?	?	?
Duong and Hurst (2016) ^[39]	+	+	?	?	+	+	?	?
Goins <i>et al.</i> (2013) ^[43]	+	+	?	?	+	?	?	?
Gonzalez <i>et al.</i> (2016) ^[44]	+	+	?	?	+	+	?	?
Gorman <i>et al.</i> (2021) ^[45]	+	?	?	?	?	?	?	?
Gras <i>et al.</i> (2019) ^[47]	+	?	?	?	?	+	?	?
Guarnizo Guzmán <i>et al.</i> (2019) ^[49]	+	+	?	?	?	+	?	?
Kwan <i>et al.</i> (2019) ^[60]	+	+	+	+	+	?	?	?
Keyhani <i>et al.</i> (2015) ^[56]	-	-	?	?	+	?	?	?
Kyriazos and Stalikas (2021)	+	+	?	?	+	+	?	?
Lauridsen <i>et al.</i> (2017) ^[62]	?	+	?	?	+	?	?	?
Levey <i>et al.</i> (2021) ^[63]	?	+	?	?	+	?	?	?
Madewell <i>et al.</i> (2016) ^[64]	-	+	?	?	+	?	+	?
Meng <i>et al.</i> (2019) ^[69]	?	+	?	?	?	+	?	?
Miller <i>et al.</i> (2021) ^[70]	?	+	?	?	+	?	+	?
Minh-Uyen and Im (2021) ^[71]	-	+	?	?	?	?	?	?
Nartova-Bochaver <i>et al.</i> (2021) ^[73]	+	+	?	?	+	?	?	?
Nguyen and Dinh (2022) ^[74]	+	+	?	?	+	?	?	?
Notario-Pacheco <i>et al.</i> (2011) ^[76]	?	+	+	?	+	?	?	?
Pretorius and Padmanabhanunni (2022) ^[77]	+	+	?	?	+	+	?	?
Rezaeipandari <i>et al.</i> (2022) ^[7]	+	+	+	?	?	?	?	?
Sharma <i>et al.</i> (2018) ^[79]	+	+	+	?	+	?	?	?
Shin <i>et al.</i> (2018) ^[81]	?	+	?	?	+	?	?	?
Smith <i>et al.</i> (2019) ^[83]	+	-	?	?	+	+	?	?
Tomyn and Weinberg (2018)	?	+	?	?	+	-	?	?
Tourunen <i>et al.</i> (2021) ^[86]	?	+	-	?	+	-	?	?
Vongsirimas <i>et al.</i> (2017) ^[90]	-	+	?	?	?	?	?	?
Waddimba <i>et al.</i> (2022) ^[92]	+	+	?	?	One hypothesis (fear of COVID-19): -Others: + hypothesis: + Known group: +	+	?	?
Wang <i>et al.</i> (2010) ^[92]	?	+	?	?	One hypothesis (MCMQ): -Others: +	?	?	?
Ye <i>et al.</i> (2017) ^[95]	+	+	?	?	One hypothesis (MCMQ): -Others: +	?	?	?
Zhang <i>et al.</i> (2021) ^[99]	+	+	?	?	+	+	?	?

CD-RISC, Connor-Davidson Resilience Scale.

Table 6
Quality assessment (CD-RISC-2-items)

Author (year)	Criteria for good measurement properties									
	Structural validity	Internal consistency	Reliability	Measurement error	Hypotheses testing for construct validity	Cross-cultural validity/invariance	Criterion validity	Responsiveness		
Kuiper et al. (2019) ^[50]	?	+	?	?	+	?	?	?		?
Jeong et al. (2015) ^[52]	?	-	?	?	+	-	?	?		?
Ni et al. (2016) ^[75]	?	+	?	?	+	?	+	?		?
Sharma et al. (2018) ^[79]	?	?	+	?	+	?	+	?		?
Vaishnavi et al. (2007) ^[89]	?	?	?	?	One hypothesis (ASEX): -Others: + Known group: +	?	?	?		+
Waddimba et al. (2022) ^[91]	?	+	?	?	One hypothesis (fear of COVID-19): -Others: +	?	-	?		?
Zhang et al. (2021) ^[98]	?	+	?	?		?	+	?		?

CD-RISC, Connor-Davidson Resilience Scale.

findings, further investigations are needed to determine why psychological resilience grasped the researchers' or journals' attention during 2016. Studies show that many countries were affected by the economic crisis before and during 2015 which significantly affected population health indicators^[103,104]. This may be one reason why resilience drew the attention of the scientific community. The findings of a systematic review study revealed that a higher prevalence of mental health problems was the outcome of periods of economic crisis. In this regard, unemployed people or those who had a precarious work situation, as well as people who faced debts and economic strain and those who had a pre-existing mental illness, were identified as vulnerable victims of economic hardship^[104]. In such a situation, it becomes important to improve people's resilience to bounce back from the adversity they face. The CD-RISC-2 is an abbreviated version of the scale that has good psychometric properties but may not capture the full range of resilience-related constructs^[88]. Here are some differences between the CD-RISC and the Brief Resilience Scale (BRS); for example, CD-RISC focuses on resources that can help individuals recover from and adapt to disruptions or stressful events, but BRS Directly measures one's ability to bounce back or be resilient. The choice of which version of the scale to use may depend on the specific research question and the available resources.

The current study findings indicated that the CD-RISC versions were assessed psychometrically among a diversity of populations, including those people who were experiencing traumatic conditions, such as patients with autism^[51], earthquake victims^[92], persons with psychological disorders^[34], as well as the general population^[87]. Considering the existing definitions of psychological resilience, it is clearly stated that resilience is people's capacity derived from individual and social assets and resources for 'bouncing back' and adapting to adversities across the lifespan^[105]. This indicated that both healthy and ill people, whether children or elderly, need resilience to improve their resistance to the challenges of life^[106]. On the other hand, the development and vast testing of the CD-RISC was conducted in response to the lack of generalizability of the other existing resilience scales^[4]. The current study findings indicated that both 25 and 10-item versions of CD-RISC were tested psychometrically almost the same; while the 25-item CD-RISC version was assessed in 44 papers, the other versions were used in 40 studies. Due to the comprehensiveness of the 25-item CD-RISC, including more aspects of resilience and considering that the CD-RISC-10 is economically and psychometrically sounder, both versions of this scale have been used in the studies in different settings^[107].

According to the COSMIN's criteria for evaluating the risk of bias, the current research findings showed limitations of included studies in assessing the three versions of CD-RISC cross-cultural and content validity as well as their stability (e.g. conducting test re-test), whereas the majority of psychometric studies of CD-RISC-25, and CD-RISC-2 rated as very good or adequate in terms of structural validity. Our findings also indicated that the included papers were qualified in term of assessing internal consistency but were weak in assessing measurement error. Assessing the methodological quality of studies included in a systematic review is crucial because ignoring it can negatively affect the trustworthiness of the results^[16,101]. As an accepted strategy, the COSMIN checklist is increasingly grasping the researchers' attention to evaluating the systematic reviews'

quality. Being aware of this consideration when assessing the studies' eligibility, the COSMIN provides evidence that determines the quality of a PROM. In terms of our findings that indicated the psychometric studies' limitations in conducting or reporting the cross-cultural validity, an insufficient explanation of the scale translation/back translation process and the pre-testing of the scale among the target population may result in low-quality evidence. The failure of the cross-cultural validity of scales may lead to misleading findings and inaccuracies that are a reflection of the constructs in each study setting^[108]. Regarding the lack of assessment of measurement error, it may lead to failure in estimating the influences of different sources of scores on variation. In other words, assessing the measurement error in each psychometric study can guide the researchers to decide about the level of trust they can place in the study findings and whether the parameters that were estimated are systematically not over- or underestimated^[101]. As a result, considering the COSMIN criteria while designing and conducting a psychometric study can be suggested to the researcher in order to improve the study's quality.

In terms of quality assessment of the included studies, the current study indicated that investigating the structural validity of the CD-RISC by researchers was mainly done based on EFA, and a lack of evaluating confirmatory factor analysis was evident. It should be mentioned that measurement is the process by which a concept relates to one or more latent variables, and the latent variables link to the observed variables, which are usually items in a questionnaire^[109]. During the development and psychometrically testing of a questionnaire to explore the underlying factors or dimensions (subscales) that explain the relationships between the items, that is called construct validity, to simplify a set of complex items (variables) using statistical techniques. Factor analysis (FA) is used as the most common method for construct validity^[110]. In fact, factor analysis is used to assess the construct validity as the heart of the design and development of a scale, questionnaire, instrument, and or tool^[111]. Construct validity by EFA provides evidence based on test content and internal structure that it tells us what the questionnaire, scale, or instrument really measures as an abstract concept^[112]. EFA is a data reduction statistical technique that reflects the theoretical structure of phenomena and summarizes its underlying variables^[113,114]. The emphasis of EFA is on the relationships among the items that use "shared covariance" to identify factors^[115,116]. It is important to mention that the extracted factors in EFA should be interpreted as explanatory and not causality. Following EFA, confirmatory factor analysis (CFA) usually applies to verifying the emerged factor structure of a set of observed variables and helps to test the hypothesis regarding how the variables are related to each other^[117]. In addition to assessing the model fit and the residual error variances of the observed variables, CFA estimates the factor loadings, variances, and covariances of the factors^[118]. Due to the requirements of conducting CFA (e.g. existing the number of factors), the COSMIN checklist guides the researchers to consider whether CFA is more appropriate^[18]. Accordingly, conducting CFA is not feasible for CD-RISC-2.

From the point of view of the quality of the available evidence, in terms of hypotheses testing for construct validity, the CD-RISC-2 was rated sufficient, and more than half of the included studies were rated as sufficient for the 25-item and 2-item versions. Hypothesis testing roughly determines whether the

relationship's direction and magnitude or difference is comparable to what can be predicted based on the structure(s) being measured. The more hypotheses that are being tried on whether the information compares to previously formulated hypotheses, the more proof is assembled to build construct validity^[120]. It has been recommended to provide clear data regarding the missing items, the sample size, the previously estimated mean differences, and correlations, as well as describe convergent validity to improve the study's quality in this aspect^[119,120].

In almost all of the studies (except three studies), the responsiveness of the CD-RISC versions was not investigated. Detecting the construct changes over time is the aim of responsiveness, and it is usually needed to use another instrument for the study samples in the same situation and time. Due to its difficulty in terms of methodology as well as the restricted introduction of responsiveness based on COSMIN, many researchers, especially clinicians, don't assess it, and reporting it remains a challenge in many studies^[121].

In the case of CD-RISC factor structure, the included studies assessed the 25-item version indicated that the factor structures have differed from the original five-factor version^[4]. It was different from 1-factor structure^[42], two-factor version^[48], three-factor version^[67], and 4-factor structure^[78]. Similarly, the CD-RISC-10-item exhibited this difference from the original version. Considering the complex nature of resilience, the mentioned variety may suggest that the process or the way this construct was understood by the study population of different ages, genders, cultures, and religious groups was somehow different. The samples of the studies were in different situations, faced various challenges, and had diverse individual capacities or environmental/social support systems^[122].

A range of different factors can influence the psychometric properties of the CD-RISC versions in the studies. The lack of a universally agreed-upon definition of resilience can make it difficult to develop relevant measurement tools and evaluate the impact of interventions and policy changes^[123]. In addition, the cultural and semantic differences between study populations can affect the validity and reliability of resilience scales^[124]. Furthermore, the psychometric properties of resilience scales may correlate with the study population's age^[125,126], and their socioeconomic status^[123].

Limitations and Strengths

The main limitation is that we included only studies that were either in Farsi or English and did not search databases such as PsychInfo, as well as searching only three international databases. The evaluation of the quality of the studies using three tools is one of the strengths of this study.

Conclusion

In this systematic review, 80 studies were evaluated for quality and risk of bias. The general result indicates the acceptability of the quality of the studies. Measurement properties such as responsiveness and criterion validity, as well as the standard error of measurement have been neglected in most studies. The evaluation of these properties in future studies can help to create more accurate evidence about the psychometric properties of the CD-RISC versions. The findings of the present study provide an image of the psychometric status of the CD-RISC versions for

researchers and clinicians in this field and can guide them in their decision to select a study tool or conduct additional psychometric studies or adopt this tool in new populations.

Ethics approval

This study was approved in ethic committee of Mazandaran University of Medical Sciences (code: IR.MAZUMS.REC.1402.264).

Consent

Informed consent was not required for this systematic review.

Source of funding

No fund was received from any organization for publishing this study.

Author contribution

H.S.N. and A.H.G. contributed in study design. A.B. and A.H.G. contributed in searching of databases. Evaluation of the quality of the studies has done by L.H. and S.H. Also, H.S.N., A.H.G., S.H., L.H., F.K.F., M.M., and F.M. wrote the manuscript. D.S.T., E.S.F., and H.S.H. critically revised the manuscript. All of the authors proved the final version of manuscript.

Conflicts of interest disclosure

The authors declare no conflict of interests.

Research registration unique identifying number (UIN)

This study wasn't submitted in Prospero. So, just the proposal of this study was approved by student research committee of Mazandaran University of medical sciences.

Guarantor

Amir Hossein Goudarzian.

Data availability

This article was derived from secondary sources (published research articles) which are cited in the reference list. No primary data are included. All data generated or analyzed during this study are included in this published article.

Provenance and peer review

Not commissioned, externally peer-reviewed.

Acknowledgements

The authors thank student research committee of Mazandaran University of Medical Sciences (Sari, Iran) for financial supports.

References

- [1] Yates TM, Masten AS. Fostering the Future: Resilience Theory and the Practice of Positive Psychology. *Positive Psychology in Practice*. John Wiley & Sons, Inc; 2004:521–539.
- [2] Yates TM, Tyrell FA, Masten AS. Resilience theory and the practice of positive psychology from individuals to societies. *Posit Psychol Pract*, 2nd ed. Wiley Online Library; 2015;44:773–88.
- [3] Knipscheer J, Sleijpen M, Frank L, *et al*. Prevalence of Potentially Traumatic Events, Other Life Events and Subsequent Reactions Indicative for Posttraumatic Stress Disorder in the Netherlands: a General Population Study Based on the Trauma Screening Questionnaire. *Int J Environ Res Public Health* 2020;17:1725.
- [4] Connor KM, Davidson JRT. Development of a new resilience scale: The Connor-Davidson Resilience Scale (CD-RISC). *Depress Anxiety* 2003; 18:76–82.
- [5] Masten AS. Resilience from a developmental systems perspective. *World Psychiatry* 2019;18:101–2.
- [6] Ayed N, Toner S, Priebe S. Conceptualizing resilience in adult mental health literature: a systematic review and narrative synthesis. *Psychol Psychother Theory Res Pract* 2019;92:299–341.
- [7] Rezaei-pandari H, Mohammadpoorasl A, Morowatisharifabad MA, *et al*. Psychometric properties of the Persian version of abridged Connor-Davidson Resilience Scale 10 (CD-RISC-10) among older adults. *BMC Psychiatry* 2022;22:493.
- [8] Sharif Nia H, She L, Froelicher ES, *et al*. Psychometric evaluation of the Connor-Davidson Resilience Scale among Iranian population. *BMC Psychiatry* 2023;23:1–9.
- [9] Lamond AJ, Depp CA, Allison M, *et al*. Measurement and predictors of resilience among community-dwelling older women. *J Psychiatr Res* 2008;43:148–54.
- [10] Salisu I, Hashim N. A critical review of scales used in resilience research. *IOSR J Business Manag* 2017;19:23–33.
- [11] Afek A, Ben-Avraham R, Davidov A, *et al*. Psychological resilience, mental health, and inhibitory control among youth and young adults under stress. *Front Psychiatry* 2021;11:608588.
- [12] Li F, Luo S, Mu W, *et al*. Effects of sources of social support and resilience on the mental health of different age groups during the COVID-19 pandemic. *BMC Psychiatry* 2021;21:16.
- [13] Xue S, Kaufman MR, Zhang X, *et al*. Resilience and Prosocial Behavior Among Chinese University Students During COVID-19 Mitigation: Testing Mediation and Moderation Models of Social Support. *Psychol Res Behav Manag* 2022;15:1531–43.
- [14] Shea BJ, Reeves BC, Wells G, *et al*. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ* 2017;358:j4008.
- [15] Page MJ, McKenzie JE, Bossuyt PM, *et al*. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Int J Surg* 2021;88:105906.
- [16] Mokkink LB, De Vet HC, Prinsen CA, *et al*. COSMIN risk of bias checklist for systematic reviews of patient-reported outcome measures. *Qual Life Res* 2018;27:1171–9.
- [17] Mokkink LB, Prinsen C, Patrick DL, *et al*. COSMIN methodology for systematic reviews of patient-reported outcome measures (PROMs). *User Manual* 2018;78:6–3.
- [18] Terwee CB, Mokkink LB, Knol DL, *et al*. Rating the methodological quality in systematic reviews of studies on measurement properties: a scoring system for the COSMIN checklist. *Qual Life Res* 2012;21:651–7.
- [19] Prinsen CA, Mokkink LB, Bouter LM, *et al*. COSMIN guideline for systematic reviews of patient-reported outcome measures. *Qual Life Res* 2018;27:1147–57.
- [20] Hahs-Vaughn DL. *Applied Multivariate Statistical Concepts*. Taylor & Francis; 2016.
- [21] Park HS, Dailey R, Lemus D. The use of exploratory factor analysis and principal components analysis in communication research. *Hum Commun Res* 2002;28:562–77.
- [22] Child D. *The Essentials of Factor Analysis*. Bloomsbury Academic; 2006.
- [23] Abdi F, Sh B, Ahadi H, *et al*. Psychometric properties of the Connor-Davidson Resilience Scale (CD-RISC) among women with breast cancer. *J Res Psychol Health* 2019;13:81–99.
- [24] Alarcón R, Cerezo MV, Hevilla S, *et al*. Psychometric properties of the Connor-Davidson Resilience Scale in women with breast cancer. *Int J Clin Health Psychol* 2020;20:81–9.

- [25] Aloba O, Ajao O, Akinsulore S, *et al.* Exploration of the psychometric properties and correlates of the 10 item connor-davidson resilience scale among family caregivers of Nigerian patients with psychiatric disorder. *Int J Ment Health Psychiatry* 2016B;4372:2.
- [26] Aloba O, Olabisi O, Aloba T. The 10-item Connor–Davidson Resilience Scale: Factorial structure, reliability, validity, and correlates among student nurses in southwestern Nigeria. *J Am Psychiatr Nurses Assoc* 2016A;22:43–51.
- [27] Anjos JF, Dos Santos MJH, Ribeiro MT, *et al.* Connor-Davidson Resilience Scale: validation study in a Portuguese sample. *BMJ Open* 2019;9:e026836.
- [28] Baek H-S, Lee K-U, Joo E-J, *et al.* Reliability and validity of the Korean version of the Connor-Davidson Resilience Scale. *Psychiatry Investig* 2010;7:109.
- [29] Bakhshayesh Eghbali B, Pourramzani A, Bahadivand Chegini AH, *et al.* Validity and Reliability of Connor-Davidson Resilience Scale in Patients with. *Multiple Scler* 2022;9:45–50.
- [30] Bezdjian S, Schneider KG, Burchett D, *et al.* Resilience in the United States Air Force: Psychometric properties of the Connor-Davidson Resilience Scale (CD-RISC). *Psychol Assess* 2017;29:479.
- [31] Bizri M, Ibrahim N, Beyrouthy C, *et al.* Psychometric properties of the Arabic version of Connor-Davidson Resilience Scale in a sample of Lebanese women. *PLoS One* 2022;17:e0269700.
- [32] Blanco V, Guisande MA, Sánchez MT, *et al.* Spanish validation of the 10-item Connor–Davidson Resilience Scale (CD-RISC 10) with non-professional caregivers. *Aging Ment Health* 2019;23:183–8.
- [33] Campbell-Sills L, Stein MB. Psychometric analysis and refinement of the connor–davidson resilience scale (CD-RISC): Validation of a 10-item measure of resilience. *J Traum Stress* 2007;20:1019–28.
- [34] Cheng C, Dong D, He J, *et al.* Psychometric properties of the 10-item Connor–Davidson Resilience Scale (CD-RISC-10) in Chinese undergraduates and depressive patients. *J Affect Disord* 2020;261: 211–20.
- [35] Coates EE, Phares V, Dedrick RF. Psychometric properties of the Connor-Davidson Resilience Scale 10 among low-income, African American men. *Psychol Assess* 2013;25:1349.
- [36] Daniel-González L, Cadena CHG, Caycho-Rodríguez T, *et al.* Validation study of the 10-item Connor-Davidson resilience scale among Mexican medical and psychology students. *Revista de Psicología y Ciencias del Comportamiento de la Unidad Académica de Ciencias Jurídicas y Sociales* 2020;11:4–18.
- [37] Derakhshanrad SA, Piven E, Rassafiani M, *et al.* Standardization of connor-davidson resilience scale in iranian subjects with cerebrovascular accident. *J Rehabil Sci Res* 2014;1:73–7.
- [38] Dominguez-Cancino KA, Calderon-Maldonado FL, Choque-Medrano E, *et al.* Psychometric Properties of the Connor-Davidson Resilience Scale for South America (CD-RISC-25SA) in Peruvian Adolescents. *Children* 2022;9:1689.
- [39] Duong C, Hurst CP. Reliability and validity of the Khmer version of the 10-item Connor-Davidson Resilience Scale (Kh-CD-RISC10) in Cambodian adolescents. *BMC Res Notes* 2016;9:1–6.
- [40] Fujikawa M, Lee E-J, Chan F, *et al.* The Connor-Davidson Resilience Scale as a positive psychology measure for people with spinal cord injuries. *Rehabil Res Policy Educ* 2013;27:213–22.
- [41] Gao J, Li Y, Wu X. Revision and validation of the Connor-Davidson Resilience Scale of coal miners in China. *Int J Ind Ergon* 2021;85: 103191.
- [42] García León MdlÁ, González Gómez A, Robles Ortega H, Padilla García, *et al.* Psychometric properties of the Connor-Davidson Resilience Scale (CD-RISC) in the Spanish population. *J Child Adolesc Trauma* 2019;14:29–40.
- [43] Goins RT, Gregg JJ, Fiske A. Psychometric properties of the Connor-Davidson resilience scale with older American Indians: The native elder care study. *Res Aging* 2013;35:123–43.
- [44] Gonzalez SP, Moore EWG, Newton M, *et al.* Validity and reliability of the Connor-Davidson Resilience Scale (CD-RISC) in competitive sport. *Psychol Sport Exerc* 2016;23:31–9.
- [45] Gorman E, Heritage B, Shepherd CC, *et al.* A psychometric evaluation of the Connor–Davidson Resilience Scale for Australian Aboriginal youth. *Aust Psychol* 2021;56:25–37.
- [46] Gouda E, Kamal D, Fouad S, *et al.* Examining the Psychometrics of the Connor-Davidson Resilience Scale (CD-RISC) among Medical Staff Members at the Faculty of Medicine, Suez Canal University, Egypt. *J Ecophysiol Occup Health* 2022;22:22–8.
- [47] Gras M-E, Font-Mayolas S, Baltasar A, *et al.* The connor-davidson resilience scale (CD-RISC) amongst young Spanish adults. *Clínica y Salud* 2019;30:73–9.
- [48] Green KT, Hayward LC, Williams AM, *et al.* Examining the Factor Structure of the Connor–Davidson Resilience Scale (CD-RISC) in a Post-9/11 U.S. Military Veteran Sample. *Assessment* 2014;21:443–51.
- [49] Guarnizo Guzmán CP, Martín G, Suárez Falcón J, *et al.* Psychometric properties of the Connor-Davidson Resilience Scale (CD-RISC) on vulnerable colombian adolescents. *Int J Psychol Psychol Ther* 2019;19: 277–89.
- [50] Guihard G, Deumier L, Alliot-Licht B, *et al.* Psychometric validation of the French version of the Connor-Davidson Resilience Scale. *L'encephale* 2018;44:40–5.
- [51] Hwang YI, Arnold S, Trollor J, *et al.* Factor structure and psychometric properties of the brief Connor–Davidson Resilience Scale for adults on the autism spectrum. *Autism* 2020;24:1572–7.
- [52] Jeong HS, Kang I, Namgung E, *et al.* Validation of the Korean version of the Connor–Davidson Resilience Scale-2 in firefighters and rescue workers. *Compr Psychiatry* 2015;59:123–8.
- [53] Jorgensen IE, Seedat S. Factor structure of the Connor-Davidson resilience scale in South African adolescents. *Int J Adolesc Med Health* 2008;20:23–32.
- [54] Jung YE, Min JA, Shin AY, *et al.* The Korean version of the Connor–Davidson Resilience Scale: an extended validation. *Stress Health* 2012; 28:319–26.
- [55] Karairmak Ö. Establishing the psychometric qualities of the Connor–Resilience Scale (CD-RISC) using exploratory and confirmatory factor analysis in a trauma survivor sample. *Psychiatry Res* 2010;179:350–6.
- [56] Keyhani M, Taghvaei D, Rajabi A, *et al.* Internal consistency and confirmatory factor analysis of the Connor-Davidson Resilience Scale (CD-RISC) among nursing female. *Iranian J Med Educ* 2015;14:857–65.
- [57] Khoshouei MS. Psychometric evaluation of the Connor-Davidson resilience scale (CD-RISC) using Iranian students. *Int J Testing* 2009;9: 60–6.
- [58] Kidd C, Romaniuk M, McLeay S, *et al.* Resilience and psychopathology in trauma-exposed Australian Veterans: An exploratory factor analysis of the Connor-Davidson Resilience Scale. *J Military Veteran Fam Health* 2019;5:27–39.
- [59] Kuiper H, van Leeuwen CC, Stolwijk-Swüste JM, *et al.* Measuring resilience with the Connor–Davidson Resilience Scale (CD-RISC): which version to choose? *Spinal Cord* 2019;57:360–6.
- [60] Kwan YH, Ng A, Lim KK, *et al.* Validity and reliability of the ten-item Connor–Davidson Resilience Scale (CD-RISC10) instrument in patients with axial spondyloarthritis (axSpA) in Singapore. *Rheumatol Int* 2019; 39:105–10.
- [61] Kyriazos T, Stalikas A. Psychometric Evidence of the 10-Item Connor-Davidson Resilience Scale (CD-RISC10, Greek Version) and the Predictive Power of Resilience on Well-Being and Distress. *Open J Soc Sci* 2021;9:280–308.
- [62] Lauridsen LS, Willert MV, Eskildsen A, *et al.* Cross-cultural adaptation and validation of the Danish 10-item Connor-Davidson Resilience Scale among hospital staff. *Scand J Public Health* 2017;45:654–7.
- [63] Levey EJ, Rondon MB, Sanchez S, *et al.* Psychometric properties of the Spanish version of the 10-item Connor Davidson Resilience Scale (CD-RISC) among adolescent mothers in Peru. *J Child Adolesc Trauma* 2021;14:29–40.
- [64] Madewell A, Ponce-Garcia E, Martin S. Data replicating the factor structure and reliability of commonly used measures of resilience: The Connor–Davidson Resilience Scale, Resilience Scale, and Scale of Protective Factors. *Data Brief* 2016;8:1387–90.
- [65] Manzano García G, Ayala Calvo JC. Psychometric properties of Connor-Davidson Resilience Scale in a Spanish sample of entrepreneurs. *Psicothema* 2013;25:245–51.
- [66] Martinez S, Jones JD, Brandt L, *et al.* Factor structure and psychometric properties of the Connor–Davidson resilience scale (CD-RISC) in individuals with opioid use disorder. *Drug Alcohol Depend* 2021;221: 108632.
- [67] McGillivray K, Ho R. Validation of the Connor Davidson Resilience Scale (CD-RISC) as applied within the Thai context. *Scholar: Human Sciences* 2016;8:178–87.
- [68] Mealer M, Schmiege SJ, Meek P. The Connor-Davidson Resilience Scale in critical care nurses: a psychometric analysis. *J Nurs Meas* 2016;24: 28–39.

- [69] Meng M, He J, Guan Y, *et al.* Factorial invariance of the 10-item Connor-Davidson resilience scale across gender among Chinese elders. *Front Psychol* 2019;10:1237.
- [70] Miller MJ, Mealer ML, Cook PF, *et al.* Psychometric assessment of the Connor-Davidson Resilience Scale for people with lower-limb amputation. *Phys Ther* 2021;101:pzab002.
- [71] Minh-Uyen VT, Im S. Psychometric examination of the Connor-Davidson resilience scale (CD-RISC-10) among Vietnamese students. *J Appl Res Higher Educ* 2021;13:325–41.
- [72] Nagle Y, Rani EK, Masood A. Validation of Connor-Davidson Resilience Scale on Indian Military Soldiers. *Eur J Psychol Res* 2021;8:9–25.
- [73] Nartova-Bochaver S, Korneev A, Bochaver K. Validation of the 10-item Connor-Davidson resilience scale: the case of Russian youth. *Front Psychiatry* 2021;12:611026.
- [74] Nguyen Q-AN, Dinh H-VT. The Validation of the 10-Item Connor-Davidson Resilience Scale and Its Correlation to Emotional Intelligence and Life Satisfaction Among Vietnamese Late Adolescents. *J Indian Assoc Child Adolesc Mental Health* 2022;18:09731342221142072.
- [75] Ni MY, Li TK, Yu NX, *et al.* Normative data and psychometric properties of the Connor-Davidson Resilience Scale (CD-RISC) and the abbreviated version (CD-RISC2) among the general population in Hong Kong. *Qual Life Res* 2016;25:111–6.
- [76] Notario-Pacheco B, Solera-Martínez M, Serrano-Parra MD, *et al.* Reliability and validity of the Spanish version of the 10-item Connor-Davidson Resilience Scale (10-item CD-RISC) in young adults. *Health Qual Life Outcomes* 2011;9:1–6.
- [77] Pretorius TB, Padmanabhanunni A. Validation of the Connor-Davidson resilience scale-10 in South Africa: item response theory and classical test theory. *Psychol Res Behav Manag* 2022;78:1235–45.
- [78] Sexton MB, Muzik M, McGinnis EG, *et al.* Psychometric characteristics of the Connor-Davidson resilience scale (CD-RISC) in postpartum mothers with histories of childhood maltreatment. *Nurs Midwifery Stud* 2016;5. doi:10.17795/NMSJOURNAL34589.
- [79] Sharma S, Pathak A, Abbott JH, *et al.* Measurement properties of the Nepali version of the Connor Davidson resilience scales in individuals with chronic pain. *Health Qual Life Outcomes* 2018;16:1–11.
- [80] She R, Yang X, Lau MM, *et al.* Psychometric properties and normative data of the 10-item Connor-Davidson Resilience Scale among Chinese adolescent students in Hong Kong. *Child Psychiatry Hum Dev* 2020;51:925–33.
- [81] Shin G-S, Choi K-S, Jeong KS, *et al.* Psychometric properties of the 10-item Connor-Davidson resilience scale on toxic chemical-exposed workers in South Korea. *Ann Occup Environ Med* 2018;30:1–7.
- [82] Singh K, Yu X-n. Psychometric evaluation of the Connor-Davidson Resilience Scale (CD-RISC) in a sample of Indian students. *J Psychol* 2010;1:23–30.
- [83] Smith KJ, Emerson DJ, Haight TD, *et al.* An examination of the psychometric properties of the Connor-Davidson Resilience Scale-10 (CD-RISC10) among accounting and business students. *J Account Educ* 2019;47:48–62.
- [84] Solano JPC, Bracher ESB, Faisal-Cury A, *et al.* Factor structure and psychometric properties of the Connor-Davidson resilience scale among Brazilian adult patients. *Sao Paulo Med J* 2016;134:400–6.
- [85] Tomin AJ, Weinberg MK. Resilience and subjective wellbeing: a psychometric evaluation in young Australian adults. *Aust Psychol* 2018;53:68–76.
- [86] Tourunen A, Siltanen S, Saajanaho M, *et al.* Psychometric properties of the 10-item Connor-Davidson resilience scale among Finnish older adults. *Aging Ment Health* 2021;25:99–106.
- [87] Tsigkaropoulou E, Douzenis A, Tsitis N, *et al.* Greek version of the Connor-Davidson resilience scale: Psychometric properties in a sample of 546 subjects. *In Vivo* 2018;32:1629–34.
- [88] Vaishnavi S, Connor K, Davidson JR. An abbreviated version of the Connor-Davidson Resilience Scale (CD-RISC), the CD-RISC2: Psychometric properties and applications in psychopharmacological trials. *Psychiatry Res* 2007;152:293–7.
- [89] Velickovic K, Rahm Hallberg I, Axelsson U, *et al.* Psychometric properties of the Connor-Davidson Resilience Scale (CD-RISC) in a non-clinical population in Sweden. *Health Qual Life Outcomes* 2020;18:1–10.
- [90] Vongsirimas N, Thanoi W, Yobas PK. Evaluating Psychometric Properties of the Connor-Davidson Resilience Scale (10-Item CD-RISC) among University Students in Thailand: การทดสอบ คุณภาพ เครื่องมือแบบ วัด ความ ความ แข็งแกร่ง และ ยึดหยุ่น ของ ชีวิต Connor-Davidson Resilience Scale (10-item CD-RISC) ใน กลุ่ม นักศึกษา ไทย. *Nursing Sci J Thailand* 2017;35:25–35.
- [91] Waddimba AC, Baker BM, Pogue JR, *et al.* Psychometric validity and reliability of the 10-and 2-item Connor-Davidson resilience scales among a national sample of Americans responding to the Covid-19 pandemic: an item response theory analysis. *Qual Life Res* 2022;31:2819–36.
- [92] Wang L, Shi Z, Zhang Y, *et al.* Psychometric properties of the 10-item Connor-Davidson Resilience Scale in Chinese earthquake victims. *Psychiatry Clin Neurosci* 2010;64:499–504.
- [93] Wu L, Tan Y, Liu Y. Factor structure and psychometric evaluation of the Connor-Davidson resilience scale in a new employee population of China. *BMC Psychiatry* 2017;17:1–7.
- [94] Xie Y, Peng L, Zuo X, *et al.* The psychometric evaluation of the Connor-Davidson resilience scale using a Chinese military sample. *PLoS One* 2016;11:e0148843.
- [95] Ye ZJ, Qiu HZ, Li PF, *et al.* Validation and application of the Chinese version of the 10-item Connor-Davidson Resilience Scale (CD-RISC-10) among parents of children with cancer diagnosis. *Eur J Oncol Nursing* 2017;27:36–44.
- [96] Yu X, Zhang J. Factor analysis and psychometric evaluation of the Connor-Davidson Resilience Scale (CD-RISC) with Chinese people. *Soc Behav Personal Int J* 2007;35:19–30.
- [97] Yu X-n, Lau JT, Mak WW, *et al.* Factor structure and psychometric properties of the Connor-Davidson Resilience Scale among Chinese adolescents. *Compr Psychiatry* 2011;52:218–24.
- [98] Zhang X, Mo Y, Yan C, *et al.* Psychometric properties of two abbreviated Connor-Davidson Resilience scales in Chinese infertile couples. *Qual Life Res* 2021;30:2405–14.
- [99] Sharif-Nia H, She L, Froelicher ES, *et al.* Resilience in Iranian adolescents during the COVID-19 pandemic: Psychometric Evaluation of the Connor-Davidson Resilience Scale. *J Child Adolesc Ment Health* 2021;33:111–22.
- [100] Levey EJ, Rondon MB, Sanchez S, *et al.* Psychometric properties of the Spanish version of the 10-item Connor Davidson Resilience Scale (CD-RISC) among adolescent mothers in Peru. *J Child Adolesc Trauma* 2019;14:29–40.
- [101] Mokkink LB, Boers M, van der Vleuten CPM, *et al.* COSMIN Risk of Bias tool to assess the quality of studies on reliability or measurement error of outcome measurement instruments: a Delphi study. *BMC Med Res Methodol* 2020;20:293.
- [102] Scimago Journal & Country Rank. Country Ranking. 2016. <https://www.scimagojr.com/countryrank.php?order=itp&ord=desc&year=2016>
- [103] Terán J-M, Varea C, Juárez S, *et al.* Social disparities in Low Birth Weight among Spanish mothers during the economic crisis (2007–2015). *Nutrición Hospitalaria* 2018;35:129–41.
- [104] Frasilho D, Matos MG, Salonna F, *et al.* Mental health outcomes in times of economic recession: a systematic literature review. *BMC Public Health* 2016;16:115.
- [105] Windle G. What is resilience? A review and concept analysis. *Rev Clin Gerontol* 2011;21:152–69.
- [106] Windle G, Bennett KM, Noyes J. A methodological review of resilience measurement scales. *Health Qual Life Outcomes* 2011;9:8.
- [107] Kavčič T, Zager Kocjan G, Dolenc P. Measurement invariance of the CD-RISC-10 across gender, age, and education: A study with Slovenian adults. *Curr Psychol* 2023;42:1727–37.
- [108] Gjersing L, Caplehorn JRM, Clausen T. Cross-cultural adaptation of research instruments: language, setting, time and statistical considerations. *BMC Med Res Methodol* 2010;10:13.
- [109] Bollen KA. Structural equations with latent variables. John Wiley & Sons; 1989.
- [110] Watson R, Thompson DR. Use of factor analysis in Journal of Advanced Nursing: literature review. *J Adv Nurs* 2006;55:330–41.
- [111] Nunnally JC. Psychometric Theory. McGraw-Hill; 1978.
- [112] Tavakol M, Wetzel A. Factor Analysis: a means for theory and instrument development in support of construct validity. *Int J Med Educ* 2020;11:245.
- [113] Loewen S, Gonulal T. Exploratory factor analysis and principal components analysis. *Advancing Quantitative Methods in Second Language Research*. Routledge; 2015:182–12.
- [114] Schreiber JB. Issues and recommendations for exploratory factor analysis and principal component analysis. *Res Soc Administr Pharm RSAP* 2021;17:1004–11.

- [115] Alavi M, Visentin DC, Thapa DK, *et al.* Cleary MJJoAN. Exploratory factor analysis and principal component analysis in clinical studies: Which one should you use?. *J Adv Nurs* 2020;76:1886–9.
- [116] Schneeweiss H, Mathes H. Factor analysis and principal components. *J Multivariate Anal* 1995;55:105–24.
- [117] Suhr DD. Exploratory or confirmatory factor analysis? SAS Institute Cary: Cary, North Carolina. 2006:200–31.
- [118] Hox JJ. Confirmatory factor analysis. *Encycl Res Methods Criminol Criminal Justice*, 1st ed. Wiley Online Library: New Jersey, USA. 2021; 158:830–2.
- [119] Mokkink LB, Terwee CB, Knol DL, *et al.* The COSMIN checklist for evaluating the methodological quality of studies on measurement properties: a clarification of its content. *BMC Med Res Methodol* 2010; 10:22.
- [120] Mokkink LB, Terwee CB, Patrick DL, *et al.* The COSMIN checklist for assessing the methodological quality of studies on measurement properties of health status measurement instruments: an international Delphi study. *Qual Life Res* 2010;19:539–49.
- [121] Angst F. The new COSMIN guidelines confront traditional concepts of responsiveness. *BMC Med Res Methodol* 2011;11:152.
- [122] Cosco TD, Kaushal A, Richards M, *et al.* Resilience measurement in later life: a systematic review and psychometric analysis. *Health Qual Life Outcomes* 2016;14:16.
- [123] Cockroft JD, Rabin J, Yockey RA, *et al.* Psychometric Properties of Scales Measuring Resilience in U.S. Latinx Populations: a Systematic Review. *Health Equity* 2023;7:148–60.
- [124] Janousch C, Anyan F, Hjerdal O, *et al.* Psychometric Properties of the Resilience Scale for Adolescents (READ) and Measurement Invariance Across Two Different German-Speaking Samples. *Front Psychol* 2020; 11:1–16.
- [125] Askeland KG, Hysing M, Sivertsen B, *et al.* Factor Structure and Psychometric Properties of the Resilience Scale for Adolescents (READ). *Assessment* 2020;27:1575–87.
- [126] Shi X, Wang S, Wang Z, *et al.* The resilience scale: factorial structure, reliability, validity, and parenting-related factors among disaster-exposed adolescents. *BMC Psychiatry* 2021;21:145.