



Universiteit
Leiden
The Netherlands

Measuring pain in aphasia: validity and reliability of the PACSLAC-D

Vries, N.J. de; Steen, J.T. van der; Achterberg, W.P.; Smaling, H.J.A.

Citation

Vries, N. J. de, Steen, J. T. van der, Achterberg, W. P., & Smaling, H. J. A. (2023).
Measuring pain in aphasia: validity and reliability of the PACSLAC-D. *Pain Management
Nursing*, 24(4), E68-E74. doi:10.1016/j.pmn.2023.03.010

Version: Publisher's Version
License: [Creative Commons CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/)
Downloaded from: <https://hdl.handle.net/1887/3761757>


Note: To cite this publication please use the final published version (if applicable).



ELSEVIER

Contents lists available at ScienceDirect

Pain Management Nursing

journal homepage: www.painmanagementnursing.org

Original Article

Measuring Pain in Aphasia: Validity and Reliability of the PACSLAC-D

Neeltje J. de Vries, M.A.^{*,†,#}, Jenny T. van der Steen, Ph.D.^{*,‡}, Wilco P. Achterberg, M.D. Ph.D.^{*,§}, Hanneke J.A. Smaling, Ph.D.^{*,§}^{*} Department of Public Health and Primary Care, Leiden University Medical Center, the Netherlands[†] TOPAZ Geriatric Rehabilitation Center Revitel, Leiden, the Netherlands[‡] Department of Primary and Community Care, Radboud university medical center, Nijmegen, the Netherlands[§] University Network for the Care sector Zuid-Holland, Leiden University Medical Center, the Netherlands

ARTICLE INFO

Article history:

Received 30 June 2022

Received in revised form 6 March 2023

Accepted 18 March 2023

ABSTRACT

Background: Post-stroke pain in patients with an inability to communicate is not systematically assessed and therefore not sufficiently treated. This stresses the need to study pain assessment instruments that do not require good communication skills.

Aim: To examine the validity and reliability of the Pain Assessment Checklist for Seniors with Limited Ability to Communicate – Dutch version (PACSLAC-D) in stroke patients with aphasia.

Method: Sixty stroke patients (mean age 79.3 years, standard deviation [SD] 8.0), of whom 27 had aphasia were observed during rest, activities of daily living (ADL), and physiotherapy using the Pain Assessment Checklist for Seniors with Limited Ability to Communicate – Dutch version (PACSLAC-D). The observations were repeated after two weeks. To examine convergent validity, correlations between the PACSLAC-D, self-report pain scales, and the clinical judgment of a health care professional (pain present yes/no) were used. To examine discriminative validity, differences in pain were investigated between rest and ADL, in patients who use pain medication and those who do not, and in patients with and without aphasia. Internal consistency and test-retest reliability were assessed to determine reliability.

Results: Convergent validity failed to meet the acceptable threshold during rest but was adequate during ADL and physiotherapy. Discriminative validity was only adequate during ADL. The internal consistency was 0.33 during rest, 0.71 during ADL, and 0.65 during physiotherapy. Test-retest reliability varied from poor during rest (intraclass correlation coefficient [ICC] = 0.07; 95% confidence interval [CI]: -0.40-0.51) to excellent during physiotherapy (ICC = 0.95; 95% CI: 0.83-0.98).

Conclusions: The PACSLAC-D captures pain in patients with aphasia who are unable to self-report, during ADL and physiotherapy, but may be less accurate during rest.

© 2023 The Authors. Published by Elsevier Inc. on behalf of American Society for Pain Management Nursing.

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

Different types of pain are common after stroke (Delpont et al., 2018), for instance headache, shoulder, and central post-stroke pain (CPSP) (Hansen et al., 2012). CPSP, for example, affects 11% of stroke patients (Liampas et al., 2020). Almost 40% of stroke survivors (n = 281) experienced pain to some degree 5 years after stroke, with 15% reporting frequent pain, and 25% reporting that their needs for pain treatment were not met (Westerlind et al., 2020). These rates are comparable with other common types of pain in older adults without stroke, who reported musculoskeletal pain (40%), peripheral neuropathic pain (40%), and chronic joint pain (Jones et al., 2016).

Self-report pain scales are considered the gold standard to measure pain, including in stroke patients (Harrison & Field, 2015). Examples of self-report pain scales are the Numerical Rating Scale (NRS) (Hjermstad et al., 2011), Visual Analogue Scale (VAS) (Heller et al., 2016), and Faces Pain Scale (FPS) (Kim & Buschmann, 2006). However, the use of self-report pain scales can be difficult for stroke patients with aphasia and other cognitive deficits. An estimated 30% of stroke patients develop aphasia (Engelter, 2006; Mitchell et al., 2020; Wu et al., 2020). Most stroke patients with aphasia or communication problems are unable to complete self-report pain scales (Schuster et al., 2020; Smith et al., 2013).

A pain observation instrument score can serve as a proxy for measuring self-reported pain in stroke patients with aphasia. Pain observation instruments are regularly used in people with demen-

[#] Address correspondence to Neeltje J. de Vries, M.A., Public Health and Primary Care, Postzone VO-P, Postbus 9600, 2300 RC Leiden.

E-mail address: n.j.de_vries@lumc.nl (N.J. de Vries).

tia who also have cognitive and communication problems (Coca & Zuniga, 2020; Haghi et al., 2020; Natavio et al., 2020; Van Dalen-Kok et al., 2021). The Pain Assessment Checklist for Seniors with Limited Ability to Communicate (PACSLAC; Kaasalainen et al., 2013) is an example of a pain observation instrument. This originally Canadian instrument consists of 60 items. The PACSLAC has shown adequate psychometric qualities for cognitively impaired older people in acute and long-term care settings (Fuchs-Lacelle & Hadjstavropoulos, 2004; Kaasalainen et al., 2013; Natavio et al., 2020; Qi et al., 2012). The PACSLAC was revised into a 31-item version, the PACSLAC-II (Chan et al., 2014; Hadjstavropoulos et al., 2014; Ruest et al., 2017). It differentiates between painful and non-painful states in older long-term care residents with dementia and older adult outpatients without dementia (Chan et al., 2014; Hadjstavropoulos et al., 2018). Several studies with translations in different languages, indicate that it is a valid and reliable observation instrument for the measurement of pain in older adults with dementia (Aubin et al., 2008; Büyükturan et al., 2018; Haghi et al., 2020; Kim et al., 2014; Takai et al., 2013; Thé et al., 2016; Van Nispen tot Pannerden et al., 2009; Zwakhalen et al., 2006).

The PACSLAC-D was developed based on the 60-item original Canadian PACSLAC instrument: (Zwakhalen et al., 2007). Zwakhalen et al. (2007) validated the reduced 24-item PACSLAC-D.

Post-stroke pain in patients with an inability to communicate is not systematically assessed and therefore not sufficiently treated (Schuster et al., 2020). This stresses the need to study the psychometric properties and feasibility of assessment instruments that do not require good communication skills. The aim of this study is to determine the validity and reliability of PACSLAC-D, an observational instrument, in stroke patients with aphasia.

Methods

Design and Study Population

This study employed a prospective observational design. Data were collected from July 2014 to December 2018. Patients who met the following selection criteria were invited by a speech and language therapist to participate in the study: ≥ 18 years old and staying at the stroke unit of a Geriatric Rehabilitation Care center in The Netherlands. Patients with dementia or delirium were not eligible. Patients who were able to communicate gave oral informed consent for participation in this study. If patients were not able to give verbal informed consent or if there were doubts about the patient's communication abilities, the legal representative also provided verbal informed consent. Stroke patients, both with and without aphasia, were included. If suspected, aphasia was diagnosed by a speech and language therapist using Token-Test (Doesborgh et al., 2003) or ScreeLing (El Hachoui et al., 2012). A score of ≥ 7 on the TokenTest or a score of ≤ 68 on ScreeLing indicates the presence of aphasia. If tests could not be taken, the clinical judgment of the speech and language therapist was decisive.

A sample size with a minimum of 50 patients is recommended for validation studies and for the analysis of reliability (De Vet et al., 2011).

Measurement Instruments

To assess the presence of pain, the PACSLAC-D, was used because of good psychometric properties in Dutch persons with dementia (Zwakhalen et al., 2007). The 24 items are related to face, resistance/defense, and social emotional/mood. The observer indicated for each item whether it was observed (1) or not (0). The total score ranges from 0 to 24, with a higher score indicating

more pain. A score of 4 or higher may indicate the presence of pain (Zwakhalen et al., 2009).

Further, two self-report pain scales were used to measure pain. The FPS consists of six vertically placed faces, with face 1 (no pain) at the bottom and face 6 (maximum pain) at the top (Kim & Buschmann, 2006). These 6 faces are assigned the following scores: 0 (no pain), 2, 4, 6, 8, and 10 (worst imaginable pain). Patients were asked to select the face that represents their experienced pain. The combined vertical NRS and VAS consists of a 10-centimeter vertical line with scores of 0 to 10, anchored by two extremes of pain: no pain (0) and extreme pain (10). Appendix A includes the FPS and the NRS/VAS combined scale.

For the clinical judgment of pain during activities of daily living (ADL), the nurse, and during physiotherapy the physiotherapist, were asked the question 'Is there pain?' Their Yes or No response was recorded by the observer as judgment nurse and judgment physiotherapist.

Assessment of Measurement Properties

Construct validity

For construct validity, the subtypes convergent and discriminative validity were determined.

Convergent validity

Moderate correlations between the PACSLAC-D and self-report pain scales were expected for the convergent validity. Similarly, moderate correlations with the clinical judgment of the presence of pain of the nurse and physiotherapist were hypothesized.

Discriminative validity

Three a-priori hypotheses were tested to examine discriminative validity: (1) more pain is expected in patients with aphasia during ADL compared with rest. In persons with dementia, more pain is observed during ADL compared to rest (Hadjstavropoulos et al., 2014; Lints-Martindale et al., 2012; Zwakhalen et al., 2006). (2) more pain is expected in patients with aphasia using pain medication than in those who use no pain medication. Persons with dementia who used pain medication have more pain than those who did not use pain medication (Rajkumar et al., 2017). Also, a study of hospitalized persons with dementia found that 60% (of $n = 108$) of persons who demonstrated pain received pain medication compared to 40% who did not receive pain medication (Boltz et al., 2021); (3) more pain is expected in patients with aphasia compared with patients without aphasia. Stroke patients with aphasia received significantly less pain medication compared with patients without aphasia and those with moderate to severe aphasia are often excluded from pain research (De Vries, Sloot, & Achterberg, 2016).

Reliability

For reliability, an acceptable internal consistency of PACSLAC-D in patients with aphasia was expected, and moderate test-retest reliability.

Procedure

After inclusion, the following sociodemographic characteristics were collected: sex, age, native language, hand dominance, stroke type, date of injury, stroke localization, analgesic medication, and the presence of aphasia. Hand dominance is related to the localization of language in the brain and, also with the localization of the stroke. In most cases, language is in the left hemisphere located and, sometimes in the right hemisphere (Carey & Johnstone, 2014; Vingerhoets, 2019).

Table 1
Patient Characteristics

		Total (n = 60)	Stroke patients with aphasia (n = 27)	Stroke patients without aphasia (n = 33)	Group comparisons t(df), p X ² (df), p or two tailed, p (Fisher's exact test)
Age (years)	Mean (SD)	79.3 (8.0)	79.3 (9.0)	79.4 (7.1)	t(58) = -0.02, p = .98
	Range	59.1-99.1	59.1-92.7	67.1-99.1	
Sex (female)	n (%)	26 (43%)	14 (52%)	20 (61%)	X ² (1) = 0.46, p = .49
Type of stroke	Ischemic, n(%)	52 (87%)	24 (89%)	28 (85%)	X ² (1) = 0.21, p = .64
	Left hemisphere	26 (43%)	21 (78%)	5 (15%)	two tailed, p = .72
	Right hemisphere	22 (37%)	3 (11%)	19 (58%)	
	Brainstem	2 (3%)	0	2 (6%)	
	Cerebellar	2 (3%)	0	2 (6%)	
	Hemorrhage, n(%)	8 (13%)	3 (11%)	5 (15%)	
	Left hemisphere	1 (2%)	0	1 (3%)	
Pain medication	Right hemisphere	1 (2%)	1 (4%)	0	
	Other	6 (10%)	2 (7%)	4 (12%)	
	Mean (SD)	1.4 (0.5)	1.3 (0.5)	1.4 (0.5)	t(58) = -1.02, p = .31
Unable to complete self-report pain scales ^b	n (%)	7 (12%)	6 (22%)	1 (3%)	two tailed, p = .04 ^a

^a p < .05^b ≥2 self-report pain scales not completed during Rest, Activities of Daily Living, and Physiotherapy. SD = standard deviation.

All patients, not blinded for aphasia, were observed by one observer on one day during rest, activities of living (ADL), and physiotherapy using the PACSLAC-D pain observation instrument for 5 to 10 minutes. The observer was a speech and language therapist with a university education level Master of Arts. Subsequently, the observer asked the patient to indicate the degree of experienced pain using the self-report pain scales FPS (Kim & Buschmann, 2006), and a combination of the NRS (Hjermstad et al., 2011), and VAS (Heller et al., 2016). 'Not applicable' was noted if the participant was unable to self-report using (one of) these scales. After the observation during ADL and physiotherapy, the nurse and physiotherapist respectively, with no knowledge of the PACSLAC-D score, were asked to use clinical judgment if the patient had experienced pain. After two weeks, the measurements were repeated.

Statistical Analysis

An overview of the characteristics of the patients was prepared using descriptive statistics. Group comparisons were obtained with *t*-test, Pearson χ^2 test or Fisher's exact test. To examine the convergent validity of PACSLAC-D in patients with aphasia, Pearson correlation coefficients were calculated between the PACSLAC-D, self-report pain scales, and the clinical judgment of the nurse and physiotherapist. To describe the strength of the correlation we used: .00-.19 = very weak; .20-.39 = weak; .40-.59 = moderate; .60-.79 = strong; .80-1.0 = very strong (Evans, 1996). Additionally, a 95% confidence interval (CI) using bootstrapping (number of samples: 1,000) of the correlations was calculated. To investigate discriminative validity, the three hypotheses were tested. First, a non-parametric Wilcoxon signed-rank test was used to examine whether more pain was observed during rest than ADL (paired test) in patients with aphasia. Second, a Kruskal-Wallis test was used to investigate if patients with aphasia who use pain medication experienced more pain than those without pain medication. Third, to examine whether patients with aphasia had more pain than patients without aphasia, a Mann-Whitney *U* test was used.

The reliability of the PACSLAC-D was examined using Cronbach's alpha. Cronbach's α -values ranging from 0.70 to 0.95 are generally considered acceptable (Bland & Altman, 1997).

In addition, test-retest reliability were assessed using intraclass correlation coefficient (ICC). Based on the 95% confident interval of

the ICC, a value between 0.50 and 0.75 indicates moderate reliability, between 0.75 and 0.90 good reliability, and higher than 0.90 excellent reliability (Koo & Li, 2016; Kunz et al., 2020). The analyses were performed using IBM SPSS Statistics version 25 for Windows, 2018.

Ethical Considerations

The study was performed in accordance with the Dutch Healthcare Quality, Complaints and Disputes Act (WKKGZ). Article 7 of this Act states that the institution should improve regular care and to that purpose should gather data. Patient participation was voluntary and took place with their consent and in compliance with data protection.

Results

This study included 60 stroke patients, of whom 43% (n = 26) were female. Their age ranged from 59 to 99 years, with a mean age of 79.3 years (SD 8.0). Of the 60 stroke patients, 27 (45%) had aphasia. Seven stroke patients (12%) were unable to complete the self-report pain scales. Six of these seven stroke patients had aphasia. The patient without aphasia had other cognitive and motor damage, including severe dysarthria due to basal nuclei stroke in the right hemisphere (Table 1).

Table 2 describes the PACSLAC-D and self-report pain scale scores during rest, ADL, and physiotherapy. All 60 patients were observed during during rest and ADL. Of these 60 stroke patients, 49 patients were observed during physiotherapy.

A small proportion of the patients (12%) could not complete all self-report pain scales, during the observations in different conditions.

Almost no pain was observed with PACSLAC-D during rest and most patients (75%) completed the self-report pain scales with the most possible score of 0. During ADL and physiotherapy more pain was observed using PACSLAC-D and some patients rate their pain with the self-report pain scales.

Significantly more pain was observed in stroke patients unable to self-report during ADL (n = 7, mean 3.0, SD 1.7) and physiotherapy (n = 4, mean 2.5, SD 1.0) compared with 53 stroke patients (88%) who were able to complete the self-report pain scales, ADL: n = 53, mean 0.54, SD 1.53; t(58) = -3.6, p < .05; physiotherapy: n = 45, mean 0.42, SD 1.03; t(47) = -3.9, p < .05. During rest,

Table 2
Descriptive Statistics of PACSLAC-D and Self-Report Pain Scales During Rest, Activities of Daily Living, and Physiotherapy

	Total (n = 60)			With aphasia (n = 27)			Without aphasia (n = 33)			Group comparisons Mann-Whitney <i>U</i> test
	N	Mean (SD)	Range	n	Mean (SD)	Range	N	Mean (SD)	Range	
Rest										
PACSLAC-D	60	0.15 (0.55)	0-3	27	0.11 (0.43)	0-2	33	0.18 (0.64)	0-3	$U = 437.00, z = -0.26, p = .79$
NRS/VAS	55	0.84 (2.13)	0-8	22	0.09 (0.30)	0	33	1.33 (2.64)	0-8	$U = 301.00, z = -1.58, p = .11$
FPS	55	0.84 (2.07)	0-8	22	0.27 (0.94)	0	33	1.21 (2.51)	0-8	$U = 304.50, z = -1.50, p = .14$
ADL										
PACSLAC-D	60	1.00 (1.71)	0-8	27	1.41 (1.70)	0-5	33	0.67 (1.67)	0-8	$U = 310.00, z = -2.37, p = .02^*$
NRS/VAS	53	1.77 (2.87)	0-9	21	1.33 (1.96)	0-7	32	2.06 (3.34)	0-9	$U = 330.00, z = -0.13, p = .90$
FPS	53	1.64 (2.66)	0-8	21	1.24 (1.61)	0-4	32	1.91 (3.16)	0-8	$U = 329.00, z = -0.15, p = .88$
Physiotherapy										
PACSLAC-D	49	0.59 (1.17)	0-4	20	0.80 (1.36)	0-4	29	0.45 (1.02)	0-4	$U = 258.00, z = -0.86, p = .39$
NRS/VAS	45	1.36 (2.58)	0-8	16	0.75 (1.30)	0-3	29	1.69 (3.03)	0-8	$U = 225.00, z = -0.21, p = .84$
FPS	45	1.24 (2.40)	0-8	16	0.75 (1.44)	0-4	29	1.52 (2.77)	0-8	$U = 216.00, z = -0.49, p = .63$

SD = standard deviation; PACSLAC-D = Pain Assessment Checklist for Seniors with Limited Ability to Communicate – Dutch version; NRS/VAS = Numeric Rating Scale/Visual Analogue Scale; FPS = Faces Pain Scale; ADL = Activities of Daily Living.

Table 3
Correlation Matrix PACSLAC-D and Self-Report Pain Scales in Patients With Aphasia

	Patients with aphasia (n = 27)	Rest		ADL		Physiotherapy			
		NRS-VAS	FPS	NRS-VAS	FPS	Judgment nurse	NRS-VAS	FPS	Judgment physiotherapist
PACSLAC-D	Pearson correlation	-0.07	-0.07	0.11	0.22	0.44 ^a	0.64 ^b	0.49	0.49
	Significance. (two-tailed)	0.76	0.77	0.65	0.35	0.05	0.01	0.054	0.054
	n	22	22	21	21	21	16	16	16
	95% CI	-0.17 - -0.05	-0.16 - -0.05	-0.27 - 0.64	-0.22 - 0.69	0.00 - 0.85	0.04 - 0.98	-0.25 - 0.94	0.21 - 1.00

^a $p < .05$.

^b $p < .01$. PACSLAC-D = Pain Assessment Checklist for Seniors with Limited Ability to Communicate – Dutch version; ADL = Activities of Daily Living; NRS/VAS = Numeric Rating Scale / Visual Analogue Scale; FPS = Faces Pain Scale; Judgment nurse = the nurse was asked to judge if any pain was present during ADL; Judgment physiotherapist = physiotherapist was asked to judge if any pain was present during physiotherapy; CI = confidence interval.

there was no difference in observed pain between patients who were unable ($n = 7$, mean 0.1, SD 0.38) and those who were able to complete self-report pain scales ($n = 53$, mean 0.2, SD 0.57); $t(58) = 0.4, p = .971$.

Convergent Validity

Table 3 shows the associations between the PACSLAC-D, self-report pain scales, and clinical judgment of pain by the nurse and physiotherapist in patients with aphasia. During rest and ADL, we found no significant correlations between PACSLAC-D and self-report pain scales. During ADL, we reported only a moderate positive correlation between PACSLAC-D and the judgment of the nurse. During physiotherapy, the PACSLAC-D was only strongly positively associated with the NRS/VAS. We found no significant correlations between the PACSLAC-D and the FPS or judgment of physiotherapist.

Discriminative Validity

No difference in pain is observed during ADL (median 1) compared with rest (median 0); $T = 25, z = -1.93, p = .053$.

Also, we found no difference in observed pain in patients with aphasia who used pain medication during rest (median = 0) and physiotherapy (median = 0) compared with those who did not use pain medication during rest (median = 0) and physiotherapy (median = 0); rest $H(1) = 0.49, p = .483$; physiotherapy $H(1) = 1.39, p = .238$. Only during ADL, significantly more pain is observed in patients with aphasia who used pain medication (median = 1) than those who did not use pain medication (median = 0); $H(1) = 6.33, p < .05$.

Table 4
Internal Consistency of PACSLAC-D Based on Observations Day 1 and 2

	Group	Cronbach's alpha
Rest	Patients with aphasia	0.33
	Patients without aphasia	0.69
ADL	Patients with aphasia	0.71
	Patients without aphasia	0.86
Physiotherapy	Patients with aphasia	0.65
	Patients without aphasia	0.73

PACSLAC-D = Pain Assessment Checklist for Seniors with Limited Ability to Communicate – Dutch version; consists of 24 items; ADL = Activities of Daily Living.

During rest, we found no difference in pain in patients with aphasia (median = 0) compared with non-aphasia patients (median = 0); $U = 437, z = -0.26, p = .792$. Significantly more pain was observed during ADL in patients with aphasia (median 1) compared with patients without aphasia (median = 0); $U = 310, z = -2.37, p < .05$. During physiotherapy, no difference in pain was observed between patients with aphasia (median = 0) and without aphasia (median = 0); $U = 258, z = -0.86, p = .388$.

Reliability

Table 4 presents the internal consistency of the PACSLAC-D in patients with aphasia and without aphasia. In patients with aphasia, Cronbach's alpha varied between 0.33 (rest) and 0.71 (ADL). In patients without aphasia, from 0.69 (rest) to 0.86 (ADL).

The test-retest reliability during rest was poor; $ICC_{consistency} = 0.07$ (95% CI: -0.40 - 0.51). By contrast, the test-retest reliability during ADL was good; $ICC_{consistency} = 0.88$ (95% CI: 0.71- 0.95) and

it was excellent during physiotherapy; $ICC_{consistency} = 0.95$ (95% CI: 0.83–0.98).

Discussion

The present study investigated the convergent and discriminative validity and reliability of PACSLAC-D in stroke patients with aphasia.

The PACSLAC-D and self-report pain scales showed poor correlations (Table 3). Van der Steen et al. (2021) also reported this finding in a study with patients with dementia who were observed with the pain observation instrument Pain Assessment in Impaired Cognition (PAIC15).

When we compared stroke patients who were unable to self-report pain to those who were able self-report their pain, more pain was observed in patients who were unable to self-report during ADL and physiotherapy. This is in line with research in persons with dementia, in which pain was observed using PACSLAC-D and where patients with pain all tended to be more severely cognitively impaired and had difficulty with self-report scales (Coca & Zuniga, 2020; Natavio et al., 2020; Zwakhalen et al., 2009).

A moderate positive correlation was found between PACSLAC-D and the clinical judgment of the nurse during ADL, and a strong positive correlation was found between PACSLAC-D and NRS/VAS during physiotherapy. Contrary to our expectations and other studies that found associations between the PACSLAC-D and FPS in cognitively impaired participants (Liu et al., 2010; Haghi et al., 2020), in this study the PACSLAC-D showed no to weak correlation with the self-report pain scales. We found only a moderate positive correlation with the judgment of the nurse during ADL, and a strong correlation with the NRS during physiotherapy. These results provide some evidence for the convergent validity of PACSLAC-D in patients with aphasia during activities, but not during rest. This may be explained by the fact that relatively few signs of pain were observed during rest, possibly because of the composition of the sample. The sample consists of patients who had no fractures, injuries or painful disorders.

The discriminating validity of the PACSLAC-D was adequate in patients with aphasia. No difference in pain was observed with the PACSLAC-D during ADL compared with rest. This result is in contrast with previous studies in which less pain is observed during rest compared with during activities (Haghi et al., 2020; Van Herk, van Dijk, Baar, Tibboel, & de Wit, 2007). Second, results were in accordance with the hypothesis that patients with aphasia who used pain medication experienced significantly more pain than patients with aphasia who did not use pain medication during ADL. Not surprisingly, the many 0 scores during rest and physiotherapy mean no significant difference was found in observed pain between both groups. A possible explanation during physiotherapy might be that the movements and exercises are more structured and guided by the physiotherapist, who may try to limit potentially painful movements while still working on therapeutic goals. Consistent with expectation in the third hypothesis, significantly more pain was observed in patients with aphasia compared with non-aphasia patients during ADL. More pain during ADL seems to be consistent with other research which found that aphasic participants score higher on body pain and general health (Cruice et al., 2010). Adequate discriminative validity of the PACSLAC-D in this study population was supported by previous relevant research of pain observation in older people with communication problems (Haghi et al., 2020; Thé et al., 2016).

The reliability of PACSLAC-D in patients with aphasia is particularly good during activities but insufficient during rest. The acceptable internal consistency during ADL and physiotherapy is in

line with studies using the PACSLAC-D in patients with dementia (Liu et al., 2010; Haghi et al., 2020; Natavio et al., 2020). Test-retest reliability was good during ADL and excellent during physiotherapy. This is in line with outcomes of test-retest reliability of PACSLAC-D in elderly with communication problems (Thé et al., 2016; Zwakhalen et al., 2006).

Limitations

Limitations of the current study include the relatively small sample size that was restricted to older stroke patients with aphasia in one geriatric rehabilitation center. This limits the generalizability of results. The order of the self-report pain scales was not randomized, the researcher who observed the patients with aphasia was not blinded and was also their speech and language therapist. Next to these limitations, the current study also has several strengths. This is the first study to examine psychometric properties of a pain observation instrument to measure pain in patients with aphasia in a clinical setting, comparing aphasia with non-aphasia patients, and in several active states.

Conclusions

The PACSLAC-D might be a useful observational instrument and alternative to screen for the presence of pain in stroke patients with aphasia, a population in which pain occurs regularly, pain is triggered by movement, and where pain management may be sub-optimal due to communication difficulties.

Implications for Nursing Education, Practice, and Research

More pain was observed in patients who were unable to self-report during potentially painful activities. This means that pain management in patients with aphasia and other communication difficulties may not be optimal, highlighting the fact that alternatives to screen for pain are essential for these patient groups. The PACSLAC-D might be a suitable alternative to screen for the absence and presence of pain in patients with aphasia who are unable to self-report during activities. The use of a pain observation instrument could help health care professionals to substantiate their opinion on whether pain is present and to evaluate whether pain interventions were successful. Notwithstanding its limitations, this study supports that a pain observation instrument might be a good alternative when self-reporting pain is not possible because of impaired cognition and/or communication problems (Coca & Zuniga, 2020; Haghi et al., 2020; Natavio et al., 2020). However, more research is required on how to measure pain in persons with aphasia in a valid and reliable manner, for example by comparing various observation instruments using larger sample sizes.

Declaration of Competing Interest


The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments


Financial support was provided by Zorgondersteuningsfonds (PROM-6) and TOPAZ.

Appendix

Faces Pain Scale (FPS)



Numeric Rating Scale / Visual Analogue Scale (NRS/VAS)



References

- Aubin, M., Verreault, R., Savoie, M., LeMay, S., Hadjistavropoulos, T., Fillion, L., Beaulieu, M., Viens, C., Bergeron, R., Vézina, L., Misson, L., & Fuchs-Lacelle, S. (2008). Validité et utilité clinique d'une grille d'observation (PACSLAC-F) pour évaluer la douleur chez des aînés atteints de démence vivant en milieu de soins de longue durée [Validity and utilities' clinic of a grid observation (PACSLAC-F) to evaluate the pain in seniors with dementia's living in the Long-Term Care]. *Canadian Journal on Aging. La Revue Canadienne Du Vieillissement*, 27(1), 45–55.
- Bland, J. M., & Altman, D. G. (1997). Cronbach's alpha. *BMJ*, 314(7080), 572.
- Boltz, M., Resnick, B., Kuzmik, A., Mogle, J., Jones, J. R., Arendacs, R., Belue, R., Cacchione, P., & Galvin, J. E. (2021). Pain Incidence, Treatment, and Associated Symptoms in Hospitalized persons with dementia. *Pain Management Nursing*, 22(2), 158–163.
- Büyükturan, Ö., Büyükturan, B., Yetiş, A., Naharcı, M., & Kırdı, N. (2018). Reliability and validity of the Turkish version of Pain Assessment Checklist for Seniors with Limited Ability to Communicate (PACSLAC-T). *Turkish Journal of Medical Sciences*, 48(4), 805–810.
- Carey, D. P., & Johnstone, L. T. (2014). Quantifying cerebral asymmetries for language in dextrals and adextrals with random-effects meta analysis. *Frontiers in Psychology*, 5, 1128.
- Chan, S., Hadjistavropoulos, T., Williams, J., & Lints-Martindale, A. (2014). Evidence-based development and initial validation of the pain assessment checklist for seniors with limited ability to communicate-II (PACSLAC-II). *Clinical Journal of Pain*, 30(9), 816–824.
- Coca, S. M., & Zuniga, R. A. A. (2020). Instruments for pain assessment in patients with advanced dementia: A systematic review of the evidence for Latin America. *Palliative & Supportive Care*, 18(6), 741–747.
- Cruice, M., Worrall, L., & Hickson, L. (2010). Health-related quality of life in people with aphasia: Implications for fluency disorders quality of life research. *Journal of Fluency Disorders*, 35(3), 173–189.
- De Vet, H., Terwee, C., Mookink, L., & Knol, D. (2011). *Measurement in Medicine: A Practical Guide (Practical Guides to Biostatistics and Epidemiology)*. Cambridge University Press.
- De Vries, N. J., Sloot, P. H., & Achterberg, W. P. (2016). Pain and pain assessment in stroke patients with aphasia: A systematic review. *Aphasiology*, 31(6), 703–719.
- Delpont, B., Blanc, C., Osseby, G. V., Hervieu-Begue, M., Giroud, M., & Bejot, Y. (2018). Pain after stroke: A review. *Revue Neurologique (Paris)*, 174(10), 671–674.
- Doesborgh, S. J., van de Sandt-Koenderman, W. M., Dippel, D. W., van Harskamp, F., Koudstaal, P. J., & Visch-Brink, E. G. (2003). Linguistic deficits in the acute phase of stroke. *Journal of Neurology*, 250(8), 977–982.
- El Hachoui, H., Sandt-Koenderman, M. W., Dippel, D. W., Koudstaal, P. J., & Visch-Brink, E. G. (2012). The ScreeLing: Occurrence of linguistic deficits in acute aphasia post-stroke. *Journal of Rehabilitation Medicine*, 44(5), 429–435.
- Engelger, S. (2006). *Aphasia als Hirnschlagsymptom: Häufigkeit und Bedeutung. [Aphasia in stroke patients: frequency and significance]*. Praxis (Bern 1994), 95(13), 489–492.
- Evans, J. D. (1996). *Straightforward statistics for the behavioral sciences*. Brooks/Cole Publishing.
- Fuchs-Lacelle, S., & Hadjistavropoulos, T. (2004). Development and preliminary validation of the Pain Assessment Checklist for Seniors With Limited Ability to Communicate (PACSLAC). *Pain Management Nursing*, 5(1), 37–49.
- Hadjistavropoulos, T., Browne, M. E., Prkachin, K. M., Taati, B., Ashraf, A., & Mihailidis, A. (2018). Pain in severe dementia: A comparison of a fine-grained assessment approach to an observational checklist designed for clinical settings. *European Journal of Pain*, 22(5), 915–925.
- Hadjistavropoulos, T., Herr, K., Prkachin, K. M., Craig, K. D., Gibson, S. J., Lukas, A., & Smith, J. H. (2014). Pain assessment in elderly adults with dementia. *Lancet Neurology*, 13(12), 1216–1227.
- Haghi, M., Fadayevatan, R., Alizadeh-Khoei, M., Kaboudi, B., Foroughan, M., & Mahdavi, B. (2020). Validation of Pain Assessment Checklist for Seniors with Limited Ability to Communicate-II (PACSLAC-II) in Iranian older adults with dementia living in nursing homes. *20(3)*, 278–287.
- Hansen, A. P., Marcussen, N. S., Klit, H., Andersen, G., Finnerup, N. B., & Jensen, T. S. (2012). Pain following stroke: A prospective study. *European Journal of Pain*, 16(8), 1128–1136.
- Harrison, R. A., & Field, T. S. (2015). Post stroke pain: Identification, assessment, and therapy. *Cerebrovasc Dis*, 39(3–4), 190–201.
- Heller, G. Z., Manuguerra, M., & Chow, R. (2016). How to analyze the Visual Analogue Scale: Myths, truths and clinical relevance. *Scandinavian Journal of Pain*, 13, 67–75.
- Hjermstad, M. J., Fayers, P. M., Haugen, D. F., Caraceni, A., Hanks, G. W., Loge, J. H., Fainsinger, R., Aass, N., & Kaasa, S. European Palliative Care Research, C. (2011). Studies comparing Numerical Rating Scales, Verbal Rating Scales, and Visual Analogue Scales for assessment of pain intensity in adults: a systematic literature review. *Journal of Pain Symptom Management*, 41(6), 1073–1093.
- Jones, M. R., Ehrhardt, K. P., Ripoll, J. G., Sharma, B., Padnos, I. W., Kaye, R. J., & Kaye, A. D. (2016). Pain in the Elderly. *Current Pain and Headache Reports*, 20(4), 23.
- Kaasalainen, S., Akhtar-Danesh, N., Hadjistavropoulos, T., Zwakhalen, S., & Verreault, R. (2013). A comparison between behavioral and verbal report pain assessment tools for use with residents in long term care. *Pain Management Nursing*, 14(4), E106–E114.
- Kim, E. J., & Buschmann, M. T. (2006). Reliability and validity of the Faces Pain Scale with older adults. *International Journal of Nursing Studies*, 43(4), 447–456.
- Kim, E. K., Kim, S. Y., Eom, M. R., Kim, H. S., & Lee, E. (2014). [Validity and reliability of the Korean version of the pain assessment checklist for seniors with

- limited ability to communicate]. *Journal of the Korean Academy of Nursing*, 44(4), 398–406.
- Koo, T. K., & Li, M. Y. (2016). A guideline of selecting and reporting intraclass correlation coefficients for reliability research. *Journal of Chiropractic Medicine*, 15(2), 155–163.
- Kunz, M., de Waal, M. W. M., Achterberg, W. P., Gimenez-Llort, L., Lobbezoo, F., Sampson, E. L., van Dalen-Kok, A. H., Defrin, R., Invitto, S., Konstantinovic, L., Oosterman, J., Petrini, L., van der Steen, J. T., Strand, L. I., de Tommaso, M., Zwakhalen, S., Husebo, B. S., & Lautenbacher, S. (2020). The Pain Assessment in Impaired Cognition scale (PAIC15): A multidisciplinary and international approach to develop and test a meta-tool for pain assessment in impaired cognition, especially dementia. *European Journal of Pain*, 24(1), 192–208.
- Liampas, A., Velidakis, N., Georgiou, T., Vadalouca, A., Varrassi, G., Hadjigeorgiou, G. M., Tsigoulis, G., & Zis, P. (2020). Prevalence and management challenges in central post-stroke neuropathic pain: A systematic review and meta-analysis. *Advances in Therapy*, 37(7), 3278–3291.
- Lints-Martindale, A. C., Hadjistavropoulos, T., Lix, L. M., & Thorpe, L. (2012). A comparative investigation of observational pain assessment tools for older adults with dementia. *Clinical Journal of Pain*, 28(3), 226–237.
- Liu, J. Y. W., Briggs, M., & Closs, S. J. (2010). The psychometric qualities of four observational pain tools (opts) for the assessment of pain in elderly people with osteoarthritic pain. *Journal of Pain and Symptom Management*, 40(4), 582–598.
- Mitchell, C., Gittins, M., Tyson, S., Vail, A., Conroy, P., Paley, L., & Bowen, A. (2020). Prevalence of aphasia and dysarthria among inpatient stroke survivors: describing the population, therapy provision and outcomes on discharge. *Aphasiology*, 35(7), 950–960.
- Natavio, T., McQuillen, E., Dietrich, M. S., Wells, N., Rhoten, B. A., Vallerand, A. H., & Monroe, T. B. (2020). A comparison of the Pain Assessment Checklist for Seniors with Limited Ability to Communicate (PACSLAC) and Pain Assessment in Advanced Dementia Scale (PAINAD). *Pain Management Nursing*, 21(6), 502–509.
- Qi, S., Diane, J., & Kay, D. (2012). The psychometric properties, feasibility and utility of behavioural observation methods in pain assessment of cognitively impaired elderly people in acute and long-term care: A systematic review. *JBI Library of Systematic Reviews*, 10(17), 977–1085.
- Rajkumar, A. P., Ballard, C., Fossey, J., Orrell, M., Moniz-Cook, E., Woods, R. T., Murray, J., Whitaker, R., Stafford, J., Knapp, M., Romeo, R., Woodward-Carlton, B., Khan, Z., Testad, I., & Corbett, A. (2017). Epidemiology of pain in people with dementia living in care homes: Longitudinal course, prevalence, and treatment implications. *Journal of the American Medical Directors Association*, 18(5), 453.e451–453.e456.
- Ruest, M., Bourque, M., Laroche, S., Harvey, M. P., Martel, M., Bergeron-Vézina, K., Apinis, C., Proulx, D., Hadjistavropoulos, T., Tousignant-Laflamme, Y., & Léonard, G. (2017). Can we quickly and thoroughly assess pain with the PACSLAC-II? A convergent validity study in long-term care residents suffering from dementia. *Pain Management Nursing*, 18(6), 410–417.
- Schuster, J., Hoyer, C., Ebert, A., & Alonso, A. (2020). Use of analgesics in acute stroke patients with inability to self-report pain: a retrospective cohort study. *BMC Neurology*, 20(1), 18.
- Smith, J. H., Bottemiller, K. L., Flemming, K. D., Cutrer, F. M., & Strand, E. A. (2013). Inability to self-report pain after a stroke: A population-based study. *Pain*, 154(8), 1281–1286.
- Takai, Y., Yamamoto-Mitani, N., Suzuki, M., Furuta, Y., Sato, A., & Fujimaki, Y. (2013). Developing and validating a Japanese version of the Assessment of Pain in Elderly People with Communication Impairment. *Archives of Gerontology and Geriatrics*, 57(3), 403–410.
- Thé, K. B., Gazoni, F. M., Cherpak, G. L., Lorenzet, I. C., Santos, L. A., Nardes, E. M., & Santos, F. C. (2016). Pain assessment in elderly with dementia: Brazilian validation of the PACSLAC scale. *Einstein (São Paulo)*, 14(2), 152–157.
- Van Dalen-Kok, A. H., Pieper, M. J. C., De Waal, M. W. M., Van der Steen, J. T., Scherder, E. J. A., & Achterberg, W. P. (2021). The impact of pain on the course of ADL functioning in patients with dementia. *Age and Ageing*, 50(3), 906–913.
- Van der Steen, J. T., Westzaan, A., Hanemaayer, K., Muhamad, M., de Waal, M. W. M., & Achterberg, W. P. (2021). Probable Pain on the Pain Assessment in Impaired Cognition (PAIC15) Instrument: Assessing sensitivity and specificity of cut-offs against three standards. *Brain Sciences*, 11(7), 869.
- Van Herk, R., van Dijk, M., Baar, F. P., Tibboel, D., & de Wit, R. (2007). Observation scales for pain assessment in older adults with cognitive impairments or communication difficulties. *Nursing Research*, 56(1), 34–43.
- Van Nispen tot Pannerden, S. C., Candel, M. J., Zwakhalen, S. M., Hamers, J. P., Curfs, L. M., & Berger, M. P. (2009). An item response theory-based assessment of the pain assessment checklist for Seniors with Limited Ability to Communicate (PACSLAC). *Journal of Pain*, 10(8), 844–853.
- Vingerhoets, G. (2019). Phenotypes in hemispheric functional segregation? Perspectives and challenges. *Phys Life Rev*, 30, 1–18. <https://doi.org/10.1016/j.plrev.2019.06.002>.
- Westerlind, E., Singh, R., Persson, H. C., & Sunnerhagen, K. S. (2020). Experienced pain after stroke: a cross-sectional 5-year follow-up study. *BMC Neurology*, 20(1). <https://doi.org/ARTN 410.1186/s12883-019-1584-z>
- Wu, C., Qin, Y. Y., Lin, Z., Yi, X. Y., Wei, X., Ruan, Y. M., & He, J. (2020). Prevalence and Impact of Aphasia among Patients Admitted with Acute Ischemic Stroke. *Journal of Stroke and Cerebrovascular Diseases*, 29(5). <https://doi.org/ARTN 10476410.1016/j.jstrokecerebrovasdis.2020.104764>
- Zwakhalen, S. M., Hamers, J. P., & Berger, M. P. (2006). The psychometric quality and clinical usefulness of three pain assessment tools for elderly people with dementia. *Pain*, 126(1-3), 210–220. <https://doi.org/10.1016/j.pain.2006.06.029>.
- Zwakhalen, S. M., Hamers, J. P., & Berger, M. P. (2007). Improving the clinical usefulness of a behavioural pain scale for older people with dementia. *Journal of Advanced Nursing*, 58(5), 493–502. <https://doi.org/10.1111/j.1365-2648.2007.04255.x>.
- Zwakhalen, S. M., Koopmans, R. T., Geels, P. J., Berger, M. P., & Hamers, J. P. (2009). The prevalence of pain in nursing home residents with dementia measured using an observational pain scale. *Eur J Pain*, 13(1), 89–93.