

University of Groningen

## CEERS Key Paper. VIII. Emission-line Ratios from NIRSpec and NIRCам Wide-Field Slitless Spectroscopy at $z > 2$

Backhaus, Bren E.; Trump, Jonathan R.; Pirzkal, Nor; Barro, Guillermo; Finkelstein, Steven L.; Haro, Pablo Arrabal; Simons, Raymond C.; Wessner, Jessica; Cleri, Nikko J.; Bagley, Micaela B.

*Published in:*  
Astrophysical Journal

*DOI:*  
[10.3847/1538-4357/ad1520](https://doi.org/10.3847/1538-4357/ad1520)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2024

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Backhaus, B. E., Trump, J. R., Pirzkal, N., Barro, G., Finkelstein, S. L., Haro, P. A., Simons, R. C., Wessner, J., Cleri, N. J., Bagley, M. B., Hirschmann, M., Nicholls, D. C., Dickinson, M., Kartaltepe, J. S., Papovich, C., Kocevski, D. D., Koekemoer, A. M., Bisigello, L., Jaskot, A. E., ... Davis, K. (2024). CEERS Key Paper. VIII. Emission-line Ratios from NIRSpec and NIRCам Wide-Field Slitless Spectroscopy at  $z > 2$ . *Astrophysical Journal*, 962(2), Article 195. <https://doi.org/10.3847/1538-4357/ad1520>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# CEERS Key Paper. VIII. Emission-line Ratios from NIRSpec and NIRCам Wide-Field Slitless Spectroscopy at $z > 2$

Bren E. Backhaus<sup>1</sup> , Jonathan R. Trump<sup>1</sup> , Nor Pirzkal<sup>2</sup> , Guillermo Barro<sup>3</sup> , Steven L. Finkelstein<sup>4</sup> , Pablo Arrabal Haro<sup>5</sup> , Raymond C. Simons<sup>1</sup> , Jessica Wessner<sup>1</sup> , Nikko J. Cleri<sup>6,7</sup> , Micaela B. Bagley<sup>4</sup> , Michaela Hirschmann<sup>8</sup> , David C. Nicholls<sup>9</sup> , Mark Dickinson<sup>5</sup> , Jeyhan S. Kartaltepe<sup>10</sup> , Casey Papovich<sup>6,7</sup> , Dale D. Kocevski<sup>11</sup> , Anton M. Koekemoer<sup>12</sup> , Laura Bisigello<sup>13,14</sup> , Anne E. Jaskot<sup>15</sup> , Ray A. Lucas<sup>12</sup> , Intae Jung<sup>12</sup> , Stephen M. Wilkins<sup>16,17</sup> , L. Y. Aaron Yung<sup>18,28</sup> , Henry C. Ferguson<sup>19</sup> , Adriano Fontana<sup>20</sup> , Andrea Grazian<sup>14</sup> , Norman A. Grogin<sup>19</sup> , Lisa J. Kewley<sup>21</sup> , Allison Kirkpatrick<sup>22</sup> , Jennifer M. Lotz<sup>23</sup> , Laura Pentericci<sup>20</sup> , Pablo G. Pérez-González<sup>24</sup> , Swara Ravindranath<sup>12</sup> , Rachel S. Somerville<sup>25</sup> , Guang Yang<sup>26</sup> , Benne W. Holwerda<sup>27</sup> , Peter Kurczynski<sup>18</sup> , Nimish P. Hathi<sup>19</sup> , Caitlin Rose<sup>10</sup> , and Kelcey Davis<sup>1</sup>

<sup>1</sup> Department of Physics, 196A Auditorium Road, Unit 3046, University of Connecticut, Storrs, CT 06269, USA

<sup>2</sup> ESA/AURA Space Telescope Science Institute, 3700 San Martin Dr. Baltimore, MD 21218, USA

<sup>3</sup> Department of Physics, University of the Pacific, Stockton, CA 90340 USA

<sup>4</sup> Department of Astronomy, The University of Texas at Austin, Austin, TX, USA

<sup>5</sup> NSF's National Optical-Infrared Astronomy Research Laboratory, 950 N. Cherry Ave., Tucson, AZ 85719, USA

<sup>6</sup> Department of Physics and Astronomy, Texas A&M University, College Station, TX 77843-4242 USA

<sup>7</sup> George P. and Cynthia Woods Mitchell Institute for Fundamental Physics and Astronomy, Texas A&M University, College Station, TX 77843-4242 USA

<sup>8</sup> Institute of Physics, Laboratory of Galaxy Evolution, Ecole Polytechnique Fédérale de Lausanne (EPFL), Observatoire de Sauverny, 1290 Versoix, Switzerland

<sup>9</sup> Research School of Astronomy and Astrophysics, Australian National University, Canberra, ACT 2600, Australia

<sup>10</sup> Laboratory for Multiwavelength Astrophysics, School of Physics and Astronomy, Rochester Institute of Technology, 84 Lomb Memorial Drive, Rochester, NY 14623, USA

<sup>11</sup> Department of Physics and Astronomy, Colby College, Waterville, ME 04901, USA

<sup>12</sup> Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218, USA

<sup>13</sup> Dipartimento di Fisica e Astronomia "G. Galilei," Università di Padova, Via Marzolo 8, I-35131 Padova, Italy

<sup>14</sup> INAF—Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, I-35122, Padova, Italy

<sup>15</sup> Department of Astronomy, Williams College, Williamstown, MA 01267, USA

<sup>16</sup> Astronomy Centre, University of Sussex, Falmer, Brighton BN1 9QH, UK

<sup>17</sup> Institute of Space Sciences and Astronomy, University of Malta, Msida MSD 2080, Malta

<sup>18</sup> Astrophysics Science Division, NASA Goddard Space Flight Center, 8800 Greenbelt Rd, Greenbelt, MD 20771, USA

<sup>19</sup> Space Telescope Science Institute, Baltimore, MD, USA

<sup>20</sup> INAF—Osservatorio Astronomico di Roma, via di Frascati 33, I-00078 Monte Porzio Catone, Italy

<sup>21</sup> Center for Astrophysics | Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138, USA

<sup>22</sup> Department of Physics and Astronomy, University of Kansas, Lawrence, KS 66045, USA

<sup>23</sup> Gemini Observatory/NSF's National Optical-Infrared Astronomy Research Laboratory, 950 N. Cherry Ave., Tucson, AZ 85719, USA

<sup>24</sup> Centro de Astrobiología (CAB), CSIC-INTA, Ctra. de Ajalvir km 4, Torrejón de Ardoz, E-28850, Madrid, Spain

<sup>25</sup> Center for Computational Astrophysics, Flatiron Institute, 162 5th Avenue, New York, NY 10010, USA

<sup>26</sup> Kapteyn Astronomical Institute, University of Groningen, P.O. Box 800, 9700 AV Groningen, The Netherlands

<sup>27</sup> Physics & Astronomy Department, University of Louisville, KY 40292, Louisville, USA

Received 2023 July 17; revised 2023 December 8; accepted 2023 December 10; published 2024 February 20

## Abstract

We use James Webb Space Telescope Near-Infrared Camera Wide Field Slitless Spectroscopy (NIRCам WFSS) and the Near-Infrared spectrograph (NIRSpec) in the Cosmic Evolution Early Release survey to measure rest-frame optical emission-line ratios of 155 galaxies at  $z > 2$ . The blind NIRCам grism observations include a sample of galaxies with bright emission lines that were not observed on the NIRSpec masks. We study the changes of the  $H\alpha$ ,  $[O III]/H\beta$ , and  $[Ne III]/[O II]$  emission lines in terms of redshift by comparing to lower-redshift SDSS, CLEAR, and MOSDEF samples. We find a significant ( $>3\sigma$ ) correlation between  $[O III]/H\beta$  with redshift, while  $[Ne III]/[O II]$  has a marginal ( $2\sigma$ ) correlation with redshift. We compare  $[O III]/H\beta$  and  $[Ne III]/[O II]$  to stellar mass and  $H\beta$  SFR. We find that both emission-line ratios have a correlation with  $H\beta$  SFR and an anticorrelation with stellar mass across the redshifts  $0 < z < 9$ . Comparison with MAPPINGS V models indicates that these trends are consistent with lower metallicity and higher ionization in low-mass and high-SFR galaxies. We additionally compare to IllustrisTNG predictions and find that they effectively describe the highest  $[O III]/H\beta$  ratios observed in our sample, without the need to invoke MAPPINGS models with significant shock ionization components.

*Unified Astronomy Thesaurus concepts:* [Emission line galaxies \(459\)](#); [Galaxy evolution \(594\)](#); [Galaxies \(573\)](#); [Active galaxies \(17\)](#)

*Supporting material:* machine-readable table

<sup>28</sup> NASA Postdoctoral Fellow.

## 1. Introduction

Galaxy emission lines provide a wealth of information about galaxy formation and physical properties. Emission lines can be used to determine a galaxy's interstellar medium (ISM)

conditions such as the metallicity, ionization, and density, as well as the physical properties such as star formation rate (Brinchmann et al. 2004; Kennicutt & Evans 2012) and dust attenuation (Cardelli et al. 1989; Calzetti & Heckman 1999; Reddy et al. 2016; Shapley et al. 2023b). One way to analyze emission lines is by comparing ratios of lines at similar wavelengths to gain information on galaxy ISM conditions. Picking emission-line ratios with similar wavelength makes the ratio less sensitive to dust attenuation. The most well-known emission-line ratio diagrams, BPT (Baldwin et al. 1981) and VO87 (Veilleux & Osterbrock 1987), compare  $[\text{O III}]\lambda 5007/\text{H}\beta$  with  $[\text{N II}]\lambda 6583/\text{H}\alpha$  or  $[\text{S II}]\lambda 6583/\text{H}\alpha$  to identify high-ionization galaxies. These diagrams make use of the strongest emission lines in rest-frame optical spectra (Baldwin et al. 1981; Veilleux & Osterbrock 1987; Kauffmann et al. 2003a; Kewley et al. 2006) to identify the dominant ionizing sources in galaxies.

Studying emission lines at different redshifts enables an understanding of how the physical conditions of galaxies change over cosmic time. At higher redshifts, the strong rest-optical lines move to the near-IR. The cosmic star formation rate density is much higher at  $z \sim 2$  than in the local Universe (Madau & Dickinson 2014). Studies have found that galaxies at  $z \sim 2$  have lower metallicities (Henry et al. 2013; Steidel et al. 2014; Papovich et al. 2022), have higher ionization in the ISM, and exhibit greater  $\alpha$ -enhancement (Steidel et al. 2016; Shapley et al. 2019) than the local Universe (Liu et al. 2008; Shapley et al. 2015; Strom et al. 2018), which is consistent with cosmic noon galaxies having higher star formation and more active galactic nuclei (AGN). Due to the higher ionization, lower metallicity, and  $\alpha$ -element enhancement of star-forming galaxies at  $z \sim 2$ , the BPT and VO87 diagrams are not effective at distinguishing star-forming galaxies from AGN (Coil et al. 2015; Backhaus et al. 2022; Cleri et al. 2023a). At redshift  $z \sim 2$ , new emission-line diagrams were established such as the OHNO diagram, comparing  $[\text{Ne III}]\lambda 3869/[\text{O II}]\lambda 3726 + 3729$  to  $[\text{O III}]/\text{H}\beta$ , to find high-ionization sources instead (Zeimann et al. 2015; Backhaus et al. 2022).

JWST now gives us access to the early Universe, where the first galaxies and black holes are being formed (e.g., Álvarez-Márquez et al. 2023; Finkelstein et al. 2023; Kokorev et al. 2023; Larson et al. 2023). This gives us the opportunity to study and understand the ISM conditions of galaxies at this early period of the Universe. JWST observations will show how these galaxies assembled and evolved into the Universe we see today. JWST near-IR spectroscopy allows us to view the rest-frame optical emission lines of galaxies at  $2 < z < 9$ .

Previous work has been done with NIRSpec spectroscopy from the JWST Early Release Observations (ERO; Pontoppidan et al. 2022) of the lensing cluster SMACS J0723.3-7327 to investigate emission lines of  $z > 5$  galaxies, but that work used a small sample of 3–6 galaxies. Schaerer et al. (2022), Trump et al. (2023), and Curti et al. (2023) studied the metallicities of these galaxies. Trump et al. (2023), Cleri et al. (2023b), and Brinchmann (2023) went further and compared emission lines at  $z > 5$  to local samples, finding higher  $([\text{Ne III}]/[\text{O II}])$  in the early galaxies compared to the local ( $z \sim 0$ ) galaxies indicating higher ionization. Additionally, analysis has been done with the Cosmic Evolution Early Release Science (CEERS) survey program. The NIRSpec spectroscopy has been used to gain a wealth of information, such as spectroscopically confirmed high-redshift galaxies (Arrabal Haro et al. 2023;

Curtis-Lake et al. 2023; Fujimoto et al. 2023), identifying and characterizing high-redshift AGN (Kocevski et al. 2023; Larson et al. 2023), studying Ly $\alpha$  emission (Jung et al. 2023; Tang et al. 2023), and studying ISM conditions as a function of galaxy properties such as stellar mass and star formation rate (Shapley et al. 2023a, 2023b). Shapley et al. (2023b) found there was not significant evolution between stellar mass and  $[\text{O III}]/\text{H}\beta$  for galaxies above  $z > 3$ , but their stellar masses alone suggest subsolar metallicity. Shapley et al. (2023b) also showed that the  $z > 5$  galaxies prefer higher  $\text{H}\alpha$  SFR when compared to the predictions of Speagle et al. (2014). The ISM conditions of galaxies at  $z > 5$  were shown to have high ionization and low metallicity ( $> 0.1 Z_{\odot}$ ) (Sanders et al. 2023). Sanders et al. (2024) make use of  $T_e$ -based metallicity to calibrate strong-line metallicity estimators for  $z > 5$  galaxies.

In this work, we use NIRCcam Wide Field Slitless Spectroscopy (WFSS) and NIRSpec Multi-Shutter Assembly (MSA) spectroscopy taken as part of CEERS to investigate the rest-frame optical emission-line evolution and galaxy properties of  $\sim 155$  galaxies at  $z > 2$ . In Section 2, we describe our data reductions and sample selection. Section 2 also establishes our comparison samples of galaxies at  $z \sim 0$ ,  $z \sim 2$ , and  $z \sim 3$ . In Section 3, we compare galaxies in our NIRSpec and NIRCcam WFSS samples. In Section 4, we use our three subsamples covering the epoch of reionization ( $z > 6$ ), cosmic noon ( $z \sim 2$ ), and the local Universe ( $z \sim 0$ ) to study how each emission-line ratio evolves with redshift. Section 5 describes the connections between galaxy properties and emission-line ratios at different redshifts. Section 6 presents the ISM conditions inferred by the emission-line ratios. We summarize our results in Section 7. In this work, we assume a  $\Lambda$  cold dark matter cosmology with  $\Omega_M = 0.3$ ,  $\Omega_{\Lambda} = 0.7$ , and  $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$  (Planck Collaboration et al. 2016).

## 2. Observational Data and Sample

### 2.1. JWST WFSS and MSA Spectroscopy

Our parent galaxy sample comes from JWST observations taken by the CEERS program, ERS-1345 (PI: Steven Finkelstein). CEERS uses NIRCcam WFSS and NIRSpec multi-object spectroscopy to cover  $\sim 100 \text{ arcmin}^2$  of the Extended Groth Strip Hubble Space Telescope (HST) legacy field (EGS; Davis et al. 2007), which is covered by the CANDELS HST survey (Grogin et al. 2011; Koekemoer et al. 2011). Our paper focuses on the four pointings with NIRCcam WFSS and six pointings with NIRSpec. CEERS has four NIRCcam WFSS pointings that partially overlap with the CEERS NIRSpec observations, allowing the emission-line measurements of both instruments to be compared, which in turn allows NIRSpec to be calibrated for slit losses (see Section 3). However, in practice, we can only perform this test for one galaxy.

The four NIRCcam WFSS pointings use the F356W filter to cover  $3.14\text{--}3.98 \mu\text{m}$ , including a suite of rest-frame optical lines for  $0 < z < 9$  galaxies with a total exposure time of 2490 s split between the two orthogonal gratings in the WFSS. The WFSS spectra have spectroscopic resolving power  $R \sim 1600$  at  $\sim 4 \mu\text{m}$ . The details of the NIRCcam WFSS are described by Greene et al. (2017).

The six NIRSpec spectroscopy pointings use the G140M/F100LP, G235M/F170LP, and G395M/F290LP filters, spanning  $1\text{--}5 \mu\text{m}$ . Four of these pointings were observed in medium

resolution,  $R \sim 1000$ , and in prism  $R \sim 100$ . Two pointings had light-leak failures in the prism observations and we use only the medium-resolution grating observations (the failed prism observations were rescheduled and observed later but are not used in this work). The details of the NIRSpec instrument are described by Jakobsen et al. (2022). Further information on the programs used in the CEERS extractions and fits can be read in Bagley et al. (2023) and Arrabal Haro et al. (2023).

### 2.1.1. Data Reduction and Sample Selection

Information on the reductions of the NIRSpec data can be found in Arrabal Haro et al. (2023, in preparation). The reduction’s main steps are as follows. The NIRSpec data is processed with the STScI JWST Calibration Pipeline v1.8.5<sup>29</sup> (Bushouse et al. 2022) and the Calibration Reference Data System (CRDS). The `calwebb_detector1` pipeline module was used to reduce the uncalibrated images by applying the correction for “snowballs” events caused by cosmic rays, the  $1/f$  noise correction, and doing a saturation check. An improved correction for the “snowball” events<sup>30</sup> is applied to the `jump` step. This step also creates the count-rate maps (CRMs).

The generated CRMs were then passed through the `calwebb_spec2` pipeline to create two-dimensional (2D) cutouts of the slitlets. The `calwebb_spec2` pipeline performs the background subtraction by using a three-nod pattern, corrects the flat fields, implements the wavelength and photometric calibrations, and resamples the 2D spectra to correct the distortion of the spectral trace.

The `calwebb_spec3` pipeline stage creates the final 2D spectra, by combining the images of the three nods. The standard pipeline `pathloss` correction is applied to correct the spectrum for source flux outside the microshutter “slit.” The 1D spectra are created by extracting from the 2D spectra using customized apertures. These custom apertures are visually defined for targets to maximize the signal-to-noise ratios (S/Ns).

The NIRSpec flux uncertainties are underestimated by a factor of  $\sim 2$ , determined by comparing the normalized median absolute deviation (NMAD) of the flux to the median of the flux uncertainty for each source. We correct for this by increasing the flux uncertainty of each spectrum by the ratio of the NMAD of the flux to the median flux uncertainty,  $\text{NMAD}(f)/\text{median}(\sigma_f)$ .

The NIRCAM WFSS data were first processed with the Stage 1 STScI pipeline to apply bias and dark corrections and perform on-the-ramp fitting to detect and remove cosmic ray impacts. A broad-filter F356W flat-field correction was applied to the data, as it was shown to reduce pixel-to-pixel variation. Finally, proper world coordinate information was added using the `AssignWCS()` Pipeline task. Spectra were extracted from these processed data sets following the Simulation Based Extraction (Pirzkal et al. 2017). We used an extraction catalog derived from the CEERS F356W mosaic. Spectral contamination was modeled and subtracted using pixel-level SEDs for each source computed from the CEERS F277W, F356W, and F444W mosaics. Finally, the dispersed background was subtracted as a combination of pre-launch synthetic dispersed F356W background and of an additional row (column) model

of the residuals in the GRISMR (GRISM C) observations. This results in the modeling and extractions of 24,000 spectra. Continuum is detected in sources down to approximately  $\sim 25.75$  AB magnitude.

The NIRCAM WFSS flux uncertainties are underestimated by a factor of  $\sim 3$  in the row dispersion and  $\sim 4$  in the column dispersion. This was determined and corrected using the same method with the ratio of NMAD of the flux to the median of the uncertainty used for the NIRSpec sample.

Our photometric redshifts used for sample selection are obtained using the same method outlined in Finkelstein et al. (2022). Briefly, the multi-band SEDs were fit using the EAZY (Brammer et al. 2008) software package. Probability distribution functions (PDFs) are created by fitting non-negative linear combinations of templates to the observed data. The templates used are a set of 12 FSPS “*tweak\_fsp\_QSF\_12\_v3*” (Conroy & Gunn 2010), and six additional templates to cover bluer colors. These bluer color templates are shown by Larson et al. (2023) to improve the photometric redshift fits for  $z > 9$  galaxies. These templates use stellar population models created with BPASS Eldridge & Stanway (2009) with 5% solar metallicities and young stellar populations,  $\log(\text{age}/\text{yr}) = 6, 6.5, 7$ .

The JWST data used in this paper can be found in MAST:10.17909/xw7n-9730.

### 2.1.2. Sample Selection

To extract the emission-line flux from both NIRSpec and NIRCAM WFSS, we first use photometric redshifts to constrain our sample to find galaxies of interest. For NIRCAM WFSS, our constraints were determined by which redshifts would have our desired emission lines land in the F356W filter. This would give redshift ranges of  $3.7 < z < 5.1$  for  $\text{H}\alpha$ , a  $5.4 < z < 7$  range for [O III], and  $7.4 < z < 9.3$  for [O II]. We increase the range of each redshift bin to account for uncertainties in the photometric redshifts, of  $3.2 < z < 5.5$  for  $\text{H}\alpha$ ,  $4.9 < z < 7.5$  for [O III], and  $6.9 < z < 9.8$  for [O II]. The photometric redshift overlap between [O III] and [O II] is due to adding  $\Delta z = 0.5$  to select galaxies. Meanwhile, for NIRSpec, we use the photometric redshift constraint of  $1.6 < z < 9$  as this covers all emission lines spanning the G140M, G235M, and G395M gratings. We determined the spectroscopic redshift for NIRCAM and NIRSpec for each source using the best-fit line center of the brightest emission line in each spectrum, usually [O III] $\lambda 5008$  or  $\text{H}\alpha$ , as they are the brightest lines in the spectra. To measure emission-line fluxes, we find the best-fit Gaussian function (and associated uncertainties) using a Levenberg–Marquardt least-squares method implemented by the `SciPy curve fit` Python code (Virtanen et al. 2020).


Our NIRCAM WFSS sample consists of 19 galaxies with [O III]/ $\text{H}\beta$  measurements and 18 galaxies with  $\text{H}\alpha$ . There are no NIRCAM WFSS [Ne III]/[O II] detections, as both emission lines were lost in the noise and had  $\text{S/N} < 1$ .

Our NIRSpec sample has a total of 118 galaxies, with 93 galaxies in the  $\text{H}\alpha$  sample, 96 galaxies in the [O III] sample, and 59 in the [O II] sample. These galaxies follow the same sample selection process as NIRCAM WFSS, requiring an  $\text{S/N} > 3$  for at least one emission line in each ratio. However, we add an additional  $\text{S/N} > 1$  requirement for the other emission line. The last sample includes [Ne III] $\lambda 3869$  and the blended [O II] $\lambda 3726 + 3729$  doublet, which is accessible up to  $z \sim 8.5$ . Figure 1 shows examples of NIRSpec 1D and 2D spectra in the bottom right panels.

<sup>29</sup> <https://jwst-pipeline.readthedocs.io/en/latest/index.html>

<sup>30</sup> <https://jwst-docs.stsci.edu/data-artifacts-and-features/snowballs-and-shower-artifacts>





**Figure 1.** Example of NIRCcam WFSS direct image (left) with 1D/2D column and row WFSS spectra for an example galaxy, 016\_B\_13569, that is in our [O III]/H $\beta$  sample. Vertical lines in the 1D spectra indicate the emission lines of interest. We also include the direct image and 1D/2D NIRSpect spectra of p5\_717 from the G395M filter (bottom).

We do not dust correct the emission lines of our samples because both H $\alpha$  and H $\beta$  are not available for all galaxies in the sample. Dust correction for CEERS NIRSpect galaxies are typically small, with a median  $E(B - V)_{\text{neb}} = 0.18$  Shapley et al. (2023b). The emission-line ratios used in this work are not affected by dust attenuation, due to the fact that the emission-line pairs are close in wavelength and are nearly equally affected by dust. We note that 62% of our [O III]/H $\beta$  galaxies between NIRSpect MSA and NIRCcam WFSS at  $z > 4$  have an SED  $A_V < 0.2$ , so we expect the H $\beta$  SFR estimates to be minimally effected by dust. Meanwhile, 48% of our H $\alpha$  galaxies at  $z > 4$  have an SED  $A_V < 0.2$ , and only three galaxies in the NIRCcam WFSS H $\alpha$  sample have SED  $A_V < 0.2$ . Galaxies affected by dust will have a lower limit for H $\beta$  or H $\alpha$  SFR.

SFR is calculated from either the H $\beta$  or H $\alpha$  emission line, depending on redshift, by following the Kennicutt & Evans (2012) SFR relation for H $\alpha$  and H $\alpha$ /H $\beta$  = 2.86 (assuming Case B recombination,  $T = 10^4$  K, and  $n_e = 10^4$  cm $^{-3}$ ; Osterbrock 1989):

$$\log(\text{SFR})[M_{\odot} \text{ yr}^{-1}] = \log[L(\text{H}\beta)] - 40.82 \quad (1)$$

$$\log(\text{SFR})[M_{\odot} \text{ yr}^{-1}] = \log[L(\text{H}\alpha)] - 41.27. \quad (2)$$

Due to the H $\beta$  and H $\alpha$  lines not being dust corrected, these SFR are a lower limit.

We create two WFSS subsamples of galaxies based on a signal-to-noise ratio of  $S/N > 3$  detection for at least one line of interest when the two orientations are coadded. The first includes galaxies with  $S/N > 3$  in H $\alpha$  at  $z \sim 4$ . The second includes galaxies with  $S/N > 3$  [O III] $\lambda$ 5007 emission lines at

$z \sim 6$ . We note there are no galaxies with [Ne III] $\lambda$ 3868 or [O II] $\lambda$ 3728 detected at  $S/N > 3$  in the CEERS NIRCcam WFSS observations, due to the low observation depth. We visually inspect the filter image, 1D, and 2D spectra of galaxies selected for these samples to ensure the emission line is detected in both orientations. An example of one galaxy inspected and included in the samples is shown in Figure 1.

The sample selection and inspection results in 18 galaxies with H $\alpha$  in the redshift range  $4 < z < 5$  and 19 galaxies with [O III] in the redshift range  $5.5 < z < 7$  from the NIRCcam WFSS.

Tables 1 and 2 present the source IDs, spectroscopic redshifts, emission-line flux, stellar mass, SED SFR, and dust attenuation measurements of the NIRCcam WFSS [O III] and H $\alpha$  galaxy samples. Our NIRSpect sample is shown in Table 3 showing the ID, redshift, and emission-line fluxes. Figures 2 and 3 show the redshift distribution, stellar mass, and SFR of both samples.

### 2.1.3. Stellar Mass

Stellar masses and dust attenuations in our NIRCcam WFSS and NIRSpect sample come from fitting the optical and NIR SEDs from the long-wavelength NIRCcam filters using FAST (Kriek et al. 2009). These models assume stellar population synthesis models used by Bruzual & Charlot (2003), following the initial mass function (IMF) defined by Chabrier (2003), a Calzetti et al. (2000) dust attenuation law, and a delayed exponential star formation history.

**Table 1**  
[O III]/H $\beta$  NIRCcam Sample

ID	R.A. (deg)	Decl. (deg)	z	[O III] Column ( $10^{-18}$ erg/s/cm $^2$ )	[O III] Row ( $10^{-18}$ erg/s/cm $^2$ )	H $\beta$ Column ( $10^{-18}$ erg/s/cm $^2$ )	H $\beta$ Row ( $10^{-18}$ erg/s/cm $^2$ )	log( $M_*$ ) [ $M_\odot$ ]	log(SFR) [ $M_\odot$ yr $^{-1}$ ]	$A_V$ (mag)
012_B_22258	214.929820	52.862205	5.635	49.37 $\pm$ 6.33	41.37 $\pm$ 5.92	6.00 $\pm$ 4.91	5.07 $\pm$ 4.91	10.05	2.16	0.8
013_A_8436	214.958411	52.843681	5.999	17.26 $\pm$ 2.02	17.06 $\pm$ 2.03	3.65 $\pm$ 2.07	2.33 $\pm$ 1.87	9.37	0.45	0.0
015_B_13732	214.987915	52.879438	6.177	44.09 $\pm$ 4.89	48.33 $\pm$ 3.98	6.03 $\pm$ 5.04	7.84 $\pm$ 3.30	9.95	1.18	0.0
015_B_19309	214.970312	52.881717	5.640	8.63 $\pm$ 3.05	4.14 $\pm$ 2.18	5.47 $\pm$ 3.73	2.80 $\pm$ 2.17	8.87	0.22	0.0
012_B_30258	214.918284	52.879359	5.645	10.95 $\pm$ 3.65	14.16 $\pm$ 3.47	2.85 $\pm$ 5.02	1.21 $\pm$ 3.91	9.20	0.65	0.0
013_A_15609	214.948684	52.856466	6.530	3.47 $\pm$ 1.29	7.49 $\pm$ 2.71	0.01 $\pm$ 1.19	1.52 $\pm$ 1.88	9.12	0.18	0.0
015_B_19805	214.958370	52.875115	6.170	34.25 $\pm$ 12.26	33.86 $\pm$ 5.80	2.43 $\pm$ 10.46	4.76 $\pm$ 6.41	9.37	0.19	0.1
016_A_13158	215.122563	52.973201	6.380	42.02 $\pm$ 5.94	49.80 $\pm$ 5.33	2.27 $\pm$ 4.11	4.32 $\pm$ 5.29	9.89	1.31	0.0
016_A_16127	215.127984	52.984951	6.664	37.96 $\pm$ 5.68	33.45 $\pm$ 4.37	0.77 $\pm$ 4.58	1.70 $\pm$ 2.64	9.98	0.81	0.0
016_A_18444	215.106518	52.975820	6.174	12.27 $\pm$ 4.72	14.37 $\pm$ 2.93	3.09 $\pm$ 4.14	0.27 $\pm$ 2.07	9.48	0.54	0.0
012_A_27683	214.989007	52.919644	6.670	11.81 $\pm$ 2.79	11.91 $\pm$ 3.97	2.01 $\pm$ 1.92	1.37 $\pm$ 2.13	9.74	0.81	0.0
013_A_10524	214.949129	52.843185	6.716	18.91 $\pm$ 4.62	16.83 $\pm$ 3.23	10.57 $\pm$ 4.10	0.52 $\pm$ 3.69	9.83	1.82	0.8
013_B_17930	214.869985	52.807034	6.749	31.76 $\pm$ 6.47	32.30 $\pm$ 4.05	6.41 $\pm$ 2.28	2.21 $\pm$ 2.68	10.19	0.73	0.0
015_B_5666	215.000956	52.865869	6.670	11.18 $\pm$ 2.92	6.66 $\pm$ 3.25	2.36 $\pm$ 2.04	1.78 $\pm$ 2.25	9.23	0.50	0.0
012_B_28896	214.921870	52.876193	5.630	14.94 $\pm$ 6.96	12.38 $\pm$ 3.83	2.35 $\pm$ 3.91	4.41 $\pm$ 1.50	9.24	0.14	0.0
015_A_16826	215.032039	52.918960	6.170	15.34 $\pm$ 3.71	14.03 $\pm$ 3.33	4.51 $\pm$ 4.90	2.12 $\pm$ 1.99	9.70	0.76	0.1
015_A_17952	215.023039	52.915309	6.165	9.08 $\pm$ 2.35	7.55 $\pm$ 3.45	0.84 $\pm$ 3.02	2.67 $\pm$ 1.51	9.25	0.31	0.0
015_B_10107	214.987324	52.868911	5.665	17.88 $\pm$ 5.82	18.44 $\pm$ 4.50	3.56 $\pm$ 7.44	7.92 $\pm$ 4.12	9.38	0.82	0.0
016_B_13823	215.090344	52.951601	5.501	18.82 $\pm$ 8.36	29.20 $\pm$ 4.27	5.69 $\pm$ 13.39	4.50 $\pm$ 4.36	9.21	0.36	0.0

**Notes.** The emission-line flux measurements for the [O III] NIRCcam WFSS sample in units of  $10^{-18}$  erg/s/cm $^2$ . Other columns show the spectroscopic redshift, stellar mass, SED SFR, and dust attenuation from the FAST SED fitting. The ID indicates the field\_panel\_WFSS ID number.

**Table 2**  
H $\alpha$  SFR NIRC*am* Sample

ID	R.A. (deg)	Decl. (deg)	$z$	H $\alpha$ Column ( $10^{-18}$ erg/s/cm $^2$ )	H $\alpha$ Row ( $10^{-18}$ erg/s/cm $^2$ )	$\log(M_*)$ [ $M_\odot$ ]	$\log(\text{SFR})$ [ $M_\odot$ ] yr $^{-1}$	$A_V$ (mag)
012_A_25344	214.977646	52.903536	4.770	8.17 $\pm$ 1.79	21.61 $\pm$ 7.47	9.84	0.66	0.0
012_A_25431	214.977566	52.903450	4.550	9.36 $\pm$ 0.85	3.35 $\pm$ 1.88	7.46	-0.07	1.3
012_A_31041	214.953995	52.907856	4.550	8.52 $\pm$ 3.03	6.49 $\pm$ 1.44	7.04	-0.53	0.8
012_A_35810	214.942294	52.919362	3.938	38.27 $\pm$ 8.28	50.02 $\pm$ 6.51	7.73	0.20	1.3
012_A_36545	214.937471	52.918283	3.938	29.55 $\pm$ 9.62	24.45 $\pm$ 10.11	10.25	0.24	0.1
012_B_26391	214.927867	52.871022	4.805	12.42 $\pm$ 1.67	10.74 $\pm$ 3.26	8.24	-1.49	0.6
012_B_29360	214.917888	52.875555	4.717	12.09 $\pm$ 2.36	12.83 $\pm$ 3.73	8.06	-1.95	0.3
013_A_19410	214.943410	52.864098	4.675	17.76 $\pm$ 5.27	16.63 $\pm$ 5.67	10.32	1.38	0.8
013_A_20628	214.941356	52.864855	4.678	8.29 $\pm$ 2.49	7.33 $\pm$ 2.36	7.86	-1.62	0.0
013_B_13408	214.894682	52.812130	4.876	13.53 $\pm$ 3.88	15.05 $\pm$ 4.52	8.08	-2.73	0.3
015_A_20196	215.022027	52.920785	4.540	14.11 $\pm$ 3.47	6.73 $\pm$ 2.95	9.03	-9.90	2.9
015_B_16325	214.978092	52.879514	4.545	12.73 $\pm$ 4.19	19.95 $\pm$ 3.85	7.48	-0.09	1.2
015_B_17048	214.985862	52.886907	4.546	9.39 $\pm$ 3.17	10.73 $\pm$ 3.36	7.20	-0.33	1.3
016_A_5887	215.151987	52.974048	4.480	29.21 $\pm$ 4.90	18.21 $\pm$ 2.09	8.42	0.61	1.9
016_A_8530	215.149544	52.978974	4.525	6.73 $\pm$ 1.59	6.29 $\pm$ 0.15	8.07	-8.80	0.7
016_B_15982	215.059038	52.936442	4.280	65.93 $\pm$ 16.13	89.81 $\pm$ 2.93	9.53	0.97	0.0
016_B_18194	215.079996	52.956800	4.745	19.97 $\pm$ 7.73	15.28 $\pm$ 5.60	8.50	-0.46	0.5
016_B_8414	215.083373	52.931987	4.117	10.45 $\pm$ 3.61	17.34 $\pm$ 3.73	8.05	-1.68	0.7

**Notes.** The H $\alpha$  flux measurements from NIRC*am* WFSS, and the spectroscopic redshift, stellar mass, SED SFR, and dust attenuation from the FAST SED fitting. The ID indicates the field\_panel\_WFSS ID number.

## 2.2. Lower-redshift Comparison Samples

We established two  $z \sim 2$  comparison samples to cover the peak of cosmic star formation (Madau & Dickinson 2014) and supermassive black hole growth (Aird et al. 2010). An SDSS  $z \sim 0$  comparison sample was also established to cover the local Universe (Kauffmann et al. 2003b, 2003c; Brinchmann et al. 2004). We acknowledge the NIRS*pec* selection was complicated and heterogeneous due to pre-JWST (HST photometry) target selection. We calculated the average emission-line luminosity from CEERS to be  $\log(L) = 41.29$  (units of erg s $^{-1}$ ), which is very similar to the average CLEAR emission-line luminosity,  $\log(L) = 41.55$  (units of erg s $^{-1}$ ), allowing a first-order comparison. These similar luminosities show it is reasonable to compare the two samples. However, the average emission-line luminosity for SDSS is  $\log(L) = 39.60$  (units of erg s $^{-1}$ ), making it deeper than the CEERS sample. Restricting the SDSS sample size based on CEERS luminosities would not provide a large enough sample for analysis. Additionally, the SDSS galaxies make use of a Kroupa IMF, while CLEAR and CEERS use a Chabrier IMF. However, as shown in Madau & Dickinson (2014), there is a small set offset of  $\text{SFR}(\text{Kroupa}) = 1.06 \text{ SFR}(\text{Chabrier})$  and  $M_*(\text{Kroupa}) = 1.08 M_*(\text{Chabrier})$ .

### 2.2.1. SDSS $z \sim 0$ Sample

The  $z \sim 0$  comparison sample is created from the Sloan Digital Sky Survey (SDSS; York et al. 2000) Data Release 10 (Ahn et al. 2014). The SDSS data set used a 2.5 m telescope at Apache Point Observatory to cover 14,555 deg $^2$  in the sky with  $R \sim 2000$  over  $3800 < \lambda < 9200 \text{ \AA}$  (Smee et al. 2013).

Emission-line measurements and redshifts for the SDSS data set are computed by Bolton et al. (2012), using a stellar template to correct the continuum for stellar absorption. Stellar masses are estimated by Montero-Dorta et al. (2016) from the broadband *ugriz* SDSS photometry using a grid of templates made from the FSPS stellar population synthesis code (Conroy et al. 2009). These templates assume a Kroupa (2001) IMF and

fit for the dust attenuation following Charlot & Fall (2000) and Calzetti et al. (2000).

The low-redshift,  $z \sim 0$ , comparison sample was selected using the same S/N > 3 line detection thresholds as for the CEERS and CLEAR samples. These selection criteria result in 284,523 galaxies with H $\alpha$ , 231,999 galaxies with [O III]/H $\beta$ , and 27,847 galaxies with [Ne III]/[O II]. We note this SDSS sample has no cuts on whether a galaxy is an AGN or SF.

### 2.2.2. CLEAR $z \sim 1.5$ Sample

Our  $z \sim 2$  comparison sample comes from the CANDELS Ly $\alpha$  Emission at Reionization (CLEAR) survey (Simons et al. 2023) and HST near-IR spectroscopy with the G102 and G141 grisms taken as part of the 3D-HST program (van Dokkum et al. 2011; Brammer et al. 2012; Momcheva et al. 2016).

We select the CLEAR comparison sample using the same S/N > 3 line detection thresholds as for the CEERS sample, visually inspecting the direct image, 1D, and 2D spectra to remove galaxies with contaminated spectra. This gives us 2890 galaxies with H $\alpha$ , 1534 galaxies with [O III], and 505 in our [O II] sample. Due to this sample's low spectral resolution ( $R \sim 100$ ), H $\alpha$  is blended with [N II] $\lambda$ 6583 + 6548. This blending causes the H $\alpha$  fluxes of CLEAR galaxies to effectively be upper limits, although most  $z \sim 2$  galaxies have [N II]/H $\alpha \ll 1$  (e.g., Shapley et al. 2015). We note this CLEAR sample makes no distinction between star-forming galaxies and AGN.

The HST data used in this paper can be found in MAST:10.17909/t9-ctff-wx60.

### 2.2.3. MOSDEF $z \sim 3$ Sample

We also compare to the stacked line-ratio measurements from the MOSFIRE Deep Evolution Field (MOSDEF) observations used in Sanders et al. (2021) at redshift  $z \sim 3.3$ . This sample was observed by the Multi-Object Spectrometer For Infrared Exploration (MOSFIRE; McLean et al. 2012) on the 10 m Keck I telescope, observing 48.5 nights over a four-year period to obtain rest-frame optical spectra of  $1.4 < z < 3.8$


**Table 3**  
NIRSpec Sample

R.A. (deg)	decl. (deg)	z	[O III] ( $10^{-18}$ ergs/s/cm $^2$ )	H $\beta$ ( $10^{-18}$ ergs/s/cm $^2$ )	H $\alpha$ ( $10^{-18}$ ergs/s/cm $^2$ )	[Ne III] ( $10^{-18}$ ergs/s/cm $^2$ )	[O II] ( $10^{-18}$ ergs/s/cm $^2$ )	$\log(M_*)$ [ $M_\odot$ ]	$\log(\text{SFR})$ [ $M_\odot$ ] yr $^{-1}$	$A_V$ (mag)
214.957160	52.872372	3.228	2.57 $\pm$ 0.31	0.54 $\pm$ 0.43	1.57 $\pm$ 0.21	1.82 $\pm$ 0.52	1.22 $\pm$ 0.47	8.91	0.18	0.0
214.959997	52.831169	4.900	3.09 $\pm$ 0.19	0.25 $\pm$ 0.12	1.20 $\pm$ 0.08	−99.00 $\pm$ 0.00	−99.00 $\pm$ 0.00	9.11	0.67	0.0
214.893181	52.882484	3.000	1.83 $\pm$ 0.22	0.78 $\pm$ 0.35	1.49 $\pm$ 0.27	2.05 $\pm$ 0.54	2.10 $\pm$ 0.59	8.80	0.06	0.0
214.943900	52.850052	5.001	65.25 $\pm$ 19.57	8.19 $\pm$ 0.88	30.26 $\pm$ 0.84	3.60 $\pm$ 0.45	11.17 $\pm$ 0.84	9.48	0.74	0.0
214.941496	52.850565	2.540	10.00 $\pm$ 4.94	5.29 $\pm$ 0.52	18.24 $\pm$ 1.76	0.56 $\pm$ 1.13	10.69 $\pm$ 0.80	9.03	0.48	0.0
214.907360	52.844535	2.010	15.23 $\pm$ 1.67	1.83 $\pm$ 1.01	5.80 $\pm$ 0.25	1.05 $\pm$ 0.77	2.53 $\pm$ 1.02	7.54	−0.86	0.0
214.898480	52.861709	1.922	1.35 $\pm$ 0.64	0.79 $\pm$ 0.58	4.22 $\pm$ 0.53	0.52 $\pm$ 229.54	4.63 $\pm$ 0.60	9.76	0.84	0.3
214.909604	52.880284	2.144	4.79 $\pm$ 0.37	1.19 $\pm$ 0.42	2.85 $\pm$ 0.18	0.36 $\pm$ 0.46	2.59 $\pm$ 0.43	8.85	0.41	0.0
214.966546	52.846672	2.136	19.14 $\pm$ 3.83	3.99 $\pm$ 0.54	14.20 $\pm$ 0.76	1.07 $\pm$ .92	11.29 $\pm$ 1.42	9.18	0.25	0.0
214.940182	52.836026	1.699	9.83 $\pm$ 0.97	6.28 $\pm$ 1.29	25.11 $\pm$ 1.09	2.14 $\pm$ 1.29	16.69 $\pm$ 2.59	10.18	1.19	0.8

**Notes.** The emission-line fluxes from the NIRSpec sample. The R.A. and decl. are given in degrees and redshifts are given for each galaxy. Emission-line fluxes are reported in units of  $10^{-18}$ ergs/s/cm $^2$ . The SFR and dust attenuation measurements are produced from SED fitting. A machine-readable version of the full table is available. When an emission line falls in a chip gap or off the spectra, a value of −99.0 is assigned.

(This table is available in its entirety in machine-readable form.)





**Figure 2.** Distribution of redshifts for each emission-line sample:  $H\alpha$  (top),  $[O\text{ III}]$  (middle), and  $[\text{Ne III}]$  and  $[O\text{ II}]$  (bottom). These are named by their respective emission lines, which have  $S/N > 3$ ; in the case of  $[\text{Ne III}]$  and  $[O\text{ II}]$ , either line may reach this requirement. The samples come from the SDSS(left), CLEAR (middle), and CEERS(right). In the right panel, the gray histograms are the NIRSpec observations and the blue ones are NIRCcam WFSS.

galaxies (Kriek et al. 2015). This survey covers the AEGIS, COSMOS, and GOODS-N fields. In order to cover multiple emission lines, two to three filters are used in the survey to observe  $H$ -band-selected galaxies. MOSDEF adopts a slit width of  $0''.7$ , which results in a spectral resolution of  $R=3400$ , 3000, 3650 and 3600 for  $Y$ ,  $J$ ,  $H$ , and  $K$ , respectively.


We use the stacked line-ratio measurements from Sanders et al. (2021), which were calculated from  $\sim 750$  galaxies at  $z \sim 2.3$  and  $\sim 375$  galaxies at  $z \sim 3.3$ . For a full description of the MOSDEF survey design and data reduction, see Kriek et al. (2015).

### 3. NIRCcam WFSS and NIRSpec MSA Comparison


The NIRCcam WFSS is a blind survey that allows us to gain access to a sample of galaxies that may be missed by targeted

NIRSpec observations, which require preselection of galaxies based on redshift and brightness. In the left panel of Figure 4, the NIRSpec  $z > 5$  galaxies are represented by gray circles, while the blue circles represent the combined column and row measurements of NIRCcam WFSS. The NIRCcam WFSS galaxies occupy a similar region of continuum measurements, but these galaxies tend to have higher  $[O\text{ III}]$  emission, due to the higher flux limits.

Due to this difference in sample selection, the NIRCcam WFSS provides an opportunity to view different types of galaxies. These galaxies also have differences in the emission-line measurements. Figure 4 also shows a comparison of the flux measurements of  $[O\text{ III}]$  and  $H\alpha$  between NIRCcam WFSS and NIRSpec. These emission lines were chosen because they have  $S/N > 3$ . NIRSpec is represented by the gray histogram and the combined measurements from both the row and column



**Figure 3.** Left: Distribution of  $H\beta$  SFR and stellar mass for our  $[O\text{ III}]/H\beta$  sample. Right: Distribution of  $H\alpha$  SFR and stellar mass for our  $H\alpha$  sample. In both the panels, galaxies observed by NIRCcam WFSS are represented by black points and arrows to represent lower limits, and galaxies observed by NIRSpec are represented by pink points. The red and gray contours represent the CLEAR and SDSS samples, respectively, with contour level indicating relative galaxy density of each sample.



**Figure 4.** Left: Comparison of the NIRSpec and NIRCcam WFSS measurements of F277W photometric flux and  $[O\text{ III}]$  flux. Comparison between emission-line flux from NIRSpec (gray histogram) and NIRCcam WFSS (blue histograms) for  $H\alpha$  emission (middle panel) and  $[O\text{ III}]$  emission (right panel). Both these emission lines were seen in both instruments. A blind survey like NIRCcam WFSS includes galaxies that have been missed in the targeted NIRSpec observations. The NIRCcam WFSS galaxies tend to have stronger emission lines, due to the higher flux limit.

dispersions for NIRCcam WFSS are shown in the blue histograms. This comparison shows that NIRCcam WFSS tends to include galaxies with brighter emission-line fluxes. This indicates that the CEERS NIRSpec observations are missing galaxies with more extreme emission lines. This should be kept in mind in our following sections when comparing NIRSpec and NIRCcam WFSS galaxies.


Of the CEERS observations, only two galaxies have been observed using both NIRSpec and NIRCcam WFSS. One galaxy is in the NIRCcam WFSS  $[O\text{ III}]/H\beta$  sample exclusively, as the  $[O\text{ III}]/H\beta$  line in the NIRSpec spectra fall into a chip gap that prevents us from measuring the emission lines. However, the other galaxy has  $H\alpha$  flux measurements of  $19.97 \pm 7.73$  (units of  $\times 10^{-18}$  erg/s/cm<sup>2</sup>) and  $15.28 \pm 5.60$  (units of  $\times 10^{-18}$  erg/s/cm<sup>2</sup>) in the NIRCcam WFSS columns and row and a  $19.39 \pm 0.72$  (units of  $\times 10^{-18}$  erg/s/cm<sup>2</sup>) measurement in NIRSpec. Though this is only a single galaxy, it shows promise that our slit loss correction in the NIRSpec measurements is working well.

#### 4. Redshift Evolution of Emission-line Galaxies

We measure the galaxy emission-line ratios with redshift by comparing CEERS galaxies with  $z \sim 2$  galaxies from CLEAR and MOSDEF and with  $z \sim 0$  galaxies from SDSS.

In Figure 5, we plot the  $[O\text{ III}]/H\beta$  emission-line ratios against redshift. The black point is the median value from the SDSS sample, the gray points are median values from two redshift bins from the CLEAR sample, and the Sanders et al. (2021) redshift 3.3 binned data are shown as pink squares. The pink circles are our NIRSpec sample and the red points are the CEERS sample. Arrows in our CEERS WFSS sample represent lower limits in the emission-line ratio, due to undetected  $H\beta$ . Other high-redshift galaxies from SMACS ERO NIRSpec observations (Trump et al. 2023) and broad-line AGN from Kocevski et al. (2023) and Larson et al. (2023) are marked as crimson squares, purple stars, and gold diamonds, respectively.

There is a 0.49 dex increase in  $[O\text{ III}]/H\beta$  between the SDSS sample and CLEAR; for more details and discussion, the reader may refer to Backhaus et al. (2022), who luminosity matched



**Figure 5.** The  $\log[\text{O III}]/\text{H}\beta$  emission-line ratio vs. redshift. The black point is the median redshift and  $\log[\text{O III}]/\text{H}\beta$  of the SDSS sample (York et al. 2000), with the error bars representing the standard deviation of the sample. The gray points and pink squares are from the CLEAR and MOSDEF samples, respectively (Sanders et al. 2021; Simons et al. 2023). The SMACS observations are marked as crimson squares. NIRSspec AGN from CEERS are purple stars. The NIRSspec and WFSS samples are represented by pink points and red points and arrows, respectively. The yellow line is the linear fit to the NIRSspec. The cyan lines represent the rolling median detection limit (MDL) to the best-fit line for the NIRSspec galaxies, indicating the range of galaxies we could see based on our S/N cutoffs. This increase of  $[\text{O III}]/\text{H}\beta$  with redshift is also shown in the linear fit line, with a slope of  $(0.06 \pm 0.02)$ .

the two samples to avoid differences in sample selection. We also find that the  $z > 5$  CEERS sample has 0.33 dex higher  $[\text{O III}]/\text{H}\beta$  than the CLEAR sample, but we note this may be due to the difference in the luminosity selection. A `linmix` linear fit to the line ratios measured from CEERS NIRSspec observations indicates a shallow but significant ( $2.3\sigma$ ) increase of  $[\text{O III}]/\text{H}\beta$  with redshift; this fit includes the lower limits. `Linmix` is a hierarchical Bayesian approach to linear regression Kelly (2007). In Guo et al. (2016), they measure a median  $[\text{O III}]/\text{H}\beta$  ratio of  $\sim 0.3$  at  $z \sim 0.6$  for  $\sim 1400$  galaxies; this matches well with the predictions from our fit line. Many of the NIRCcam WFSS line ratios are lower limits, due to undetected  $\text{H}\beta$  lines. These limits are generally consistent with the measured NIRSspec line ratios, but they are also consistent with a steeper increase of  $[\text{O III}]/\text{H}\beta$  with redshift.


The median detection limits (MDL) are created by taking the upper and lower limit for each line ratio of each galaxy. The rolling MDL are the median difference between the detection limits and the linear fit. This line represents the lowest signal that can be observed with a  $1\sigma$  detection of the line flux. We note that our NIRSspec MSA data at high redshifts are well separated from the rolling MDL, indicating there is no selection bias for our high-redshift sample. Thus, the small increase of  $[\text{O III}]/\text{H}\beta$  ratio with redshift is not likely to be a simple result of a changing detection limit with redshift.

Figure 6 shows the  $[\text{Ne III}]/[\text{O II}]$  ratio with redshift, using the same notation as Figure 5. The  $[\text{Ne III}]/[\text{O II}]$  ratio also increases with redshift. There is a 0.34 dex increase between the SDSS and CLEAR samples. This is different from the results from Backhaus et al. (2022), who found  $[\text{Ne III}]/[\text{O II}]$  had a 0.2 dex difference between SDSS and CLEAR. This may be because Backhaus et al. (2022) required an  $S/N > 1$  in both  $[\text{O II}]$  and  $[\text{Ne III}]$ , whereas we only require  $S/N > 3$  for  $[\text{O II}]$  for the SDSS, CLEAR, and CEERS samples. There is also a 0.37 dex increase between CLEAR and galaxies with  $z > 5$ .


This is smaller than the 0.5 dex increase between CLEAR and SMACS reported by Trump et al. (2023), but Trump et al. (2023) only had five galaxies above  $z > 5$ . Similarly to Figure 5, this fit is to the CEERS NIRSspec data. There is only a marginal ( $2.5\sigma$ ) correlation between the  $[\text{Ne III}]/[\text{O II}]$  ratio and redshift in the CEERS samples, with a slope of  $0.05 \pm 0.02$ . The increase of  $[\text{Ne III}]/[\text{O II}]$  at  $z > 2$  is further emphasized when looking at the work of Pharo et al. (2023), who found that the stacked  $[\text{Ne III}]/[\text{O II}]$  ratio of a representative low-mass ( $\log(M_*/M_\odot) < 9$ ) star-forming galaxies at  $z \sim 0.7$  is the same as the stacked  $[\text{Ne III}]/[\text{O II}]$  ratio of local ( $z \sim 0$ ) galaxies, while the  $z \sim 2$  measurements from HST are higher than their stacked  $z \sim 0.7$  measurements. This indicates that  $[\text{Ne III}]/[\text{O II}]$  does not evolve substantially from between  $0 < z < 1$  but has a jump around  $z \sim 2$ .

The rolling MDL indicates the CEERS  $z > 2$  sample may be affected by the detection limit caused by the  $S/N > 1$  detection threshold for  $[\text{Ne III}]$ , such that there might exist  $z > 1.5$  galaxies with lower  $[\text{Ne III}]/[\text{O II}]$  that are undetected. On the other hand, the measured line ratios are well-separated from the upper detection limit and there appears to be a genuine lack of high- $[\text{Ne III}]/[\text{O II}]$  galaxies at  $z > 2$ .

We show the relationship between  $\text{H}\alpha$  SFR with redshift in Figure 7. This sample is not dust corrected, as we do not have a pair of hydrogen lines in all observations and the SED-based  $A_V$  may be unreliable for estimating the nebular attenuation. We note that the  $\text{H}\alpha$  NIRCcam WFSS galaxies have higher  $A_V$  measured by their SED fitting than the other samples. This would push the galaxies to higher  $\text{H}\alpha$  SFR. We fit a line to the CEERS NIRSspec galaxies and see a significant ( $6\sigma$ ) and strong correlation of  $\text{H}\alpha$  SFR with redshift. This trend is further emphasized by the 1.5 dex increase in  $\text{H}\alpha$  SFR between SDSS and CLEAR galaxies and a smaller 0.5 dex increase between CLEAR and  $z > 5$  galaxies. We see a 0.7 dex difference between the NIRCcam WFSS and NIRSspec galaxies between



**Figure 6.** The  $\log[\text{Ne III}]/[\text{O II}]$  emission-line ratio vs. redshift. The black point is the median redshift and  $\log[\text{Ne III}]/[\text{O II}]$  of the SDSS sample York et al. (2000). The gray point is a median value from CLEAR Simons et al. (2023). The pink points and red squares are from the CEERS NIRSpec and SMACS samples, respectively. The NIRSpec  $[\text{Ne III}]/[\text{O II}]$  line ratio has a  $2.5\sigma$  slope with redshift,  $(0.05 \pm 0.02)$ , as shown by the yellow best-fit line.



**Figure 7.** The  $\text{H}\alpha$  SFR vs. redshift. The black point is the median value of the SDSS sample (York et al. 2000). The gray point is a median value from CLEAR (Simons et al. 2023). The pink points and red circles are from the CEERS NIRSpec and NIRCam WFSS samples, respectively. The cyan and green lines represent the lower detection limits of the NIRSpec and NIRCam WFSS data, respectively. There is a 1.5 dex increase in the median  $\text{H}\alpha$  SFR measurements from SDSS to CLEAR and a 0.5 dex increase in the median  $\text{H}\alpha$  SFR measurements between CLEAR and  $z > 5$  galaxies. The yellow best-fit line to the NIRSpec  $\text{H}\alpha$  SFR has a  $8\sigma$  slope with redshift,  $(0.24 \pm 0.03)$ .

$3.8 < z < 5$ , due to the different detection limits of each instrument.

The relationships between emission lines and galaxy properties and ISM conditions are further explored in Sections 5 and 6 of the paper.

## 5. Emission-line Properties with Galaxy Stellar Mass and SFR

We will now investigate how the emission-line ratios correlate to galaxy properties such as stellar mass and SFR.


Figure 8 relates the  $[\text{O III}]/\text{H}\beta$  ratio to  $\text{H}\beta$  SFR and stellar mass. Figure 9 similarly compares  $[\text{Ne III}]/[\text{O II}]$  to  $\text{H}\beta$  SFR and stellar mass. Equation (1) was used to derive  $\text{H}\beta$  SFR; this is a proxy for the unobscured galaxy SFR. Here, our  $z \sim 0$  SDSS and  $z \sim 2$  CLEAR samples are represented by the gray and red contours, respectively. Our NIRCam WFSS galaxies

are represented by black points for well-measured galaxies and black arrow for lower limits. NIRSpec galaxies are shown as pink points. In Figure 8, we include the median value of the Illustris simulations as the purple line. We calculate a best-fit line using the CEERS NIRSpec  $z > 5$  sample and subsamples of SDSS galaxies and CLEAR galaxies the same size as the CEERS sample.


We additionally compare our observed  $[\text{O III}]/\text{H}\beta$  to the simulated ratios of Hirschmann et al. (2023a). These model line ratios are built on the IllustrisTNG simulations (reference), applying both Cloudy (Ferland et al. 2013) and MAPPINGS V (Sutherland et al. 2018) photoionization models to the simulated galaxies. These simulated galaxies include a mix of nebular emission contributions from star-forming H II regions, post-AGB stars, shocks, and AGN narrow-line regions. The details of these model line ratios are described by Hirschmann et al. (2017) and Hirschmann et al. (2023b). We compare to model line ratios from simulated  $z \sim 6$  galaxies with  $\log(M_*/M_\odot) \sim 9$  and  $\log(\text{SFR}/[M_\odot/\text{yr}]) \sim 0.5$ , matching the median stellar mass and SFR of the observed galaxies. The ranges of line ratios from these simulated galaxies are shown as purple shading in Figure 8.

The  $[\text{O III}]/\text{H}\beta$  emission-line ratio has a significant ( $>3\sigma$ ) correlation with  $\text{H}\beta$  SFR, with a slope of  $0.2 \pm 0.04$ , and a significant anticorrelation with stellar mass, with a slope of  $-0.41 \pm 0.09$ . Our  $z > 5$  NIRCam WFSS and NIRSpec samples are on average 0.5 dex higher in  $\text{H}\beta$  SFR than the CLEAR galaxies. The IllustrisTNG simulated line ratios are effective at reproducing the highest  $[\text{O III}]/\text{H}\beta$  ratios observed in our sample, but most of our observed galaxies have lower  $[\text{O III}]/\text{H}\beta$  ratios than the simulated galaxies of similar galaxy mass and SFR. We attempted to do multiple linear regression on our sample to detangle the relationship between the  $[\text{O III}]/\text{H}\beta$  emission line with redshift,  $\text{H}\beta$  SFR, and stellar mass; however, due to the sample size and the differences between the NIRSpec and NIRCam WFSS samples, this produced unreliable results. A larger sample size would be needed for further analysis.





**Figure 8.** Left: The relationships between the [O III]/H $\beta$  emission-line ratio and the galaxy H $\beta$  luminosity and H $\beta$  SFR. Right: The relationships between the [O III]/H $\beta$  emission-line ratio and stellar mass. In both panels, the gray and red contours are the SDSS and CLEAR samples, respectively. The black points and arrows are [O III]/H $\beta$  measurements and limits from NIRCам WFSS observations. The yellow line is fit to the CEERS  $z > 5$  sample and randomly selected SDSS and CLEAR galaxies matching the size of the CEERS sample. All the IllustrisTNG simulations are shown as the purple region.



**Figure 9.** Left: The relationships between the [Ne III]/[O II] emission-line ratio and the galaxy H $\beta$  luminosity and H $\beta$  SFR. Right: The relationships between the [Ne III]/[O II] emission-line ratio and stellar mass. In both panels, the gray and red contours are the SDSS and CLEAR sample, respectively. The black points and arrows are [O III]/H $\beta$  measurements and limits from NIRCам WFSS observations. The yellow line is fit to the CEERS  $z > 5$  sample and randomly selected SDSS and CLEAR galaxies matching the size of the CEERS sample. There is a significant ( $>3\sigma$ ) relationship between [Ne III]/[O II] and H $\beta$  SFR, and a marginal relationship ( $2.8\sigma$ ) with stellar mass.


The [Ne III]/[O II] ratio has a significant correlation with H $\beta$  SFR, with a slope of  $0.23 \pm 0.04$ , which is similar to [O III]/H $\beta$ . Similarly to [O III]/H $\beta$ , [Ne III]/[O II] has a marginal ( $2\sigma$ ) anticorrelation to stellar mass with a slope of  $-0.2 \pm 0.07$ , but this is about three times less significant than what was found between stellar mass and [O III]/H $\beta$ .

The anticorrelation between  $\log([\text{O III}]/\text{H}\beta)$  and stellar mass has been shown in Dickey et al. (2016) and Kashino et al. (2019), and it is due to lower metallicity and higher ionization in galaxies with higher specific star formation rates (sSFR). This was also shown in simulations done in Hirschmann et al. (2017) and Hirschmann et al. (2023b).

## 6. ISM Conditions at $z > 5$

We compare our measurements of [O III]/H $\beta$  and [Ne III]/[O II] to theoretical models to infer the physical conditions of

the ISM. The models we compare to come from the work of Kewley et al. (2019), which uses both the Starburst99 (Leitherer et al. 1999) models of stellar ionizing spectra and the MAPPINGS V ionization code (Sutherland et al. 2018). The Starburst99 model spectra includes mass loss and uses a Salpeter (1955) IMF. The atomic data used in MAPPINGS V come from the CHIANTI 8 database (Dere et al. 1997; Del Zanna et al. 2015), which includes the effects of excitation, dust depletion, recombination, and photoionization in the model H II regions. The ‘‘Pressure Models’’ of Kewley et al. (2019) describe synthetic emission-line spectra created from different combinations of pressure  $\log(P/k)$ , ionization  $\log(q)$ , and metallicity  $Z/Z_{\odot}$ . These models interpolate between Starburst99 models and CHIANTI 8 data in order to match the grid. We use the following values of ionization and metallicity:




**Figure 10.** Top Left: Comparison of  $[\text{O III}]/\text{H}\beta$  measurements of our sample to MAPPINGS V models of ISM ionization and metallicities. The gray histogram represents the distribution of the NIRSpect measurements of  $[\text{O III}]/\text{H}\beta$ , while the blue histogram represents the NIRCcam WFSS sample; neither of the CEERS samples has a measured metallicity. We also include the four SMACS galaxies that have metallicity measurements using the coronal  $[\text{O III}]\lambda 4364$ . Top Right: Comparison of  $[\text{Ne III}]/[\text{O II}]$  measurements of our sample to MAPPINGS V models of the ISM’s ionization and metallicity. The gray histogram represents the distribution of the NIRSpect measurements of  $[\text{Ne III}]/[\text{O II}]$ . Bottom: Distribution of the CEERS  $z > 5$  sample where the metallicities are derived using the strong-line calibrations provided in Sanders et al. (2024).

1. Ionization  $\log(q) = [7, 8, 9]$ , units of  $\text{cm s}^{-1}$ ; and
2. Metallicity  $Z/Z_{\odot} = [0.05, 0.2, 0.4, 1.0]$ .

Due to neither emission-line ratio varying significantly with pressure, we choose  $\log P = 8$  for the models. We also note the  $Z/Z_{\odot} = 0.05$  bin is extrapolated in the Starburst99 input spectra, which causes those synthetic spectra to be the least certain of the theoretical predictions.

In Figure 10, we compare our high-redshift sample to the MAPPINGS V models. The gray and blue histograms in both the  $[\text{O III}]/\text{H}\beta$  and  $[\text{Ne III}]/[\text{O II}]$  top panels represent the NIRSpect and NIRCcam WFSS emission-line ratio distribution, respectively. The four SMACS galaxies that have metallicity measurements are also included as red squares. These are compared to the MAPPINGS V models that are represented by the three colored lines, where each color is a different ionization and each point is a different metallicity. The bottom panels are individual galaxy measurements, where metallicities are derived using the strong-line calibrations from Sanders et al. (2024). The  $[\text{O III}]/\text{H}\beta$  ratios cover a range of moderate to high ionization. The  $[\text{Ne III}]/[\text{O II}]$  ratios are similarly well-matched to MAPPINGS V models for moderate to high ionization.

Due to the degeneracy between ionization and metallicity in the MAPPINGS V model prediction, it is difficult to infer ISM metallicity from the  $[\text{O III}]/\text{H}\beta$  ratio in Figure 10. To mitigate this, we also make use of the OHNO diagram in Figure 11, to give us a more narrow range of ISM conditions and highlight the evolution of ISM conditions with redshift. The gray contours represent the SDSS sample; the galaxies that are in the AGN region of the diagram have higher ionization. The black points are the CLEAR sample from CLEAR. We include the measurements from the SMACS galaxies from Trump et al. (2023) and the two NIRSpect AGN from Kocevski et al. (2023) as red squares and purple stars, respectively. The  $z = 8.7$  AGN observed by Larson et al. (2023) is shown as a yellow star. Our sample is shown as pink circles representing the NIRSpect sample above  $z > 5$  and a pink bar for NIRCcam WFSS. This bar represents the range of  $[\text{O III}]/\text{H}\beta$  covered by the WFSS sample, as no  $[\text{Ne III}]/[\text{O II}]$  measurements were detected for these galaxies. The MAPPINGS V models are represented the same way as in Figure 10. The inset arrow shows the typical scale and direction of  $\sim 1$  dex changes in metallicity. From this, we can see our  $z > 5$  sample prefers a moderate to high ionization,  $\log(q) = 8, 9$ , with a moderate metallicity,  $Z/Z_{\odot} = 0.2, 0.4$ . The MAPPINGS V models indicate that



**Figure 11.** Theoretical OHNO line ratio predictions from the MAPPINGS V models (colored lines) compared to the low-redshift SDSS galaxies (gray contours), cosmic noon CLEAR galaxies (black points), SMACS galaxies (red stars), CEERS AGN (purple stars), and CEERS NIRSpec galaxies (pink points). Due to the constraint of the filter, only  $[\text{O III}]/\text{H}\beta$  can be observed for the NIRCcam WFSS observations. These galaxies are represented by the histogram in the right-hand panel, the gray histogram represents the NIRCcam WFSS measurements with a lower limit, and red histogram are galaxies with an  $S/N$  measurement  $> 3$  for both emission lines. The arrow on this histogram indicates that the true  $[\text{O III}]/\text{H}\beta$  values may be higher. The red histogram in the panel represents the well-measured galaxies from the NIRCcam WFSS, this sample is also represented by the pink bar across the panel. Model ionization is indicated by the color of the line, and metallicity decreases from left to right for each set of connected model points. Inset vectors indicate the direction and amplitude of 1 dex increases in metallicity and pressure. The black dashed line is the empirical AGN/SF dividing line defined for  $z \sim 2$  galaxies in Backhaus et al. (2022). The CEERS data set is best described by moderate/high ionization and a moderate metallicity, while CLEAR is described by lower ionization and more moderate metallicity.

ionization increases from the  $z \sim 0$  SDSS sample to the  $z \sim 2$  CLEAR sample to the  $z > 5$  CEERS sample. We note that the three AGN sources have a higher  $[\text{Ne III}]/[\text{O II}]$  ratio and are best described by MAPPINGS V models with higher ionization.

## 7. Summary

In this work, we studied optical emission-line ratios from  $z \sim 0$  to  $z \sim 9$  using SDSS, CLEAR, and CEERS data sets. We have used NIRCcam WFSS to define two samples measuring  $\text{H}\alpha$  of 18 galaxies at  $3.9 < z < 4.9$  and  $[\text{O III}]/\text{H}\beta$  of 19 galaxies at  $5.5 < z < 6.7$ . This sample was found by first setting a constraint on the photometric redshifts, before visually inspecting for the emission lines. We also have three samples using NIRSpec observations to give additional measurements of  $\text{H}\alpha$  in 93 galaxies,  $[\text{O III}]/\text{H}\beta$  of 96 galaxies, and  $[\text{Ne III}]/[\text{O II}]$  for 59 galaxies. The NIRSpec sample is selected with a  $S/N > 3$  in the emission lines of interest, before being visually inspected.

We studied these emission lines and summarize our results as follows:

1. Our CEERS samples cover a redshift range of  $2 < z < 9$  and show a marginal ( $3\sigma$ ) correlation between  $[\text{O III}]/\text{H}\beta$  with redshift,  $0.07 \pm 0.03$ . Additionally, there is a marginal  $2\sigma$  correlation with  $[\text{Ne III}]/[\text{O II}]$  with redshift,  $0.05 \pm 0.02$ . When looking at the  $\text{H}\alpha$  SFR, a very strong and significant correlation, having a slope of  $0.18 \pm 0.03$ , with redshift was found.
2. We see a 0.33 dex increase in  $[\text{O III}]/\text{H}\beta$ , a 0.37 dex increase in  $[\text{Ne III}]/[\text{O II}]$ , and a 0.5 dex increase with  $\text{H}\alpha$  SFR when comparing our CEERS  $z > 5$  sample to our  $z \sim 2$  sample from CLEAR.
3. We show  $[\text{O III}]/\text{H}\beta$  and  $[\text{Ne III}]/[\text{O II}]$  both have significant correlations with  $\text{H}\beta$  SFR, with slopes of  $0.2 \pm 0.04$  and  $0.23 \pm 0.04$ , respectively.  $[\text{O III}]/\text{H}\beta$  is also shown to have a significant anticorrelation with stellar mass with a slope of  $-0.41 \pm 0.09$ , while  $[\text{Ne III}]/[\text{O II}]$  has a marginal anticorrelation with a slope of  $-0.2 \pm 0.07$ .
4. The IllustrisTNG simulations match the measurements of our highest  $[\text{O III}]/\text{H}\beta$  values when we compare them to our observed galaxies at similar redshift, SFR, and stellar masses.
5. When comparing our high-redshift  $z > 5$  sample to MAPPINGS V model spectra, we found they are best described by high ionization with moderate metallicity. Comparing the CEERS line ratios with  $z \sim 0$  and  $z \sim 2$  samples indicates that the ISM ionization increases with increasing redshift.

Larger samples of  $z > 5$  galaxies with JWST spectroscopy are needed to better disentangle the relationship between ISM conditions and galaxy properties at cosmic dawn. The NIRCcam WFSS observations are very useful as a blind survey that includes galaxies missed by targeted NIRSpec observations, but the single-filter NIRCcam WFSS observations in CEERS results in a limited wavelength range that includes only a single emission-line ratio for these galaxies. Multi-filter NIRCcam WFSS observations would provide broader wavelength

coverage and more effectively probe the ISM conditions for blind and representative samples of emission-line galaxies in the early Universe.

### Acknowledgments

We acknowledge the work of our colleagues in the CEERS collaboration and everyone involved in the JWST mission. B.E. B. and J.R.T. acknowledge support from NASA grants JWST-ERS-01345 and JWST-AR-01721, and NSF grant CAREER-1945546.

*Software:* AstroPy (Astropy Collaboration et al. 2013), Matplotlib (Hunter 2007), NumPy (van der Walt et al. 2011), SciPy (Virtanen et al. 2020), Linmix Kelly 2007, eazy-py (Brammer et al. 2008).

### ORCID iDs

Bren E. Backhaus <https://orcid.org/0000-0001-8534-7502>  
Jonathan R. Trump <https://orcid.org/0000-0002-1410-0470>  
Nor Pirzkal <https://orcid.org/0000-0003-3382-5941>  
Guillermo Barro <https://orcid.org/0000-0001-6813-875X>  
Steven L. Finkelstein <https://orcid.org/0000-0001-8519-1130>  
Pablo Arrabal Haro <https://orcid.org/0000-0002-7959-8783>  
Raymond C. Simons <https://orcid.org/0000-0002-6386-7299>  
Jessica Wessner <https://orcid.org/0000-0002-4390-1816>  
Nikko J. Cleri <https://orcid.org/0000-0001-7151-009X>  
Micaela B. Bagley <https://orcid.org/0000-0002-9921-9218>  
Michaela Hirschmann <https://orcid.org/0000-0002-3301-3321>  
David C. Nicholls <https://orcid.org/0000-0003-0892-5203>  
Mark Dickinson <https://orcid.org/0000-0001-5414-5131>  
Jeyhan S. Kartaltepe <https://orcid.org/0000-0001-9187-3605>  
Casey Papovich <https://orcid.org/0000-0001-7503-8482>  
Dale D. Kocevski <https://orcid.org/0000-0002-8360-3880>  
Anton M. Koekemoer <https://orcid.org/0000-0002-6610-2048>  
Laura Bisigello <https://orcid.org/0000-0003-0492-4924>  
Anne E. Jaskot <https://orcid.org/0000-0002-6790-5125>  
Ray A. Lucas <https://orcid.org/0000-0003-1581-7825>  
Intae Jung <https://orcid.org/0000-0003-1187-4240>  
Stephen M. Wilkins <https://orcid.org/0000-0003-3903-6935>  
L. Y. Aaron Yung <https://orcid.org/0000-0003-3466-035X>  
Henry C. Ferguson <https://orcid.org/0000-0001-7113-2738>  
Adriano Fontana <https://orcid.org/0000-0003-3820-2823>  
Andrea Grazian <https://orcid.org/0000-0002-5688-0663>  
Norman A. Grogin <https://orcid.org/0000-0001-9440-8872>  
Lisa J. Kewley <https://orcid.org/0000-0001-8152-3943>  
Allison Kirkpatrick <https://orcid.org/0000-0002-1306-1545>  
Jennifer M. Lotz <https://orcid.org/0000-0003-3130-5643>  
Laura Pentericci <https://orcid.org/0000-0001-8940-6768>  
Pablo G. Pérez-González <https://orcid.org/0000-0003-4528-5639>  
Swara Ravindranath <https://orcid.org/0000-0002-5269-6527>  
Rachel S. Somerville <https://orcid.org/0000-0002-6748-6821>  
Guang Yang <https://orcid.org/0000-0001-8835-7722>  
Benne W. Holwerda <https://orcid.org/0000-0002-4884-6756>  
Peter Kurczynski <https://orcid.org/0000-0002-8816-5146>  
Nimish P. Hathi <https://orcid.org/0000-0001-6145-5090>

Caitlin Rose <https://orcid.org/0000-0002-8018-3219>  
Kelcey Davis <https://orcid.org/0000-0001-8047-8351>

### References

- Ahn, C. P., Alexandroff, R., Allende Prieto, C., et al. 2014, *ApJS*, 211, 17  
Aird, J., Nandra, K., Laird, E. S., et al. 2010, *MNRAS*, 401, 2531  
Álvarez-Márquez, J., Colina, L., Crespo Gómez, A., et al. 2023, arXiv:2309.06319  
Arrabal Haro, P., Dickinson, M., Finkelstein, S. L., et al. 2023, *Natur*, 622, 7984  
Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33  
Backhaus, B. E., Trump, J. R., Cleri, N. J., et al. 2022, *ApJ*, 926, 161  
Bagley, M. B., Finkelstein, S. L., Koekemoer, A. M., et al. 2023, *ApJL*, 946, L12  
Baldwin, J. A., Phillips, M. M., & Terlevich, R. 1981, *PASP*, 93, 5  
Bolton, A. S., Schlegel, D. J., Aubourg, É., et al. 2012, *AJ*, 144, 144  
Brammer, G. B., van Dokkum, P. G., & Coppi, P. 2008, *ApJ*, 686, 1503  
Brammer, G. B., van Dokkum, P. G., Franx, M., et al. 2012, *ApJS*, 200, 13  
Brinchmann, J. 2023, *MNRAS*, 525, 2087  
Brinchmann, J., Charlot, S., White, S. D. M., et al. 2004, *MNRAS*, 351, 1151  
Bruzual, G., & Charlot, S. 2003, *MNRAS*, 344, 1000  
Bushouse, H., Eisenhamer, J., Dencheva, N., et al. 2022, JWST Calibration Pipeline v1.8.5, Zenodo, doi:10.5281/zenodo.7429939  
Calzetti, D., Armus, L., Bohlin, R. C., et al. 2000, *ApJ*, 533, 682  
Calzetti, D., & Heckman, T. M. 1999, *ApJ*, 519, 27  
Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245  
Chabrier, G. 2003, *PASP*, 115, 763  
Charlot, S., & Fall, S. M. 2000, *ApJ*, 539, 718  
Cleri, N. J., Yang, G., Papovich, C., et al. 2023a, *ApJ*, 948, 112  
Cleri, N. J., Olivier, G. M., Hutchison, T. A., et al. 2023b, *ApJ*, 953, 10  
Coil, A. L., Aird, J., Reddy, N., et al. 2015, *ApJ*, 801, 35  
Conroy, C., & Gunn, J. E. 2010, *ApJ*, 712, 833  
Conroy, C., Gunn, J. E., & White, M. 2009, *ApJ*, 699, 486  
Curti, M., D'Eugenio, F., Carniani, S., et al. 2023, *MNRAS*, 518, 425  
Curtis-Lake, E., Carniani, S., Cameron, A., et al. 2023, *NatAs*, 7, 622  
Davis, M., Guhathakurta, P., Konidaris, N. P., et al. 2007, *ApJL*, 660, L1  
Del Zanna, G., Dere, K. P., Young, P. R., Landi, E., & Mason, H. E. 2015, *A&A*, 582, A56  
Dere, K. P., Landi, E., Mason, H. E., Monsignor Fossi, B. C., & Young, P. R. 1997, *A&AS*, 125, 149  
Dickey, C. M., van Dokkum, P. G., Oesch, P. A., et al. 2016, *ApJL*, 828, L11  
Eldridge, J. J., & Stanway, E. R. 2009, *MNRAS*, 400, 1019  
Ferland, G. J., Porter, R. L., van Hoof, P. A. M., et al. 2013, *RMxAA*, 49, 137  
Finkelstein, S. L., Bagley, M., Song, M., et al. 2022, *ApJ*, 928, 52  
Finkelstein, S. L., Bagley, M. B., Ferguson, H. C., et al. 2023, *ApJL*, 946, L13  
Fujimoto, S., Kohno, K., Ouchi, M., et al. 2023, arXiv:2303.01658  
Greene, T. P., Kelly, D. M., Stansberry, J., et al. 2017, *JATIS*, 3, 035001  
Grogin, N. A., Kocevski, D. D., Faber, S. M., et al. 2011, *ApJS*, 197, 35  
Guo, Y., Koo, D. C., Lu, Y., et al. 2016, *ApJ*, 822, 103  
Henry, A., Scarlata, C., Domínguez, A., et al. 2013, *ApJL*, 776, L27  
Hirschmann, M., Charlot, S., Feltre, A., et al. 2017, *MNRAS*, 472, 2468  
Hirschmann, M., Charlot, S., & Somerville, R. S. 2023a, *MNRAS*, 526, 3504  
Hirschmann, M., Charlot, S., Feltre, A., et al. 2023b, *MNRAS*, 526, 3610  
Hunter, J. D. 2007, *CSE*, 9, 90  
Jakobsen, P., Ferruit, P., Alves de Oliveira, C., et al. 2022, *A&A*, 661, A80  
Jung, I., Finkelstein, S. L., Arrabal Haro, P., et al. 2023, arXiv:2304.05385  
Kashino, D., Silverman, J. D., Sanders, D., et al. 2019, *ApJS*, 241, 10  
Kauffmann, G., Heckman, T. M., Tremonti, C., et al. 2003a, *MNRAS*, 346, 1055  
Kauffmann, G., Heckman, T. M., White, S. D. M., et al. 2003b, *MNRAS*, 341, 33  
Kauffmann, G., Heckman, T. M., White, S. D. M., et al. 2003c, *MNRAS*, 341, 54  
Kelly, B. C. 2007, *ApJ*, 665, 1489  
Kennicutt, R. C., & Evans, N. J. 2012, *ARA&A*, 50, 531  
Kewley, L. J., Groves, B., Kauffmann, G., & Heckman, T. 2006, *MNRAS*, 372, 961  
Kewley, L. J., Nicholls, D. C., Sutherland, R., et al. 2019, *ApJ*, 880, 16  
Kocevski, D. D., Barro, G., McGrath, E. J., et al. 2023, *ApJL*, 946, L14  
Koekemoer, A. M., Faber, S. M., Ferguson, H. C., et al. 2011, *ApJS*, 197, 36  
Kokorev, V., Fujimoto, S., Labbé, I., et al. 2023, *ApJL*, 957, L7  
Kriek, M., van Dokkum, P. G., Labbé, I., et al. 2009, *ApJ*, 700, 221



- Kriek, M., Shapley, A. E., Reddy, N. A., et al. 2015, *ApJS*, **218**, 15
- Kroupa, P. 2001, *MNRAS*, **322**, 231
- Larson, R. L., Finkelstein, S. L., Kocevski, D. D., et al. 2023, *ApJL*, **953**, L29
- Leitherer, C., Schaerer, D., Goldader, J. D., et al. 1999, *ApJS*, **123**, 3
- Liu, X., Shapley, A. E., Coil, A. L., Brinchmann, J., & Ma, C.-P. 2008, *ApJ*, **678**, 758
- Madau, P., & Dickinson, M. 2014, *ARA&A*, **52**, 415
- McLean, I. S., Steidel, C. C., Epps, H. W., et al. 2012, *Proc. SPIE*, **8446**, 84460J
- Momcheva, I. G., Brammer, G. B., van Dokkum, P. G., et al. 2016, *ApJS*, **225**, 27
- Montero-Dorta, A. D., Bolton, A. S., Brownstein, J. R., et al. 2016, *MNRAS*, **461**, 1131
- Osterbrock, D. E. 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei* (Sausalito, CA: Univ. Science Books)
- Papovich, C., Simons, R. C., Estrada-Carpenter, V., et al. 2022, *ApJ*, **937**, 22
- Pharo, J., Guo, Y., Koo, D. C., Forbes, J. C., & Guhathakurta, P. 2023, *ApJL*, **946**, L5
- Pirzkal, N., Malhotra, S., Ryan, R. E., et al. 2017, *ApJ*, **846**, 84
- Planck Collaboration, Ade, P. A. R., Aghanim, N., et al. 2016, *A&A*, **594**, A13
- Pontoppidan, K. M., Barrientes, J., Blome, C., et al. 2022, *ApJL*, **936**, L14
- Reddy, N. A., Steidel, C. C., Pettini, M., & Bogosavljević, M. 2016, *ApJ*, **828**, 107
- Salpeter, E. E. 1955, *ApJ*, **121**, 161
- Sanders, R. L., Shapley, A. E., Topping, M. W., Reddy, N. A., & Brammer, G. B. 2023, *ApJ*, **955**, 54
- Sanders, R. L., Shapley, A. E., Topping, M. W., Reddy, N. A., & Brammer, G. B. 2024, *ApJ*, **962**, 24
- Sanders, R. L., Shapley, A. E., Jones, T., et al. 2021, *ApJ*, **914**, 19
- Schaerer, D., Marques-Chaves, R., Barrufet, L., et al. 2022, *A&A*, **665**, L4
- Shapley, A. E., Reddy, N. A., Sanders, R. L., Topping, M. W., & Brammer, G. B. 2023a, *ApJL*, **950**, L1
- Shapley, A. E., Sanders, R. L., Reddy, N. A., Topping, M. W., & Brammer, G. B. 2023b, *ApJ*, **954**, 157
- Shapley, A. E., Reddy, N. A., Kriek, M., et al. 2015, *ApJ*, **801**, 88
- Shapley, A. E., Sanders, R. L., Shao, P., et al. 2019, *ApJL*, **881**, L35
- Simons, R. C., Papovich, C., Momcheva, I. G., et al. 2023, *ApJS*, **266**, 13
- Smee, S. A., Gunn, J. E., Uomoto, A., et al. 2013, *AJ*, **146**, 32
- Speagle, J. S., Steinhardt, C. L., Capak, P. L., & Silverman, J. D. 2014, *ApJS*, **214**, 15
- Steidel, C. C., Strom, A. L., Pettini, M., et al. 2016, *ApJ*, **826**, 159
- Steidel, C. C., Rudie, G. C., Strom, A. L., et al. 2014, *ApJ*, **795**, 165
- Strom, A. L., Steidel, C. C., Rudie, G. C., Trainor, R. F., & Pettini, M. 2018, *ApJ*, **868**, 117
- Sutherland, R., Dopita, M., Binette, L., & Groves, B. 2018, MAPPINGS V: Astrophysical plasma modeling code, Astrophysics Source Code Library, ascl:1807.005
- Tang, M., Stark, D. P., Chen, Z., et al. 2023, *MNRAS*, **526**, 1657
- Trump, J. R., Arrabal Haro, P., Simons, R. C., et al. 2023, *ApJ*, **945**, 35
- van der Walt, S., Colbert, S. C., & Varoquaux, G. 2011, *CSE*, **13**, 22
- van Dokkum, P. G., Brammer, G., Fumagalli, M., et al. 2011, *ApJL*, **743**, L15
- Veilleux, S., & Osterbrock, D. E. 1987, *ApJS*, **63**, 295
- Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, *NatMe*, **17**, 261
- York, D. G., Adelman, J., Anderson, J. E. J., et al. 2000, *AJ*, **120**, 1579
- Zeimann, G. R., Ciardullo, R., Gebhardt, H., et al. 2015, *ApJ*, **798**, 29