


PAPER



Cite this: DOI: 10.1039/d4dt01376j

Ammonium halide selective ion pair recognition and extraction with a chalcogen bonding heteroditopic receptor†

Andrew Docker,^a Yuen Cheong Tse,^a Hui Min Tay,^b Zongyao Zhang^b and Paul D. Beer *^b

The first example of a heteroditopic receptor capable of cooperative recognition and extraction of ammonium salt (NH₄X) ion-pairs is described. Consisting of a bidentate 3,5-bis-tellurotriazole chalcogen bond donor binding cleft, the appendage of benzo-15-crown-5 (B15C5) substituents to the tellurium centres facilitates binding of the ammonium cation *via* a co-facial bis-B15C5 sandwich complex, which serves to switch on chalcogen bonding-mediated anion binding potency. Extensive quantitative ion-pair recognition ¹H NMR titration studies in CD₃CN/CDCl₃ (1 : 1, v/v) solvent media reveal impressive ion-pair binding affinities towards a variety of ammonium halide, nitrate and thiocyanate salts, with the heteroditopic receptor displaying notable ammonium halide salt selectivity. The prodigious solution phase NH₄X recognition also translates to efficient solid–liquid and liquid–liquid extraction capabilities.

Received 10th May 2024,

Accepted 6th June 2024

DOI: 10.1039/d4dt01376j

rsc.li/dalton

Introduction

The ammonium cation (NH₄⁺) is implicated in pivotal roles across various domains from biological metabolism^{1–4} to environmental monitoring,^{5–9} and industrial processes.¹⁰ Indeed, in biotic systems NH₄⁺ is a crucial intermediate in nitrogen metabolism, amino acid synthesis and pH regulation,¹¹ whilst, from an anthropogenic perspective NH₄⁺ constitutes a key component in agricultural fertiliser, and a crucial reagent in chemical manufacturing.¹² Considering the importance of ammonium, it is surprising that more effort has not been directed towards designing receptors capable of its molecular recognition,¹³ wherein the majority of reports to date primarily rely on tripodal pyrazolyl- or cryptand-based host systems.^{14–21} The development of heteroditopic receptors for ion pair recognition *via* the simultaneous binding of a cation and an anion has proven a powerful strategy in augmenting ion affinity and selectivity profiles of supramolecular host systems.^{22–24} Typically such ditopic hosts target alkali metal salts, employing a crown ether motif for metal cation binding covalently linked to hydrogen bond donors for recognition of the counter-anion guest species.

In recent decades the supramolecular toolbox for anion recognition has been expanded to include sigma (σ)-hole type interactions,^{25,26} with halogen bonding (XB)^{27–30} and chalcogen bonding (ChB)^{31–34} monotopic host systems commonly exhibiting remarkable anion binding affinity enhancements and unique selectivity behaviours relative to more traditionally employed hydrogen bonding based receptors.^{35–41} Despite this, the integration of sigma (σ)-hole interactions into heteroditopic host structural design remains rare.^{42–44} In light of these advantages and paucity of receptors targeting ammonium ion-pairs,⁴⁵ we sought to apply our recent report of a ChB heteroditopic ion-pair receptor,⁴⁶ **1-ChB^{PFP}**, consisting of a 3,5-bis-tellurotriazole nitro-benzene central scaffold functionalised with electron-deficient perfluorophenyl substituents and benzo-15-crown-5 appended telluro-triazoles (Fig. 1), for the purpose of ammonium ion-pair (NH₄X) recognition. Herein, we report to the best of our knowledge the first example of a heteroditopic receptor capable of cooperative solution phase recognition of NH₄X ion-pairs, where extensive quantitative ion-pair affinity measurements demonstrate considerable selectivity towards ammonium halides. Crucially, the anion affinity of the receptor relies on NH₄⁺ complexation *via* an intramolecular co-facial sandwich complex by the B15C5 units which not only conformationally preorganises the tellurotriazole ChB donors, but also switches on Te σ-hole Lewis acidity for anion recognition. This prodigious cooperative ion-pair recognition behaviour of **1-ChB^{PFP}** is further exploited for successful ammonium salt solid–liquid and liquid–liquid extraction purposes.

^aYusuf Hamied Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, UK

^bChemistry Research Laboratory, Department of Chemistry, University of Oxford, Mansfield Road, Oxford OX1 3TA, UK. E-mail: paul.beer@chem.ox.ac.uk

† Electronic supplementary information (ESI) available. CCDC 2155965. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d4dt01376j>




Fig. 1 Ammonium salt (NH_4X) binding chalcogen bonding heteroditopic receptor $1\text{-ChB}^{\text{PFP}}$.

Results and discussion

Anion and ion-pair recognition studies

It is well known that B15C5 is capable of forming 2 : 1 host-guest stoichiometric complexes with alkali metal cations K^+ ,

Rb^+ , Cs^+ ,^{47–49} and similar sandwich complex formation has also been reported with NH_4^+ .⁵⁰ Motivated by this, we sought to determine whether $1\text{-ChB}^{\text{PFP}}$ could bind an ammonium cation in an analogous intramolecular manner between the two pendant B15C5 units and thereby potentially function as a receptor for NH_4X ion-pairs (Fig. 2a). To this end, a qualitative ^1H NMR titration experiment was initially conducted, wherein to a $\text{CD}_3\text{CN}/\text{CDCl}_3$ (1 : 1, v/v) solution of $1\text{-ChB}^{\text{PFP}}$ was added an equimolar amount of solid NH_4PF_6 . Upon comparison of the ^1H NMR spectra of pre- and post-ammonium salt addition, dramatic perturbations and broadening of the resonances associated with the crown ether aromatic and methylene regions, namely signals c, d, e, f and h respectively were observed relative to the free receptor (Fig. 2b). Specifically, the dramatic *ca.* 1 and 0.5 ppm upfield shifts of CH_2 signals f and h of the polyether chain, evidence a strong shielding effect from a proximal aromatic ring current, supporting the formation of a cofacial NH_4^+ bis-B15C5 sandwich complex and is wholly consistent with previously reported diagnostic chemical shift changes associated with this type of complex.^{47–49} The sequential addition of further equivalents of NH_4PF_6 elicited no further changes in the ^1H NMR spectrum, indicative of 1 : 1 $\text{NH}_4^+ : 1\text{-ChB}^{\text{PFP}}$ complex stoichiometry and the association of ammonium cation to the B15C5 units is of considerable




Fig. 2 (a) NH_4X $1\text{-ChB}^{\text{PFP}}$ binding equilibria. ^1H NMR titration experiments of (b) NH_4PF_6 and $1\text{-ChB}^{\text{PFP}}$ (c) TBABr and $1\text{-ChB}^{\text{PFP}}$ in the presence of 1 equivalent of NH_4PF_6 ($\text{CD}_3\text{CN}/\text{CDCl}_3$ 1 : 1 (v/v), 500 MHz, 298 K).

strength ($K_a > 10^4 \text{ M}^{-1}$). Furthermore, the relatively minor perturbations observed in the nitro phenyl signals a and b suggest minimal perturbation of the ChB binding cleft, as anticipated from the non-coordinating nature of the hexafluorophosphate counter-anion.

Encouraged by this strong evidence for NH_4^+ sandwich complexation, we investigated the ammonium salt ion-pair recognition properties of $1\text{-ChB}^{\text{PFP}}$. To this end, a series of ^1H NMR anion titration experiments were conducted on a $\text{CD}_3\text{CN}/\text{CDCl}_3$ (1 : 1, v/v) solution of $1\text{-ChB}^{\text{PFP}}$ in the presence of equimolar NH_4PF_6 . The addition of increasing equivalents of tetrabutylammonium halide, nitrate and thiocyanate salts all induced progressive downfield shifts of the heteroditopic receptor's internal aromatic proton signal b, providing strong evidence for the participation of the tellurotriazole ChB donors mediating the anion recognition process (a representative example for bromide is shown in Fig. 2c). During the course of anion addition, it was noted that characteristic features of the NH_4^+ bis-B15C5 sandwich complex in the ^1H NMR spectrum persisted, indicating that the anion binding and cation binding events occur concomitantly *i.e.* genuine ion-pair binding. Monitoring proton b, Bindfit⁵¹ analysis of the resulting anion-induced chemical shift perturbation isotherm titration data (Fig. 3) determined 1 : 1 stoichiometric host/guest apparent association constants (K_a)⁵² for a range of halides and polyatomic anions. Table 1 shows the co-bound ammonium complex receptor ($1\text{-ChB}^{\text{PFP}}\cdot\text{NH}_4^+$) displays strong halide affinities, with a particularly impressive affinity for chloride with $K_a(\text{Cl}^-) = 2530 \text{ M}^{-1}$, notably greater than both $K_a(\text{Br}^-)$ and $K_a(\text{I}^-)$ by at least a factor of two. Interestingly, relative to the halides the affinities for nitrate and thiocyanate are considerably diminished ($K_a(\text{NO}_3^-) = 357 \text{ M}^{-1}$ and $K_a(\text{SCN}^-) = 294 \text{ M}^{-1}$). It is also noteworthy that whilst anion affinity trends in simple acyclic HB based receptors are typically governed by the anion's inherent basicity, usually correlated with $\text{p}K_a$ values, this is not observed for $1\text{-ChB}^{\text{PFP}}$, and empirically appears to be a consistent feature of sigma-hole based anion receptors.⁵³




Fig. 3 Anion-binding isotherms for $1\text{-ChB}^{\text{PFP}}$ in the presence of 1 equivalent of NH_4PF_6 ($\text{CD}_3\text{CN}/\text{CDCl}_3$ 1 : 1 (v/v), 500 MHz, 298 K).

Table 1 Anion association constants for $1\text{-ChB}^{\text{PFP}}$ from ^1H NMR titration experiments (1 : 1 $\text{CD}_3\text{CN}/\text{CDCl}_3$ (v/v), 500 MHz, 298 K)

Anion association constant (K_a , M^{-1}) of $1\text{-ChB}^{\text{PFP}}$ in the presence of equimolar NH_4PF_6 ^{a,b}	Anion $\text{p}K_a$	
Cl^-	2530	−8.0
Br^-	1140	−9.0
I^-	1130	−10
NO_3^-	357	−1.3
SCN^-	294	4.0

^a Determined from Bindfit analysis, monitoring signal b, error <5%.

^b Anions added as their tetrabutylammonium salts.

Interestingly, the co-bound ammonium complex receptor $1\text{-ChB}^{\text{PFP}}\cdot\text{NH}_4^+$ halide association constant values are of significantly larger magnitude than those determined with the potassium complex $1\text{-ChB}^{\text{PFP}}\cdot\text{K}^+$ ⁴⁶ with a particularly notable 2-fold enhancement in chloride affinity. Whilst the exact origin of this enhancement is not definitively known, it is postulated the $\text{N-H}\cdots\text{O}$ hydrogen bonding interactions formed between NH_4^+ and the crown ether oxygens more effectively electronically polarise the Te ChB donor centres than a potassium cation thereby raising anion binding potency. Importantly, in the absence of NH_4PF_6 , $1\text{-ChB}^{\text{PFP}}$ exhibited no measurable anion binding affinity, thereby confirming the crucial role of bis-B15C5 sandwich bound NH_4^+ in switching on ChB mediated anion recognition *via* favourable proximal electrostatic interactions and preorganised through bond polarisation of the Te sigma-hole donors.

Solid state single-crystal X-ray diffraction study of NH_4Br ion-pair complex

Further insight into the ammonium halide salt ion-pair recognition mode of $1\text{-ChB}^{\text{PFP}}$ in the solid state was provided by single crystal diffraction X-ray analysis of $1\text{-ChB}^{\text{PFP}}$ complexed with NH_4Br (Fig. 4). Consistent with the ^1H NMR solution phase evidence, the ammonium cation is complexed *via* a cova-




Fig. 4 Solid-state structure of $1\text{-ChB}^{\text{PFP}}$ complexed with NH_4Br (solvent molecules and hydrogen atoms, except those of NH_4^+ , are omitted for clarity). Grey = carbon, blue = nitrogen, red = oxygen, light green = fluorine, orange = tellurium and dark red = bromine.

cial B15C5 sandwich complex. However, in contrast to the structure of alkali metal cation (M^+) complexes previously obtained for $1\text{-ChB}^{\text{PFP}}$,⁴⁶ there are notable differences. Specifically, in the M^+ crystal structures the orientation of the crown ether rings appear relatively symmetric, typically allowing all five oxygen atoms of each B15C5 unit to coordinate the cation. In contrast, the somewhat skewed conformations of the B15C5 units observed in the ammonium bromide complex of $1\text{-ChB}^{\text{PFP}}$ seem to suggest a less symmetric NH_4^+ complexation mode.⁴⁶ This is presumably due to the formation of directional hydrogen bonds between the ammonium and crown ether oxygens; $\text{N-H}\cdots\text{O}$. The Br^- counteranion is shown to be chelated, moderately asymmetrically, by bifurcated chalcogen bond formation, exhibiting short $\text{Te}\cdots\text{Br}$ contacts of 3.241 Å and 3.583 Å, corresponding to contractions in their van der Waal radii of 83% and 92% respectively. Determination of the $\text{C-Te}\cdots\text{Br}$ angles; 174° and 167° reveals a commonly observed preference for ChB interactions in which the preferential bonding geometry approaches linearity.

Ammonium salt solid-liquid and liquid-liquid ion-pair extraction studies

Motivated by the impressive ion-pair affinity of $1\text{-ChB}^{\text{PFP}}$ for ammonium salts, as evidenced by quantitative solution phase ^1H NMR binding studies, attention was directed toward investigating the heteroditopic receptor's potential as an extraction agent for NH_4X salts under solid-liquid (SLE) and liquid-liquid extraction (LLE) conditions. In a typical SLE experiment, a CDCl_3 solution of $1\text{-ChB}^{\text{PFP}}$ was exposed to a 5-fold molar excess microcrystalline solid sample of NH_4X ($\text{X} = \text{Cl}^-$, Br^- , I^- , NO_3^- , SCN^-) and stirred for 10 minutes (Fig. 5a). With the

exception of NH_4Cl , the ^1H NMR spectrum of the resultant post-extraction solution in general revealed dramatic changes relative to the pre-extraction spectrum and closely resembled spectroscopic features observed during the solution phase ion-pair titration experiments with NH_4PF_6 and TBAX (Fig. 5b). Specifically, in the case of NH_4Br , NH_4I , NH_4NO_3 and NH_4SCN , dramatic broadening of the methylene and aromatic signals of B15C5 units, indicative of the NH_4^+ sandwich complexation binding mode and downfield perturbations of the internal aromatic signal b, consistent with ChB mediated anion binding, were observed. It was also noted a new signal appeared presenting as a triplet consistent with the heteronuclear spin-spin coupling between ^1H and ^{14}N of NH_4^+ , further confirming successful extraction of these NH_4X salts into CDCl_3 . The extraction efficiency for each NH_4X salt was estimated by integration of the N-H signals of NH_4^+ relative to signal a of $1\text{-ChB}^{\text{PFP}}$, thereby determining the ratio of NH_4X extracted by $1\text{-ChB}^{\text{PFP}}$, the results of which are summarised in Table 2, together with graphical presentation shown in Fig. 6. The efficiency of a host to perform SLE is a subtle balance

Table 2 Solid-liquid extraction efficiencies of $1\text{-ChB}^{\text{PFP}}$

Ammonium salt	NH_4Cl	NH_4Br	NH_4I	NH_4NO_3	NH_4SCN
Extraction ^a	0%	85%	>95%	~5% ^b	>95%
$\Delta H_f/\text{kJ mol}^{-1}$	705	647	608	646	597

^a Determined from relative integration of proton signals a of $1\text{-ChB}^{\text{PFP}}$ and of NH_4^+ , error estimated at $\pm 5\%$. ^b Despite evident extraction of NH_4NO_3 from the post SLE ^1H NMR spectrum the very low signal intensity of the co-extracted NH_4^+ precluded precise extraction percentage determination.




Fig. 5 (a) Representative SLE equilibrium of ammonium salt (NH_4X) by $1\text{-ChB}^{\text{PFP}}$ and cartoon representation of the SLE process (b) Pre and post-SLE ^1H NMR spectra (CDCl_3 , 500 MHz, 298 K), with various ammonium salts.




Fig. 6 Plot showing extraction efficiency (blue) and lattice energy ΔH_L (red) versus NH_4X ($\text{X} = \text{Cl}, \text{Br}, \text{I}, \text{NO}_3$).

between two competing factors: the affinity of the host for the ion-pair and the lattice enthalpy (ΔH_L) of the salt to be extracted. In the former, a higher ion-pair binding affinity usually translates to improved SLE (or LLE) extraction performance, whilst in the latter a larger ΔH_L is energetically unfavourable to extraction performance. Inspection of Table 2 reveals that whilst NH_4I and NH_4SCN are extracted with the highest efficiencies >95%, as expected on the basis of their low ΔH_L values, interestingly, in stark contrast, the extraction efficiencies observed for NH_4Br and NH_4NO_3 , are 85% and ~5% respectively. Considering their near identical lattice enthalpies; $\Delta H_L(\text{NH}_4\text{Br}) = 647 \text{ kJ mol}^{-1}$ and $\Delta H_L(\text{NH}_4\text{NO}_3) = 646 \text{ kJ mol}^{-1}$, it is apparent the 3-fold enhancement in solution phase ion-pair affinity for $K_a(\text{Br}^-)$ relative to $K_a(\text{NO}_3^-)$ notably translates to an improved extraction capability providing strong evidence for **1-ChB^{PPF}** functioning as a genuine ion-pair receptor. However, despite **1-ChB^{PPF}** exhibiting the largest NH_4Cl ion-pair affinity from solution phase experiments (Table 2), no evidence of SLE was noted, suggesting the appreciable magnitude of $\Delta H_L(\text{NH}_4\text{Cl})$ dominates in this case.

Liquid-liquid extraction (LLE) experiments were undertaken for ammonium chloride, bromide and iodide. In a typical LLE experiment a CDCl_3 solution of **1-ChB^{PPF}** (2 mM) was exposed to an ammonium halide D_2O solution (4 M), stirred vigorously for 30 minutes and ^1H NMR spectrum of the post-extraction CDCl_3 organic phase recorded. As for the SLE experiment, while for NH_4Cl no receptor proton perturbations were observed, in the case of NH_4Br and NH_4I a comparison of the pre- and post-extraction spectra revealed successful extraction of the ammonium halides as evidenced by similar proton perturbations to those observed in the SLE experiments (Fig. S7†). However, unlike in the SLE experiment, the signal corresponding to the co-extracted NH_4^+ was significantly broadened or not observable, presumably due to deuterium-proton isotope exchange from the deuterated aqueous source phase, which prevented quantitative determination of LLE efficiencies.

Conclusions

In summary, the unprecedented cooperative ion-pair recognition of ammonium salt ion-pair species (NH_4X) is achieved by a heteroditopic receptor **1-ChB^{PPF}**. Exploiting the bis-telluro-triazole ChB donor framework, wherein the Te-centres are directly appended with B15C5 units, co-facial intramolecular bis-B15C5 NH_4^+ sandwich complex formation not only preorganises the receptor's ChB donor groups, but also effectively serves to enhance and switch on the Lewis acidity of the Te-centres for anion recognition. Quantitative ^1H NMR binding studies demonstrate prodigious NH_4X ion-pair binding properties, highlighting a significant selectivity preference for ammonium halide salts over NH_4NO_3 and NH_4SCN . Solid state X-ray structural analysis of the **1-ChB^{PPF}**· NH_4Br ion-pair complex supports the postulated bis-crown ether NH_4^+ sandwich binding mode, concomitant with strong $\text{ChB}\cdots\text{Br}^-$ interactions driving anion coordination. The notable solution phase NH_4X ion-pair binding properties of **1-ChB^{PPF}** are also reflected in the heteroditopic receptor's efficient solid-liquid and liquid-liquid extraction NH_4Br and NH_4I salt capabilities. Importantly, these results highlight the exciting potential of sigma (σ)-hole based heteroditopic host structural design for the future development of ammonium salt selective ion-pair recognition applications.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

A. D. thanks EPSRC for studentship funding (Grant reference number EP/N509711/1), Y. C. T. thanks the Croucher Foundation for a scholarship. H. M. T. acknowledges the Clarendon Fund and the Oxford Australia Scholarships Fund for a Postgraduate Research Scholarship.

References

- 1 S. S. Mohiuddin and D. Khattar, in *StatPearls*, StatPearls Publishing, Treasure Island (FL), 2024.
- 2 R. G. Wetzel, in *Limnology*, ed. R.G. Wetzel, Academic Press, San Diego, 3rd edn, 2001, pp. 205–237.
- 3 D. S. Dimski, *J. Vet. Intern. Med.*, 1994, **8**, 73–78.
- 4 S. M. Howitt and M. K. Udvardi, *Biochim. Biophys. Acta, Biomembr.*, 2000, **1465**, 152–170.
- 5 G. D. Boardman, S. M. Starbuck, D. B. Hudgins, X. Li and D. D. Kuhn, *Environ. Toxicol.*, 2004, **19**, 134–142.
- 6 D. Li, X. Xu, Z. Li, T. Wang and C. Wang, *Trends Anal. Chem.*, 2020, **127**, 115890.
- 7 R. Temino-Boes, R. Romero-López and I. Romero, *Water*, 2019, **11**, 2143.

- 8 R. Temino-Boes, R. García-Bartual, I. Romero and R. Romero-Lopez, *J. Environ. Manage.*, 2021, **282**, 111739.
- 9 J. Walker, D. Nelson and V. P. Aneja, *Environ. Sci. Technol.*, 2000, **34**, 3527–3534.
- 10 B. L. Bodirsky, A. Popp, H. Lotze-Campen, J. P. Dietrich, S. Rolinski, I. Weindl, C. Schmitz, C. Müller, M. Bonsch, F. Humpeöder, A. Biewald and M. Stevanovic, *Nat. Commun.*, 2014, **5**, 3858.
- 11 I. D. Weiner and J. W. Verlander, *Physiol. Rev.*, 2017, **97**, 465–494.
- 12 G. J. Leigh, in *Catalysts for Nitrogen Fixation: Nitrogenases, Relevant Chemical Models and Commercial Processes*, ed. B. E. Smith, R. L. Richards and W. E. Newton, Springer Netherlands, Dordrecht, 2004, pp. 33–54.
- 13 A. Späth and B. König, *Beilstein J. Org. Chem.*, 2010, **6**, 32.
- 14 A. Pazik and A. Skwierawska, *Sens. Actuators, B*, 2014, **196**, 370–380.
- 15 T. M. Jonah, L. Mathivathanan, A. N. Morozov, A. M. Mebel, R. G. Raptis and K. Kavallieratos, *New J. Chem.*, 2017, **41**, 14835–14838.
- 16 N. Koch, W. Seichter and M. Mazik, *Tetrahedron*, 2015, **71**, 8965–8974.
- 17 S. Y. Jon, J. Kim, M. Kim, S.-H. Park, W. S. Jeon, J. Heo and K. Kim, *Angew. Chem., Int. Ed.*, 2001, **40**, 2116–2119.
- 18 A. Rueda-Zubiaurre, N. Herrero-García, M. del Rosario Torres, I. Fernández and J. Osío Barcina, *Chem. – Eur. J.*, 2012, **18**, 16884–16889.
- 19 K.-M. Park, H. J. Kim, S.-H. Moon, J. J. Vittal, J. H. Jung and S. S. Lee, *New J. Chem.*, 2010, **34**, 603–606.
- 20 B. Metz, J. M. Rosalky and R. Weiss, *J. Chem. Soc., Chem. Commun.*, 1976, 533b–5534.
- 21 X. Wang, O. Shyshov, M. Hanževački, C. M. Jäger and M. von Delius, *J. Am. Chem. Soc.*, 2019, **141**, 8868–8876.
- 22 Q. He, G. I. Vargas-Zúñiga, S. H. Kim, S. K. Kim and J. L. Sessler, *Chem. Rev.*, 2019, **119**, 9753–9835.
- 23 A. J. McConnell, A. Docker and P. D. Beer, *ChemPlusChem*, 2020, **85**, 1824–1841.
- 24 S. Roelens, A. Vacca, O. Francesconi and C. Venturi, *Chem. – Eur. J.*, 2009, **15**, 8296–8302.
- 25 J. Y. C. Lim and P. D. Beer, *Chem*, 2018, **4**, 731–783.
- 26 M. S. Taylor, *Coord. Chem. Rev.*, 2020, **413**, 213270.
- 27 M. G. Sarwar, B. Dragisic, S. Sagoo and M. S. Taylor, *Angew. Chem., Int. Ed.*, 2010, **49**, 1674–1677.
- 28 B. Chowdhury, S. Sinha and P. Ghosh, *Chem. – Eur. J.*, 2016, **22**, 18051–18059.
- 29 C. J. Massena, N. B. Wageling, D. A. Decato, E. Martin Rodriguez, A. M. Rose and O. B. Berryman, *Angew. Chem., Int. Ed.*, 2016, **55**, 12398–12402.
- 30 R. Tepper, B. Schulze, M. Jäger, C. Friebe, D. H. Scharf, H. Görls and U. S. Schubert, *J. Org. Chem.*, 2015, **80**, 3139–3150.
- 31 A. Docker, C. H. Guthrie, H. Kuhn and P. D. Beer, *Angew. Chem., Int. Ed.*, 2021, **60**, 21973–21978.
- 32 G. E. Garrett, E. I. Carrera, D. S. Seferos and M. S. Taylor, *Chem. Commun.*, 2016, **52**, 9881–9884.
- 33 S. Akbaba, T. Steinke, L. Vogel, E. Engelage, M. Erdelyi and S. M. Huber, *Chem. – Eur. J.*, 2024, e202400608.
- 34 A. Docker, T. G. Johnson, H. Kuhn, Z. Zhang and M. J. Langton, *J. Am. Chem. Soc.*, 2023, **145**, 2661–2668.
- 35 A. Docker, X. Shang, D. Yuan, H. Kuhn, Z. Zhang, J. J. Davis, P. D. Beer and M. J. Langton, *Angew. Chem., Int. Ed.*, 2021, **60**, 19442–19450.
- 36 A. J. Taylor, A. Docker and P. D. Beer, *Chem. – Asian J.*, 2023, **18**, e202201170.
- 37 A. Docker, Y. C. Tse, H. M. Tay, A. J. Taylor, Z. Zhang and P. D. Beer, *Angew. Chem., Int. Ed.*, 2022, **61**, e202214523.
- 38 A. Borissov, I. Marques, J. Y. C. Lim, V. Félix, M. D. Smith and P. D. Beer, *J. Am. Chem. Soc.*, 2019, **141**, 4119–4129.
- 39 D. Mungalpara, S. Stegmüller and S. Kubik, *Chem. Commun.*, 2017, **53**, 5095–5098.
- 40 T. G. Johnson, A. Docker, A. Sadeghi-Kelishadi and M. J. Langton, *Chem. Sci.*, 2023, **14**, 5006–5013.
- 41 A. Docker, J. G. Stevens and P. D. Beer, *Chem. – Eur. J.*, 2021, **27**, 14600–14604.
- 42 P. Sabater, F. Zapata, B. López, I. Fernández, A. Caballero and P. Molina, *Dalton Trans.*, 2018, **47**, 15941–15947.
- 43 T. Bunchuay, A. Docker, U. Eiamprasert, P. Surawatanawong, A. Brown and P. D. Beer, *Angew. Chem., Int. Ed.*, 2020, **59**, 12007–12012.
- 44 A. Mele, P. Metrangolo, H. Neukirch, T. Pilati and G. Resnati, *J. Am. Chem. Soc.*, 2005, **127**, 14972–14973.
- 45 Z. Kokan and M. J. Chmielewski, *J. Am. Chem. Soc.*, 2018, **140**, 16010–16014.
- 46 A. Docker, I. Marques, H. Kuhn, Z. Zhang, V. Félix and P. D. Beer, *J. Am. Chem. Soc.*, 2022, **144**, 14778–14789.
- 47 L. A. Lapkina, A. A. Sinelshchikova, K. P. Birin, V. E. Larchenko, M. S. Grigoriev, A. Yu. Tsivadze and Y. G. Gorbunova, *Inorg. Chem.*, 2021, **60**, 1948–1956.
- 48 A. G. Martynov, M. A. Polovkova, G. S. Bereznoy, A. A. Sinelshchikova, F. M. Dolgushin, K. P. Birin, G. A. Kirakosyan, Y. G. Gorbunova and A. Yu. Tsivadze, *Inorg. Chem.*, 2020, **59**, 9424–9433.
- 49 S. P. Gromov, A. I. Vedernikov, N. A. Lobova, L. G. Kuz'mina, S. S. Basok, Y. A. Strelenko, M. V. Alfimov and J. A. K. Howard, *New J. Chem.*, 2011, **35**, 724–737.
- 50 K. M. Doxsee, P. E. Francis and T. J. R. Weakley, *Tetrahedron*, 2000, **56**, 6683–6691.
- 51 <https://supramolecular.org>.
- 52 The determined anion binding constants (K_a) are apparent as the 1:1 stoichiometric host–guest binding model of Bindfit assumes quantitative and complete NH_4^+ complexation, which is justified by the observed experimental results.
- 53 L. Chen, S. N. Berry, X. Wu, E. N. W. Howe and P. A. Gale, *Chem*, 2020, **6**, 61–141.