



Citation for published version:

Lambert, J, Taylor, A, Streeter, A, Greaves, C, Ingram, WM, Dean, S, Jolly, K, Mutrie, N, Price, L & Campbell, J 2023, 'Adding web-based support to exercise referral schemes improves symptoms of depression in people with elevated depressive symptoms: A secondary analysis of the e-coachER randomised controlled trial', *Mental Health and Physical Activity*, vol. 25, 100535. <https://doi.org/10.1016/j.mhpa.2023.100535>

DOI:

[10.1016/j.mhpa.2023.100535](https://doi.org/10.1016/j.mhpa.2023.100535)

Publication date:

2023

Document Version

Peer reviewed version

[Link to publication](#)

Publisher Rights

CC BY-NC-ND

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Background

Depression and anxiety are common mood disorders, with 7.8% of people in the UK estimated to meet clinical diagnostic criteria (NICE, 2018) with a substantial proportion of mental health care taking place within primary care (Park & Zarate, 2019). Depression is also co-morbid with a range of chronic physical conditions including type 2 diabetes, hypertension, osteoarthritis, and obesity (Pilling et al., 2009), so promoting physical activity to patients has multiple benefits (Cooney, 2018). For patients with depression, physical activity is an evidence-based intervention, which is just as effective as anti-depressants and psychological therapies, with the added benefit of treating many chronic physical conditions (Cooney et al., 2013; Heissel et al., 2023; Rebar et al., 2015; Stubbs et al., 2017). Physical activity has also been shown to be an effective treatment for anxiety (Heissel et al., 2023).

Providing support to increase physical activity within primary care is challenging, especially among patients with poor mental health. Patients with poor mental health often have unique barriers to engaging in physical activity (e.g., low mood, pain, fear, low self-efficacy (perceived ability to be active) and low motivation (Busch et al., 2016; Rebar & Taylor, 2017). Behavioural and psychological support to increase physical activity within primary care can be effective (Harris et al., 2015) but often GPs are unlikely to have enough time to deliver the support needed (Lowe et al., 2022), especially if patients have multiple physical and mental health conditions. Group-based physical activity is also currently recommended by the National Institute for Health and Care Excellence (NICE, 2022) for depression, but not for anxiety (NICE, 2011).

Exercise Referral Schemes (ERS), historically hosted in leisure centres and gyms and more recently through behavioural community support, are common within UK primary care for patients with physical and mental health conditions. There is currently no single model for ERSs in the UK, but the predominant mode of delivery involves 10–12 weeks of structured, supervised exercise at a local exercise facility or a counselling approach to support patients to engage in a variety of types of PA (Taylor et al., 2020). A recent study found that at least

half of those attending exercise referral schemes were referred for low mood (Taylor et al., 2021). Exercise referral schemes are also potentially effective for people with mental health conditions: A systematic review of randomised controlled trials (RCTs) found large effects size in favour of exercise referral schemes for depression (pooled standardised mean difference -0.82 , 95% CI -1.28 to -0.35) when compared to usual care (Pavey et al., 2011). However, another recent review found that uptake and adherence to exercise referral schemes tended to be lower for those referred for mental health conditions (Tomlinson-Perez et al., 2022). One way to overcome the challenge of adherence in people with mental health conditions is through theory-informed approaches.

Self Determination Theory (SDT), a theory of human motivation, posits that increasing a person's sense of competence, autonomy, and relatedness will increase their autonomous motivation. SDT also states that more autonomously motivated individuals are more likely to persevere and achieve their desired goals whilst experiencing more positive mental health outcomes (Deci & Ryan, 2012; Rouse et al., 2011). SDT has been tested extensively, with systematic reviews showing a consistent positive relationship between more autonomous forms of motivation and well-being. Exercise referral schemes grounded in SDT have also been shown to generate improvements in mental health outcomes (Duda et al., 2014).

Web-based interventions, when complementing usual ERS are potentially effective at supporting short-term improvements in physical activity in a range of clinical conditions (Taylor et al., 2021). Web-based interventions promoting physical activity may also improve symptoms of depression and anxiety in people with depression in the community (Lambert et al., 2018), but the evidence is fairly limited (Carneiro et al., 2022) and none examined the effects of augmenting an exercise scheme with web-based support. Web-based interventions promoting physical activity offer a potentially inexpensive and scalable option for people with depression and anxiety (Carneiro et al., 2022). For a group of people who may be referred to an exercise programme, and who may find it more difficult to maintain engagement, an augmented virtual support system that fosters improvements in self-

determination constructs may be particularly valuable. Therefore, understanding the effects of existing web-based interventions targeting physical activity on depression and anxiety may offer important insights into whether a web-based physical activity intervention could offer augmented support, particularly for people with depression.

Exercise referral schemes provide an ideal context to evaluate the effects of web-based support to increase physical activity for people with depression and anxiety. The e-coachER RCT investigated whether adding interactive weekly web-based behavioural support to usual exercise referral schemes could increase accelerometer-measured moderate to vigorous physical activity (MVPA) over exercise referral schemes alone at 12 months post-randomisation. The e-coachER intervention, compared with exercise referral schemes alone showed only a small non-significant effect on accelerometer-measured MVPA at 12 months (Taylor et al., 2021). However, e-coachER led to some short-term changes in most process outcomes, some of which also appeared to mediate e-coachER effects on changes in MVPA (Lambert et al., 2022).

The research questions for this secondary analysis of people with elevated symptoms of depression within the e-coachER trial were:

1. Does e-coachER + exercise referral improve symptoms of depression and anxiety compared to exercise referral alone at 4 and 12 months?
2. Does e-coachER + exercise referral increase accelerometer-measured and self-reported MVPA compared to exercise referral schemes alone at 4 and 12 months?
3. Are improvements in symptoms of depression and anxiety at 4 and 12 months mediated by changes in accelerometer-measured and self-reported MVPA at 4 months?
4. Are improvements in symptoms of depression and anxiety at 4 and 12 months mediated by changes in the e-coachER processes (as defined by the logic model) at 4 months?

Methods

Study design and participants

This secondary analysis utilises data from participants in the e-coachER multi-centre RCT with a score of ≥ 8 on the Hospital Anxiety and Depression Scale (HADS), indicating at least mild depression. The e-coachER RCT recruited 450 adults from Greater Glasgow, Birmingham or Plymouth and adjacent rural areas, who had been or were about to be referred by a primary care practitioner to their local ERS. Participants registered with a general practitioner (GP) surgery with one or more of the following: obesity (body mass index (BMI), 30–40), a diagnosis of hypertension, prediabetes, type 2 diabetes, lower limb osteoarthritis or having a history of treatment for depression were eligible for inclusion in the trial. Participants also needed to be categorised as ‘inactive’ or ‘moderately inactive’ (i.e. some activity but < 1 hour per week and in a sedentary occupation, according to the General Practice Physical Activity Questionnaire (GPPAQ) (Ahmad et al., 2015). Participants were excluded if they did not meet the eligibility criteria for their local ERS, had an unstable, severe and enduring mental health condition, were being treated for an alcohol or drug addiction that may have limited their involvement with the study, or were unable to use written materials in English.

Participants were randomly allocated 1:1 to either usual exercise referral schemes alone (control arm) or usual exercise referral schemes plus e-coachER (intervention arm) using a secure, password-protected web-based system created and managed by the Peninsula Clinical Trials Unit (CTU). Randomisation was stratified by site. The participant’s perceived main reason for their referral to the exercise referral schemes (i.e., chronic condition) and self-reported IT literacy/confidence (using a 10-point scale) were used as minimisation variables to ensure balanced participant characteristics in each group.

The randomisation sequence was generated by researchers in the University of Plymouth Medical Statistics department independent of the e-coachER RCT. The trial statistician,

health economist and analyst handling the accelerometer data conducted their analyses blinded to allocation. Exercise referral practitioners were not aware of which group participants were randomised. Those involved with the qualitative process evaluation were not blinded. Due to the nature of the intervention, it was not possible to blind participants. The protocol for the study has been published elsewhere (Ingram et al., 2018). The study was approved by the National Research Ethics Committee North West—Preston (15/NW/0347).

Participants in both arms of the trial were offered the usual primary care ERS. Participants randomised to the intervention arm were also offered the e-coachER package in addition to their usual ERS. The intervention has previously been described in detail elsewhere (Ingram et al., 2018; Taylor et al., 2020, 2021). In brief, e-coachER was a web-based support package using the LifeGuide© platform (<https://www.lifeguideonline.org/>), informed by SDT, with behaviour change techniques (BCTs) specifically selected to target the basic psychological needs of autonomy, competence and relatedness (Teixeira et al., 2020). Participants allocated to the intervention group received a small box through the mail containing a user guide for accessing the e-coachER web-based support system, a pedometer (step-counter) and a fridge magnet with tear-off sheets to record weekly step counts or minutes of MVPA. Participants were able to use the e-coachER intervention throughout the entire duration of the study. See Appendix 1 for the e-coachER logic model. Participants were sent an accelerometer and questionnaire booklet by post, and a prepaid envelope to return to Peninsula CTU at baseline, 4 months and 12 months post-randomisation. Weekly minutes of MVPA, recorded in bouts of at least 10 minutes were collected using GENEActiv accelerometers (Activinsights; <https://www.geneactiv.org/>), over 1 week at 4 months and 12 months post-randomisation.

Outcomes

The focus of the present study was the change in symptoms and depression and anxiety self-reported by participants using the HADS. The HADS is a 14-item, 4-point scale, used to

identify possible and probable cases of anxiety and depression among patients in nonpsychiatric hospital clinics. The HADS is divided into both anxiety and depression subscales. A score of 8-10 indicates mild depression or anxiety and 11 to 21 indicates moderate to severe depression or anxiety (Stern, 2014). The HADS performs well in assessing the symptom severity and clinical presence of anxiety and depression in primary care patients and the general population. Mediators included baseline to 4-month change in accelerometer-measured physical activity and process variables. Accelerometer measured physical activity was assessed by counting the number of weekly minutes of MVPA (both in 10-minute bouts and continuous), using GENEActiv accelerometers. Self-reported MVPA over one week was measured using the 7-day recall of PA (7-day Physical Activity Recall questionnaire) at 4 and 12 months (Blair et al., 1985). Process variables were selected to capture key psychological processes for changing physical activity behaviour, as specified by the underlying logic model (Appendix 1). They reflected theoretical mechanisms of change and enactment (i.e., participant use of the BCTs in day-to-day settings). Briefly, items were derived from extensive reviews of the literature to ensure they matched the theoretical constructs specified by the logic model, but also were fit for purpose within a randomised trial and were acceptable and easy to understand according to our Public and Patient Involvement advisory group. Process variables included the following:

Perceived importance to be physically active (single item, 11-point scale). Example question: “Doing at least 30 min of moderate intensity physical activity (e.g., brisk walk) on at least 5 days a week is important to me”

Perceived confidence to be physically active (single item, 11-point scale). Example question: “I am confident that I can do at least 30 min of moderate intensity physical activity (e.g., brisk walk) on at least 5 days a week”

Perceived competence in being regularly physically active (4 item, 5-point scale). Example question: “I am able to meet the challenge of being physically active regularly”

Autonomous in decisions about PA (4 items, 5-point scale). Example question: “I feel free to be physically active in my own way”

Availability of support (3 item, 5-point scale). Example question: “There are others in my life with whom I can be physically active”

Frequency of support (3 item, 5-point scale). Example question: “In the last 30 days how often did others discuss physical activity with me?”

Action planning (5 item, 5-point scale). Example question: “In the last 30 days I have regularly made weekly plans for when to be physically active”

Self-monitoring (2 item, 5-point scale). Example question: “In the last 30 days I have consistently monitored the amount of physical activity I do”

A full report of the outcome and process measures is reported elsewhere (Lambert et al., 2022; Taylor et al., 2020). The participant’s perception of the reason for referral to the exercise referral schemes (i.e., clinical condition) was collected. If multiple conditions were stated, the participant’s most important reason for referral was used as a minimisation variable. The level of engagement in the online e-coachER intervention was captured from the LifeGuide© platform.

Statistical analysis

All models were adjusted for baseline outcome values and stratification variables (by site) and minimisation variables (participant’s perception of the main medical reason for referral to the ERS and IT literacy level) age and gender. Baseline descriptive statistics were presented by the randomisation arm. To investigate the effect of the intervention on the HADS and PA outcomes at 4 and 12 months, according to aims 1 and 2, between-group differences were compared, using complete case data following intention-to-treat (ITT) principles, applying linear mixed-effects models to allow for clustering by the centre. This was followed by secondary per-protocol analysis using a complier average causal effect (CACE) approach,

with robust standard errors clustered by centre, to examine the respective impact of participants registering for e-coachER and of participants completing ‘Step 5’ (i.e., setting a goal and reviewing a goal online). For the mediation analyses in aims 3 and 4, linear mixed effects models, clustered by the centre, were fitted and the product of coefficients in the path analysis (Alwin & Hauser, 1975) (figure 1) of the mediating effects of accelerometer-measured MVPA (both in ≥ 10 -minute bouts and continuous), self-reported MVPA and process variables at 4 months. Mediation analysis was not contingent on the significance of results from the main analysis, as a mediated effect could still be possible without a significant net effect of the intervention on the primary outcome (Cerin & MacKinnon, 2009). Confidence intervals for the mediated path were estimated through the bootstrap resampling method, with 1000 replications. Only those with valid accelerometer data were included in the analysis for the mediating effect of accelerometer-measured physical activity.




Figure 1 A priori path model for testing mediation effects

Results

At baseline, 205 of the 450 patients randomised to the trial had a score of ≥ 8 on the HADS-D indicating at least mild depression, 69 had a score ranging from 11 to 14 indicating moderate depression and 36 had a score ranging from 15 to 21 indicating severe depression. For anxiety, 178 had a score of ≥ 8 on the HADS-A, indicated mild anxiety, 73 had a score ranging from 11 to 14 indicating moderate anxiety and 52 had a score ranging from 15 to 21 indicating severe anxiety. Of the baseline sample, 138 (67%) provided follow-up HADS scores at 4 months and 126 (61%) provided follow-up HADS scores at 12 months. The groups were well balanced at baseline with no statistically significant differences (Table 1). For accelerometer-measured, bouted physical activity, 187 (91%) provided usable accelerometer data at baseline, 108 (53%) at 4 months and 100 (49%) at 12 months. For self-reported physical activity, 205 provided usable data at baseline, 154 (75%) at 4 months and 138 (67%) at 12 months. Those who did not attend their first ERS session were significantly less likely to have 4-month follow-up data ($p > 0.05$). Overall mean age and BMI was 47.6 years (SD 12.7) and 32.7kg/m² (SD 4.4) respectively. Descriptive data for the whole sample (N=450) have been described previously (Taylor et al., 2020, 2021).

Table 1. Baseline demographic and clinical characteristics

	Intervention			Control			Total		
	N	Mean/n	SD/%	N	Mean/n	SD/%	N	Mean/n	SD/%
Gender	95			110			205		
Female		68	71.58		73	66.36		141	68.78
Male		27	28.42		37	33.64		64	31.22
Age (years)	95	47.33	12.75	110	47.77	12.71	205	47.57	12.70
BMI	95	32.36	4.41	110	33.01	4.47	205	32.71	4.44
IT confidence	95			110			205		
High		76	80.00		94	85.45		170	82.93
Low		19	20.00		16	14.55		35	17.07
GPPAQ Score	95			110			205		
2		69	72.63		66	60.00		135	65.85
3		26	27.37		44	40.00		70	34.15
Yes		65	69.15		84	77.78		149	73.76
No		29	30.85		24	22.22		53	26.24

Notes: N varies due to invalid wear-time for PA, or non-completion of a full set of measures. *Exercise referral scheme

There was a significant between-group difference of -1.36, (95% Confidence Interval (CI): -2.55 to -0.18, $p < 0.05$) for depression at 4 months post-randomisation in favour of the e-coachER arm (table 2). There were no significant differences in anxiety at 4 months or for depression and anxiety at 12 months (Table 2). There were no significant differences in physical activity at any time point (Table 3).

For the CACE analysis, greater between group differences in depression at 4-months were observed for participants who registered for e-coachER (-1.98, 95% CI: -2.55 to -1.40, $p < 0.001$) or completed a goal review (-2.89, 95% CI: -3.70 to -2.09, $p < 0.001$). Greater between group differences in self-reported physical activity at 12-months were also observed for participants who registered for e-coachER (125.54, 95% CI: 84.47 to 166.60, $p < 0.001$) or

completed a goal review (199.18, 95% CI: 132.13 to 266.24, $p < 0.001$). See appendix 2 for full CACE analysis.

Table 2. Summary of HADS outcome data at baseline and 4- and 12 months follow-up and estimated between-group differences

	Intervention			Control			Coefficient (95% CI)	P
	N	Mean	SD	N	Mean	SD		
HADS-D								
Baseline	95	11.66	3.23	110	11.28	2.80		
Month 4	57	9.35	4.58	81	10.31	4.34	-1.36 (-2.55 to -0.18)	0.024
Month 12	51	10.32	5.04	75	9.93	4.57	-0.10 (-1.57 to 1.37)	0.893
HADS-A								
Baseline	95	12.14	3.86	110	11.40	4.04		
Month 4	57	10.46	4.50	81	10.94	4.34	-1.00 (-2.21 to 0.21)	0.105
Month 12	51	11.33	4.44	75	10.83	4.34	-0.06 (-1.39 to 1.27)	0.933

Table 3. Summary of MVPA measured by accelerometer at 4 and 12 months and estimated between group-differences

	Intervention			Control			Coefficient (95% CI)	P
	N	Mean	SD	N	Mean	SD		
Bouted								
MVPA								
Baseline	86	26.28	44.88	97	17.30	35.78		
Month 4	42	25.47	42.57	66	21.89	32.08	1.71 (-11.03 to 14.44)	0.793
Month 12	41	16.95	39.77	59	13.73	29.42	2.92 (-10.69 to 16.52)	0.674
Continuous								
MVPA								
Baseline	89	387.29	265.21	98	332.40	231.71		
Month 4	43	379.30	207.98	67	341.52	264.79	1.36 (-49.25 to 51.96)	0.958
Month 12	44	333.16	263.46	59	328.45	224.11	-26.45 (-96.18 to 43.29)	0.457
Self-reported								
MVPA								
Baseline	95	160.59	338.12	110	167.91	296.35		
Month 4	64	244.13	338.82	90	333.12	579.82	-17.5 (-163.31 to 128.31)	0.814
Month 12	59	222.69	442.43	79	182.90	420.58	84.7 (-55.92 to 225.38)	0.238

Notes: N varies due to invalid wear-time for PA, or non-completion of a full set of measures.

Mediation analyses revealed no mediation effect of change in weekly minutes of accelerometer-measured MVPA or self-reported MVPA on depression and anxiety at 4- and 12 months post-randomisation (Appendices 3 to 6).

Intervention effects on confidence, competence and self-monitoring at 4 months significantly mediated the effect of e-coachER in lowering depression at 4 months (mediated effect, Table 4). All remaining processes were not significant. There were direct effects of e-coachER on depression scores at 4 months when controlling for changes in all the process

measures apart from confidence and competence (direct C' -path in Table 8). There were no significant mediating effects or direct effects at 12 months for any of the process measures (Appendix 7).

Table 4. Mediation effects for intervention effects on processes at 4 months on depression at 4 months

Mediators	N	A path		B path		Mediated effect		C' path	
		β (SE)	P	β (SE)	P	β (SE)	95% CI	β (SE)	P
Importance	132	0.55 (0.43)	0.203	-0.04 (0.10)	0.672	-0.02 (0.08)	-0.24, 0.12	-1.35 (0.62)	0.030
Confidence	133	1.08 (0.51)	0.033	-0.37 (0.09)	0	-0.40 (0.20)	-0.82, -0.04	-0.98 (0.59)	0.097
Competence	127	1.37 (0.62)	0.027	-0.24 (0.08)	0.002	-0.33 (0.21)	-0.82, 0.00	-0.92 (0.63)	0.146
Autonomy	130	0.48 (0.60)	0.424	-0.23 (0.07)	0.002	-0.11 (0.15)	-0.41, 0.16	-1.23 (0.57)	0.031
Freq of support	134	0.69 (0.59)	0.247	-0.01 (0.08)	0.875	-0.01 (0.08)	-0.16, 0.16	-1.36 (0.62)	0.028
Avail of support	133	0.19 (0.48)	0.693	-0.06 (0.10)	0.555	-0.01 (0.06)	-0.13, 0.12	-1.37 (0.62)	0.026
Action planning	125	0.48 (0.85)	0.573	-0.15 (0.04)	0	-0.07 (0.13)	-0.34, 0.20	-1.37 (0.60)	0.022
Self-monitoring	131	0.86 (0.36)	0.016	-0.34 (0.11)	0.002	-0.29 (0.16)	-0.65, -0.02	-1.40 (0.58)	0.015

Notes: N varies due to lack of valid wear-time for PA, or non-completion of full set of measures.

No direct effects of e-coachER on anxiety scores at 4 or 12 months were observed when controlling for changes in the process measures (c' -path, Table 5 and Appendix 8). Change in competence and self-monitoring at 4 months significantly mediated the effect of e-

coachER in reducing anxiety scores at 4 (mediated effect, Table 5) but not 12 months (mediated effect, Appendix 8).

Table 5. Mediation effects for intervention effects on processes at 4 months on anxiety at 4 months

Mediators	N	A path		B path		Mediated effect		C' path	
		β (SE)	P	β (SE)	P	β (SE)	95%CI	β (SE)	P
Importance	132	0.55 (0.43)	0.203	-0.24 (0.11)	0.025	-0.13 (0.13)	-0.42, 0.07	-0.96 (0.62)	0.125
Confidence	133	1.08 (0.51)	0.033	-0.17 (0.10)	0.09	-0.18 (0.15)	-0.49, 0.07	-0.81 (0.63)	0.201
Competence	127	1.37 (0.62)	0.027	-0.24 (0.08)	0.002	-0.33 (0.19)	-0.77, -0.02	-0.75 (0.63)	0.238
Autonomy	130	0.48 (0.60)	0.424	-0.10 (0.08)	0.21	-0.05 (0.08)	-0.23, 0.10	-1.05 (0.61)	0.083
Freq of support	134	0.69 (0.59)	0.247	-0.11 (0.08)	0.153	-0.08 (0.09)	-0.31, 0.07	-1.03 (0.63)	0.102
Avail of support	133	0.19 (0.48)	0.693	-0.08 (0.10)	0.456	-0.01 (0.07)	-0.19, 0.12	-1.19 (0.62)	0.055
Action planning	125	0.48 (0.85)	0.573	-0.09 (0.04)	0.036	-0.04 (0.09)	-0.26, 0.13	-1.11 (0.60)	0.066
Self-monitoring	131	0.86 (0.36)	0.016	-0.30 (0.11)	0.009	-0.26 (0.16)	-0.61, -0.02	-1.03 (0.61)	0.091

Notes: N varies due to lack of valid wear-time for PA, or non-completion of full set of measures

Discussion

When added to usual ERS, e-coachER led to significant between-group improvements in symptoms of depression at 4 months (which became larger for participants who registered versus didn't or completed an e-coachER goal review versus didn't), but not at the 12-month follow-up. However, no significant differences were observed for anxiety at 4 or 12 months. No differences were observed for any of the MVPA outcomes. However, differences in self-reported MVPA at 12 months for participants who registered or completed an e-coachER

goal review. MVPA did not mediate the relationship between e-coachER and depression or anxiety and 4 or 12 months follow-up. However, self-reported competence and confidence to engage in physical activity and self-reported use of self-monitoring of physical activity positively partially mediated the effects of e-coachER on the reduction in depression symptoms at 4 months. Competence and use of self-monitoring positively and partially mediated changes in symptoms of anxiety at 4-month follow-up. These results suggest that adding a web-based behavioural support programme to exercise referral schemes yields additional mental health benefits for patients and that these benefits are partly derived from changes in patients' competence and confidence to engage in physical activity and self-reported use of self-monitoring of physical activity, irrespective of whether changes in MVPA were achieved. The present study is the first to evaluate whether adding a web-based behavioural support programme to exercise referral schemes can lead to changes in symptoms of depression and anxiety. Furthermore, the present study is one of the few to evaluate the mediational effects of accelerometer-measured physical activity on depression or anxiety in the context of an RCT.

Relation to other literature

Previous RCTs of web-based behavioural support for MVPA have also observed improvements in depression but not changes in self-reported or accelerometer-measured MVPA. For example, one RCT of 48 people with depression found that a web-based supported intervention targeting physical activity led to changes in depression but not self-reported physical activity (Ström et al., 2013). Similarly, a 3-arm RCT (n=501) observed significant reductions in depression and stress (at 3 months) and anxiety (at 9 months) but no changes in accelerometer-measured MVPA when comparing web-based video-tailored and text-tailored physical activity intervention to a control group (Vandelanotte et al., 2022). However, Vandelanotte et al., (2022) did not solely include people with elevated baseline depression symptoms. Another study compared the effects of an individually tailored, mailed print PA intervention for Latinas (n=266) on symptoms of depression vs a wellness control at

12 months. They found no direct effects of the intervention on symptoms of depression. However, improvements in self-reported MVPA at 12 months appeared to mediate intervention effects on depression at 12 months (Mendoza-Vasconez et al., 2019). This is contrary to our findings despite employing similar behavioural strategies (e.g., goal-setting and self-monitoring). One reason for this could be the differences in the mode of delivery. For example, the e-coachER study used a printed, tailored approach which addressed specific PA barriers identified by Latinas in focus groups and the literature (Marcus et al., 2013). To note, none of the above studies took place within ERS, limiting their comparability with e-coachER.

Our findings also build on previous research investigating the effect of exercise referral schemes grounded in SDT on mental health. A previous cluster RCT compared exercise referral, grounded in SDT vs exercise referral alone and found significant decreases in anxiety but not depression at 6 months in favour of the intervention arm (Duda et al., 2014). However, no differences in participant perceptions of autonomy support from the health and fitness advisor (HFA). The authors attributed the lack of findings to a possible lack of training and support for the HFAs assigned to the SDT arm (Duda et al., 2014). Lack of training can directly impact delivery fidelity, as outlined in the treatment fidelity framework provided by the NIH's Behavioral Change Consortium (BCC) (Borrelli et al., 2005). Therefore, this lack of training may have led to a lack of delivery fidelity to the protocol from the HFA which led to sub-optimal outcomes. In e-coachER, the SDT content was standardised, fixed, and delivered directly to the participant, avoiding this breakdown in fidelity (Watkins et al., 2016) and potentially explaining why we observed further changes in key processes related to SDT and mental health.

There are several possible reasons explaining why we observed reductions in depression but not MVPA. First, increases in perceived autonomy/developing a sense of competence, control or relatedness might be more important than changes in physical activity for improving mood (Nyström et al., 2015). SDT posits that if a person's basic psychological

need for competence, autonomy and relatedness are met, they should experience improvements in mental health via increased autonomous motivation (Deci & Ryan, 2012). The present study found that increases in self-monitoring, competence and confidence towards physical activity mediated the effects of e-coachER on improvements in depression and anxiety at 4 months. This aligns with previous studies which have consistently found self-efficacy to be a strong candidate mechanism for the relationship between physical activity and depression (Haller et al., 2018; Howarter et al., 2014; Kandola et al., 2019; Miller et al., 2019; White et al., 2009). However, self-efficacy refers to confidence in carrying out a behaviour under challenging circumstances, rather than just competence to carry out a behaviour (Rodgers et al., 2014). Previous research has shown that people who demonstrate greater internalisation of their reasons for engaging with PA also exhibit more positive mental health outcomes (Fortier et al., 2012). This is consistent with SDT self-determination theory which shows that autonomously regulated behaviours can translate into enhanced psychological wellness (Teixeira et al., 2012) with less autonomous regulations (e.g., identified regulation) only predicting wellbeing contingent on behavioural performance (Burton et al., 2006). It could be that participant's reasons for engaging in PA in the e-coachER arm were more internalised, leading to decreases in depression and anxiety. It may therefore be more important to focus on activities that are intrinsically motivating to the patient, rather than focusing on frequency, intensity and type of MVPA (Nyström et al., 2015).

Second, e-coachER could be improving symptoms through the same mechanism of behavioural activation (BA). BA reduces depression through increasing exposure to positive reinforcement and reducing negative reinforcement in the environment (Hopko et al., 2003; Lejuez et al., 2001). BA is also compatible with SDT sharing many techniques including providing a rationale, graded tasks, self-monitoring and goal setting, action planning and goal review (Lambert et al., 2017; Teixeira et al., 2020). e-coachER may have supported people to increase their engagement in routine pleasurable and necessary activities that

don't necessarily involve MVPA. For example, a 2-arm pilot RCT using BA to promote PA to people with depression found significant reductions in depression and anxiety at 2 months despite no observed changes in MVPA (Lambert et al., 2018). In another 5-arm RCT, encouraging people to gradually increase their degree of PA over weekly treatment modules (using self-monitoring and providing a rationale for their depression) led to decreases in depressive symptoms when compared with a waiting list control group (Nyström et al., 2017).

Third, the present analysis did not account for the co-dependence of sedentary behaviour and physical activity. A prospective cohort study of 60,235 UK Biobank participants used compositional data analysis methods to assess the impact of replacing daily time spent in sedentary behaviours with sleep, light, or moderate to vigorous physical activity on depression and anxiety symptoms. A 12.5%, 7.6% and 1.3% reduction in depressive symptoms at 2-year follow-up was found when replacing 60 minutes of sedentary behaviour with MVPA, sleep, or light PA, respectively (Kandola et al., 2021). As there is a finite period in the day, it could be that, rather than increasing MVPA, e-coachER led to changes in depression by replacing sedentary behaviour with MVPA, light activity or sleep.

Strengths and limitations

This study had several strengths. First, this secondary analysis was part of a large RCT with a large overall sample size and included people with a range of health conditions, making the findings relevant to patients presenting at primary care who might fit the criteria for exercise referral. Second, the use of accelerometer-measured MVPA is a key strength considering the trial participants were not blinded to their assignment to usual ERS or augmented e-coachER. Third, it followed patients after 4 and 12 months. Fourth it employed mediation analysis to investigate the indirect effects of e-coachER on depression and anxiety via processes of change and MVPA.

Several limitations need to be acknowledged. First, e-coachER struggled with loss to follow-up at 4 and 12 months limiting the level of confidence in the findings. The e-coachER qualitative analysis also showed that participants reported IT-related difficulties and a lack of support from the exercise referral schemes which may have impacted their confidence to use e-coachER (Taylor et al., 2020). Second, we did not use multiple imputation analysis to address the missing data since multiple imputation in trials with a large amount of missing data may introduce bias (Jakobsen et al., 2017). Instead, since we have reported an exploratory analysis, a complete case analysis was used. Third, the sub-sampling from the main trial could have introduced some selection bias. However, both groups were reasonably well matched in key demographic and clinical characteristics.

Directions for future research

This study provides support for SDT underpinning web-based interventions to reduce mental health issues, at least in the short term. Future research could replicate this work by employing a more person-centred approach focussing on people with elevated depression in the intervention design process (e.g., by using the person-based approach) (Yardley et al., 2015). This could enhance the effects of the intervention by making it more specific and relevant for people with depression. Another avenue for future research could be to use compositional data analysis methods on existing RCTs which have collected accelerometer-measured physical activity to see whether displacement of sedentary time with other activities leads to changes in mental health.

In summary, for participants with depression, adding web-based support to usual ERS leads to significantly greater reductions in depression at 4 months compared to ERS alone, but the effects were not sustained at 12 months. The intervention had no effect on physical activity within the study. However, changes in depression can be influenced by changing people's motivational regulations toward physical activity. e-coachER may represent a low-cost option to further reduce depression in the context of ERS. Furthermore, this research provides evidence that web-based interventions promoting physical activity, but without a specific

focus on mental health, have the potential to alleviate depressive symptoms, even in the absence of increasing physical activity. Simply engaging in processes linked to self-determination (e.g., gaining a sense of competence and control) and physical activity can reduce depression. These findings have clear implications for future web-based interventions highlighting the need to capture mental health outcomes and design them to target people with mental health issues. Our mediational analyses also provide insight into how key theoretical components embedded within the intervention can impact depression and anxiety over the short term, further extending the applicability of SDT beyond behavioural engagement into the realm of health and wellbeing (Teixeira et al., 2012). Further research is needed to help to understand how web-based physical activity interventions can be used to facilitate longer-term benefits for people with a mental health condition.

Role of the funding source

This study was funded by the National Institute for Health Research (NIHR), Health Technology Assessment Programme (grant reference: 13/25/20). The funder of the study had no role in study design, data collection, data analysis, data interpretation, or writing of the report. This study is funded by the NIHR Health Technology Assessment Programme (grant reference: 13/25/20). The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care. Sarah Dean's time is partially supported by the National Institute for Health Research (NIHR) Applied Research Collaboration South West Peninsula. Kate Jolly is part-funded by NIHR Applied Research Collaboration (ARC) West Midlands.

References

Ahmad, S., Harris, T., Limb, E., Kerry, S., Victor, C., Ekelund, U., Iliffe, S., Whincup, P., Beighton, C., Ussher, M., & Cook, D. G. (2015). Evaluation of reliability and validity of the General Practice

- Physical Activity Questionnaire (GPPAQ) in 60–74 year old primary care patients. *BMC Family Practice*, 16(1), 113. <https://doi.org/10.1186/s12875-015-0324-8>
- Alwin, D. F., & Hauser, R. M. (1975). The decomposition of effects in path analysis. *American Sociological Review*, 40(1), 37. <https://doi.org/10.2307/2094445>
- Blair, S., Haskell, W., Paffenbarger Jr, R., Vranizan, K., Farquhar, J., & Wood, P. (1985). Assessment of habitual physical activity by a seven-day recall in a community survey and controlled experiments. *American Journal of Epidemiology*, 122(5), 794–804.
- Borrelli, B., Sepinwall, D., Ernst, D., Bellg, A. J., Czajkowski, S., Breger, R., DeFrancesco, C., Levesque, C., Sharp, D. L., Ogedegbe, G., Resnick, B., & Orwig, D. (2005). A new tool to assess treatment fidelity and evaluation of treatment fidelity across 10 years of health behavior research. *Journal of Consulting and Clinical Psychology*, 73(5), 852–860. <https://doi.org/10.1037/0022-006X.73.5.852>
- Burton, K. D., Lydon, J. E., D'Alessandro, D. U., & Koestner, R. (2006). The differential effects of intrinsic and identified motivation on well-being and performance: Prospective, experimental, and implicit approaches to self-determination theory. *Journal of Personality and Social Psychology*, 91(4), 750–762. <https://doi.org/10.1037/0022-3514.91.4.750>
- Busch, A. M., Ciccolo, J. T., Puspitasari, A. J., Nosrat, S., Whitworth, J. W., & Stults-Kolehmainen, M. A. (2016). Preferences for exercise as a treatment for depression. *Mental Health and Physical Activity*, 10, 68–72. <https://doi.org/10.1016/j.mhpa.2015.12.004>
- Carneiro, L., Rosenbaum, S., Ward, P. B., Clemente, F. M., Ramirez-Campillo, R., Monteiro-Júnior, R. S., Martins, A., & Afonso, J. (2022). Web-based exercise interventions for patients with depressive and anxiety disorders: A systematic review of randomized controlled trials. *Brazilian Journal of Psychiatry*, 44(3), 331–341. <https://doi.org/10.1590/1516-4446-2021-2026>

- Cerin, E., & MacKinnon, D. P. (2009). A commentary on current practice in mediating variable analyses in behavioural nutrition and physical activity. *Public Health Nutrition*, *12*(8), 1182–1188. <https://doi.org/10.1017/S1368980008003649>
- Cooney, G. (2018). Exercise and mental health: A complex and challenging relationship. *The Lancet Psychiatry*, *5*(9), 692–693. [https://doi.org/10.1016/S2215-0366\(18\)30291-8](https://doi.org/10.1016/S2215-0366(18)30291-8)
- Cooney, G., Dwan, K., Greig, C. A., Lawlor, D. A., Rimer, J., Waugh, F. R., McMurdo, M., & Mead, G. E. (2013). Exercise for depression. *Cochrane Database of Systematic Reviews*. <https://doi.org/10.1002/14651858.CD004366.pub6>
- Deci, E. L., & Ryan, R. M. (2012). Self-determination theory. In *Handbook of theories of social psychology*, Vol. 1 (pp. 416–436). Sage Publications Ltd. <https://doi.org/10.4135/9781446249215.n21>
- Duda, J. L., Williams, G. C., Ntoumanis, N., Daley, A., Eves, F. F., Mutrie, N., Rouse, P. C., Lodhia, R., Blamey, R. V., & Jolly, K. (2014). Effects of a standard provision versus an autonomy supportive exercise referral programme on physical activity, quality of life and well-being indicators: A cluster randomised controlled trial. *International Journal of Behavioral Nutrition and Physical Activity*, *11*(1), 10. <https://doi.org/10.1186/1479-5868-11-10>
- Fortier, M. S., Duda, J. L., Guerin, E., & Teixeira, P. J. (2012). Promoting physical activity: Development and testing of self-determination theory-based interventions. *International Journal of Behavioral Nutrition and Physical Activity*, *9*(1), 20. <https://doi.org/10.1186/1479-5868-9-20>
- Haller, N., Lorenz, S., Pfirmann, D., Koch, C., Lieb, K., Dettweiler, U., Simon, P., & Jung, P. (2018). Individualized web-based exercise for the treatment of depression: Randomized controlled trial. *JMIR Mental Health*, *5*(4), e10698–e10698. PubMed. <https://doi.org/10.2196/10698>
- Harris, T., Kerry, S. M., Victor, C. R., Ekelund, U., Woodcock, A., Iliffe, S., Whincup, P. H., Beighton, C., Ussher, M., Limb, E. S., David, L., Brewin, D., Adams, F., Rogers, A., & Cook, D. G. (2015). A primary care nurse-delivered walking intervention in older adults: PACE (pedometer

- accelerometer consultation evaluation)-Lift cluster randomised controlled trial. *PLOS Medicine*, 12(2), e1001783. <https://doi.org/10.1371/journal.pmed.1001783>
- Heissel, A., Heinen, D., Brokmeier, L. L., Skarabis, N., Kangas, M., Vancampfort, D., Stubbs, B., Firth, J., Ward, P. B., Rosenbaum, S., Hallgren, M., & Schuch, F. (2023). Exercise as medicine for depressive symptoms? A systematic review and meta-analysis with meta-regression. *British Journal of Sports Medicine*, bjsports-2022-106282. <https://doi.org/10.1136/bjsports-2022-106282>
- Hopko, D. R., Lejuez, C. W., Ruggiero, K. J., & Eifert, G. H. (2003). Contemporary behavioral activation treatments for depression: Procedures, principles, and progress. *Clinical Psychology Review*, 23(5), 699–717. [https://doi.org/10.1016/S0272-7358\(03\)00070-9](https://doi.org/10.1016/S0272-7358(03)00070-9)
- Howarter, A. D., Bennett, K. K., Barber, C. E., Gessner, S. N., & Clark, J. M. R. (2014). Exercise self-efficacy and symptoms of depression after cardiac rehabilitation: Predicting changes over time using a piecewise growth curve analysis. *Journal of Cardiovascular Nursing*, 29(2). https://journals.lww.com/jcnjournal/Fulltext/2014/03000/Exercise_Self_efficacy_and_Symp_toms_of_Depression.10.aspx
- Ingram, W., Webb, D., Taylor, R. S., Anokye, N., Yardley, L., Jolly, K., Mutrie, N., Campbell, J. L., Dean, S. G., Greaves, C., Steele, M., Lambert, J., McAdam, C., Jane, B., King, J., Jones, R. B., Little, P., Woolf, A., Erwin, J., ... Taylor, A. (2018). Multicentred randomised controlled trial of an augmented exercise referral scheme using web-based behavioural support in individuals with metabolic, musculoskeletal and mental health conditions: Protocol for the e-coachER trial. *BMJ Open*, 8(9), e022382. <https://doi.org/10.1136/bmjopen-2018-022382>
- Jakobsen, J. C., Gluud, C., Wetterslev, J., & Winkel, P. (2017). When and how should multiple imputation be used for handling missing data in randomised clinical trials – a practical guide with flowcharts. *BMC Medical Research Methodology*, 17(1), 162. <https://doi.org/10.1186/s12874-017-0442-1>

- Kandola, A., Ashdown-Franks, G., Hendrikse, J., Sabiston, C. M., & Stubbs, B. (2019). Physical activity and depression: Towards understanding the antidepressant mechanisms of physical activity. *Neuroscience & Biobehavioral Reviews*, *107*, 525–539.
<https://doi.org/10.1016/j.neubiorev.2019.09.040>
- Kandola, A., del Pozo Cruz, B., Osborn, D. P. J., Stubbs, B., Choi, K. W., & Hayes, J. F. (2021). Impact of replacing sedentary behaviour with other movement behaviours on depression and anxiety symptoms: A prospective cohort study in the UK Biobank. *BMC Medicine*, *19*(1), 133.
<https://doi.org/10.1186/s12916-021-02007-3>
- Lambert, J., Greaves, C. J., Farrand, P., Haase, A. M., & Taylor, A. (2017). Development of a web-based intervention (eMotion) based on behavioural activation to promote physical activity in people with depression. *Mental Health and Physical Activity*, *13*, 120–136.
<https://doi.org/10.1016/j.mhpa.2017.10.003>
- Lambert, J., Greaves, C. J., Farrand, P., Price, L., Haase, A. M., & Taylor, A. (2018). Web-based intervention using behavioral activation and physical activity for adults with depression (the eMotion study): Pilot randomized controlled trial. *Journal of Medical Internet Research*, *20*(7), e10112. <https://doi.org/10.2196/10112>
- Lambert, J., Taylor, A., Streeter, A., Greaves, C., Ingram, W. M., Dean, S., Jolly, K., Mutrie, N., Taylor, R. S., Yardley, L., Price, L., & Campbell, J. (2022). A process evaluation, with mediation analysis, of a web-based intervention to augment primary care exercise referral schemes: The e-coachER randomised controlled trial. *International Journal of Behavioral Nutrition and Physical Activity*, *19*(1), 128. <https://doi.org/10.1186/s12966-022-01360-7>
- Lejuez, C. W., Hopko, D. R., & Hopko, S. D. (2001). A brief behavioral activation treatment for depression: Treatment manual. *Behavior Modification*, *25*(2), 255–286.
- Lowe, A., Myers, A., Quirk, H., Blackshaw, J., Palanee, S., & Copeland, R. (2022). Physical activity promotion by GPs: A cross-sectional survey in England. *BJGP Open*, *6*(3), BJGPO.2021.0227.
<https://doi.org/10.3399/BJGPO.2021.0227>


- Marcus, B. H., Dunsiger, S. I., Pekmezi, D. W., Larsen, B. A., Bock, B. C., Gans, K. M., Marquez, B., Morrow, K. M., & Tilkemeier, P. (2013). The Seamos Saludables study: A randomized controlled physical activity trial of Latinas. *American Journal of Preventive Medicine, 45*(5), 598–605. <https://doi.org/10.1016/j.amepre.2013.07.006>
- Mendoza-Vasconez, A. S., Marquez, B., Linke, S., Arredondo, E. M., & Marcus, B. H. (2019). Effect of physical activity on depression symptoms and perceived stress in Latinas: A mediation analysis. *Mental Health and Physical Activity, 16*, 31–37. <https://doi.org/10.1016/j.mhpa.2019.03.001>
- Miller, K. J., Mesagno, C., McLaren, S., Grace, F., Yates, M., & Gomez, R. (2019). Exercise, mood, self-efficacy, and social support as predictors of depressive symptoms in older adults: Direct and interaction effects. *Frontiers in Psychology, 10*. <https://www.frontiersin.org/articles/10.3389/fpsyg.2019.02145>
- NICE. (2011). *Generalised anxiety disorder and panic disorder in adults: Management*.
- NICE. (2018). Depression in adults: Treatment and management. *NICE Guideline: Short Version Draft for Second Consultation*.
- NICE. (2022). Depression in adults: Treatment and management. *NICE Guideline: Short Version Draft for Second Consultation*.
- Nyström, M. B. T., Neely, G., Hassmén, P., & Carlbring, P. (2015). Treating Major Depression with Physical Activity: A Systematic Overview with Recommendations. *Cognitive Behaviour Therapy, 44*(4), 341–352. <https://doi.org/10.1080/16506073.2015.1015440>
- Nyström, M. B. T., Stenling, A., Sjöström, E., Neely, G., Lindner, P., Hassmén, P., Andersson, G., Martell, C., & Carlbring, P. (2017). Behavioral activation versus physical activity via the internet: A randomized controlled trial. *Journal of Affective Disorders, 215*, 85–93. <https://doi.org/10.1016/j.jad.2017.03.018>
- Park, L. T., & Zarate, C. A. (2019). Depression in the Primary Care Setting. *New England Journal of Medicine, 380*(6), 559–568. <https://doi.org/10.1056/NEJMcp1712493>

- Pavey, T. G., Taylor, A., Fox, K. R., Hillsdon, M., Anokye, N., Campbell, J. L., Foster, C., Green, C., Moxham, T., Mutrie, N., Searle, J., Trueman, P., & Taylor, R. S. (2011). Effect of exercise referral schemes in primary care on physical activity and improving health outcomes: Systematic review and meta-analysis. *BMJ*, *343*, d6462. <https://doi.org/10.1136/bmj.d6462>
- Pilling, S., Anderson, I., Goldberg, D., Meader, N., & Taylor, C. (2009). Depression in adults, including those with a chronic physical health problem: Summary of NICE guidance. *Bmj*, *339*.
- Rebar, A. L., Stanton, R., Geard, D., Short, C., Duncan, M. J., & Vandelanotte, C. (2015). A meta-meta-analysis of the effect of physical activity on depression and anxiety in non-clinical adult populations. *Health Psychology Review*, *9*(3), 366–378. <https://doi.org/10.1080/17437199.2015.1022901>
- Rebar, A. L., & Taylor, A. (2017). Physical activity and mental health; it is more than just a prescription. *Mental Health and Physical Activity*, *13*, 77–82. <https://doi.org/10.1016/j.mhpa.2017.10.004>
- Rodgers, W. M., Markland, D., Selzler, A.-M., Murray, T. C., & Wilson, P. M. (2014). Distinguishing Perceived Competence and Self-Efficacy: An Example From Exercise. *Research Quarterly for Exercise and Sport*, *85*(4), 527–539. <https://doi.org/10.1080/02701367.2014.961050>
- Rouse, P. C., Ntoumanis, N., Duda, J. L., Jolly, K., & Williams, G. C. (2011). In the beginning: Role of autonomy support on the motivation, mental health and intentions of participants entering an exercise referral scheme. *Psychology & Health*, *26*(6), 729–749. <https://doi.org/10.1080/08870446.2010.492454>
- Stern, A. F. (2014). The Hospital Anxiety and Depression Scale. *Occupational Medicine*, *64*(5), 393–394. <https://doi.org/10.1093/occmed/kqu024>
- Ström, M., Uckelstam, C.-J., Andersson, G., Hassmén, P., Umefjord, G., & Carlbring, P. (2013). Internet-delivered therapist-guided physical activity for mild to moderate depression: A randomized controlled trial. *PeerJ*, *1*, e178. <https://doi.org/10.7717/peerj.178>

- Stubbs, B., Vancampfort, D., Rosenbaum, S., Firth, J., Cosco, T., Veronese, N., Salum, G. A., & Schuch, F. B. (2017). An examination of the anxiolytic effects of exercise for people with anxiety and stress-related disorders: A meta-analysis. *Psychiatry Research*, *249*, 102–108.
<https://doi.org/10.1016/j.psychres.2016.12.020>
- Taylor, A., Taylor, R. S., Ingram, W., Dean, S. G., Jolly, K., Mutrie, N., Lambert, J., Yardley, L., Streeter, A., Greaves, C., McAdam, C., Price, L., Anokye, N. K., & Campbell, J. (2021). Randomised controlled trial of an augmented exercise referral scheme using web-based behavioural support for inactive adults with chronic health conditions: The e-coachER trial. *British Journal of Sports Medicine*, *55*, 444–450.
- Taylor, A., Taylor, R. S., Ingram, W. M., Anokye, N., Dean, S., Jolly, K., Mutrie, N., Lambert, J., Yardley, L., Greaves, C., King, J., McAdam, C., Steele, M., Price, L., Streeter, A., Charles, N., Terry, R., Webb, D., Campbell, J., ... Cavanagh, C. (2020). Adding web-based behavioural support to exercise referral schemes for inactive adults with chronic health conditions: The e-coachER RCT. *Health Technology Assessment*, *24*(63), 1–106. <https://doi.org/10.3310/hta24630>
- Teixeira, P. J., Carraça, E. V., Markland, D., Silva, M. N., & Ryan, R. M. (2012). Exercise, physical activity, and self-determination theory: A systematic review. *International Journal of Behavioral Nutrition and Physical Activity*, *9*(1), 78. <https://doi.org/10.1186/1479-5868-9-78>
- Teixeira, P. J., Marques, M. M., Silva, M. N., Brunet, J., Duda, J. L., Haerens, L., La Guardia, J., Lindwall, M., Lonsdale, C., Markland, D., Michie, S., Moller, A. C., Ntoumanis, N., Patrick, H., Reeve, J., Ryan, R. M., Sebire, S. J., Standage, M., Vansteenkiste, M., ... Hagger, M. S. (2020). A classification of motivation and behavior change techniques used in self-determination theory-based interventions in health contexts. *Motivation Science*, *6*(4), 438–455.
<https://doi.org/10.1037/mot0000172>
- Tomlinson-Perez, S., Machaczek, K. K., Firth, J., Pollard, N., Meda, G., Keddie, E., & Goyder, E. (2022). Evaluation of the uptake, retention and effectiveness of exercise referral schemes for the

- management of mental health conditions in primary care: A systematic review. *BMC Public Health*, 22(1), 249. <https://doi.org/10.1186/s12889-022-12638-7>
- Vandelanotte, C., Duncan, M. J., Plotnikoff, R. C., Rebar, A., Alley, S., Schoeppe, S., To, Q., Mummery, W. K., & Short, C. E. (2022). Impact of a web-based personally tailored physical activity intervention on depression, anxiety, stress and quality of life: Secondary outcomes from a randomized controlled trial. *Mental Health and Physical Activity*, 23, 100477. <https://doi.org/10.1016/j.mhpa.2022.100477>
- Watkins, E., Newbold, A., Tester-Jones, M., Javaid, M., Cadman, J., Collins, L. M., Graham, J., & Mostazir, M. (2016). Implementing multifactorial psychotherapy research in online virtual environments (IMPROVE-2): Study protocol for a phase III trial of the MOST randomized component selection method for internet cognitive-behavioural therapy for depression. *BMC Psychiatry*, 16(1), 345. <https://doi.org/10.1186/s12888-016-1054-8>
- White, K., Kendrick, T., & Yardley, L. (2009). Change in self-esteem, self-efficacy and the mood dimensions of depression as potential mediators of the physical activity and depression relationship: Exploring the temporal relation of change. *Mental Health and Physical Activity*, 2(1), 44–52. <https://doi.org/10.1016/j.mhpa.2009.03.001>
- Yardley, L., Ainsworth, B., Arden-Close, E., & Muller, I. (2015). The person-based approach to enhancing the acceptability and feasibility of interventions. *Pilot and Feasibility Studies*, 1(1), 37. <https://doi.org/10.1186/s40814-015-0033-z>

Appendix 1 The e-coachER Logic model



Appendix 2. Complier-average casual effect (CACE) analysis of all outcomes at 4 and 12 months

	Coefficient	95% CI Lower	95% CI Upper	P value
HADS-D Month 4				
Registered	-1.98	-2.55	-1.40	0.000
Step 5	-2.89	-3.70	-2.09	0.000
HADS-D Month 12				
Registered	-0.24	-0.62	0.13	0.205
Step 5	-0.36	-0.88	0.16	0.170
HADS-A Month 4				
Registered	-1.37	-3.20	0.45	0.140
Step 5	-2.02	-4.76	0.73	0.150
HADS-A Month 12				
Registered	-0.07	-1.23	1.08	0.900
Step 5	-0.11	-1.84	1.62	0.899
Bouted MVPA month 4				
Registered	2.09	-10.75	14.93	0.750
Step 5	3.08	-15.47	21.63	0.745
Bouted MVPA month 12				
Registered	3.51	-1.45	8.46	0.166
Step 5	5.04	-2.07	12.15	0.165
Continuous MVPA month 4				
Registered	-5.45	-18.77	7.87	0.423
Step 5	-7.85	-26.00	10.30	0.397
Continuous MVPA month 12				
Registered	-34.28	-89.37	20.81	0.223
Step 5	-48.87	-125.58	27.85	0.212
Self-reported MVPA month 4				
Registered	-33.20	-171.00	104.60	0.637
Step 5	-51.01	-256.22	154.20	0.626
Self-reported MVPA month 12				
Registered	125.54	84.47	166.60	0.000
Step 5	199.18	132.13	266.24	0.000

Appendix 3. Mediation effects of changes in accelerometer measured and self-reported MVPA at 4 months on the intervention effect on depression at 4 months

Mediators	N	A path		B path		Mediated effect		C' path	
		β (SE)	P	β (SE)	P	β (SE)	95%CI	β (SE)	P
Bouted	100	1.71	0.793	0.00	0.803	0.00	-0.17, 0.20	-1.38	0.047
MVPA		(6.50)		(0.01)		(0.09)		(0.70)	
Continuous	104	1.36	0.958	0.00	0.919	0.00	-0.22, 0.22	-1.55	0.027
MVPA		(25.82)		(0.00)		(0.10)		(0.70)	
Self-reported	306	22.77	0.602	0.00	0.098	-0.02	-0.20, 0.03	-1.09	0.002
MVPA		(43.61)		(0.00)		(0.06)		(0.36)	

N varies due to invalid wear-time for PA, or non-completion of a full set of measures.

Appendix 4. Mediation effects of changes in accelerometer measured and self-reported MVPA at 4 months on the intervention effect on depression at 12 months.

Mediators	N	A path		B path		Mediated effect		C' path	
		β (SE)	P	β (SE)	P	β (SE)	95%CI	β (SE)	P
Bouted	89	1.705	0.793	0.000	0.982	0.000	(-0.251,0.197)	-0.147	0.874
MVPA		(6.499)		(0.012)		(0.107)		(0.926)	
Continuous	92	1.356	0.958	-0.003	0.401	-0.004	(-0.329,0.247)	-0.222	0.809
MVPA		(25.819)		(0.003)		(0.130)		(0.921)	
Self-reported	275	22.766	0.602	-0.001	0.11	-0.023	(-0.209,0.034)	-0.107	0.798
MVPA		(43.607)		(0.000)		(0.065)		(0.419)	

Notes: N varies due to invalid wear-time for PA, or non-completion of a full set of measures.

Appendix 5. Mediation effects of changes in accelerometer measured and self-reported MVPA at 4 months on the intervention effect on anxiety at 4 months.

Mediators	N	A path		B path		Mediated effect		C' path	
		β (SE)	P	β (SE)	P	β (SE)	95%CI	β (SE)	P
Bouted	100	1.705	0.793	-0.000	0.964	0.000	(-0.194,0.149)	-0.670	0.357
MVPA		(6.499)		(0.009)		(0.084)		(0.728)	
Continuous	104	1.356	0.958	-0.001	0.782	-0.001	(-0.165,0.153)	-0.857	0.223
MVPA		(25.819)		(0.003)		(0.078)		(0.702)	
Self-reported	138	-17.497	0.814	0.000	0.625	0.000	(-0.157,0.042)	-0.988	0.11
MVPA		(74.394)		(0.001)		(0.052)		(0.618)	

Notes: N varies due to invalid wear-time for PA, or non-completion of a full set of measures.

Appendix 6. Mediation effects of changes in accelerometer measured and self-reported MVPA at 4 months on the intervention effect on anxiety at 12 months.

Mediators	N	A path		B path		Mediated effect		C' path	
		β (SE)	P	β (SE)	P	β (SE)	95%CI	β (SE)	P
Bouted	89	1.705	0.793	-0.007	0.475	-0.012	(-0.309,0.135)	0.270	0.728
MVPA		(6.499)		(0.010)		(0.103)		(0.777)	
Continuous	92	1.356	0.958	-0.004	0.117	-0.005	(-0.437,0.307)	-0.042	0.956
MVPA		(25.819)		(0.003)		(0.158)		(0.758)	
Self-reported	119	-17.497	0.814	0.000	0.962	0.000	(-0.167,0.144)	0.083	0.909
MVPA		(74.394)		(0.001)		(0.066)		(0.722)	

Notes: N varies due to invalid wear-time for PA, or non-completion of a full set of measures.

Appendix 7. Mediation effects for intervention effects on processes at 4 months on depression at 12 months

Mediators	N	A path		B path		Mediated effect		C' path	
		β (SE)	P	β (SE)	P	β (SE)	95%CI	β (SE)	P
Importance	112	0.545 (0.428)	0.203	-0.136 (0.134)	0.308	-0.074 (0.112)	(-0.350,0.120)	-0.368 (0.824)	0.655
Confidence	113	1.075 (0.505)	0.033	-0.410 (0.124)	0.001	-0.441 (0.265)	(-1.024,0.018)	0.218 (0.802)	0.785
Competence	108	1.370 (0.621)	0.027	-0.242 (0.100)	0.016	-0.332 (0.221)	(-0.873,0.001)	0.343 (0.808)	0.671
Autonomy	111	0.481 (0.602)	0.424	-0.292 (0.103)	0.004	-0.140 (0.197)	(-0.551,0.246)	-0.132 (0.798)	0.868
Freq of support	113	0.685 (0.592)	0.247	-0.095 (0.101)	0.347	-0.065 (0.113)	(-0.324,0.129)	-0.283 (0.804)	0.725
Avail of support	112	0.187 (0.475)	0.693	-0.077 (0.133)	0.56	-0.014 (0.089)	(-0.173,0.197)	-0.241 (0.808)	0.766
Action planning	105	0.480 (0.851)	0.573	-0.122 (0.061)	0.045	-0.059 (0.114)	(-0.318,0.181)	-0.408 (0.821)	0.619
Self-monitoring	111	0.863 (0.358)	0.016	-0.256 (0.152)	0.093	-0.221 (0.181)	(-0.611,0.084)	-0.274 (0.811)	0.735

Notes: N varies due to non-completion of full set of measures.

Appendix 8. Mediation effects for intervention effects on processes at 4 months on anxiety at 12 months

Mediators	N	A path		B path		Mediated effect		C' path	
		β (SE)	P	β (SE)	P	β (SE)	95%CI	β (SE)	P
Importance	112	0.545 (0.428)	0.203	-0.117 (0.119)	0.328	-0.064 (0.098)	(-0.319,0.072)	-0.150 (0.724)	0.836
Confidence	113	1.075 (0.505)	0.033	-0.121 (0.114)	0.287	-0.130 (0.155)	(-0.509,0.119)	0.044 (0.738)	0.953
Competence	108	1.370 (0.621)	0.027	-0.079 (0.092)	0.392	-0.108 (0.145)	(-0.438,0.147)	0.236 (0.736)	0.748
Autonomy	111	0.481 (0.602)	0.424	-0.119 (0.096)	0.214	-0.057 (0.099)	(-0.265,0.136)	0.148 (0.745)	0.843
Freq of support	113	0.685 (0.592)	0.247	-0.070 (0.090)	0.439	-0.048 (0.097)	(-0.267,0.132)	-0.146 (0.725)	0.841
Avail of support	112	0.187 (0.475)	0.693	-0.122 (0.119)	0.308	-0.023 (0.082)	(-0.193,0.165)	-0.069 (0.720)	0.923
Action planning	105	0.480 (0.851)	0.573	-0.041 (0.054)	0.448	-0.020 (0.063)	(-0.158,0.100)	-0.197 (0.725)	0.786
Self-monitoring	111	0.863 (0.358)	0.016	-0.050 (0.143)	0.729	-0.043 (0.155)	(-0.359,0.282)	-0.118 (0.739)	0.873

Notes: N varies due to non-completion of full set of measures.