
Association of Tooth Loss in Chronic Periodontitis Patients with And Without Mellitus

PAVITHRA H DAVE¹, BALAJI GANESH S^{2*}, MAHESH RAMAKRISHNAN³

¹Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai 77

²Senior Lecturer, Department of Periodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai 77

³Reader, Department of Pedodontics and Preventive Dentistry, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai 77

*Corresponding Author

Email ID: 151501037.sdc@saveetha.com¹, balajiganeshs.sdc@saveetha.com², mahesh@saveetha.com³

Abstract: Tooth loss is a negative public health outcome that substantially affects not only the general and oral health status of people, but also their quality of life. Diabetes mellitus is one of the most widespread epidemics experienced worldwide; hence the aim of this study is to find the association of tooth loss in chronic periodontitis patients with and without diabetes mellitus. This cross sectional study consisted of a sample size of 547 patients of which 270 were diabetic patients and 277 were non diabetic patients and were distributed into two groups. After collection of data, it was tabulated with parameters such as name of the patient, age, gender, systemic condition, type of periodontal disease and tooth loss. The cast sheets of patients were obtained from the patient management software. The data of each patient was obtained and tabulated. The most common missing teeth were found to be upper anteriors and it had a higher prevalence of tooth loss in the non-diabetic group (26.82%) than in the diabetic group (18.43%). Association between tooth loss and systemic conditions was done using Chi square test p value- 0.008 and was found to be statistically significant. Tooth loss was higher in the age group of 61-70 years for the diabetic group and in the age group of 50-60 years for the non-diabetic group. The association between diabetes mellitus and periodontal disease is a well known fact. Therefore oral health education programs should be promoted in patients, which aids in early diagnosis and prevention of severe periodontal diseases.

Keywords: Adults; Diabetes; Gender; Periodontitis; Tooth loss, innovative

INTRODUCTION

Tooth loss and edentulism both are considered poor health outcomes that have a negative impact on a person's quality of life (Gerritsen et al., 2010). The prevalence of tooth loss increases with age and is a major problem for people 60 years and older. Adverse consequences of tooth loss include difficulty in chewing and speaking, esthetic dissatisfaction, and social stigma (Beltrán-Aguilar et al., 2005). Partial or total loss of permanent teeth is the result of a multifactorial process involving biological developments such as caries, periodontal disease, pulp pathology, trauma, and oral cancer, non-biological factors related to dental procedures, and sociodemographic and socioeconomic factors (Isabel Jimenez-Gayosso, 2015). The World Health Organization considers tooth loss a negative public health outcome that substantially affects not only the general and oral health status of people but also their quality of life. However, despite its predominance as part of the dental component of public health, tooth loss is often neglected (Steele et al., 2004). Researchers across the world have systematically reported the relevance of certain conditions, such as caries and periodontal disease, as major causes of tooth loss (Medina-Solis et al., 2013) (Kassebaum et al., 2014).

Oral health status plays an important role in the nutrition of children, adults, and older adults, with tooth loss linked to changes in taste, food preferences, and nutritional deficiency (Johnson, 2005) Salient among the negative effects of tooth loss are reduced chewing performance, limited food selection, weight loss, deficient communication, low self-esteem, and diminished well-being (Yoshida et al., 2005). Most dental problems share several risk factors, particularly unhealthy diets, and tobacco consumption, alcohol consumption, orally with other noncommunicable diseases, such as cardiovascular conditions, cancer, diabetes, and respiratory disorders (Marcenes et al., 2003). In 2010, the World Health Organization stated that this generated similar patterns of inequality among different population groups concerning oral and systemic illnesses. Those with diabetes

prevalence of the periodontal disease among patients with diabetes ranges from 22% through 68% for gingivitis and from 25% through 98% for periodontitis, and the prevalence rates of gingivitis and periodontitis in those without diabetes are markedly lower (Lee et al., 2015). Patients with type 2 diabetes are as much as 2.6 times more at risk of developing periodontal disease than those without diabetes. This appears to indicate that diabetes is an important modulator of periodontal inflammation and a leading cause of tooth loss. In a literature review, researchers established a strong association between periodontal disease and diabetes (Lewis et al., 2016). Increased life expectancy without a concomitant improvement in the quality of life directly affects health expenditure and is becoming an urgent public health problem throughout the world (Minaya-Sánchez et al., 2017).

Periodontitis is a multifactorial disease, with the primary etiological agent being plaque microflora. Research has also shown that periodontal microflora is similar to that found in atheromatous plaques (Priyanka et al., 2017). Cytokines play an important role in the pathogenesis and progression of periodontitis and the levels of TNF α , endothelins, IL - 21 and varies in chronic and aggressive periodontitis (Khalid et al., 2016, 2017; Mootha et al., 2016; Varghese et al., 2015). Periodontal disease can be a risk factor for cardiac disease and COPD (Mootha et al., 2016; Priyanka et al., 2017; Ramesh et al., 2019; Ramesh, Sheeja S. Varghese, et al., 2016) and it is important to manage the periodontal disease effectively and treat osseous defects for implant treatment (Ramesh et al., 2017). Platelet-rich fibrin and growth factors offer an advantage over other forms of treatment (Panda et al., 2014; Ravi et al., 2017). Researchers have attempted various regenerative methods (Avinash et al., 2017; Kavarthapu and Thamaraiselvan, 2018; Panda et al., 2014; Ravi et al., 2017) like PRF, growth factors and stem cells to salvage the destruction of periodontal tissues in chronic as well as aggressive forms of periodontitis (Thamaraiselvan et al., 2015). The treatment modalities such as antimicrobial therapy has provided added advantage in treating periodontitis cases. (Ramesh, Sheeja Saji Varghese, et al., 2016) (Ramamurthy and Mg, 2018). Our department is passionate about research we have published numerous high quality articles in this domain over the past years (Kavitha et al., 2014), (Praveen et al., 2001),(Devi and Gnanavel, 2014), (Putchala et al., 2013), (Vijayakumar et al., 2010), (Lekha et al., 2014a, 2014b) (Danda, 2010) (Danda, 2010) (Parthasarathy et al., 2016) (Gopalakannan et al., 2012), (Rajendran et al., 2019), (Govindaraju et al., 2017), (P. Neelakantan et al., 2015), (PradeepKumar et al., 2016), (Sajan et al., 2011), (Lekha et al., 2014a), (Neelakantan et al., 2013), (Patil et al., 2017), (Jeevanandan and Govindaraju, 2018), (Abdul Wahab et al., 2017), (Eapen et al., 2017), (Menon et al., 2018), (Wahab et al., 2018), (Vishnu Prasad et al., 2018), (Uthrakumar et al., 2010), (Ashok et al., 2017), (Prasanna Neelakantan et al., 2015). The aim of our study was to find the association of tooth loss in chronic periodontitis patients with and without diabetes mellitus reported to a private dental college.

MATERIALS AND METHOD

Study Setting

This was a cross sectional study of patients who had periodontitis and undergone tooth loss. It revolved around a university setting having patients visiting the Department of Periodontics of Saveetha Dental College and Hospital, Chennai, Tamil Nadu in between June 2019-April 2020. The approval for this study was obtained from the Institutional Ethical Committee (SDC/SIHEC/2020/DIASDATA/0619-0320).

Data Collection & Tabulation

The sample size consisted of 547 patients of which 270 were diabetic patients and 277 were non diabetic patients and were distributed into two groups. After collection of data, it was tabulated with parameters – name of the patient, age, gender, systemic condition, type of periodontal disease and tooth loss.

Statistical Analysis

After further verification of data by an external reviewer, it was imported to SPSS software developed by IBM for statistical analysis. Frequency, percentage of parameters was employed in the analysis. Chi square test was used to detect the association between gender, age, region of tooth loss. p value less than 0.05 was considered statistically significant.

RESULTS AND DISCUSSION

In this study, we conclude that there is a higher prevalence of tooth loss in chronic periodontitis patients in both the diabetic and non-diabetic groups. Tooth loss was higher in the age group of 61-70 years for the diabetic group and in the age group of 50-60 years for the non-diabetic group.

Graph 1 shows association of tooth loss with the systemic condition, upper anteriors had a higher prevalence of tooth loss in the non diabetic (26.82%) when compared to diabetic group (18.43%). Association was done using Chi square test p value- 0.008 and hence the two parameters, tooth loss and systemic conditions were found to be statistically significant. From graph 1 we infer that in the diabetic group, the highest incidence of tooth loss was seen in upper anteriors (18.43%) followed by upper molars (10.22%), lower anteriors (10.04%), lower molars

(6.20%), upper premolars (3.28%) and lower premolars (1.28%). In the non-diabetic group, the highest incidence of tooth loss was seen in upper anteriors (26.82%) followed by lower anteriors (8.03%), upper molars (7.85%), lower molars (4.01%), upper premolars (3.28%) and lower premolars (0.55%).

Graph 2 shows association of ages and tooth loss among diabetic patients, upper anteriors had a higher prevalence of tooth loss in the age group of 61-70 years (17.04%). Association was done using Chi square test p value- 0.003 and hence the two parameters, age groups and tooth loss were found to be statistically significant.

Graph 3 shows association of gender and tooth loss among diabetic patients, upper anteriors had a higher prevalence of tooth loss in male population (23.33%) when compared with the female population (14.07%). Association was done using Chi square test p value- 0.323 and hence the two parameters, gender and tooth loss were found to be not statistically significant.

Graph 4 shows association of types of periodontal disease and tooth loss among diabetic patients, upper anteriors had a higher prevalence of tooth loss in generalised chronic periodontitis (34.81%) when compared with localised chronic periodontitis (2.59%). Association was done using Chi square test p value- 0.133 and hence the two parameters, type of periodontal disease and tooth loss were found to be not statistically significant.

Graph 5 shows association of age groups and tooth loss among non-diabetic patients, upper anteriors had a higher prevalence of tooth loss in the age group 50-60 years (41.52%). Association was done using chi square test p value- 0.076 and hence the two parameters, age groups and tooth loss were not found to be statistically significant.

Graph 6 shows the association between gender and tooth loss among non-diabetic patients, upper anteriors had a higher prevalence of tooth loss in male population (31.41%) when compared with the female population (21.66%). Association was done using chi square test p value- 0.607 and hence the two parameters, gender and tooth loss were not found to be statistically significant.

Graph 7 shows the association between type of periodontal disease and tooth loss among non-diabetic patients, upper anteriors had a higher prevalence of tooth loss in generalised chronic periodontitis (44.4%) when compared with localised chronic periodontitis (8.66%). Association was done using chi square test p value- 0.353 and hence the two parameters, type of periodontal disease and tooth loss were not found to be statistically significant.

We undertook this study to ascertain whether a relationship existed between tooth loss and diabetes while adjusting for other variables. In this sample of Chennai adults, we observed a connection not only between tooth loss and diabetes but also between tooth loss and other kinds of variables such as age and gender. It has been observed that chronic diseases, such as diabetes, are associated with oral diseases. Researchers have also reported that tooth loss among middle-aged adults with chronic diseases is higher compared with the systemically healthy population, heightened risk of developing periodontitis among people with diabetes, which, along with cavities, is a risk factor for tooth loss and the most common reason for tooth extraction in the population (Mealey, 2006). This may be a result of the heightened risk is seen as closely linked to glycemic control and severity of the periodontal disease.

The study results by López et al showed that diabetes is associated independently with both edentulism and tooth loss in the U.S. population (Medina-Solís et al., 2006). The findings are consistent with the results of two other large studies conducted in the United States and Mexico in which investigators also reported greater tooth loss among people with diabetes. The researchers in the U.S. study used self-reported data regarding tooth loss and diabetes from the CDC's Behavioral Risk Factor Surveillance System to explore the association between diabetes and tooth loss among people 18 years and older (Kapp et al., 2007).

The global prevalence of diabetes in adults (> 18 years) increased from 4.7% in 1980 to 8.5% in 2014, and the World Health Organization predicts that it will reach 439 million people, nearly 10% of adults, in 2030. The results of this and our study and other research efforts in the United States and throughout the world indicate an association between diabetes and tooth loss; people with diabetes are more likely to lose more teeth than people without diabetes. The impact of diabetes on oral health has therefore acquired particular relevance for dental clinical practice, 45-48 bringing to light the need to better understand the mechanisms underlying the oral health implications of diabetes and to implement comprehensive treatment strategies for the prevention of possible complications (D'Aiuto et al., 2017).

Periodontitis is the second-largest health problem and the advanced periodontal disease affects 10 to 15% of the population worldwide (Silva et al., 2010). Recognition of the mounting evidence on the relationship between oral and systemic health will confront dentists, physicians, and other healthcare workers with the importance of working together. Nowhere is this more important than in the early identification of individuals with undiagnosed diabetes and the co-management of oral and overall health in patients with diabetes. There is sufficient evidence of the bidirectional relationship between diabetes and periodontal disease to formulate guidelines for screening undiagnosed diabetes and the co-management of patients with diabetes in the clinical practice of dentistry and dental hygiene. There is also sufficient evidence regarding the role periodontal disease plays in increasing systemic inflammation to suggest that non-dental healthcare providers should screen patients for periodontal disease.

For dental and non-dental practitioners who embrace the chance to become more actively involved in this vital area of healthcare, this new and exciting level of clinical follow is definite to benefit patients and be professionally pleasing and rewarding. Drumond-Santana *et al.* (2007) found a 49.1% prevalence of periodontitis, in a population of diabetic individuals in a public hospital, which is higher than the value found in the previous literature (de Pinho et al., 2012). The reason may be due to the sample assessed being restricted to a general hospital, which may reflect lesser proximity to the actual situation of the periodontal health of the diabetic population.

In the present study, tooth loss was more common in male diabetic patients and non diabetic patients when compared with females. This corroborates other studies reporting that periodontal problems are more prevalent and more severe among diabetic men. Such a finding implies that public healthcare services need to actively seek changes in clinic hours and other measures that address care for men with diabetes, as males seek health services less often than women. Our study has limitations that must be taken into account for an adequate interpretation of its results such as geographic limitation, confounding factors, and does not represent all the ethnic groups or populations from around the world. Future scope of the study includes extensive research to find the role of systemic diseases in the prevalence of periodontitis in diverse populations which can help in further diagnosis and treatment planning.

CONCLUSION

Within the limitations of the study, we conclude that the most common missing teeth were found to be upper anteriors and it had a higher prevalence of tooth loss in the non-diabetic group (26.82%) than in the diabetic group (18.43%). Tooth loss was higher in the age group of 61-70 years for the diabetic group and in the age group of 50-60 years for the non-diabetic group. The association between diabetes mellitus and periodontal disease is a well known fact. Therefore oral health education programs should be promoted in patients, which aids in early diagnosis and prevention of severe periodontal diseases.

Author Contributions

First author (Pavithra H Dave) performed the analysis, interpretation and wrote the manuscript. Second author (Dr.Balaji Ganesh S) contributed to conception, data design, analysis, interpretation and critically revised the manuscript. Third author (Dr.Mahesh) participated in the study and revised the manuscript. All the three authors have discussed the results and contributed to the final manuscript.

Conflicts of Interest

The authors declare no conflicts of interest.


REFERENCES

1. Abdul Wahab PU, Senthil Nathan P, Madhulaxmi M, et al. (2017) Risk Factors for Post-operative Infection Following Single Piece Osteotomy. *Journal of maxillofacial and oral surgery* 16(3): 328–332.
2. Ashok BS, Ajith TA and Sivanesan S (2017) Hypoxia-inducible factors as neuroprotective agent in Alzheimer's disease. *Clinical and experimental pharmacology & physiology* 44(3): 327–334.
3. Avinash K, Malaippan S and Dooraiswamy JN (2017) Methods of Isolation and Characterization of Stem Cells from Different Regions of Oral Cavity Using Markers: A Systematic Review. *International journal of stem cells* 10(1): 12–20.
4. Beltrán-Aguilar ED, Barker LK, Canto MT, et al. (2005) Surveillance for dental caries, dental sealants, tooth retention, edentulism, and enamel fluorosis--United States, 1988-1994 and 1999-2002. *Morbidity and mortality weekly report. Surveillance summaries* 54(3): 1–43.
5. D'Aiuto F, Gable D, Syed Z, et al. (2017) Evidence summary: The relationship between oral diseases and diabetes. *British dental journal* 222(12): 944–948.
6. Danda AK (2010) Comparison of a single noncompression miniplate versus 2 noncompression miniplates in the treatment of mandibular angle fractures: a prospective, randomized clinical trial. *Journal of oral and maxillofacial surgery: official journal of the American Association of Oral and Maxillofacial Surgeons*


- 68(7): 1565–1567.
7. de Pinho AMS, Borges CM, de Abreu MHNG, et al. (2012) Impact of periodontal disease on the quality of life of diabetics based on different clinical diagnostic criteria. *International journal of dentistry* 2012: 986412.
 8. Devi VS and Gnanavel BK (2014) Properties of Concrete Manufactured Using Steel Slag. *Procedia Engineering* 97: 95–104.
 9. Eapen BV, Baig MF and Avinash S (2017) An Assessment of the Incidence of Prolonged Postoperative Bleeding After Dental Extraction Among Patients on Uninterrupted Low Dose Aspirin Therapy and to Evaluate the Need to Stop Such Medication Prior to Dental Extractions. *Journal of maxillofacial and oral surgery* 16(1): 48–52.
 10. Gerritsen AE, Allen PF, Witter DJ, et al. (2010) Tooth loss and oral health-related quality of life: a systematic review and meta-analysis. *Health and quality of life outcomes* 8: 126.
 11. Gopalakannan S, Senthilvelan T and Ranganathan S (2012) Modeling and Optimization of EDM Process Parameters on Machining of Al 7075-B4C MMC Using RSM. *Procedia Engineering* 38: 685–690.
 12. Govindaraju L, Neelakantan P and Gutmann JL (2017) Effect of root canal irrigating solutions on the compressive strength of tricalcium silicate cements. *Clinical oral investigations* 21(2): 567–571.
 13. Isabel Jimenez-Gayosso S (2015) Socioeconomic inequalities in the use of oral health services (USSB) ever in life by mexican school children 6-12 years old. *GACETA. ACAD NACIONAL MEDICINA*
 14. Jeevanandan G and Govindaraju L (2018) Clinical comparison of Kedo-S paediatric rotary files vs manual instrumentation for root canal preparation in primary molars: a double blinded randomised clinical trial. *European Archives of Paediatric Dentistry*. DOI: 10.1007/s40368-018-0356-6.
 15. Johnson CSJ (2005) Psychosocial correlates of nutritional risk in older adults. *Canadian journal of dietetic practice and research: a publication of Dietitians of Canada = Revue canadienne de la pratique et de la recherche en dietetique: une publication des Dietetistes du Canada* 66(2): 95–97.
 16. Kapp JM, Boren SA, Yun S, et al. (2007) Diabetes and tooth loss in a national sample of dentate adults reporting annual dental visits. *Preventing chronic disease* 4(3): A59.
 17. Kassebaum NJ, Bernabé E, Dahiya M, et al. (2014) Global Burden of Severe Tooth Loss: A Systematic Review and Meta-analysis. *Journal of dental research* 93(7 Suppl): 20S–28S.
 18. Kavarthapu A and Thamaraiselvan M (2018) Assessing the variation in course and position of inferior alveolar nerve among south Indian population: A cone beam computed tomographic study. *Indian journal of dental research: official publication of Indian Society for Dental Research* 29(4): 405–409.
 19. Kavitha M, Subramanian R, Narayanan R, et al. (2014) Solution combustion synthesis and characterization of strontium substituted hydroxyapatite nanocrystals. *Powder Technology* 253: 129–137.
 20. Khalid W, Vargheese SS, Lakshmanan R, et al. (2016) Role of endothelin-1 in periodontal diseases: A structured review. *Indian journal of dental research: official publication of Indian Society for Dental Research* 27(3): 323–333.
 21. Khalid W, Vargheese SS, Sankari M, et al. (2017) Comparison of Serum Levels of Endothelin-1 in Chronic Periodontitis Patients Before and After Treatment. *Journal of clinical and diagnostic research: JCDR* 11(4): ZC78–ZC81.
 22. Lee C-Y, Chang Y-Y, Shieh T-Y, et al. (2015) Reasons for permanent tooth extractions in Taiwan. *Asia-Pacific journal of public health / Asia-Pacific Academic Consortium for Public Health* 27(2): NP2350–7.
 23. Lekha L, Kanmani Raja K, Rajagopal G, et al. (2014a) Schiff base complexes of rare earth metal ions: Synthesis, characterization and catalytic activity for the oxidation of aniline and substituted anilines. *Journal of organometallic chemistry* 753: 72–80.
 24. Lekha L, Kanmani Raja K, Rajagopal G, et al. (2014b) Synthesis, spectroscopic characterization and antibacterial studies of lanthanide(III) Schiff base complexes containing N, O donor atoms. *Journal of Molecular Structure*. DOI: 10.1016/j.molstruc.2013.10.014.
 25. Lewis A, Kitson A and Harvey G (2016) Improving oral health for older people in the home care setting: An exploratory implementation study. *Australasian journal on ageing* 35(4): 273–280.
 26. Marcenes W, Steele JG, Sheiham A, et al. (2003) The relationship between dental status, food selection, nutrient intake, nutritional status, and body mass index in older people. *Cadernos de saude publica* 19(3): 809–816.
 27. Mealey BL (2006) Periodontal disease and diabetes. A two-way street. *Journal of the American Dental Association* 137 Suppl: 26S–31S.
 28. Medina-Solís CE, Pérez-Núñez R, Maupomé G, et al. (2006) Edentulism among Mexican adults aged 35 years and older and associated factors. *American journal of public health* 96(9): 1578–1581.
 29. Medina-Solis CE, Pontigo-Loyola AP, Pérez-Campos E, et al. (2013) Principal reasons for extraction of permanent tooth in a sample of Mexicans adults. *Revista de investigacion clinica; organo del Hospital de Enfermedades de la Nutricion* 65(2): 141–149.
 30. Menon S, Ks SD, R S, et al. (2018) Selenium nanoparticles: A potent chemotherapeutic agent and an

- elucidation of its mechanism. *Colloids and surfaces. B, Biointerfaces* 170: 280–292.
31. Minaya-Sánchez M, Medina-Solís CE, Casanova-Rosado JF, et al. (2017) Prevalence of Functional Dentition in a Group of Mexican Adult Males. *Puerto Rico health sciences journal* 36(3): 146–151.
 32. Mootha A, Malaiappan S, Jayakumar ND, et al. (2016) The Effect of Periodontitis on Expression of Interleukin-21: A Systematic Review. *International journal of inflammation* 2016: 3507503.
 33. Neelakantan P, Grotra D and Sharma S (2013) Retreatability of 2 mineral trioxide aggregate-based root canal sealers: a cone-beam computed tomography analysis. *Journal of endodontia* 39(7): 893–896.
 34. Neelakantan P., Cheng CQ, Mohanraj R, et al. (2015) Antibiofilm activity of three irrigation protocols activated by ultrasonic, diode laser or Er:YAG laser in vitro. *International endodontic journal* 48(6): 602–610.
 35. Neelakantan Prasanna, Sharma S, Shemesh H, et al. (2015) Influence of Irrigation Sequence on the Adhesion of Root Canal Sealers to Dentin: A Fourier Transform Infrared Spectroscopy and Push-out Bond Strength Analysis. *Journal of endodontia* 41(7): 1108–1111.
 36. Panda S, Jayakumar ND, Sankari M, et al. (2014) Platelet rich fibrin and xenograft in treatment of intrabony defect. *Contemporary clinical dentistry* 5(4): 550–554.
 37. Parthasarathy M, Isaac Joshua Ramesh Lalvani J, Dhinesh B, et al. (2016) Effect of hydrogen on ethanol-biodiesel blend on performance and emission characteristics of a direct injection diesel engine. *Ecotoxicology and environmental safety* 134(Pt 2): 433–439.
 38. Patil SB, Durairaj D, Suresh Kumar G, et al. (2017) Comparison of Extended Nasolabial Flap Versus Buccal Fat Pad Graft in the Surgical Management of Oral Submucous Fibrosis: A Prospective Pilot Study. *Journal of maxillofacial and oral surgery* 16(3): 312–321.
 39. PradeepKumar AR, Shemesh H, Jothilatha S, et al. (2016) Diagnosis of Vertical Root Fractures in Restored Endodontically Treated Teeth: A Time-dependent Retrospective Cohort Study. *Journal of endodontia* 42(8): 1175–1180.
 40. Praveen K, Narayanan V, Muthusekhar MR, et al. (2001) Hypotensive anaesthesia and blood loss in orthognathic surgery: a clinical study. *The British journal of oral & maxillofacial surgery* 39(2): 138–140.
 41. Priyanka S, Kaarthikeyan G, Nadathur JD, et al. (2017) Detection of cytomegalovirus, Epstein-Barr virus, and Torque Teno virus in subgingival and atheromatous plaques of cardiac patients with chronic periodontitis. *Journal of Indian Society of Periodontology* 21(6): 456–460.
 42. Putchala MC, Ramani P, Herald J. Sherlin, et al. (2013) Ascorbic acid and its pro-oxidant activity as a therapy for tumours of oral cavity – A systematic review. *Archives of Oral Biology*. DOI: 10.1016/j.archoralbio.2013.01.016.
 43. Rajendran R, Kunjusankaran RN, Sandhya R, et al. (2019) Comparative Evaluation of Remineralizing Potential of a Paste Containing Bioactive Glass and a Topical Cream Containing Casein Phosphopeptide-Amorphous Calcium Phosphate: An in Vitro Study. *Pesquisa Brasileira em Odontopediatria e Clínica Integrada*. DOI: 10.4034/pboci.2019.191.61.
 44. Ramamurthy J and Mg V (2018) COMPARISON OF EFFECT OF HIORA MOUTHWASH VERSUS CHLORHEXIDINE MOUTHWASH IN GINGIVITIS PATIENTS: A CLINICAL TRIAL. *Asian J Pharm Clin Res* 11(7): 84–88.
 45. Ramesh A, Varghese Sheeja S., Jayakumar ND, et al. (2016) Chronic obstructive pulmonary disease and periodontitis – unwinding their linking mechanisms. *Journal of oral biosciences / JAOB, Japanese Association for Oral Biology* 58(1): 23–26.
 46. Ramesh A, Varghese Sheeja Saji, Doraiswamy JN, et al. (2016) Herbs as an antioxidant arsenal for periodontal diseases. *Journal of intercultural ethnopharmacology* 5(1): 92–96.
 47. Ramesh A, Ravi S and Kaarthikeyan G (2017) Comprehensive rehabilitation using dental implants in generalized aggressive periodontitis. *Journal of Indian Society of Periodontology* 21(2): 160–163.
 48. Ramesh A, Vellayappan R, Ravi S, et al. (2019) Esthetic lip repositioning: A cosmetic approach for correction of gummy smile - A case series. *Journal of Indian Society of Periodontology* 23(3): 290–294.
 49. Ravi S, Malaiappan S, Varghese S, et al. (2017) Additive Effect of Plasma Rich in Growth Factors With Guided Tissue Regeneration in Treatment of Intrabony Defects in Patients With Chronic Periodontitis: A Split-Mouth Randomized Controlled Clinical Trial. *Journal of periodontology* 88(9): 839–845.
 50. Sajan D, Udaya Lakshmi K, Erdogdu Y, et al. (2011) Molecular structure and vibrational spectra of 2,6-bis(benzylidene)cyclohexanone: a density functional theoretical study. *Spectrochimica acta. Part A, Molecular and biomolecular spectroscopy* 78(1): 113–121.
 51. Silina V and Kalda R (2018) Challenges for clinical practice and research in family medicine in reducing the risk of chronic diseases. Notes on the EGPRN Spring Conference 2017 in Riga. *The European journal of general practice* 24(1): 112–117.
 52. Silva AM, Vargas AMD, Ferreira EFE, et al. (2010) Periodontitis in individuals with diabetes treated in the public health system of Belo Horizonte, Brazil. *Revista brasileira de epidemiologia = Brazilian journal of epidemiology* 13(1): 118–125.


53. Steele JG, Sanders AE, Slade GD, et al. (2004) How do age and tooth loss affect oral health impacts and quality of life? A study comparing two national samples. *Community dentistry and oral epidemiology* 32(2): 107–114.
54. Thamaraiselvan M, Elavarasu S, Thangakumaran S, et al. (2015) Comparative clinical evaluation of coronally advanced flap with or without platelet rich fibrin membrane in the treatment of isolated gingival recession. *Journal of Indian Society of Periodontology* 19(1): 66–71.
55. Uthrakumar R, Vesta C, Raj CJ, et al. (2010) Bulk crystal growth and characterization of non-linear optical bithiourea zinc chloride single crystal by unidirectional growth method. *Current applied physics: the official journal of the Korean Physical Society* 10(2): 548–552.
56. Varghese SS, Thomas H, Jayakumar ND, et al. (2015) Estimation of salivary tumor necrosis factor-alpha in chronic and aggressive periodontitis patients. *Contemporary clinical dentistry* 6(Suppl 1): S152–6.
57. Vijayakumar GNS, Devashankar S, Rathnakumari M, et al. (2010) Synthesis of electrospun ZnO/CuO nanocomposite fibers and their dielectric and non-linear optic studies. *Journal of alloys and compounds* 507(1): 225–229.
58. Vishnu Prasad S, Kumar M, Ramakrishnan M, et al. (2018) Report on oral health status and treatment needs of 5-15 years old children with sensory deficits in Chennai, India. *Special care in dentistry: official publication of the American Association of Hospital Dentists, the Academy of Dentistry for the Handicapped, and the American Society for Geriatric Dentistry* 38(1): 58–59.
59. Wahab PUA, Madhulaxmi M, Senthilnathan P, et al. (2018) Scalpel Versus Diathermy in Wound Healing After Mucosal Incisions: A Split-Mouth Study. *Journal of oral and maxillofacial surgery: official journal of the American Association of Oral and Maxillofacial Surgeons* 76(6): 1160–1164.
60. Yoshida M, Morikawa H, Yoshikawa M, et al. (2005) Eight-year mortality associated with dental occlusion and denture use in community-dwelling elderly persons. *Gerodontology* 22(4): 234–237.




Graph 1: Bar graph reveals the association between tooth loss and systemic conditions. X- axis represents the systemic condition and Y- axis represents percentage of tooth loss. It was found that upper anteriors had a higher prevalence of tooth loss in the non diabetic (26.82%) and diabetic group (18.43%). Pearson chi square value- 15.569, p value- 0.008 <0.05 and hence the two parameters tooth loss and systemic conditions were found to be statistically significant, with significant increase in loss of upper anteriors among the non diabetic group.




Graph 2: Bar graph reveals the association between age groups and tooth loss among diabetic patients. X- axis represents the age groups and Y- axis represents percentage of tooth loss. It was found that upper anteriors had a higher prevalence of tooth loss in the age group 61-70 years (17.04%). Pearson chi square value- 26.521, p value- 0.003 <0.05 and hence the two parameters, age groups and tooth loss were found to be statistically significant.




Graph 3: Bar graph reveals the association between gender and tooth loss among diabetic patients. X- axis represents the gender and Y- axis represents percentage of tooth loss. It was found that upper anteriors had a higher prevalence of tooth loss in male population (23.33%) when compared with the female population (14.07%). Pearson chi square value- 5.834, p value- 0.323 >0.05 and hence the two parameters, gender and tooth loss were not found to be statistically significant.




Graph 4: Bar graph reveals the association between type of periodontal disease and tooth loss among diabetic patients. X- axis represents the type of periodontal disease and Y- axis represents percentage of tooth loss. It was found that upper anteriors had a higher prevalence of tooth loss in generalised chronic periodontitis (34.81%) when compared with localised chronic periodontitis (2.59%). Pearson chi square value- 8.459, p value- 0.133 >0.05 and hence the two parameters, type of periodontal disease and tooth loss were not found to be statistically significant.



Graph 5: Bar graph reveals the association between age groups and tooth loss among non-diabetic patients. X- axis represents the age groups and Y- axis represents percentage of tooth loss. It was found that upper anteriors had a higher prevalence of tooth loss in the age group 50-60 years (41.52%). Pearson chi square value- 23.408, p value- 0.076 >0.05 and hence the two parameters, age groups and tooth loss were not found to be statistically significant.



Graph 6: Bar graph reveals the association between gender and tooth loss among non-diabetic patients. X- axis represents the gender and Y- axis represents percentage of tooth loss. It was found that upper anteriors had a higher prevalence of tooth loss in male population (31.41%) when compared with the female population (21.66%). Pearson chi square value- 3.611, p value- 0.607 >0.05 and hence the two parameters, gender and tooth loss were not found to be statistically significant.



Graph 7: Bar graph reveals the association between type of periodontal disease and tooth loss among non-diabetic patients. X- axis represents the type of periodontal disease and Y- axis represents percentage of tooth loss. It was found that upper anteriors had a higher prevalence of tooth loss in generalised chronic periodontitis (44.4%) when compared with localised chronic periodontitis (8.66%). Pearson chi square value- 5.544, p value- 0.353 >0.05 and hence the two parameters, type of periodontal disease and tooth loss were not found to be statistically significant.