



THE AGA KHAN UNIVERSITY

eCommons@AKU

Internal Medicine, East Africa

Medical College, East Africa

5-2024

Psychiatric morbidity among patients on treatment for tuberculosis at a tertiary referral hospital in Western Kenya

Robina Momanyi

Edith Kwobah

Philip Owiti

Henry Nyamogoba

Lukoye Atwoli

Follow this and additional works at: https://ecommons.aku.edu/eastafrica_fhs_mc_intern_med



Part of the [Psychiatry Commons](#), and the [Psychiatry and Psychology Commons](#)

RESEARCH ARTICLE

Psychiatric morbidity among patients on treatment for tuberculosis at a tertiary referral hospital in Western Kenya

Robina Kerubo Momanyi^{1*}, Edith Kamaru Kwobah², Philip Owiti³, Henry Nyamogoba⁴, Lukoye Atwoli^{1,5,6}

1 Department of Mental Health and Behavioural Sciences, School of Medicine Moi University, Kenya, **2** Department of Mental Health, Moi Teaching and Referral Hospital, Kenya, **3** National Tuberculosis Program, Ministry of Health, Kenya, **4** Department of Medical Microbiology and Parasitology, School of Medicine Moi University, Kenya, **5** Department of Medicine, Medical College East Africa, Aga Khan University, Kenya, **6** Brain and Mind Institute Aga Khan University, Kenya

* robinamomanyi@gmail.com



Abstract

Background

Mental disorders account for nine percent of the overall global burden of disease and are among the top ten leading causes of disability. Mental illness and tuberculosis share risk factors including poverty, overcrowding, stigma, poor nutrition, substance use and retro-viral disease co-infection. Presence of mental illness in tuberculosis delays health-seeking, affects drug adherence, increases cost of treatment, prolongs disease duration, lowers quality of life, and increases mortality. Early diagnosis, linkage, and treatment of psychiatric morbidity among patients with tuberculosis would improve outcomes for both. This study thus aimed to determine the prevalence and factors associated with psychiatric morbidity among patients on treatment for tuberculosis at a low- middle- income country.

Methods

A cross-sectional study carried out at the tuberculosis clinic at Moi Teaching and Referral Hospital (MTRH), Eldoret, Kenya. 367 participants on TB treatment were interviewed using Mini-International Neuropsychiatric Interview (MINI) tool. The key outcome was presence of psychiatric illness. Pearson's Chi-square and logistic regression were used to assess relationships at bivariate and multivariate level respectively.

Results

Majority of the respondents were male (61.3%) and overall median age was 33 years. About half of participants (48.5%) had at least one psychiatric illness. Common disorders were alcohol use disorder (30.3%), depression (23.4%), substance use disorder (12.8%) and suicidality (8.2%). Odds of 'any psychiatric illness' were increased by being male (aOR = 1.92; $P = 0.04$), being separated or divorced (aOR = 6.86; $P = 0.002$), using alcohol (aOR = 3.2; $P < 0.001$), having been previously treated for tuberculosis (aOR = 2.76; $P = 0.01$), having

OPEN ACCESS

Citation: Momanyi RK, Kwobah EK, Owiti P, Nyamogoba H, Atwoli L (2024) Psychiatric morbidity among patients on treatment for tuberculosis at a tertiary referral hospital in Western Kenya. PLOS ONE 19(5): e0302744. <https://doi.org/10.1371/journal.pone.0302744>

Editor: Petros Isaakidis, Médecins Sans Frontières (MSF), SOUTH AFRICA

Received: July 25, 2023

Accepted: April 11, 2024

Published: May 15, 2024

Copyright: © 2024 Momanyi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its [Supporting Information](#) files.

Funding: The author(s) received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

other medical comorbidities (aOR = 4.2; $P = 0.004$) and family history of mental illness (aOR = 2.4; $P = 0.049$).

Conclusion

Almost half of the patients on treatment for tuberculosis had at least one psychiatric illness. Introduction of protocols for screening for mental illness and integration of mental health services with tuberculosis care would aid prompt diagnosis, referral, and quality of care.

Introduction

Mental disorders account for nine percent of the overall global burden of disease and are listed among the top ten leading causes of disability. Tuberculosis (TB) is the second leading cause of death from a single infectious disease with an estimated 1.6 million deaths in 2021 [1]. Patients on treatment for TB are at higher risk for mental disorders as they have various triggers and shared risk factors including stigma associated with TB and HIV in case of co-infection, overcrowding, poor nutrition, smoking, alcohol use, lower income and isolation in case of multi-drug resistant tuberculosis [2–5]. Physical weakness caused by TB leads to decreased work productivity, absence from work and later job insecurity and financial instability that could be precursors or triggers of mental illness [6, 7]. Additionally, certain anti-tuberculosis drugs like isoniazid and ethambutol have side effects that include mood fluctuation, psychosis, seizures, mental confusion, dizziness and disorientation [8] and whether this relates to eventual psychiatric morbidity is yet to be ascertained.

Research on the prevalence of mental illness among patients on treatment for TB has mostly looked at depression and anxiety with a range of 35–84% for depression and 36–65% for anxiety in Ethiopia, Pakistan, India, Cameroon and South Africa [9–13]. A study from Kenya, found a prevalence of depression at 31% among 51 TB patients [14]. Studies in Zambia, Thailand and India of patients on treatment for TB found a range of 20%–30% for alcohol use disorder [15–18] while substance use disorder ranged from 3%–39% in Pakistan, Peru and Ethiopia [19–21]. Prevalence of mental illness varies depending on region and screening tool used but is higher when compared to that in the general population [22–24]. Few studies have explored psychiatric morbidity as a whole in the region and specifically in Kenya. As we aim for multisectoral collaboration to combat the TB epidemic, and as TB care teams work on holistic patient-centered care, the psychological distress and associated mental illness related to TB, cannot be ignored as it may burden the patient and affect treatment outcomes.

The presence of mental illness in TB delays health-seeking, affects drug adherence, increases cost of treatment, prolongs disease duration, lowers quality of life, and increases mortality [2, 3]. The potential for mental illness to impair adherence to TB medication increases risk of TB transmission and the development of multidrug resistance [25] and as such is a threat to public health. Mental illness affecting compliance to both anti-TB medication and anti-psychotics would lead to relapse of mental illness and further re-infection with TB [7, 25]. Knowing the prevalence, and type of patients at higher risk of mental illness among patients on treatment for TB in high TB burden countries, is crucial in directing policy to integrate mental health and TB care. Within this context, the study sought to determine the prevalence of psychiatric illness among patients on treatment for TB in Kenya. The researchers also sought to establish socio-demographic and clinical determinants, associated with psychiatric morbidity among patients on treatment for TB.

Methods

Study design

This was a cross-sectional study conducted between March 2019 and March 2020.

Study setting

The study was done at the TB clinic of Moi Teaching and Referral Hospital (MTRH)—the second largest public referral hospital in Kenya. The hospital, located in the western part of the country has a wide catchment population of approximately 24 million and serves patients from both urban and rural setups. Kenya had an estimated TB incidence of 133,000 in 2021 and is ranked among the high TB and TB/HIV burden countries by the World Health Organization [1]. The TB incidence was 146,000 in 2018 prior to this study [26] and 136,000 in 2019 [27].

Study population

All adult patients enrolled for TB treatment at the TB clinic at MTRH were eligible for recruitment into the study. On average from the TB clinic records, 400 patients were seen at the clinic every month with about 50–80 of them being newly-diagnosed with TB. Others were patients who had come for revisits and drug refill. This data was obtained in August, 2018 at the time of proposal development and was the monthly average for the preceding seven months.

Sample size

Sample size was derived from Fishers formula where p was the estimated proportion of TB patients with mental illness of 31% (19), $q = 1 - p$ (i.e. 0.69), resulting in a sample size of 329 participants. Assuming a 10% non-response the sample size was increased to a minimum of 362 respondents.

Eligibility criteria

All adult patients with TB enrolled at the MTRH TB clinic were included. Patients who were physically severely ill or in distress or hospitalized were excluded from the study.

Data collection

Recruitment of study participants and data collection were done simultaneously between 1st August, 2019 and 28th February, 2020. Interviewer -administered questionnaires were used. The first questionnaire covered socio-demographics (age, gender, marital status, level of education, occupation) and other clinical and TB characteristics including history of cigarette smoking or alcohol use, type of TB, patient type, comorbid medical illness, phase of TB treatment (initiation or continuation), prior diagnosis of psychiatric illness and family history of mental illness. The second questionnaire was the Mini International Neuropsychiatric Interview (MINI) based on Version 7 of Diagnostic and Statistical Manual 5th edition (DSM-5) [28]. MINI has sixteen modules spanning a wide range of mental illness including major depressive disorder, suicidality, bipolar disorder, post-traumatic stress disorder, alcohol and substance use disorders, psychotic disorder, eating disorders, generalized anxiety disorder and antisocial personality disorder. Its validity and reliability have been assessed across multiple settings with good results [28, 29]. It has been used previously in Kenya after adaptation and adoption [30–32]. Every third patient who met eligibility criteria was approached and if they gave consent they were interviewed. This was repeated until the sample size was achieved. To

get the interval of every 3rd patient, we divided the average of 1200 patients expected in 3 months by the sample size $1200/367 = 3.26$. This was rounded to every 3rd patient to approximate the total expected patients in the duration. The data collection however took 6 months because after the initial 3 months most of the patients were revisits who had already been interviewed and hence a smaller pool to draw from.

Analysis

The data was cleaned and analyzed using Stata v.14 (*StataCorp College Station, TX, USA*). Recategorization was carried out for some variables like age, smoking, alcohol use and the psychiatric illness. Discrete variables were summarized as frequencies and percentages and displayed as tables while continuous variables were represented with median and interquartile ranges (IQR) and categorized during further analyses.

The main outcome was presence of any psychiatric illness(es) and these were presented with their 95% confidence intervals (CI). Associations between the outcomes and the socio-demographic and clinical characteristics were assessed using Pearson's χ^2 with the strengths of association presented as odds ratio (OR) with their 95% CI. Multivariable logistic regression was used to identify factors independently associated with the main outcomes. All the socio-demographic and clinical exposure variables were included in the multivariable models unless there was collinearity e.g., HIV, diabetes and hypertension which are related with 'any comorbidity'. The level of significance was set at $P < 0.05$.

Ethical considerations

Ethical approval and permission were obtained from the Moi University/ Moi Teaching and Referral Hospital Institutional Research and Ethics Committee (IREC), FAN: IREC 3276 and MTRH management respectively. Participation was entirely voluntary and all enrolled participants gave voluntary written informed consent. Anonymity was preserved by de-identification of data. Authors did not have access to information that could identify individual participants during or after data collection as only study numbers were used. Patients assessed to have serious mental health illnesses, including suicidality were referred to the mental health unit of the hospital.

Results

Socio-demographic and clinical characteristics of study participants

Of the 367 patients interviewed, majority ($n = 225, 61\%$) were males with an overall median age of 33 years (IQR: 25–45). Slightly more than half of the patients (52%) were aged between 18–34 years. The proportion married ($n = 155$) was comparable to those single ($n = 153$) at 42% each, while 43 of the respondents (11.7%) were either divorced or separated. Majority ($n = 354, 96\%$) of the participants had at least a primary level of education. In fact, almost a third ($n = 105, 28\%$) had attained a tertiary level of education. Half ($n = 184$) of the patients worked in the informal sector. Half ($n = 183$) of the patients reported previous or current use of alcohol. Sociodemographic characteristics are shown in [Table 1](#).

Most of the patients interviewed ($n = 310, 85\%$) were new (taking anti-TBs for the first time) while slightly more than half ($n = 201, 55\%$) were in the initiation phase of treatment (first two months of treatment and on a regimen of 4 drugs). Most of the patients ($n = 284, 77\%$) had pulmonary type TB, of which 89% was bacteriologically confirmed (sputum positive). Of the 83 patients with extrapulmonary TB, 8 had TB meningitis. Two-fifths ($n = 146$) of the respondents had at least one comorbidity, of which the commonest was HIV at 30%

Table 1. Socio-demographic and clinical characteristics of patients on treatment for tuberculosis.

Characteristic	Sub- category	n	(%)
Sex	Male	225	(61.3%)
	Female	142	(38.7%)
Age, years, median (IQR)		33	(25–45)
	18 -34yrs	190	(51.8%)
	35 -49yrs	116	(31.6%)
	50 -64yrs	48	(13.1%)
	65+yrs	13	(3.5%)
Marital status	Single	153	(41.7%)
	Married	155	(42.2%)
	Divorced /separated	43	(11.7%)
	Widowed	16	(4.4%)
Education	No formal education	13	(3.8%)
	Primary	117	(32.1%)
	Secondary	129	(35.4%)
	Tertiary	105	(28.9%)
Occupation	Formal	52	(14.2%)
	Informal	184	(50.1%)
	Unemployed	72	(19.6%)
	Student	59	(16.1%)
Smoking	Never	274	(74.6%)
	Previously	77	(21%)
	Current	16	(4.4%)
Alcohol use	Never	183	(49.8%)
	Previously	157	(42.8%)
	Current	27	(7.4%)
Patient type	Newly diagnosed with TB	310	(84.5)
	Previously treated	57	(15.5)
TB type	PTB Bacteriologically confirmed	252	(68.7)
	PTB clinically diagnosed	32	(8.7)
	Extra-pulmonary TB	83	(22.6)
Treatment phase	Initiation	201	(54.8)
	Continuation	166	(45.2)
Comorbidities	Any comorbidity	146	(39.8)
	HIV	110	(30)
	Hypertension	17	(4.6)
	Diabetes	18	(4.9)
	Others	16	(4.3)
Psychiatric diagnosis	Never	358	(97.6)
	Previously	6	(1.6)
	Current	3	(0.8)
Psychiatric Family history	Yes	40	(10.9)
	No	327	(89.1)

TB–tuberculosis; PTB–pulmonary TB

<https://doi.org/10.1371/journal.pone.0302744.t001>

followed by diabetes (4.9%) and hypertension (4.6%). Other comorbidities included asthma, anaemia, renal disease, and cardiac disease. As regards mental illness, 358 (97.6%) of the respondents had never had a psychiatric diagnosis before while 40 (11%) had a family history of psychiatric illness. The clinical characteristics are seen in [Table 1](#).

Prevalence of psychiatric illnesses

Of the respondents, 48.5% ($n = 178$) had at least one psychiatric disorder (95% CI: 43.3–53.8). The most common psychiatric disorder among patients on treatment for TB was alcohol use disorder (AUD) at a prevalence of 30.3%. Major depressive disorder had a prevalence of 23.4% followed by substance use disorder at 12.8% and suicidality at 8.2%. Other psychiatric illnesses that were less common added up to 14.6% as in [Table 2](#).

Determinants associated with ‘any psychiatric illness’

Of the 178 respondents with psychiatric illness, males were almost twice as likely to have any psychiatric illness compared to females (aOR 1.92, 95% CI 1.03–3.56). Patients aged 65 years and above had 88% reduced odds of any psychiatric illness as compared to those 18–34 years old (aOR 0.12, 95% CI 0.02–0.62). Being separated or divorced had almost seven times increased odds (aOR 6.86, 95% CI 2.03–23.2) of being associated with any psychiatric illness compared to the married. History of alcohol use was associated with three times the likelihood of psychiatric morbidity (aOR 3.2, 95% CI 1.68–6.08), while being previously treated for TB (aOR 2.76, 95% CI 1.26–6.04) and family history of psychiatric illness (aOR 2.40, 95% CI 1.00–5.76) was associated with more than double the likelihood. Having extra-pulmonary TB was associated with half the odds of any psychiatric illness (aOR 0.51, 95% CI 0.27–0.98). Other factors like education, occupation, treatment phase and presence of comorbidities were not significantly associated with any psychiatric morbidity as shown in [Table 3](#).

Determinants associated with specific psychiatric disorders

Being separated or divorced (aOR 2.44, 95% CI 1.06–5.62) or having been previously treated for TB (aOR 2.03, 95% CI 1.00–4.1) was associated with increased odds of major depressive

Table 2. Prevalence of specific psychiatric disorders.

<i>Psychiatric disorder</i>	<i>n (%)</i>	<i>(95% CI)</i>
<i>Alcohol use disorder</i>	111 (30.3%)	(25.6–35.2)
<i>Major depressive disorder</i>	86 (23.4%)	(19.2–28.1)
<i>Substance use disorder</i>	47 (12.8%)	(9.6–16.7)
<i>Suicidality</i>	30 (8.2%)	(5.6–11.5)
<i>Suicide behaviour disorder</i>	9 (2.5%)	(1.1–4.6)
<i>Mood disorder with psychotic features</i>	9 (2.5%)	(1.1–4.6)
<i>Antisocial personality disorder</i>	7 (1.9%)	(0.8–3.9)
<i>Psychotic disorder</i>	6 (1.6%)	(0.6–3.5)
<i>Panic disorder</i>	5 (1.4%)	(0.4–3.2)
<i>Obsessive compulsive disorder</i>	4 (1.1%)	(0.3–2.8)
<i>Bipolar disorder</i>	4 (1.1%)	(0.3–2.8)
<i>Generalized anxiety disorder</i>	3 (0.8%)	(0.2–2.4)
<i>Post-traumatic stress disorder</i>	2 (0.5%)	(0.1–2.0)
<i>Manic episode</i>	1 (0.3%)	(0.0–1.5)
<i>Hypomanic episode</i>	1 (0.3%)	(0.0–1.5)
<i>Agoraphobia</i>	1 (0.3%)	(0.0–1.5)
<i>Social anxiety disorder</i>	1 (0.3%)	(0.0–1.5)

<https://doi.org/10.1371/journal.pone.0302744.t002>

Table 3. Determinants associated with 'any psychiatric illnesses'.

Characteristic	Sub-category	Yes		No		Bivariate		Multivariate		
		n	(%)	n	(%)	cOR	p-value	aOR	95% CI	p-value
Gender	Female	47	(33.1)	95	(66.9)	1		1		
	Male	131	(58.2)	94	(41.8)	2.82	<0.001	1.92	1.03–3.56	0.039
Age (yrs)	18–34	73	(38.4)	117	(61.6)	1		1		
	35–49	72	(62.1)	44	(37.9)	2.62	<0.001	0.86	0.43–1.71	0.66
	50–64	30	(62.5)	18	(37.5)	2.67	<0.001	1.14	0.44–2.96	0.78
	65+	3	(23.1)	10	(76.9)	0.48	0.28	0.12	0.02–0.62	0.011
Education	No formal education	7	(53.9)	6	(46.1)	1		1		
	Primary	71	(60.7)	46	(39.3)	1.32	0.63	0.60	0.14–2.65	0.51
	Secondary	66	(51.2)	63	(48.8)	0.90	0.85	0.68	0.15–3.00	0.61
	Tertiary	31	(29.5)	74	(70.5)	0.36	0.09	0.30	0.06–1.44	0.13
Occupation	Formal	24	(46.1)	28	(53.9)	1		1		
	Informal	102	(55.4)	82	(44.6)	1.45	0.24	0.76	0.34–1.73	0.52
	Unemployed	40	(55.6)	32	(44.4)	1.46	0.30	1.35	0.53–3.45	0.53
	Student	12	(20.3)	47	(79.7)	0.30	0.01	0.49	0.16–1.46	0.20
Marital status	Married	78	(50.3)	77	(49.7)	1		1		
	Single	55	(36)	98	(64)	0.55	0.01	0.96	0.50–1.87	0.91
	Separated /divorced	39	(90.7)	4	(9.3)	9.61	<0.001	6.86	2.03–23.2	0.002
	Widowed	6	(37.5)	10	(62.5)	0.59	0.33	0.67	0.18–2.50	0.56
Comorbidities	Without	94	(42.5)	127	(57.5)	1		1		
	With	84	(57.5)	62	(42.5)	1.83	0.01	1.41	0.79–2.53	0.25
HIV status	No	113	(44)	144	(56)	1		1		
	Yes	65	(59.1)	45	(40.9)	1.84	0.01			
Smoking	Never	105	(38.3)	169	(61.7)	1		1		
	Yes	73	(78.5)	20	(21.5)	5.87	<0.001	1.51	0.72–3.19	0.28
Alcohol use	Never	49	(26.8)	134	(73.2)	1		1		
	Yes	129	(70.1)	55	(29.9)	6.41	<0.001	3.20	1.68–6.08	<0.0001
Patient type	New	136	(43.9)	174	(56.1)	1		1		
	Previously treated	42	(73.7)	15	(26.3)	3.58	<0.001	2.76	1.26–6.04	0.01
Type of TB	PTB bacteriological	132	(52.4)	120	(47.6)	1		1		
	PTB clinically diagnosed	16	(50)	16	(50)	0.91	0.80	0.62	0.24–1.62	0.33
	Extra-PTB	30	(36.1)	53	(63.9)	0.51	0.01	0.51	0.27–0.98	0.04
Treatment phase	Initiation	92	(45.8)	109	(54.2)	1		1		
	Continuation	86	(51.8)	80	(48.2)	1.27	0.25	1.20	0.71–2.02	0.50
Family psychiatric history	No	152	(46.5)	175	(53.5)	1		1		
	Yes	26	(65)	14	(35)	2.14	0.03	2.40	1.00–5.76	0.05

PTB–pulmonary TB; Extra-PTB–extrapulmonary TB

<https://doi.org/10.1371/journal.pone.0302744.t003>

disorder while age 65 years and above was associated with reduced odds (aOR 0.09, 95% CI 0.01–0.89) (Table 4).

Being male (aOR 7.98, 95% CI 3.5–18.2), separated or divorced (aOR 3.51, 95% CI 1.3–9.65) or having history of smoking (aOR 5.03, 95% CI 2.5–10.2) was associated with increased odds of alcohol use disorder while being a student (aOR 0.02, 95% CI 0.0–0.25) and having clinically diagnosed TB (aOR 0.10, 95% CI 0.03–0.4) were associated with decreased odds (Table 5).

Being male (aOR 16.5, 95% CI 1.9–145.2), having a history of smoking (aOR 3.57, 95% CI 1.41–9.03) or alcohol use (aOR 7.69, 95% CI 1.42–41.5) or family history of psychiatric illness

Table 4. Characteristics of TB patients with major depressive disorder.

Characteristic	Sub-category	Yes		No		Bivariate		Multivariate		
		n	(%)	n	(%)	cOR	p-value	aOR	95% CI	p-value
Gender	Female	38	(26.8)	104	(73.2)	1		1		
	Male	47	(20.9)	178	(79.1)	0.72	0.19	0.86	0.45–1.66	0.66
Age (yrs)	18–34	40	(21.0)	150	(79.0)	1		1		
	35–49	33	(28.4)	83	(71.6)	1.49	0.14	0.77	0.38–1.56	0.47
	50–64	11	(22.9)	37	(77.1)	1.11	0.78	0.56	0.21–1.49	0.24
	65+	1	(7.7)	12	(92.3)	0.31	0.27	0.09	0.01–0.89	0.04
Education	No formal education	5	(38.5)	8	(61.5)	1		1		
	Primary	34	(29.1)	83	(70.9)	0.66	0.49	0.45	0.11–1.95	0.29
	Secondary	31	(24.0)	98	(76.0)	0.51	0.26	0.42	0.10–1.85	0.24
	Tertiary	14	(13.3)	91	(86.7)	0.25	0.03	0.24	0.05–1.17	0.08
Occupation	Formal	8	(15.4)	44	(84.6)	1		1		
	Informal	42	(22.8)	142	(77.2)	1.63	0.25	1.01	0.40–2.57	0.98
	Unemployed	29	(40.3)	43	(59.7)	3.71	0.004	2.51	0.93–6.80	0.07
	Student	6	(10.2)	53	(89.8)	0.62	0.41	0.52	0.15–1.87	0.32
Marital status	Married	31	(20.0)	124	(80.0)	1		1		
	Single	30	(19.6)	123	(80.4)	0.98	0.93	1.02	0.51–2.06	0.95
	Separated /divorced	21	(48.8)	22	(51.2)	3.82	<0.001	2.44	1.06–5.62	0.04
	Widowed	3	(18.8)	13	(81.2)	0.92	0.91	0.95	0.22–4.04	0.94
Comorbidities	Without	43	(19.5)	178	(80.5)	1		1		
	With	42	(28.8)	104	(71.2)	1.67	0.04	1.46	0.80–2.66	0.21
Smoking	Never	61	(22.3)	213	(77.7)	1		1		
	Yes	24	(25.8)	69	(74.2)	1.21	0.48	0.95	0.43–2.06	0.89
Alcohol use	Never	40	(21.9)	143	(78.1)	1		1		
	Yes	45	(24.5)	139	(75.5)	1.15	0.56	0.92	0.45–1.87	0.81
Patient type	New	65	(21.0)	245	(79.0)	1		1		
	Previously treated	20	(35.1)	37	(64.9)	2.04	0.02	2.03	1.00–4.10	0.05
Type of TB	PTB bacteriological	57	(22.6)	195	(77.4)	1		1		
	PTB clinically diagnosed	12	(37.5)	20	(62.5)	2.05	0.07	1.80	0.72–4.51	0.21
	Extra-PTB	16	(19.3)	67	(80.7)	0.82	0.52	0.91	0.46–1.82	0.80
Treatment phase	Initiation	49	(24.4)	152	(75.6)	1		1		
	Continuation	36	(21.7)	130	(78.3)	0.86	0.54	0.88	0.51–1.53	0.65
Family psychiatric history	No	70	(21.4)	257	(78.6)	1		1		
	Yes	15	(37.5)	25	(62.5)	2.20	0.03	1.78	0.81–3.91	0.15

PTB–pulmonary TB; Extra-PTB–extrapulmonary TB

<https://doi.org/10.1371/journal.pone.0302744.t004>

(aOR 3.97, 95% CI 1.32–12.0) were statistically associated with substance use disorder (Table 6).

Having other medical comorbidities like hypertension, diabetes and HIV among others was fourfold associated with suicidality (aOR 4.2, 95% CI 1.57–11.3) after adjustments.

Discussion

Prevalence of psychiatric morbidity

This study found that about half of the patients on treatment for TB (48.5%) had at least one mental illness, almost all of which was prior undiagnosed (97.6%). The most common

Table 5. Characteristics of TB patients with alcohol use disorder.

Characteristic	Sub-category	Yes		No		Bivariate		Multivariate		
		n	(%)	n	(%)	cOR	p-value	aOR	95%CI	p-value
Gender	Female	12	(8.4)	130	(91.6)	1		1		
	Male	99	(44)	126	(56)	8.5	<0.001	7.98	3.5–18.2	<0.001
Age (yrs)	18–34	37	(19.5)	153	(80.5)	1		1		
	35–49	51	(44)	65	(56)	3.24	<0.001	0.87	0.39–1.9	0.73
	50–64	21	(43.8)	27	(56.2)	3.21	0.001	0.83	0.3–2.3	0.73
	65+	2	(15.4)	11	(84.6)	0.75	0.71	0.26	0.04–2.0	0.20
Education	No formal education	3	(23.1)	10	(76.9)	1		1		
	Primary	50	(42.7)	67	(57.3)	2.49	0.18	1.09	0.22–5.4	0.91
	Secondary	40	(31)	89	(67)	1.50	0.55	1.10	0.22–5.5	0.90
	Tertiary	16	(15.2)	89	(84.8)	0.60	0.47	0.32	0.06–1.8	0.06
Occupation	Formal	19	(36.5)	33	(63.5)	1		1		
	Informal	70	(38)	114	(62)	1.07	0.84	0.42	0.2–1.04	0.06
	Unemployed	21	(29.2)	51	(70.8)	0.72	0.39	0.53	0.18–1.5	0.24
	Student	1	(1.7)	58	(98.3)	0.03	0.001	0.02	0.0–0.25	0.002
Marital status	Married	55	(35.5)	100	(64.5)	1		1		
	Single	26	(17)	127	(83)	0.37	<0.001	0.94	0.42–2.1	0.89
	Sep /divorced	26	(60.5)	17	(39.5)	2.78	0.004	3.51	1.3–9.65	0.02
	Widowed	4	(25)	12	(75)	0.61	0.41	0.98	0.21–4.4	0.98
Comorbidities	Without	59	(26.7)	162	(73.3)	1		1		
	With	52	(35.6)	94	(64.4)	1.52	0.07	1.39	0.72–2.7	0.32
HIV status	No	68	(26.5)	189	(73.5)	1		1		
	Yes	43	(39.1)	67	(60.9)	1.78	0.02			
Smoking	Never	48	(17.5)	226	(82.5)	1		1		
	Yes	63	(67.7)	30	(32.3)	9.89	<0.001	5.03	2.5–10.2	<0.001
Patient type	New	81	(26.1)	229	(73.9)	1		1		
	Previously treated	30	(52.6)	27	(47.4)	3.14	<0.001	2.16	0.96–4.8	0.06
Type of TB	PTB bacteriological confirmed	87	(34.5)	165	(65.5)	1		1		
	PTB clinically diagnosed	5	(15.6)	27	(84.4)	0.35	0.04	0.10	0.03–0.4	0.001
	Extra-PTB	19	(22.9)	64	(77.1)	0.56	0.05	0.65	0.31–1.4	0.26
Treatment phase	Initiation	57	(28.4)	144	(71.6)	1		1		
	Continuation	54	(32.5)	112	(67.5)	1.22	0.39	0.85	0.46–1.6	0.61
Family psychiatric history	No	97	(29.7)	230	(70.3)	1		1		
	Yes	14	(35)	26	(65)	1.28	0.49	0.79	0.31–2.0	0.62

PTB–pulmonary TB; Extra-PTB–extrapulmonary TB

<https://doi.org/10.1371/journal.pone.0302744.t005>

disorders were alcohol use disorder (in one-third of the patients), depression (in one-in-four patients) while substance use disorder and suicidality were each prevalent in about one-in-ten patients. Generalized anxiety disorder and post-traumatic disorder were among the other less common disorders with both combined affecting one-in-a-hundred patients.

The 48.5% prevalence of any mental illness is higher compared to findings of a study in Pakistan with a rate of 31%, Nigeria at 32% and a pooled rate of 34% in a meta-analysis [19, 33, 34]. It is however lower than 82% rate in South Africa [35]. Prevalence of mental illness among patients on treatment for TB is anticipated to be higher due to shared risk factors including alcohol use, smoking, overcrowding, stigma, association with HIV, psychosocial and economic stressors or due to compromised immunity and neglected self-care associated with mental

Table 6. Characteristics of TB patients with substance use disorder.

Characteristic	Sub-category	Yes		No		Bivariate		Multivariate		
		n	(%)	n	(%)	cOR	p-value	aOR	95%CI	p-value
Gender	Female	1	(0.7)	141	(99.3)	1		1		
	Male	46	(20.4)	179	(79.6)	36.2	0.00	16.5	1.9–145.2	0.01
Age (yrs)	18–34	16	(8.4)	174	(91.6)					
	35–49	22	(19)	94	(81)	2.55	0.00	0.79	0.27–2.32	0.66
	50–64	9	(18.8)	39	(81.2)	2.51	0.04	1.14	0.29–4.45	0.85
	65+	0	(0)	13	(100)	-		-		
Education	No formal education	1	(7.7)	12	(92.3)	1		1		
	Primary	25	(21.4)	92	(78.6)	3.26	0.28	1.23	0.81–18.7	0.88
	Secondary	15	(11.6)	114	(88.4)	1.58	0.67	0.78	0.05–12.0	0.05
	Tertiary	4	(3.8)	101	(96.2)	0.48	0.52	0.22	0.01–4.41	0.33
Occupation	Formal	5	(9.6)	47	(90.4)	1		1		
	Informal	33	(17.9)	151	(82.1)	2.05	0.16	0.81	0.24–2.78	0.74
	Unemployed	7	(9.7)	65	(90.3)	1.01	0.98	0.62	0.13–2.96	0.55
	Student	2	(3.4)	57	(96.6)	0.33	0.2	0.71	0.09–5.84	0.75
Marital status	Married	19	(12.3)	136	(87.7)	1		1		
	Single	13	(8.5)	140	(91.5)	0.66	0.28	2.12	0.66–6.88	0.21
	Sep /divorced	13	(30.2)	30	(69.8)	3.1	0.00	2.38	0.82–6.87	0.11
	Widowed	2	(12.5)	14	(87.5)	1.02	0.98	2.17	0.27–17.5	0.47
Comorbidities	Without	28	(12.7)	193	(87.3)	1		1		
	With	19	(13)	127	(87)	1.03	0.92	1.03	0.43–2.44	0.95
Smoking	Never	11	(4)	263	(96)	1		1		
	Yes	36	(38.7)	57	(61.3)	15.1	0.00	3.57	1.41–9.03	0.01
Alcohol use	Never	2	(1.1)	181	(98.9)	1		1		
	Yes	45	(24.5)	139	(75.5)	29.3	0.00	7.69	1.42–41.5	0.02
Patient type	New	34	(11)	276	(89)	1		1		
	Previously treated	13	(22.8)	44	(77.2)	2.4	0.02	0.90	0.33–2.43	0.84
Type of TB	PTB bacteriological confirmed	35	(13.9)	217	(86.1)	1		1		
	PTB clinically diagnosed	5	(15.6)	27	(84.4)	1.15	0.79	1.03	0.25–4.2	0.97
	Extra-PTB	7	(8.4)	76	(91.6)	0.57	0.20	1.04	0.36–2.99	0.94
Treatment phase	Initiation	25	(12.4)	176	(87.6)	1		1		
	Continuation	22	(13.2)	144	(86.8)	1.08	0.82	0.89	0.38–2.05	0.78
Family psychiatric history	No	36	(11)	291	(89)	1		1		
	Yes	11	(27.5)	29	(72.5)	3.07	0.00	3.97	1.32–12.0	0.01

PTB–pulmonary TB; Extra-PTB–extrapulmonary TB

<https://doi.org/10.1371/journal.pone.0302744.t006>

illness [25, 36]. Vulnerability, marginalization, stigma, and discrimination are often faced by people affected by TB and these could predispose to mental illness.

The low rate of a prior diagnosis of mental illness showed that there is a major treatment gap between diagnosed and undiagnosed mental illness. Similar findings were reported in another Kenyan study where of the 42% of subjects with symptoms of mental illness, only 4% had any psychiatric illness documented [31]. Health seeking for mental illness is usually affected by the associated stigma and the non-medical explanations given to mental illness. Additionally, there is lack of screening at primary health settings given that there is no diagnostic test for mental illness but rather clinical assessment or use of screening tools.

The assessed prevalence of alcohol use disorder is similar to a pooled average of 30% in a meta-analysis of 28 studies in tuberculosis patients [15], but double that previously reported for the general Kenyan population at 13.6% [37]. There's a strong association between heavy alcohol use and TB as alcohol weakens the immune system by interfering with the integrity and microbiomes of the gastrointestinal tract, decreasing and damaging the epithelial cells and by disruption of the ciliary function of the upper airways thus ultimately increasing the risk of TB [38]. Heavy alcohol use has been linked to higher reinfection rate, poor treatment compliance and development of drug resistant TB [15]. Also people with chronic alcohol use often have poor nutritional status predisposing to activation of latent TB [3].

Prevalence of depression was lower than that in other studies with a range from 31% - 54% [9, 13, 14, 25, 39–41], likely because of the different tools used (screening versus diagnostic). Nevertheless, many studies have consistently reported higher prevalence of depression among patients with TB as compared to the general population [2, 42], and could be attributed to the psychological stressors reported by the patients including stigma, association with HIV, drug-burden especially if co-infected, long duration of treatment, lost or reduced revenue, fear of infecting people close to them, isolation from their families and increased alcohol consumption [5, 36, 43].

Prevalence of substance use was 12.8% in this study and this was a combination of all non-alcohol substances including khat/miraa, bhang, tobacco, 'kuber' and glue. This rate was comparable to that in a Kenyan survey with a prevalence of 13.4% [23]. Prevalence of suicidality was at 8.2% for suicidal ideation and 2.5% for suicidal attempt/ behavior. This was comparable to a study in South Africa where 9% reported suicidal ideation while 3.1% had previous suicide attempts [18]. A different study in Ethiopia had higher rates of 17.3% for suicidal ideation and 7.5% for suicide attempts [44].

Determinants associated with psychiatric morbidity

The odds of psychiatric morbidity were increased in males, those who were separated or divorced, those with history of alcohol use, smoking, medical comorbidities, or previous history of TB and those who reported a family history of mental illness.

Males were more likely to have 'any' psychiatric illness or alcohol and substance use disorders, similar to findings of a meta-analysis [15] and several other studies in Africa and Asia [16–18, 45]. Culturally, social drinking among men is more tolerated compared to the same in females while it could be the start or gateway to alcohol or substance use disorders. Males, those with history of alcohol use and family psychiatric history had higher odds of substance use disorder. Those with a history of cigarette smoking even though not currently smoking still had higher odds of having substance use disorder. The gender and alcohol use association were replicated in a different study in Ethiopia where males were seven times and those with history of alcohol use were twice as likely to have substance use disorder.

Conversely, females are reported to have higher rates of depression but not alcohol or substance use disorders [11, 40, 46]. Gender differences could be as a result of males having more of externalizing symptoms as opposed to females who would present with more of internalizing symptoms [47].

This study reported lower rates of 'any psychiatric illness' and depression in those aged 65 +. This finding of decreased rates of depression in the older age was different from other studies. A study in Ethiopia reported that for every 14 years increase in age, the risk of having depression increased by 19% [48] and another study in Nigeria reported that depression was significantly more prevalent in older patients with tuberculosis compared with their family members [5]. The difference could be due to the lower numbers in this age group given that

their total number was 13, only three had any psychiatric illness and only 1 of the 13 had depression. Thus, with the lower numbers chance findings may not be ruled out.

This study found that being separated or divorced was associated with increased odds of any psychiatric illness, depression and alcohol use disorder, which were similar to other studies [10, 16, 41, 44]. Van Rensburg *et al.* in a systematic review of 100 articles also reported similar results [7]. A meta-analysis, however, reported that respondents who were single had higher odds of alcohol use disorder [15]. Those who are single, separated or divorced may lack good social support and may be at higher risk of maladaptive coping including alcohol use while on the other hand addiction may lead to change in marital status [28].

Having other medical comorbidities like hypertension, diabetes and HIV among others was fourfold associated with suicidality in this study. This was similar to studies in Ethiopia and South Africa that showed that having more than one other chronic illness and being HIV positive were positive predictors for suicidal ideation and suicide attempt [18, 44].

Other studies have similarly reported increased odds of psychiatric morbidity in TB patients who were smokers and tobacco users [17, 41], those with familial history of mental illness [11, 49], been previously treated for TB [5, 15, 38, 45, 50, 51], or with other comorbidities [8, 11, 43, 44, 52]. Substance use disorders and TB share common risk factors, including undernutrition, HIV infection, concomitant alcohol use and smoking [53]. Smoking is also considered a gateway drug to other drugs with shared risk factors for both smoking and alcohol use including genetic predisposition and family history. Studies have also reported genetic predisposition of anxiety, alcohol use disorder and depression [12, 22, 54, 55]. However, few studies have attempted to investigate the relationship of family history and mental illness among patients on treatment for TB in our setting.

The relationship between previous treatment and mental illness could be bi-directional as patients with uncontrolled mental illness could have poor adherence to anti-TB medication thus leading to retreatment as treatment interrupters or loss to follow-up. Further, because of their living conditions they could have re-infection of TB and need retreatment even after cure or treatment completion [3]. On the other hand, retreatment of TB can be a stressor given the type of drugs, long duration of treatment, drug burden and antecedent stigma associated with HIV thus predisposing to mental illness especially depression or alcohol use as a coping mechanism, albeit maladaptive.

Limitations of the study

The main strength of this study is that the sample was socio-demographically diverse, having been drawn from a population at the second largest TB treatment program in Kenya and probably in East Africa. Secondly, the study employed a diagnostic tool (MINI) that has been validated and adapted for use in this setting. Thirdly the study is generalizable to patients on treatment for TB in similar settings based on sampling method and sample size being representative. It can, however, not be generalized to a primary care setting as this was a tertiary hospital and as such the sample may be different in a community setting.

The cross-sectional design was a limitation as causal relationships between the compared variables cannot be inferred. Much as TB could lead to or increase the risk of mental illness, this study could not establish the proportional increase in size. This thus remains an area of further research. However, the cross-sectional design was useful in bringing out the prevalence of psychiatric morbidity among patients on treatment for tuberculosis in Kenya.

Another limitation was social desirability bias due to self-reporting nature of the interviewer-administered questionnaire. Some of the sections like alcohol and substance use and suicidal ideation may attract socially acceptable responses. This would consequently cause an

under-reporting of some symptoms and as such true prevalence may even be higher than reported. To mitigate against this, the interviewer employed principles of reassurance, non-judgmental attitude, and assurance of confidentiality.

Lastly, residence was not included in the socio-demographics questionnaire and considering that it impacts on socio-economics, it would additionally influence psychological and emotional well-being that would affect mental health. This was not included in this study and can be an area of research. Also, the study results found no association of females and depression which was unexpected. Further research would be needed to explain why this was different in this population.

Conclusions

In conclusion, this study found a high prevalence of psychiatric morbidity with most common disorders being alcohol use disorder, depression, substance use disorder and suicidality. Factors associated with increased odds of having ‘any psychiatric morbidity’ were being male, being separated or divorced, history of alcohol use or smoking, family history of psychiatric illness, having medical comorbidities and being previously treated for TB.

We thus recommend for integrated and bi-directional screening, diagnosis, referral and management of comorbid TB and mental illness, akin to the TB-HIV collaborative efforts. Longitudinal studies are recommended to show causal associations and to follow up outcomes of patients with TB and psychiatric comorbidities.

Supporting information

S1 Data.
(CSV)

Acknowledgments

We acknowledge Moi university, School of Medicine and the department of Mental Health and Behavioural sciences for their input during the development of this research article. We also acknowledge TB clinic staff at MTRH for a conducive environment for data collection and the participants who consented to be part of this work.

Author Contributions

Conceptualization: Robina Kerubo Momanyi, Edith Kamaru Kwobah, Philip Owiti, Henry Nyamogoba, Lukoye Atwoli.

Data curation: Robina Kerubo Momanyi, Philip Owiti.

Formal analysis: Philip Owiti.

Investigation: Robina Kerubo Momanyi.

Methodology: Robina Kerubo Momanyi, Edith Kamaru Kwobah, Philip Owiti, Lukoye Atwoli.

Project administration: Robina Kerubo Momanyi.

Resources: Robina Kerubo Momanyi.

Supervision: Edith Kamaru Kwobah, Henry Nyamogoba, Lukoye Atwoli.

Validation: Robina Kerubo Momanyi, Edith Kamaru Kwobah, Philip Owiti, Lukoye Atwoli.

Visualization: Robina Kerubo Momanyi, Edith Kamaru Kwobah, Philip Owiti, Henry Nyamogoba, Lukoye Atwoli.

Writing – original draft: Robina Kerubo Momanyi.

Writing – review & editing: Robina Kerubo Momanyi, Edith Kamaru Kwobah, Philip Owiti, Henry Nyamogoba, Lukoye Atwoli.

References

1. World Health Organization. Global Tuberculosis Report 2022.
2. Bender A, Guruge S, Hyman I, Janjua M. Tuberculosis, and common mental disorders: International lessons for Canadian immigrant health. *Can J Nurs Res*. 2012; 44(4):56–75. PMID: [23448075](#)
3. Doherty AM, Kelly J, McDonald C, O'Dwyer AM, Keane J, Cooney J. A review of the interplay between tuberculosis and mental health. *Gen Hosp Psychiatry [Internet]*. 2013; 35(4):398–406. Available from: <https://doi.org/10.1016/j.genhosppsy.2013.03.018> PMID: [23660587](#)
4. Mason PH, Sweetland AC, Fox GJ, Halovic S, Nguyen TA, Marks GB. Tuberculosis and mental health in the Asia-Pacific. *Australas Psychiatry*. 2016; 24(6):553–5. <https://doi.org/10.1177/1039856216649770> PMID: [27206468](#)
5. Oladimeji O, Ushie BA, Udoh EE, Oladimeji KE, Ige OM, Obasanya O, et al. Psychosocial wellbeing of patients with multidrug resistant tuberculosis voluntarily confined to long-term hospitalisation in Nigeria. *BMJ Glob Heal [Internet]*. 2016; 1(3): e000006. Available from: <http://gh.bmj.com/lookup/doi/https://doi.org/10.1136/bmjgh-2015-000006> PMID: [28588950](#)
6. Tola HH, Shojaeizadeh D, Garmaroudi G, Tol A, Yekaninejad MS, Ejeta LT, et al. Psychological distress and its effect on tuberculosis treatment outcomes in Ethiopia. *Glob Health Action*. 2015; 8(1). <https://doi.org/10.3402/gha.v8.29019> PMID: [26610316](#)
7. Janse Van Rensburg A, Dube A, Curran R, Ambaw F, Murdoch J, Bachmann M, et al. Comorbidities between tuberculosis and common mental disorders: A scoping review of epidemiological patterns and person-centred care interventions from low-to-middle income and BRICS countries. Vol. 9, *Infectious Diseases of Poverty*. 2020. <https://doi.org/10.1186/s40249-019-0619-4> PMID: [31941551](#)
8. Amreen, Rizvi N. Frequency of Depression and Anxiety among Tuberculosis Patients. *J Tuberc Res [Internet]*. 2016; 04(04):183–90. Available from: <http://www.scirp.org/journal/PaperDownload.aspx?DOI=10.4236/jtr.2016.44021>
9. Kumar K, Kumar A, Chandra P, Kansal H. A study of prevalence of depression and anxiety in patients suffering from tuberculosis. *J Fam Med Prim Care [Internet]*. 2016; 5(1):150. Available from: <http://www.jfmpc.com/text.asp?2016/5/1/150/184641> <https://doi.org/10.4103/2249-4863.184641> PMID: [27453861](#)
10. Rahul Mandaknalli BG. Prevalence Depression in Tuberculosis Patients in a Tertiary Care Hospital. *Sch J App Med Sci*. 2015; 3(6D):2445–8.
11. Kehbila J, Ekabe CJ, Aminde LN, Noubiap JJJ, Fon PN, Monekosso GL. Prevalence, and correlates of depressive symptoms in adult patients with pulmonary tuberculosis in the Southwest Region of Cameroon. *Infect Dis Poverty [Internet]*. 2016; 5(1):51. Available from: <http://idpjournal.biomedcentral.com/articles/10.1186/s40249-016-0145-6> PMID: [27268138](#)
12. Peltzer K, Naidoo P, Matseke G, Louw J, Mchunu G, Tutshana B. Prevalence of psychological distress and associated factors in tuberculosis patients in public primary care clinics in South Africa. *BMC Psychiatry [Internet]*. 2012; 12(1):89. Available from: <http://bmcp psychiatry.biomedcentral.com/articles/10.1186/1471-244X-12-89> PMID: [22839597](#)
13. Duko B, Gebeyehu A, Ayano G. Prevalence and correlates of depression and anxiety among patients with tuberculosis at Wolaita Sodo University Hospital and Sodo Health Center, Wolaita Sodo, South Ethiopia, Cross sectional study. *BMC Psychiatry [Internet]*. 2015; 15(1):214. Available from: <http://bmcp psychiatry.biomedcentral.com/articles/10.1186/s12888-015-0598-3>
14. Lee E. Prevalence of Depression among Active TB and DEPRESSION AMONG ACTIVE TB AND TB / HIV PATIENTS IN KISUMU. 2015;1–35. Available from: http://digitalcollections.sit.edu/isp_collection
15. Necho M, Tsehay M, Seid M, Zenebe Y, Belete A, Gelaye H. Prevalence and associated factors for alcohol use disorder among tuberculosis patients: a systematic review and meta- analysis study. 2021;1–15.
16. Connell RO, Chishinga N, Kinyanda E, Patel V, Ayles H, Helen A. Prevalence and Correlates of Alcohol Dependence Disorder among TB and HIV Infected Patients in Zambia. 2013; 8(9):1–9.

17. Thapa P, Kamath R, Bk S, Monteiro A, Vc S. Prevalence and Associated Factors of Alcoholism among Tuberculosis Patients in Udipi Taluk, Karnataka, India: A Cross Sectional Study. 2014; 12(3):177–81. PMID: [26032055](#)
18. Laprawat S, Peltzer K, Pansila W, Tansakul C, Africa S. Alcohol use disorder and tuberculosis treatment: A longitudinal mixed method study in Thailand.: 1–5.
19. Islam AT, Hoque A. Pattern of psychiatric illness among tuberculosis Patients an analysis in a tertiary care hospital of Bangladesh. 2015; 1:763–6.
20. Scuffell J, Boccia D, Velarde FG, Leon SR, Raviola G, Lecca L, et al. Mental disorders, and drug/alcohol use in patients commencing extensively drug-resistant tuberculosis treatment. *Public Heal Action* [Internet]. 2017; 7(3):237–9. Available from: <http://www.ingentaconnect.com/content/10.5588/pha.17.0044> PMID: [29201659](#)
21. Id MS, Tolessa O, Tesfaye M, Adorjan K, Krahl W, Tesfaye E, et al. Magnitude and predictors of khat use among patients with tuberculosis in Southwest Ethiopia: A longitudinal study. 2020;1–12.
22. Kessler RC, Aguilar-gaxiola S, Alonso J, Chatterji S, Lee S, Ormel J, et al. *World Mental Health (WMH) Surveys*. 2011; 18(1):23–33.
23. Kamenderi M, Muteti J, Okioma V, Kimani S. Status of Drugs and Substance Abuse among the General Population in Kenya. 2021; 1:54–9.
24. Sweetland A, Oquendo M, Wickramaratne P, Weissman M, Wainberg M. Depression: A silent driver of the global tuberculosis epidemic. *World Psychiatry*. 2014; 13(3):325–6. <https://doi.org/10.1002/wps.20134> PMID: [25273311](#)
25. Ambaw F, Mayston R, Hanlon C, Alem A. Depression among patients with tuberculosis: determinants, course, and impact on pathways to care and treatment outcomes in a primary care setting in southern Ethiopia—a study protocol. *BMJ Open* [Internet]. 2015; 5(7):e007653. Available from: <http://bmjopen.bmj.com/lookup/doi/10.1136/bmjopen-2015-007653> PMID: [26155818](#)
26. Ministry of Health. Annual Report 2018. 2018. p. 1–78.
27. WHO. Tuberculosis Report. Vol. XLIX, Baltimore Health News. 2020. 8 p.
28. Lecrubier Y, Sheehan D V., Weiller E, Amorim P, Bonora I, Sheehan KH, et al. The Mini International Neuropsychiatric Interview (MINI). A short diagnostic structured interview: Reliability and validity according to the CIDI. *Eur Psychiatry*. 1997; 12(5):224–31.
29. Sheehan D V, Sheehan KH, Shytle RD, Janavs J, Bannon Y, Rogers JE, et al. Reliability and validity of the Mini International Neuropsychiatric Interview for Children and Adolescents (MINI-KID). *J Clin Psychiatry*. 2010; 71(3):313–26. <https://doi.org/10.4088/JCP.09m05305whi> PMID: [20331933](#)
30. Aillon JL, Ndeti DM, Khasakhala L, Ngari WN, Achola HO, Akinyi S, et al. Prevalence, types, and comorbidity of mental disorders in a Kenyan primary health centre. *Soc Psychiatry Psychiatr Epidemiol*. 2014; 49(8):1257–68. <https://doi.org/10.1007/s00127-013-0755-2> PMID: [23959589](#)
31. Ndeti DM, Khasakhala LI, Kuria MW, Mutiso VN, Ongecha-Owuor FA, Kokonya DA. The prevalence of mental disorders in adults in different level general medical facilities in Kenya: A cross-sectional study. *Ann Gen Psychiatry*. 2009; 8:1–8. <https://doi.org/10.1186/1744-859X-8-1> PMID: [19144164](#)
32. Kwobah E, Epstein S, Mwangi A, Litzelman D, Atwoli L. PREVALENCE of psychiatric morbidity in a community sample in Western Kenya. *BMC Psychiatry* [Internet]. 2017; 17(1):1–6. Available from: <https://doi.org/10.1186/s12888-017-1202-9> PMID: [28100210](#)
33. Adeyemi, Akeem akinloye. Psychiatric Morbidity and Quality of Life in Patients with Pulmonary Tuberculosis, HIV/AIDS, and Healthy Individuals: A Comparative Study; A dissertation
34. Njie GJ, Khan A. Prevalence of Tuberculosis and Mental Disorders Comorbidity: A Systematic Review and Meta-analysis. *J Immigr Minor Heal*. 2022; 24(6):1550–6. <https://doi.org/10.1007/s10903-021-01312-6> PMID: [34796457](#)
35. Thungana Y, Wilkinson R, Zingela Z. Comorbidity of mental ill-health in tuberculosis patients under treatment in a rural province of South Africa: a cross-sectional survey. *BMJ Open*. 2022; 12(11):1–8.
36. Hansel NN, Wu AW, Chang B, Diette GB. Quality of life in tuberculosis: patient and provider perspectives. *Qual Life Res*. 2004; 13(3):639–52. <https://doi.org/10.1023/B:QURE.0000021317.12945.f0> PMID: [15130027](#)
37. NATIONAL SURVEY ON THE STATUS OF DRUGS AND SUBSTANCE. 2022;
38. Veerakumar AM, Kumar S, Sarkar S. Author's personal copy ScienceDirect Alcohol use disorders among pulmonary tuberculosis patients under RNTCP in urban Pondicherry, India.
39. Vega P, Sweetland A, Acha J, Castillo H, Guerra D, Fawzi MCS, et al. Psychiatric issues in the management of patients with multidrug-resistant tuberculosis SUMMARY. 2004; 8(July 2003):749–59.
40. Husain MO, Dearman SP, Chaudhry IB, Rizvi N, Waheed W. The relationship between anxiety, depression, and illness perception in tuberculosis patients in Pakistan. *Clin Pract Epidemiol Ment Health*

[Internet]. 2008; 4(February 2008):4. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2288599&tool=pmcentrez&rendertype=abstract>

41. Paula A, Sc M, Lazzari TK, Sc M, Silva DR, Ph D. Health-Related Quality of Life, Depression and Anxiety in Hospitalized Patients with Tuberculosis. 2017; 3536:69–76.
42. Qayyum S, Rizvi R. Prevalence of depression in TB patients in comparison to non-tuberculous family contacts visiting the DOTS clinic in tertiary care hospital and its correlation with disease pattern. 2008; 17:3–5. *Infect Dis J Pakistan*. 2008; 17:3–5.
43. Deribew A, Tesfaye M, Hailmichael Y, Apers L, Abebe G, Duchateau L, et al. Common mental disorders in TB/HIV co-infected patients in Ethiopia. *BMC Infect Dis*. 2010; 10(1):201. <https://doi.org/10.1186/1471-2334-10-201> PMID: 20618942
44. Molla A, Mengesha A, Derajew H, Kerebih H. Suicidal Ideation, Attempt, and Associated Factors among Patients with Tuberculosis in Ethiopia: A Cross-Sectional Study. 2020; 2019.
45. Peltzer K, Louw J, Mchunu G, Naidoo P, Matseke G. Hazardous and Harmful Alcohol Use and Associated Factors in Tuberculosis Public Primary Care Patients in South Africa. 2012;3245–57.
46. Aamir S, Aisha A. Co-morbid anxiety, and depression among pulmonary tuberculosis patients. *J Coll Physicians Surg Pakistan*. 2010; 20(10):703–4. <https://doi.org/10.2010/JCPSP.703704> PMID: 20943121
47. Wilhelm KA. Gender and mental health. *Aust N Z J Psychiatry*. 2014; 48(7):603–5. <https://doi.org/10.1177/0004867414538678> PMID: 24973270
48. Ambaw F, Mayston R, Hanlon C, Alem A. Burden and presentation of depression among newly diagnosed individuals with TB in primary care settings in Ethiopia. 2017;1–10.
49. Mohammedhusein M, Alenko A, Tessema W, Mamaru A. Prevalence and associated factors of depression and anxiety among patients with pulmonary tuberculosis attending treatment at public health facilities in southwest Ethiopia. *Neuropsychiatr Dis Treat*. 2020; 16:1095–104. <https://doi.org/10.2147/NDT.S249431> PMID: 32636628
50. Adem A, Tesfaye M, Adem M. The Prevalence and Pattern of Depression in Patients with Tuberculosis on Follow-up at Jimma University Specialized Hospital and Jimma Health Center. *Med Sci*. 2014; 3(1):955–68.
51. Singh L, Pardal P, Prakash J. Psychiatric morbidity in patients of pulmonary tuberculosis-an observational study. *Ind Psychiatry J [Internet]*. 2015; 24(2):168. Available from: <http://www.industrialpsychiatry.org/text.asp?2015/24/2/168/181722> <https://doi.org/10.4103/0972-6748.181722> PMID: 27212822
52. Peltzer K, Naidoo P, Matseke G, Louw J, McHunu G, Tutshana B. Prevalence of post-traumatic stress symptoms and associated factors in tuberculosis (TB), TB retreatment and/or TB-HIV co-infected primary public health-care patients in three districts in South Africa. *Psychol Heal Med*. 2013; 18(4):387–97. <https://doi.org/10.1080/13548506.2012.726364> PMID: 23061988
53. World Health Organization. END TB Global Tuberculosis Report 2017. 2017. 147 p.
54. Kamau JW, Kuria W, Mathai M, Atwoli L, Kangethe R. Psychiatric morbidity among HIV-infected children and adolescents in a resource-poor Kenyan urban community. *AIDS Care*. 2012; 24(7):836–42. <https://doi.org/10.1080/09540121.2011.644234> PMID: 22292795
55. Woollett N, Cluver L, Bandeira M, Brahmhatt H. Identifying risks for mental health problems in HIV positive adolescents accessing HIV treatment in Johannesburg. *J Child Adolesc Ment Health*. 2017; 29(1):11–26. <https://doi.org/10.2989/17280583.2017.1283320> PMID: 28287023