



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

K-stability of Fano 3-folds of Picard rank 3 and degree 20

Citation for published version:

Denisova, E 2024, 'K-stability of Fano 3-folds of Picard rank 3 and degree 20', *Annali dell' Università di Ferrara*. <https://doi.org/10.1007/s11565-024-00516-6>

Digital Object Identifier (DOI):

[10.1007/s11565-024-00516-6](https://doi.org/10.1007/s11565-024-00516-6)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Annali dell' Università di Ferrara

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



K-STABILITY OF FANO 3-FOLDS OF PICARD RANK 3 AND DEGREE 20

ELENA DENISOVA

ABSTRACT. We prove K-stability of smooth Fano 3-folds of Picard rank 3 and degree 20 that satisfy very explicit generality condition.

CONTENTS

1. Introduction	1
2. The Proof	2
Appendix A. Polarized δ -invariant via Kento Fujita's formulas	5
A.1. Polarized δ -invariant on Del Pezzo surface of degree 4 with \mathbb{A}_1 singularity.	6
A.2. Polarized δ -invariant on Del Pezzo surface of degree 4 with two \mathbb{A}_1 singularities.	14
A.3. Polarized δ -invariant on Del Pezzo surface of degree 4 with \mathbb{A}_2 singularity	23
References	25

1. INTRODUCTION

Let $S = \mathbb{P}^1 \times \mathbb{P}^1$, let C be a smooth curve in S of degree $(5, 1)$, and let $\eta: C \rightarrow \mathbb{P}^1$ be the morphism induced by the projection $S \rightarrow \mathbb{P}^1$ to the first factor. Then η is a finite morphism of degree five, and we may assume that the points $([1 : 0], [0 : 1])$ and $([0 : 1], [1 : 0])$ are among its ramifications points. This assumption implies that the curve C is given by

$$u(x^5 + a_1x^4y + a_2x^3y^2 + a_3x^2y^3) = v(y^5 + b_1xy^4 + b_2x^2y^3 + b_3x^3y^2)$$

for some $a_1, a_2, a_3, b_1, b_2, b_3$, where $([u : v], [x : y])$ are coordinates on S . Note that the ramification index of the point $([1 : 0], [0 : 1])$ can be computed as follows:

$$\begin{cases} 2 \text{ if } a_3 \neq 0, \\ 3 \text{ if } a_3 = 0 \text{ and } a_2 \neq 0, \\ 4 \text{ if } a_3 = a_2 = 0 \text{ and } a_1 \neq 0, \\ 5 \text{ if } a_3 = a_2 = a_1 = 0. \end{cases}$$

Likewise, we can compute the ramification index of the point $([0 : 1], [1 : 0])$. We may assume that

- $([1 : 0], [0 : 1])$ has the largest ramification index among all ramifications points of η
- the ramification index of the point $([0 : 1], [1 : 0])$ is the second largest index.

If both these indices are 5, then $a_1 = a_2 = a_3 = b_1 = b_2 = b_3 = 0$, the morphism η does not have other ramification points, and the equation of the curve C simplifies as

$$ux^5 = vy^5.$$

In this case, we have $\text{Aut}(S, C) \cong \mathbb{C}^* \rtimes \mathbb{Z}/2\mathbb{Z}$. In all other cases, this group is finite [6, Corollary 2.7]. Now, we consider embedding $S \hookrightarrow \mathbb{P}^1 \times \mathbb{P}^2$ given by

$$([u : v], [x : y]) \mapsto ([u : v], [x^2 : xy : y^2]),$$

Throughout this paper, all varieties are assumed to be projective and defined over \mathbb{C} .

and identify S and C with their images in $\mathbb{P}^1 \times \mathbb{P}^2$. Let $\pi: X \rightarrow \mathbb{P}^1 \times \mathbb{P}^2$ be the blow up of the curve C . Then X is a smooth Fano threefold in the deformation family № 3.5 in the Mori–Mukai list and every smooth member of this family can be obtained in this way. We know from [2, Section 5.14], that

- X is K-stable if the numbers $a_1, a_2, a_3, b_1, b_2, b_3$ are general enough,
- X is K-polystable if $a_1 = a_2 = a_3 = b_1 = b_2 = b_3 = 0$.

However, for some $a_1, a_2, a_3, b_1, b_2, b_3$, the threefold X is not K-polystable.

Example 1. If $(a_1, a_2, a_3) = (0, 0, 0) \neq (b_1, b_2, b_3)$, then X is not K-polystable [2, Lemma 7.6].

Note also that it follows from the proof of [6, Lemma 8.7] that $\text{Aut}(X) \cong \text{Aut}(S, C)$. In particular, we conclude the group $\text{Aut}(X)$ is finite if and only if $(a_1, a_2, a_3, b_1, b_2, b_3) \neq (0, 0, 0, 0, 0, 0)$. In this case, the threefold X is K-polystable if and only if it is K-stable. Moreover, we have

Conjecture 1 ([2]). *The Fano threefold X is K-stable if and only if $(a_1, a_2, a_3) \neq (0, 0, 0)$.*

Geometrically, this conjecture says that the following two conditions are equivalent:

- (1) the threefold X is K-stable,
- (2) the morphism $\eta: C \rightarrow \mathbb{P}^1$ does not have ramification points of ramification index five.

The goal of this paper is to prove the following (slightly weaker) result:

Theorem. *If all ramification points of η have ramification index two, then X is K-stable.*

Let $\text{pr}_1: \mathbb{P}^1 \times \mathbb{P}^2 \rightarrow \mathbb{P}^1$ be the projection to the first factor and $\phi_1 = \text{pr}_1 \circ \pi$. Then ϕ_1 is a fibration into del Pezzo surfaces of degree four, and Theorem and Conjecture 1 can be restated as follows:

Main Theorem. *If every singular fiber of ϕ_1 has only singular points of type \mathbb{A}_1 , then X is K-stable.*

Conjecture 2. *The Fano threefold X is K-stable if and only if every singular fiber of ϕ_1 has only singular points of type $\mathbb{A}_1, \mathbb{A}_2$ or \mathbb{A}_3 .*

Acknowledgments: I am grateful to my supervisor Professor Ivan Cheltsov for the introduction to the topic and his continuous support.

2. THE PROOF

To prove **Main Theorem**, we suppose that each singular fiber of the fibration ϕ_1 has one or two singular points of type \mathbb{A}_1 . Note that this fiber is a del Pezzo surface of degree 4 with Du Val singularities. We know ([8, 10]) that the Fano threefold X is K-stable if and only if for every prime divisor \mathbf{F} over X we have

$$\beta(\mathbf{F}) = A_X(\mathbf{F}) - S_X(\mathbf{F}) > 0$$

where $A_X(\mathbf{F})$ is the log discrepancy of the divisor \mathbf{F} , and

$$S_X(\mathbf{F}) = \frac{1}{(-K_X)^3} \int_0^\infty \text{vol}(-K_X - u\mathbf{F}) du.$$

To show this, we fix a prime divisor \mathbf{F} over X . Then we set $Z = C_X(\mathbf{F})$. If Z is an irreducible surface, then it follows from [9] that $\beta(\mathbf{F}) > 0$, see also [2, Theorem 3.17]. Therefore, we may assume that

- either Z is an irreducible curve in X ,
- or Z is a point in X .

In both cases, we fix a point $O \in Z$. Let \bar{T} be the fiber of ϕ_1 which contains O . Then \bar{T} is a del Pezzo surface with at most Du Val singularities. Set

$$\tau(\bar{T}) = \sup \left\{ u \in \mathbb{R}_{>0} \mid \text{the divisor } -K_X - u\bar{T} \text{ is pseudo-effective} \right\}$$

For $u \in [0, \tau(\bar{T})]$ let $P(u)$ be the positive part of the Zariski decomposition of the divisor $-K_X - u\bar{T}$, and let $N(u)$ be its negative part. Then we have

$$P(u) = \begin{cases} -K_X - u\bar{T} & \text{if } u \in [0, 1], \\ -K_X - u\bar{T} - (u-1)\tilde{S} & \text{if } u \in [1, 2], \end{cases} \quad \text{and } N(u) = \begin{cases} 0 & \text{if } u \in [0, 1], \\ (u-1)\tilde{S} & \text{if } u \in [1, 2], \end{cases}$$

which gives

$$S_X(\bar{T}) = \frac{1}{20} \int_0^2 P(u)^3 du = \frac{69}{80} < 1$$

Now, for every prime divisor F over the surface \bar{T} , we set

$$S(W_{\bullet,\bullet}^{\bar{T}}; F) = \frac{3}{(-K_X)^3} \int_0^\tau \text{ord}_F(N(u)|_{\bar{T}}) (P(u)|_{\bar{T}})^2 du + \frac{3}{(-K_X)^3} \int_0^\tau \int_0^\infty \text{vol}(P(u)|_{\bar{T}} - vF) dv du.$$

Then, following [1, 2], we let

$$\delta_O(\bar{T}, W_{\bullet,\bullet}^{\bar{T}}) = \inf_{\substack{F/\bar{T} \\ O \in C_{\bar{T}}(F)}} \frac{A_T(F)}{S(W_{\bullet,\bullet}^{\bar{T}}; F)},$$

where the infimum is taken by all prime divisors over the surface \bar{T} whose center on \bar{T} contains O . Then it follows from [1, 2] that

$$\frac{A_X(\mathbf{F})}{S_X(\mathbf{F})} \geq \min \left\{ \frac{1}{S_X(\bar{T})}, \delta_O(\bar{T}, W_{\bullet,\bullet}^{\bar{T}}) \right\}.$$

Therefore, if $\beta(\mathbf{F}) \leq 0$, then $\delta_O(\bar{T}, W_{\bullet,\bullet}^{\bar{T}}) \leq 1$.

Let's prove that $\delta_O(\bar{T}, W_{\bullet,\bullet}^{\bar{T}}) > 1$. To estimate $\delta_O(\bar{T}, W_{\bullet,\bullet}^{\bar{T}})$, we set $\bar{D} = P(u)|_{\bar{T}}$. We have

$$\bar{D} = \begin{cases} -K_{\bar{T}} & \text{if } u \in [0, 1], \\ -K_{\bar{T}} - (u-1)\bar{C}_2 & \text{if } u \in [1, 2], \end{cases}$$

where $\bar{C}_2 := \tilde{S}|_{\bar{T}}$. Then \bar{D} is ample for $u \in [0, 2)$, and

$$\bar{D}^2 = \begin{cases} 4 & \text{if } u \in [0, 1], \\ 5 - u^2 & \text{if } u \in [1, 2]. \end{cases}$$

We denote \tilde{S} to be the proper transform on X of the surface S . By Lemma [2, 5.68] and Lemma[2, 5.69] we have

Lemma 1. *If $O \in \tilde{S}$ then $\delta_O(X) > 1$.*

Lemma 2. *If \bar{T} is smooth then $\delta_O(X) > 1$.*

Thus, to prove **Main Theorem**, we may assume that $O \notin \tilde{S}$ and \bar{T} is singular. Recall that

$$\delta_O(\bar{T}, \bar{D}) = \inf_{\substack{F/\bar{T} \\ O \in C_{\bar{T}}(F)}} \frac{A_{\bar{T}}(F)}{S_{\bar{D}}(F)} \text{ where } S_{\bar{D}}(F) = \frac{1}{\bar{D}^2} \int_0^\tau \text{vol}(\bar{D} - vF) dv$$

where $\tau = \tau(F)$ is the pseudo-effective threshold of F with respect to \overline{D} . Usually $\delta_O(\overline{T}, -K_{\overline{T}})$ is denoted by $\delta_O(\overline{T})$.

Note that since $O \notin \widetilde{S}$ then for any divisor F over \overline{T} then we get

$$\begin{aligned}
S(W_{\bullet,\bullet}^{\overline{T}}; F) &= \frac{3}{(-K_X)^3} \left(\int_0^\tau (P(u)^2 \cdot \overline{T}) \cdot \text{ord}_O(N(u)|_{\overline{T}}) du + \int_0^\tau \int_0^\infty \text{vol}(P(u)|_{\overline{T}} - vF) dv du \right) = \\
&= \frac{3}{20} \int_0^\tau \int_0^\infty \text{vol}(P(u)|_{\overline{T}} - vF) dv du = \\
&= \frac{3}{20} \left(\int_0^1 \int_0^\infty \text{vol}(-K_{\overline{T}} - vF) dv du + \int_1^2 \int_0^\infty \text{vol}(-K_{\overline{T}} - (u-1)\overline{C}_2 - vF) dv du \right) = \\
&= \frac{3}{20} \left(\int_0^\infty \text{vol}(-K_{\overline{T}} - vF) dv + \int_0^\infty \text{vol}(-K_T - (u-1)\overline{C}_2 - vF) dv \right) \leq \\
&= \frac{3}{20} \left(\int_0^\infty \text{vol}(-K_{\overline{T}} - vF) dv + \int_0^\infty \text{vol}(-K_{\overline{T}} - vF) dv \right) = \\
&= \frac{3}{10} \left(\int_0^\infty \text{vol}(-K_{\overline{T}} - vF) dv \right) = \frac{6}{5} \left(\frac{1}{4} \int_0^\infty \text{vol}(-K_{\overline{T}} - vF) dv \right) = \\
&= \frac{6}{5} S_{\overline{T}}(F) \leq \frac{6}{5} \cdot \frac{A_{\overline{T}}(F)}{\delta_O(\overline{T})}
\end{aligned}$$

Thus, if $\delta_O(\overline{T}) > 6/5$, then $\delta_O(\overline{T}, W_{\bullet,\bullet}^{\overline{T}}) > 1$. To estimate $\delta_O(\overline{T}, W_{\bullet,\bullet}^{\overline{T}})$ in the case when $\delta_O(\overline{T}) \leq 6/5$, we define the following positive continuous function on $[1, 2]$:

$$f(u) := \begin{cases} \frac{15 - 3u^2}{16 + 3u - 9u^2 + 2u^3} & \text{if } u \in [1, a], \\ \frac{15 - 3u^2}{11 - u^3} & \text{if } u \in [a, 2] \end{cases}$$

where a is a root of $3u^3 - 9u^2 + 3u + 5$ on $[1, 2]$. More precisely, $a \in [1.355, 1.356]$. In the appendix we prove that for each O such that $\delta_O(\overline{T}) \leq \frac{6}{5}$ we have $\delta_O(\overline{T}, \overline{D}) \geq f(u)$ for every $u \in [1, 2]$. So we obtain

$$\begin{aligned}
S(W_{\bullet,\bullet}^{\overline{T}}; F) &= \frac{3}{(-K_X)^3} \int_1^2 \int_0^\tau \text{vol}(P(u)|_{\overline{T}} - vF) dv du + \frac{3}{(-K_X)^3} \int_0^1 \int_0^\tau \text{vol}(P(u)|_{\overline{T}} - vF) dv du \leq \\
&\leq \frac{3}{20} \left(\int_1^2 \frac{(5 - u^2)}{\delta_O(\overline{T}, \overline{D})} du \right) A_{\overline{T}}(F) + \frac{3}{20} \cdot \frac{4A_{\overline{T}}(F)}{\delta_O(\overline{T})} \leq \frac{3}{20} \left(\int_1^2 \frac{(5 - u^2)}{f(u)} du \right) A_{\overline{T}}(F) + \frac{3}{5} A_{\overline{T}}(F) \leq \\
&\leq \frac{3}{20} \left(\int_1^{1.356} (5 - u^2) \frac{16 + 3u - 9u^2 + 2u^3}{15 - 3u^2} du + \int_{1.355}^2 (5 - u^2) \frac{11 - u^3}{15 - 3u^2} du \right) A_{\overline{T}}(F) + \frac{3}{5} A_{\overline{T}}(F) \leq \\
&\leq \frac{99}{100} A_{\overline{T}}(F)
\end{aligned}$$

Thus $\frac{A_{\overline{T}}(F)}{S(W_{\bullet,\bullet}^{\overline{T}}; F)} \geq \frac{100}{99}$ for every prime divisor F over \overline{T} whose support on F contains O , so that $\delta_O(W^{\overline{T}}, F) \geq \frac{100}{99}$, which implies $\beta(F) > 0$ and X is K -stable.

Remark 1. If O were a singular point of type \mathbb{A}_2 , this approach would not work, because as is shown in Appendix A.3 there exists a curve \overline{C} on \overline{T} containing O such that $\delta_O(\overline{T}, \overline{D}) = \frac{u^3 - 6u^2 + 19}{15 - 3u^2}$ so we get that

$$S(W^{\overline{T}}; \overline{C}) \leq \frac{3}{20} \left(\int_1^2 \frac{(5 - u^2)}{\delta_P(\overline{T}, \overline{C})} du \right) A_{\overline{T}}(\overline{C}) + \frac{3}{5} A_{\overline{T}}(\overline{C}) = \frac{83}{80} A_{\overline{T}}(\overline{C})$$

so $\frac{A_{\overline{T}}(F)}{S(W_{\bullet,\bullet}^{\overline{T}}; \overline{C})} < 1$ and we do not get a contradiction.

APPENDIX A. POLARIZED δ -INVARIANT VIA KENTO FUJITA'S FORMULAS

Let us use notations from Section 2. Recall that \overline{T} is a Du Val del Pezzo surface, and the blow up π induces a birational morphism $v : \overline{T} \rightarrow \mathbb{P}^2$. We assume that \overline{T} is singular so v is a weighted blow up. We have the following commutative diagram

$$\begin{array}{ccc} T & & \\ \sigma \swarrow \quad \searrow \eta & & \\ \overline{T} & \xrightarrow{v} & \mathbb{P}^2 \end{array}$$

Suppose that $u \in [1, 2]$. Recall that $\overline{D} = -K_{\overline{T}} - (1 - u)\overline{C}_2$. Observe that \overline{C}_2 is contained in the smooth locus of the surface \overline{T} . Let C_2 be the strict transform of the curve \overline{C}_2 on the surface T , set $D = -K_T - (1 - u)C_2$. Note that $D = \sigma^*(\overline{D})$ so the divisor D is big and nef for $u \in [1, 2]$. Recall that

$$\delta_O(\overline{T}, \overline{D}) = \inf_{\substack{F/\overline{T} \\ O \in C_{\overline{T}}(F)}} \frac{A_{\overline{T}}(F)}{S_D(F)}$$

where the infimum is run over all prime divisor F over \overline{T} such that $O \in C_{\overline{T}}(F)$. For every point $P \in T$, we also define

$$\delta_P(T, D) = \inf_{\substack{E/T \\ P \in C_T(E)}} \frac{A_T(E)}{S_D(E)}$$

where the infimum is run over all prime divisor E over T such that $P \in C_T(E)$. Since $D = \sigma^*(\overline{D})$ and $K_T = \sigma^*(K_{\overline{T}})$, we have

$$\delta_O(\overline{T}, \overline{D}) = \inf_{P: O = \sigma(P)} \delta_P(T, D)$$

So, to estimate $\delta_O(\overline{T}, \overline{D})$ it is enough to estimate $\delta_P(T, D)$ for P all points P such that $\sigma(P) = O$. Let \mathcal{C} be a smooth curve on T containing P . Set

$$\tau(\mathcal{C}) = \sup \left\{ v \in \mathbb{R}_{\geq 0} \mid \text{the divisor } -K_T - v\mathcal{C} \text{ is pseudo-effective} \right\}.$$

For $v \in [0, \tau]$, let $P(v)$ be the positive part of the Zariski decomposition of the divisor $-K_T - \mathcal{C}$, and let $N(v)$ be its negative part. Then we set

$$S(W_{\bullet,\bullet}^{\mathcal{C}}; P) = \frac{2}{D^2} \int_0^{\tau(\mathcal{C})} h_D(v) dv,$$

where

$$h_D(v) = (P(v) \cdot \mathcal{C}) \times (N(v) \cdot \mathcal{C})_P + \frac{(P(v) \cdot \mathcal{C})^2}{2}.$$

It follows from [1, 2] that:

$$\delta_P(T, D) \geq \min \left\{ \frac{1}{S_D(\mathcal{C})}, \frac{1}{S(W_{\bullet,\bullet}^{\mathcal{C}}, P)} \right\}.$$


We will estimate $\delta_P(T, D)$ in the following using the notations above for a suitable choice of the curve \mathcal{C} , $\tau(\mathcal{C})$, $P(v)$ and $N(v)$ later in special cases.

A similar approach was taken in [3] and [5].

A.1. Polarized δ -invariant on Del Pezzo surface of degree 4 with \mathbb{A}_1 singularity. Suppose that \bar{T} has one singular point and this point is a singular point of type \mathbb{A}_1 . Then η is a blow up of \mathbb{P}^2 at points P_1, P_2, P_3 and P_4 in general position and a point P_5 which belongs to the exceptional divisor corresponding to P_4 and no other negative curve. Suppose $\mathbf{E} := L_{14} \cup L_{24} \cup L_{24} \cup E_5$. By [7, Section 6.2] we have:

$$\delta_P(T) = \begin{cases} 1 & \text{if } P \in E_4, \\ 6/5 & \text{if } P \in \mathbf{E} \setminus E_4, \\ 4/3 & \text{if } P \text{ belongs to two curves in } \{E_1, E_2, E_3, L_{12}, L_{13}, L_{23}, L_{45}, C_2\}, \\ 18/13 & \text{if } P \text{ belongs to exactly one curve in } \{E_1, E_2, E_3, L_{12}, L_{13}, L_{23}, L_{45}, C_2\} \setminus \mathbf{E}, \\ 3/2 & \text{otherwise} \end{cases}$$

where E_1, E_2, E_3, E_4, E_5 are exceptional divisors corresponding to P_1, P_2, P_3, P_4, P_5 respectively, C_2 is a strict transform of a (-1) -curve coming from the conic on \mathbb{P}^2 , L_{ij} are strict transforms of the lines passing through P_i and P_j for $(i, j) \in \{(1, 2), (1, 3), (1, 4), (2, 3), (2, 4), (3, 4)\}$ and L_{45} a strict transform of a (-1) -curve coming from a line on \mathbb{P}^2 . The dual graph of (-1) and (-2) -curves is given in the following picture:



Lemma A.1. Suppose P is a point on T and $D = -K_T - (u-1)C_2$ with $D^2 = 5 - u^2$ then

$$\delta_P(T, D) = \begin{cases} \frac{15 - 3u^2}{16 + 3u - 9u^2 + 2u^3} & \text{for } P \in E_4 \setminus E_5 \text{ and } u \in [1, 2] \\ \frac{15 - 3u^2}{11 - u^3} & \text{for } P \in E_5 \setminus E_4 \text{ and } u \in [1, 2] \\ \frac{15 - 3u^2}{3u^3 - 12u^2 + 6u + 13} & \text{for } P \in L_{14} \setminus (E_4 \cup E_5 \cup E_1) \text{ and } u \in [1, 2] \end{cases}$$

and

$$\delta_P(T, D) \geq \begin{cases} \frac{15 - 3u^2}{3u^3 - 12u^2 + 6u + 13} & \text{for } P = E_4 \cap E_5 \text{ and } u \in [1, a] \\ \frac{15 - 3u^2}{11 - u^3} & \text{for } P = E_4 \cap E_5 \text{ and } u \in [a, 2] \end{cases}$$

and

$$\delta_P(T, D) \geq \begin{cases} \frac{15 - 3u^2}{3u^3 - 12u^2 + 6u + 13} & \text{for } P = L_{14} \cap E_1 \text{ and } u \in [1, b] \\ \frac{2(15 - 3u^2)}{19 - 2u^3} & \text{for } P = L_{14} \cap E_1 \text{ and } u \in [b, 3/2] \\ \frac{15 - 3u^2}{3u^3 - 18u^2 + 27u - 4} & \text{for } P = L_{14} \cap E_1 \text{ and } u \in [3/2, 2] \end{cases}$$

where a is a root of $3u^3 - 9u^2 + 3u + 5$ on $[1, 2]$, b is a root of $8u^3 - 24u^2 + 12u + 7$ on $[1, 3/2]$. Note that $a \in [1.355, 1.356]$, $b \in [1.261, 1.262]$.

Proof. Step 1. Suppose $P \in E_4$. In this case we set $\mathcal{C} = E_4$. Then $\tau(\mathcal{C}) = 3 - u$. The Zariski Decomposition of the divisor $D - vE_4$ is given by:

$$P(v) = \begin{cases} -K_T - (u-1)C_2 - vE_4 & \text{for } v \in [0, 2-u] \\ -K_T - (u-1)C_2 - vE_4 - (u+v-2)E_5 & \text{for } v \in [2-u, 1] \\ -K_T - (u-1)C_2 - vE_4 - (u+v-2)E_5 - (v-1)(L_{14} + L_{24} + L_{34}) & \text{for } v \in [1, 3-u] \end{cases}$$

and

$$N(v) = \begin{cases} 0 & \text{for } v \in [0, 2-u] \\ (u+v-2)E_5 & \text{for } v \in [2-u, 1] \\ (u+v-2)E_5 + (v-1)(L_{14} + L_{24} + L_{34}) & \text{for } v \in [1, 3-u] \end{cases}$$

Moreover,

$$P(v)^2 = \begin{cases} 5 - u^2 - 2v^2 & \text{for } v \in [0, 2-u] \\ 9 + 2uv - 4u - 4v - v^2 & \text{for } v \in [2-u, 1] \\ 2(2-v)(3-u-v) & \text{for } v \in [1, 3-u] \end{cases}$$

and

$$P(v) \cdot \mathcal{C} = \begin{cases} 2v & \text{for } v \in [0, 2-u] \\ 2-u+v & \text{for } v \in [2-u, 1] \\ 5-u-2v & \text{for } v \in [1, 3-u] \end{cases}$$

Thus,

$$\begin{aligned} S_D(\mathcal{C}) &= \frac{1}{5-u^2} \left(\int_0^{2-u} 5 - u^2 - 2v^2 dv + \int_{2-u}^1 9 + 2uv - 4u - 4v - v^2 dv + \right. \\ &\quad \left. + \int_1^{3-u} 2(2-v)(3-u-v) dv \right) = \frac{16 + 3u - 9u^2 + 2u^3}{15 - 3u^2} \end{aligned}$$

Thus, $\delta_P(T, D) \leq \frac{15-3u^2}{16+3u-9u^2+2u^3}$ for $P \in E_4$. Note that we have:

- if $P \in E_4 \setminus (E_5 \cup L_{14} \cup L_{24} \cup L_{34})$

$$h_D(v) = \begin{cases} 2v^2 & \text{for } v \in [0, 2-u] \\ \frac{(2-u+v)^2}{2} & \text{for } v \in [2-u, 1] \\ \frac{(5-u-2v)^2}{2} & \text{for } v \in [1, 3-u] \end{cases}$$

- if $P = E_4 \cap E_5$

$$h_D(v) = \begin{cases} 2v^2 & \text{for } v \in [0, 2-u] \\ \frac{(2-u+v)(u+3v-2)}{2} & \text{for } v \in [2-u, 1] \\ \frac{(u+1)(5-u-2v)}{2} & \text{for } v \in [1, 3-u] \end{cases}$$

- if $P \in E_4 \cap (L_{14} \cup L_{24} \cup L_{34})$

$$h_D(v) = \begin{cases} 2v^2 & \text{for } v \in [0, 2-u] \\ \frac{(2-u+v)^2}{2} & \text{for } v \in [2-u, 1] \\ \frac{(3-u)(5-u-2v)}{2} & \text{for } v \in [1, 3-u] \end{cases}$$

So we have

- if $P \in E_4 \setminus (E_5 \cup L_{14} \cup L_{24} \cup L_{34})$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^C; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} 2v^2 dv + \int_{2-u}^1 \frac{(2-u+v)^2}{2} dv + \int_1^{3-u} \frac{(5-u-2v)^2}{2} dv \right) = \\ &= \frac{9+6u-9u^2+2u^3}{15-3u^2} \leq \frac{16+3u-9u^2+2u^3}{15-3u^2} \end{aligned}$$

- if $P = E_4 \cap E_5$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^C; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} 2v^2 dv + \int_{2-u}^1 \frac{(2-u+v)(u+3v-2)}{2} dv + \right. \\ &\quad \left. + \int_1^{3-u} \frac{(u+1)(5-u-2v)}{2} dv \right) = \frac{11-u^3}{15-3u^2} \end{aligned}$$

- if $P \in E_4 \cap (L_{14} \cup L_{24} \cup L_{34})$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^C; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} 2v^2 dv + \int_{2-u}^1 \frac{(2-u+v)^2}{2} dv + \right. \\ &\quad \left. + \int_1^{3-u} \frac{(3-u)(5-u-2v)}{2} dv \right) = \\ &= \frac{13+3u^3-12u^2+6u}{15-3u^2} \leq \frac{16+3u-9u^2+2u^3}{15-3u^2} \end{aligned}$$

We obtain that

$$\delta_P(T, D) = \frac{15-3u^2}{16+3u-9u^2+2u^3} \text{ for } P \in E_4 \setminus E_5 \text{ and } u \in [1, 2]$$

and

$$\delta_P(T, D) \geq \begin{cases} \frac{15-3u^2}{16+3u-9u^2+2u^3} & \text{for } P = E_4 \cap E_5 \text{ and } u \in [1, a] \\ \frac{15-3u^2}{11-u^3} & \text{for } P = E_4 \cap E_5 \text{ and } u \in [a, 2] \end{cases}$$

where a is a root of $3u^3 - 9u^2 + 3u + 5$ on $[1, 2]$. Note that $a \in [1.355, 1.356]$.

Step 2. Suppose $P \in E_5$. In this case we set $\mathcal{C} = E_5$. Then $\tau(\mathcal{C}) = 2$. The Zariski Decomposition of the divisor $D - vE_5$ is given by:

$$P(v) = \begin{cases} -K_T - (u-1)C_2 - vE_5 - \frac{v}{2}E_4 & \text{for } v \in [0, 1] \\ -K_T - (u-1)C_2 - vE_5 - \frac{v}{2}E_4 - (v-1)L_{45} & \text{for } v \in [1, u] \\ -K_T - (u-1)C_2 - vE_5 - \frac{v}{2}E_4 - (v-1)L_{45} - (v-u)C_2 & \text{for } v \in [u, 2] \end{cases}$$

and

$$N(v) = \begin{cases} \frac{v}{2}E_4 & \text{for } v \in [0, 1] \\ \frac{v}{2}E_4 + (v-1)L_{45} & \text{for } v \in [1, u] \\ \frac{v}{2}E_4 + (v-1)L_{45} + (v-u)C_2 & \text{for } v \in [u, 2] \end{cases}$$

Moreover,

$$P(v)^2 = \begin{cases} 5 - 4v + 2uv - u^2 - v^2/2 & \text{for } v \in [0, 1] \\ 6 - 6v + v^2/2 + 2uv - u^2 & \text{for } v \in [1, u] \\ \frac{3(2-v)^2}{2} & \text{for } v \in [u, 2] \end{cases}$$

and

$$P(v) \cdot \mathcal{C} = \begin{cases} 2 - u + v/2 & \text{for } v \in [0, 1] \\ 3 - u - v/2 & \text{for } v \in [1, u] \\ 3 - 3v/2 & \text{for } v \in [u, 2] \end{cases}$$

Thus,

$$\begin{aligned} S_D(\mathcal{C}) &= \frac{1}{5-u^2} \left(\int_0^1 5 - 4v + 2uv - u^2 - v^2/2 dv + \int_1^u 6 - 6v + v^2/2 + 2uv - u^2 dv + \right. \\ &\quad \left. + \int_u^2 \frac{3(2-v)^2}{3} dv \right) = \frac{11-u^3}{15-3u^2} \end{aligned}$$

Thus, $\delta_P(T, D) \leq \frac{15-3u^2}{11-u^3}$ for $P \in E_5$. Note that we have:

- if $P \in E_5 \setminus (E_4 \cup C_2 \cup L_{45})$ then

$$h_D(v) = \begin{cases} \frac{(2-u+v/2)^2}{2} & \text{for } v \in [0, 1] \\ \frac{(3-u-v/2)^2}{2} & \text{for } v \in [1, u] \\ \frac{(3-3v/2)^2}{2} & \text{for } v \in [u, 2] \end{cases}$$

- if $P = E_5 \cap C_2$ then

$$h_D(v) = \begin{cases} \frac{(2-u+v/2)^2}{2} & \text{for } v \in [0, 1] \\ \frac{(3-u-v/2)^2}{2} & \text{for } v \in [1, u] \\ \frac{3(2-v)(6-4u+v)}{8} & \text{for } v \in [u, 2] \end{cases}$$

- if $P = E_5 \cap L_{45}$ then

$$h_D(v) = \begin{cases} \frac{(2-u+v/2)^2}{2} & \text{for } v \in [0, 1] \\ \frac{(6-2u-v)(2-2u+3v)}{8} & \text{for } v \in [1, u] \\ \frac{3(2-v)(v+2)}{8} & \text{for } v \in [u, 2] \end{cases}$$

So we have

- if $P \in E_5 \setminus (E_4 \cup C_2 \cup L_{45})$ then

$$S_D(W_{\bullet,\bullet}^C; P) = \frac{2}{5-u^2} \left(\int_0^1 \frac{(2-u+v/2)^2}{2} dv + \int_1^u \frac{(3-u-v/2)^2}{2} dv + \int_u^2 \frac{(3-3v/2)^2}{2} dv \right) = \frac{21+6u-18u^2+5u^3}{2(15-3u^2)} \leq \frac{11-u^3}{15-3u^2}$$

- if $P = E_5 \cap C_2$ then

$$S_D(W_{\bullet,\bullet}^C; P) = \frac{2}{5-u^2} \left(\int_0^1 \frac{(2-u+v/2)^2}{2} dv + \int_1^u \frac{(3-u-v/2)^2}{2} dv + \int_u^2 \frac{3(2-v)(6-4u+v)}{8} dv \right) = \frac{45-30u+2u^3}{2(15-3u^2)} \leq \frac{11-u^3}{15-3u^2}$$

- if $P = E_5 \cap L_{45}$ then

$$S_D(W_{\bullet,\bullet}^C; P) = \frac{2}{5-u^2} \left(\int_0^1 \frac{(2-u+v/2)^2}{2} dv + \int_1^u \frac{(6-2u-v)(2-2u+3v)}{8} dv + \int_u^2 \frac{3(2-v)(v+2)}{8} dv \right) = \frac{26-12u^2+3u^3}{2(15-3u^2)} \leq \frac{11-u^3}{15-3u^2}$$

We obtain that

$$\delta_P(T, D) = \frac{15-3u^2}{11-u^3} \text{ for } P \in E_5 \setminus E_4 \text{ and } u \in [1, 2].$$

Step 3.1. Suppose $P \in L_{14} \cup L_{24} \cup L_{34}$ and $u \in [1, 3/2]$. In this case we set $\mathcal{C} = L_{14}$. Then $\tau(\mathcal{C}) = 3 - u$. Without loss of generality, we can assume that $P \in L_{14}$. The Zariski Decomposition of the divisor $D - vL_{14}$ is given by:

$$P(v) = \begin{cases} D - vL_{14} - \frac{v}{2}E_4 & \text{for } v \in [0, 2-u] \\ D - vL_{14} - \frac{v}{2}E_4 - (u+v-2)E_1 & \text{for } v \in [2-u, 1] \\ D - vL_{14} - \frac{v}{2}E_4 - (u+v-2)E_1 - (v-1)L_{23} & \text{for } v \in [1, 4-2u] \\ D - vL_{14} - (u+v-2)(E_1 + E_4) - (v-1)L_{23} - (2u+v-4)E_5 & \text{for } v \in [4-2u, 3-u] \end{cases}$$

and

$$N(v) = \begin{cases} \frac{v}{2}E_4 & \text{for } v \in [0, 2-u] \\ \frac{v}{2}E_4 + (u+v-2)E_1 & \text{for } v \in [2-u, 1] \\ \frac{v}{2}E_4 + (u+v-2)E_1 + (v-1)L_{23} & \text{for } v \in [1, 4-2u] \\ (u+v-2)(E_1 + E_4) + (v-1)L_{23} + (2u+v-4)E_5 & \text{for } v \in [4-2u, 3-u] \end{cases}$$

Moreover

$$P(v)^2 = \begin{cases} 5-2v-v^2/2-u^2 & \text{for } v \in [0, 2-u] \\ 9-4u-6v+v^2/2+2uv & \text{for } v \in [2-u, 1] \\ \frac{(v-2)(3v+4u-10)}{2} & \text{for } v \in [1, 4-2u] \\ 2(u+v-3)^2 & \text{for } v \in [4-2u, 3-u] \end{cases}$$

and

$$P(v) \cdot \mathcal{C} = \begin{cases} v/2+1 & \text{for } v \in [0, 2-u] \\ 3-u-v/2 & \text{for } v \in [2-u, 1] \\ 4-u-3v/2 & \text{for } v \in [1, 4-2u] \\ 2(3-u-v) & \text{for } v \in [4-2u, 3-u] \end{cases}$$

Thus,

$$S_D(\mathcal{C}) = \frac{1}{5-u^2} \left(\int_0^{2-u} 5 - 2v - v^2/2 - u^2 dv + \int_{2-u}^1 9 - 4u - 6v + v^2/2 + 2uv dv + \right. \\ \left. + \int_1^{4-2u} \frac{(v-2)(3v+4u-10)}{2} dv + \int_{4-2u}^{3-u} 2(u+v-3)^2 dv \right) = \frac{3u^3 - 12u^2 + 6u + 13}{15 - 3u^2}$$

Thus, $\delta_P(T, D) \leq \frac{15-3u^2}{3u^3-12u^2+6u+13}$ for $P \in L_{14}$. Note that we have:

- if $P \in L_{14} \setminus (E_4 \cup E_1 \cup L_{23} \cup E_5)$ then

$$h_D(v) = \begin{cases} \frac{(v/2+1)^2}{2} & \text{for } v \in [0, 2-u] \\ \frac{(3-u-v/2)^2}{2} & \text{for } v \in [2-u, 1] \\ \frac{(4-u-3v/2)^2}{2} & \text{for } v \in [1, 4-2u] \\ 2(3-u-v)^2 & \text{for } v \in [4-2u, 3-u] \end{cases}$$

- if $P = L_{14} \cap E_1$ then

$$h_D(v) = \begin{cases} \frac{(v/2+1)^2}{2} & \text{for } v \in [0, 2-u] \\ \frac{(6-2u-v)(2u+3v-2)}{8} & \text{for } v \in [2-u, 1] \\ \frac{(8-2u-3v)(2u+v)}{8} & \text{for } v \in [1, 4-2u] \\ (3-u-v) & \text{for } v \in [4-2u, 3-u] \end{cases}$$

- if $P = L_{14} \cap L_{23}$ then

$$h_D(v) = \begin{cases} \frac{(v/2+1)^2}{2} & \text{for } v \in [0, 2-u] \\ \frac{(3-u-v/2)^2}{2} & \text{for } v \in [2-u, 1] \\ \frac{(8-2u-3v)(4-2u+v)}{8} & \text{for } v \in [1, 4-2u] \\ 2(2-u)(3-u-v) & \text{for } v \in [4-2u, 3-u] \end{cases}$$

So we have

- if $P \in L_{14} \setminus (E_4 \cup E_1 \cup L_{23} \cup E_5)$ then

$$S_D(W_{\bullet,\bullet}^{\mathcal{C}}; P) = \frac{2}{5-u^2} \left(\int_0^{2-u} \frac{(v/2+1)^2}{2} dv + \int_{2-u}^1 \frac{(3-u-v/2)^2}{2} dv + \right. \\ \left. + \int_1^{4-2u} \frac{(4-u-3v/2)^2}{2} dv + \int_{4-2u}^{3-u} 2(3-u-v)^2 dv \right) = \\ = \frac{21 - u^3 - 6u}{2(15 - 3u^2)} \leq \frac{3u^3 - 12u^2 + 6u + 13}{15 - 3u^2}$$

- if $P = L_{14} \cap E_1$ then

$$S_D(W_{\bullet,\bullet}^{\mathcal{C}}; P) = \frac{2}{5-u^2} \left(\int_0^{2-u} \frac{(v/2+1)^2}{2} dv + \int_{2-u}^1 \frac{(6-2u-v)(2u+3v-2)}{8} dv + \right. \\ \left. + \int_1^{4-2u} \frac{(8-2u-3v)(2u+v)}{8} dv + \int_{4-2u}^{3-u} (3-u-v) dv \right) = \frac{19 - 2u^3}{2(15 - 3u^2)}$$

- if $P = L_{14} \cap L_{23}$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^C; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} \frac{(v/2+1)^2}{2} dv + \int_{2-u}^1 \frac{(3-u-v/2)^2}{2} dv + \right. \\ &\quad \left. + \int_1^{4-2u} \frac{(8-2u-3v)(4-2u+v)}{8} dv + \int_{4-2u}^{3-u} 2(2-u)(3-u-v) dv \right) = \\ &= \frac{26-12u^2+3u^3}{2(15-3u^2)} \leq \frac{3u^3-12u^2+6u+13}{15-3u^2} \end{aligned}$$

We obtain that

$$\delta_P(T, D) = \frac{15-3u^2}{3u^3-12u^2+6u+13} \text{ for } P \in L_{14} \setminus (E_4 \cup E_5 \cup E_1) \text{ and } u \in [1, 3/2]$$

and

$$\delta_P(T, D) \geq \begin{cases} \frac{15-3u^2}{3u^3-12u^2+6u+13} & \text{for } P = L_{14} \cap E_1 \text{ and } u \in [1, b] \\ \frac{2(15-3u^2)}{19-2u^3} & \text{for } P = L_{14} \cap E_1 \text{ and } u \in [b, 3/2] \end{cases}$$

where b is a root of $8u^3 - 24u^2 + 12u + 7$ on $[1, 3/2]$. Note that $b \in [1.261, 1.262]$.

Step 3.2. Suppose $P \in L_{14} \cup L_{24} \cup L_{34}$ and $u \in [3/2, 2]$. In this case we set $\mathcal{C} = L_{14}$. Then $\tau(\mathcal{C}) = 3 - u$. Without loss of generality, we can assume that $P \in L_{14}$. The Zariski Decomposition of the divisor $D - vL_{14}$ is given by:

$$P(v) = \begin{cases} D - vL_{14} - \frac{v}{2}E_4 & \text{for } v \in [0, 2-u] \\ D - vL_{14} - \frac{v}{2}E_4 - (u+v-2)E_1 & \text{for } v \in [2-u, 4-2u] \\ D - vL_{14} - (u+v-2)(E_1+E_4) - (2u+v-4)E_5 & \text{for } v \in [4-2u, 1] \\ D - vL_{14} - (u+v-2)(E_1+E_4) - (v-1)L_{23} - (2u+v-4)E_5 & \text{for } v \in [1, 3-u] \end{cases}$$

and

$$N(v) = \begin{cases} \frac{v}{2}E_4 & \text{for } v \in [0, 2-u] \\ \frac{v}{2}E_4 + (u+v-2)E_1 & \text{for } v \in [2-u, 4-2u] \\ (u+v-2)(E_1+E_4) + (2u+v-4)E_5 & \text{for } v \in [4-2u, 1] \\ (u+v-2)(E_1+E_4) + (v-1)L_{23} + (2u+v-4)E_5 & \text{for } v \in [1, 3-u] \end{cases}$$

Moreover

$$P(v)^2 = \begin{cases} 5-2v-v^2/2-u^2 & \text{for } v \in [0, 2-u] \\ 9-4u-6v+v^2/2+2uv & \text{for } v \in [2-u, 4-2u] \\ 2u^2+4uv+v^2-12u-10v+17 & \text{for } v \in [4-2u, 1] \\ 2(u+v-3)^2 & \text{for } v \in [1, 3-u] \end{cases}$$

and

$$P(v) \cdot \mathcal{C} = \begin{cases} 1+v/2 & \text{for } v \in [0, 2-u] \\ 3-u-v/2 & \text{for } v \in [2-u, 4-2u] \\ 5-2u-v & \text{for } v \in [4-2u, 1] \\ 2(3-u-v) & \text{for } v \in [1, 3-u] \end{cases}$$

Thus,

$$\begin{aligned} S_D(\mathcal{C}) &= \frac{1}{5-u^2} \left(\int_0^{2-u} 5-2v-v^2/2-u^2 dv + \int_{2-u}^{4-2u} 9-4u-6v+v^2/2+2uv dv + \right. \\ &\quad \left. + \int_{4-2u}^1 2u^2+4uv+v^2-12u-10v+17 dv + \int_1^{3-u} 2(u+v-3)^2 dv \right) = \frac{3u^3-12u^2+6u+13}{15-3u^2} \end{aligned}$$

Thus, $\delta_P(T, D) \leq \frac{15-3u^2}{3u^3-12u^2+6u+13}$ for $P \in L_{14}$. Note that we have:

- if $P \in L_{14} \setminus (E_4 \cup E_1 \cup L_{23} \cup E_5)$ then

$$h_D(v) = \begin{cases} \frac{(1+v/2)^2}{2} & \text{for } v \in [0, 2-u] \\ \frac{(3-u-v/2)^2}{2} & \text{for } v \in [2-u, 4-2u] \\ \frac{(5-2u-v)^2}{2} & \text{for } v \in [4-2u, 1] \\ 2(3-u-v)^2 & \text{for } v \in [1, 3-u] \end{cases}$$

- if $P = L_{14} \cap E_1$ then

$$h_D(v) = \begin{cases} \frac{(1+v/2)^2}{2} & \text{for } v \in [0, 2-u] \\ \frac{(6-2u-v)(2u+3v-2)}{8} & \text{for } v \in [2-u, 4-2u] \\ \frac{(v+1)(5-2u-v)}{2} & \text{for } v \in [4-2u, 1] \\ 2(3-u-v) & \text{for } v \in [1, 3-u] \end{cases}$$

- if $P = L_{14} \cap L_{23}$ then

$$h_D(v) = \begin{cases} \frac{(1+v/2)^2}{2} & \text{for } v \in [0, 2-u] \\ \frac{(3-u-v/2)^2}{2} & \text{for } v \in [2-u, 4-2u] \\ \frac{(5-2u-v)^2}{2} & \text{for } v \in [4-2u, 1] \\ 2(2-u)(3-u-v) & \text{for } v \in [1, 3-u] \end{cases}$$

- if $P \in L_{14} \setminus (E_4 \cup E_1 \cup L_{23} \cup E_5)$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^C; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} \frac{(v/2+1)^2}{2} dv + \int_{2-u}^{4-2u} \frac{(3-u-v/2)^2}{2} dv + \right. \\ &\quad \left. + \int_{4-2u}^1 \frac{(5-2u-v)^2}{2} dv + \int_1^{3-u} 2(3-u-v)^2 dv \right) = \\ &= \frac{7u^3 - 36u^2 + 48u - 6}{2(15-3u^2)} \leq \frac{3u^3 - 12u^2 + 6u + 13}{15-3u^2} \end{aligned}$$

- if $P = L_{14} \cap E_1$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^C; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} \frac{(v/2+1)^2}{2} dv + \int_{2-u}^{4-2u} \frac{(6-2u-v)(2u+3v-2)}{8} dv + \right. \\ &\quad \left. + \int_{4-2u}^1 \frac{(v+1)(5-2u-v)}{2} dv + \int_1^{3-u} 2(3-u-v) dv \right) = \frac{3u^3 - 18u^2 + 27u - 4}{15-3u^2} \end{aligned}$$

- if $P = L_{14} \cap L_{23}$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^C; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} \frac{(v/2+1)^2}{2} dv + \int_{2-u}^{4-2u} \frac{(3-u-v/2)^2}{2} dv + \right. \\ &\quad \left. + \int_{4-2u}^1 \frac{(5-2u-v)^2}{2} dv + \int_1^{3-u} 2(2-u)(3-u-v) dv \right) = \\ &= \frac{3u^3 - 12u^2 + 26}{2(15-3u^2)} \leq \frac{3u^3 - 12u^2 + 6u + 13}{15-3u^2} \end{aligned}$$

We obtain that

$$\delta_P(T, D) = \frac{15-3u^2}{3u^3-12u^2+6u+13} \text{ for } P \in L_{14} \setminus (E_1 \cup E_4 \cup E_5) \text{ and } u \in [3/2, 2]$$

and

$$\delta_P(T, D) \geq \frac{15 - 3u^2}{3u^3 - 18u^2 + 27u - 4} \text{ for } P = L_{14} \cap E_1 \text{ and } u \in [3/2, 2]$$

□

Corollary A.1. *Let P be a point in T that is contained in $L_{12} \cup L_{24} \cup L_{34} \cup E_4 \cup E_5$ then*

$$\delta_P(T, D) \geq \begin{cases} \frac{15 - 3u^2}{16 + 3u - 9u^2 + 2u^3} & \text{for } u \in [1, a], \\ \frac{15 - 3u^2}{11 - u^3} & \text{for } u \in [a, 2] \end{cases}$$


Corollary A.2. *Suppose O is a point on a del Pezzo surface \bar{T} with \mathbb{A}_1 singularity and $\delta_O(T) \leq \frac{6}{5}$ then*

$$\delta_O(\bar{T}, \bar{D}) \geq \begin{cases} \frac{15 - 3u^2}{16 + 3u - 9u^2 + 2u^3} & \text{for } u \in [1, a], \\ \frac{15 - 3u^2}{11 - u^3} & \text{for } u \in [a, 2] \end{cases}$$

A.2. Polarized δ -invariant on Del Pezzo surface of degree 4 with two \mathbb{A}_1 singularities. Suppose that \bar{T} has two singular points and these points are singular point of type \mathbb{A}_1 . Then η is a blow up of \mathbb{P}^2 at points P_1, P_2 , and P_4 in general position and after that blowing up a point P_3 which belongs to the exceptional divisor corresponding to P_2 and a point P_5 which belongs to the exceptional divisor corresponding to P_4 and no other negative curve. By [7, Section 6.2] we have:

$$\delta_P(T) = \begin{cases} 1 & \text{if } P \in (E_2 \cup E_4 \cup L_{24}), \\ 6/5 & \text{if } P \in (E_3 \cup E_5 \cup L_{12} \cup L_{14}) \setminus (E_2 \cup E_4), \\ 4/3 & \text{if } P \in (C_2 \cap E_1) \cup (L_{23} \cap L_{45}), \\ 18/13 & \text{if } P \in (C_2 \cup E_1 \cup L_{23} \cup L_{45}) \setminus ((C_2 \cap E_1) \cup (L_{23} \cap L_{45}) \cup (E_3 \cup E_5 \cup L_{12} \cup L_{14})), \\ 3/2, & \text{otherwise} \end{cases}$$

where E_1, E_2, E_3, E_4, E_5 are exceptional divisors corresponding to P_1, P_2, P_3, P_4, P_5 respectively, C_2 is a strict transform of a (-1) -curve coming from the conic on \mathbb{P}^2 , L_{ij} are strict transforms of the lines passing through P_i and P_j for $(i, j) \in \{(1, 2), (1, 4)\}$ and L_{45} and L_{23} are strict transforms of a (-1) -curve coming from lines passing through P_2 and P_4 respectively on \mathbb{P}^2 . The dual graph of (-1) and (-2) -curves is given in the following picture:



Lemma A.2. Suppose P is a point on T and $D = -K_T - (u-1)C_2$ with $D^2 = 5 - u^2$ then

$$\delta_P(T, D) = \begin{cases} \frac{15 - 3u^2}{16 + 3u - 9u^2 + 2u^3} \text{ for } P \in (E_2 \cup E_4) \setminus (E_3 \cup E_5) \text{ and } u \in [1, 2], \\ \frac{15 - 3u^2}{11 - u^3} \text{ for } P \in (E_3 \cup E_5) \setminus (E_2 \cup E_4) \text{ and } u \in [1, 2], \\ \frac{15 - 3u^2}{u^3 - 6u^2 + 6u + 5} \text{ for } P \in L_{24} \setminus (E_2 \cup E_4) \text{ and } u \in [1, 2], \\ \frac{15 - 3u^2}{3u^3 - 12u^2 + 6u + 13} \text{ for } P \in L_{14} \setminus (E_4 \cup E_5 \cup E_1) \text{ and } u \in [1, 2], \end{cases}$$

and

$$\delta_P(T, D) \geq \begin{cases} \frac{15 - 3u^2}{16 + 3u - 9u^2 + 2u^3} \text{ for } P \in \{E_2 \cap E_3, E_4 \cap E_5\} \text{ and } u \in [1, a], \\ \frac{15 - 3u^2}{11 - u^3} \text{ for } P \in \{E_2 \cap E_3, E_4 \cap E_5\} \text{ and } u \in [a, 2] \end{cases}$$

and

$$\delta_P(T, D) \geq \begin{cases} \frac{15 - 3u^2}{3u^3 - 12u^2 + 6u + 13} \text{ for } P = L_{14} \cap E_1 \text{ and } u \in [1, b], \\ \frac{2(15 - 3u^2)}{19 - 2u^3} \text{ for } P = L_{14} \cap E_1 \text{ and } u \in [b, 3/2], \\ \frac{15 - 3u^2}{3u^3 - 18u^2 + 27u - 4} \text{ for } P = L_{14} \cap E_1 \text{ and } u \in [3/2, 2], \end{cases}$$

where a is a root of $3u^3 - 9u^2 + 3u + 5$ on $[1, 2]$, b is a root of $8u^3 - 24u^2 + 12u + 7$ on $[1, 3/2]$. Note that $a \in [1.355, 1.356]$, $b \in [1.261, 1.262]$.

Proof. Step 1. Suppose $P \in E_2 \cup E_4$. Without loss of generality we can assume that $P \in E_4$. In this case we set $\mathcal{C} = E_4$. Then $\tau(\mathcal{C}) = 3 - u$. The Zariski Decomposition of the divisor $D - vE_4$ is given by:

$$P(v) = \begin{cases} -K_T - (u-1)C_2 - vE_4 \text{ for } v \in [0, 2-u] \\ -K_T - (u-1)C_2 - vE_4 - (u+v-2)E_5 \text{ for } v \in [2-u, 1] \\ -K_T - (u-1)C_2 - vE_4 - (u+v-2)E_5 - (v-1)(L_{14} + 2L_{24} + E_2) \text{ for } v \in [1, 3-u] \end{cases}$$

and

$$N(v) = \begin{cases} 0 \text{ for } v \in [0, 2-u] \\ (u+v-2)E_5 \text{ for } v \in [2-u, 1] \\ (u+v-2)E_5 + (v-1)(L_{14} + 2L_{24} + E_2) \text{ for } v \in [1, 3-u] \end{cases}$$

Moreover,

$$P(v)^2 = \begin{cases} 5 - u^2 - 2v^2 \text{ for } v \in [0, 2-u] \\ 9 + 2uv - 4u - 4v - v^2 \text{ for } v \in [2-u, 1] \\ 2(2-v)(3-u-v) \text{ for } v \in [1, 3-u] \end{cases}$$

and

$$P(v) \cdot \mathcal{C} = \begin{cases} 2v \text{ for } v \in [0, 2-u] \\ 2-u+v \text{ for } v \in [2-u, 1] \\ 5-u-2v \text{ for } v \in [1, 3-u] \end{cases}$$

Thus,

$$S_D(\mathcal{C}) = \frac{1}{5-u^2} \left(\int_0^{2-u} 5 - u^2 - 2v^2 dv + \int_{2-u}^1 9 + 2uv - 4u - 4v - v^2 dv + \int_1^{3-u} 2(2-v)(3-u-v) dv \right) = \frac{16+3u-9u^2+2u^3}{15-3u^2}$$

Thus, $\delta_P(T, D) \leq \frac{15-3u^2}{16+3u-9u^2+2u^3}$ for $P \in E_4$. Note that we have:

- if $P \in E_4 \setminus (E_5 \cup L_{14} \cup L_{24} \cup L_{34})$

$$h_D(v) = \begin{cases} 2v^2 & \text{for } v \in [0, 2-u] \\ \frac{(2-u+v)^2}{2} & \text{for } v \in [2-u, 1] \\ \frac{(5-u-2v)^2}{2} & \text{for } v \in [1, 3-u] \end{cases}$$

- if $P = E_4 \cap E_5$

$$h_D(v) = \begin{cases} 2v^2 & \text{for } v \in [0, 2-u] \\ \frac{(2-u+v)(u+3v-2)}{2} & \text{for } v \in [2-u, 1] \\ \frac{(u+1)(5-u-2v)}{2} & \text{for } v \in [1, 3-u] \end{cases}$$

- if $P \in E_4 \cap (L_{14} \cup L_{24})$

$$h_D(v) \leq \begin{cases} 2v^2 & \text{for } v \in [0, 2-u] \\ \frac{(2-u+v)^2}{2} & \text{for } v \in [2-u, 1] \\ \frac{(5-u-2v)(1-u+2v)}{2} & \text{for } v \in [1, 3-u] \end{cases}$$

So we have

- if $P \in E_4 \setminus (E_5 \cup L_{14} \cup L_{24} \cup L_{34})$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^{\mathcal{C}}; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} 2v^2 dv + \int_{2-u}^1 \frac{(2-u+v)^2}{2} dv + \int_1^{3-u} \frac{(5-u-2v)^2}{2} dv \right) = \\ &= \frac{9+6u-9u^2+2u^3}{15-3u^2} \leq \frac{16+3u-9u^2+2u^3}{15-3u^2} \end{aligned}$$

- if $P = E_4 \cap E_5$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^{\mathcal{C}}; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} 2v^2 dv + \int_{2-u}^1 \frac{(2-u+v)(u+3v-2)}{2} dv + \right. \\ &\quad \left. + \int_1^{3-u} \frac{(u+1)(5-u-2v)}{2} dv \right) = \frac{11-u^3}{15-3u^2} \end{aligned}$$

- if $P \in E_4 \cap (L_{14} \cup L_{24})$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^{\mathcal{C}}; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} 2v^2 dv + \int_{2-u}^1 \frac{(2-u+v)^2}{2} dv + \int_1^{3-u} \frac{(5-u-2v)(1-u+2v)}{2} dv \right) = \\ &= \frac{2u^3-6u^2+8}{15-3u^2} \leq \frac{16+3u-9u^2+2u^3}{15-3u^2} \end{aligned}$$

We obtain that

$$\delta_P(T, D) = \frac{15-3u^2}{16+3u-9u^2+2u^3} \text{ for } P \in E_4 \setminus E_5 \text{ and } u \in [1, 2]$$

and

$$\delta_P(T, D) \geq \begin{cases} \frac{15 - 3u^2}{16 + 3u - 9u^2 + 2u^3} & \text{for } P = E_4 \cap E_5 \text{ and } u \in [1, a] \\ \frac{15 - 3u^2}{11 - u^3} & \text{for } P = E_4 \cap E_5 \text{ and } u \in [a, 2] \end{cases}$$

where a is a root of $3u^3 - 9u^2 + 3u + 5$ on $[1, 2]$. Note that $a \in [1.355, 1.356]$.

Step 2. Suppose $P \in E_3 \cup E_5$. In this case we set $\mathcal{C} = E_5$. Then $\tau(\mathcal{C}) = 2$. Without loss of generality we can assume that $P \in E_5$. The Zariski Decomposition of the divisor $D - vE_5$ is given by:

$$P(v) = \begin{cases} -K_T - (u - 1)C_2 - vE_5 - \frac{v}{2}E_4 & \text{for } v \in [0, 1] \\ -K_T - (u - 1)C_2 - vE_5 - \frac{v}{2}E_4 - (v - 1)L_{45} & \text{for } v \in [1, u] \\ -K_T - (u - 1)C_2 - vE_5 - \frac{v}{2}E_4 - (v - 1)L_{45} - (v - u)C_2 & \text{for } v \in [u, 2] \end{cases}$$

and

$$N(v) = \begin{cases} \frac{v}{2}E_4 & \text{for } v \in [0, 1] \\ \frac{v}{2}E_4 + (v - 1)L_{45} & \text{for } v \in [1, u] \\ \frac{v}{2}E_4 + (v - 1)L_{45} + (v - u)C_2 & \text{for } v \in [u, 2] \end{cases}$$

Moreover,

$$P(v)^2 = \begin{cases} 5 - 4v + 2uv - u^2 - v^2/2 & \text{for } v \in [0, 1] \\ 6 - 6v + v^2/2 + 2uv - u^2 & \text{for } v \in [1, u] \\ \frac{3(2-v)^2}{2} & \text{for } v \in [u, 2] \end{cases}$$

and

$$P(v) \cdot \mathcal{C} = \begin{cases} 2 - u + v/2 & \text{for } v \in [0, 1] \\ 3 - u - v/2 & \text{for } v \in [1, u] \\ 3 - 3v/2 & \text{for } v \in [u, 2] \end{cases}$$

Thus,

$$S_D(\mathcal{C}) = \frac{1}{5 - u^2} \left(\int_0^1 5 - 4v + 2uv - u^2 - v^2/2 dv + \int_1^u 6 - 6v + v^2/2 + 2uv - u^2 dv + \int_u^2 \frac{3(2-v)^2}{3} dv \right) = \frac{11 - u^3}{15 - 3u^2}$$

Thus, $\delta_P(T, D) \leq \frac{15 - 3u^2}{11 - u^3}$ for $P \in E_5$. Note that we have:

- if $P \in E_5 \setminus (E_4 \cup C_2 \cup L_{45})$ then

$$h_D(v) = \begin{cases} \frac{(2-u+v/2)^2}{2} & \text{for } v \in [0, 1] \\ \frac{(3-u-v/2)^2}{2} & \text{for } v \in [1, u] \\ \frac{(3-3v/2)^2}{2} & \text{for } v \in [u, 2] \end{cases}$$

- if $P = E_5 \cap C_2$ then

$$h_D(v) = \begin{cases} \frac{(2-u+v/2)^2}{2} & \text{for } v \in [0, 1] \\ \frac{(3-u-v/2)^2}{2} & \text{for } v \in [1, u] \\ \frac{3(2-v)(6-4u+v)}{8} & \text{for } v \in [u, 2] \end{cases}$$

- if $P = E_5 \cap L_{45}$ then

$$h_D(v) = \begin{cases} \frac{(2-u+v/2)^2}{2} & \text{for } v \in [0, 1] \\ \frac{(6-2u-v)(2-2u+3v)}{8} & \text{for } v \in [1, u] \\ \frac{3(2-v)(v+2)}{8} & \text{for } v \in [u, 2] \end{cases}$$

So we have

- if $P \in E_5 \setminus (E_4 \cup C_2 \cup L_{45})$ then

$$S_D(W_{\bullet,\bullet}^C; P) = \frac{2}{5-u^2} \left(\int_0^1 \frac{(2-u+v/2)^2}{2} dv + \int_1^u \frac{(3-u-v/2)^2}{2} dv + \int_u^2 \frac{(3-3v/2)^2}{2} dv \right) = \frac{21+6u-18u^2+5u^3}{2(15-3u^2)} \leq \frac{11-u^3}{15-3u^2}$$

- if $P = E_5 \cap C_2$ then

$$S_D(W_{\bullet,\bullet}^C; P) = \frac{2}{5-u^2} \left(\int_0^1 \frac{(2-u+v/2)^2}{2} dv + \int_1^u \frac{(3-u-v/2)^2}{2} dv + \int_u^2 \frac{3(2-v)(6-4u+v)}{8} dv \right) = \frac{45-30u+2u^3}{2(15-3u^2)} \leq \frac{11-u^3}{15-3u^2}$$

- if $P = E_5 \cap L_{45}$ then

$$S_D(W_{\bullet,\bullet}^C; P) = \frac{2}{5-u^2} \left(\int_0^1 \frac{(2-u+v/2)^2}{2} dv + \int_1^u \frac{(6-2u-v)(2-2u+3v)}{8} dv + \int_u^2 \frac{3(2-v)(v+2)}{8} dv \right) = \frac{26-12u^2+3u^3}{2(15-3u^2)} \leq \frac{11-u^3}{15-3u^2}$$

We obtain that

$$\delta_P(T, D) = \frac{15-3u^2}{11-u^3} \text{ for } P \in (E_3 \cup E_5) \setminus (E_2 \cup E_4) \text{ and } u \in [1, 2].$$

Step 3. Suppose $P \in L_{24}$. In this case we set $\mathcal{C} = L_{24}$. Then $\tau(\mathcal{C}) = 3 - u$. The Zariski Decomposition of the divisor $D - vL_{24}$ is given by:

$$P(v) = \begin{cases} D - vL_{24} - \frac{v}{2}(E_2 + E_4) & \text{for } v \in [0, 4-2u] \\ D - vL_{24} - (u+v-2)(E_2 + E_4) - (2u+v-4)(E_3 + E_5) & \text{for } v \in [4-2u, 3-u] \end{cases}$$

and

$$N(v) = \begin{cases} \frac{v}{2}(E_2 + E_4) & \text{for } v \in [0, 4-2u] \\ (u+v-2)(E_2 + E_4) + (2u+v-4)(E_3 + E_5) & \text{for } v \in [4-2u, 3-u] \end{cases}$$

Moreover,

$$P(v)^2 = \begin{cases} -u^2 - 2v + 5 & \text{for } v \in [0, 4-2u] \\ (u+v-3)(3u+v-7) & \text{for } v \in [4-2u, 3-u] \end{cases}$$

and

$$P(v) \cdot \mathcal{C} = \begin{cases} 1 & \text{for } v \in [0, 4-2u] \\ 5-2u-v & \text{for } v \in [4-2u, 3-u] \end{cases}$$

Thus,

$$\begin{aligned} S_D(\mathcal{C}) &= \frac{1}{5-u^2} \left(\int_0^{4-2u} -u^2 - 2v + 5 dv + \int_{4-2u}^{3-uu} (u+v-3)(3u+v-7) dv \right) = \\ &= \frac{4u^3 - 15u^2 + 6u + 17}{15-3u^2} \end{aligned}$$

Thus, $\delta_P(T, D) \leq \frac{15-3u^2}{4u^3-15u^2+6u+17}$ for $P \in L_{24}$. If $P \in L_{24} \setminus (E_2 \cup E_4)$ then

$$h_D(v) = \begin{cases} \frac{1}{2} & \text{for } v \in [0, 4-2u] \\ \frac{(5-2u-v)^2}{2} & \text{for } v \in [4-2u, 3-u] \end{cases}$$

So for $P \in L_{24} \setminus (E_2 \cup E_4)$ we have

$$\begin{aligned} S_D(W_{\bullet,\bullet}^{\mathcal{C}}; P) &= \frac{2}{5-u^2} \left(\int_0^{4-2u} \frac{1}{2} dv + \int_{4-2u}^{3-u} \frac{(5-2u-v)^2}{2} dv = \right. \\ &= \frac{u^3 - 6u^2 + 6u + 5}{15 - 3u^2} \leq \frac{4u^3 - 15u^2 + 6u + 17}{15 - 3u^2} \end{aligned}$$

We obtain that

$$\delta_P(T, D) = \frac{15 - 3u^2}{u^3 - 6u^2 + 6u + 5} \text{ for } P \in L_{24} \setminus (E_2 \cup E_4) \text{ and } u \in [1, 2].$$

Step 4.1. Suppose $P \in L_{12} \cup L_{14}$ and $u \in [1, 3/2]$. In this case we set $\mathcal{C} = L_{14}$. Then $\tau(\mathcal{C}) = 3 - u$. Without loss of generality, we can assume that $P \in L_{14}$. The Zariski Decomposition of the divisor $D - vL_{14}$ is given by:

$$P(v) = \begin{cases} D - vL_{14} - \frac{v}{2}E_4 \text{ for } v \in [0, 2-u] \\ D - vL_{14} - \frac{v}{2}E_4 - (u+v-2)E_1 \text{ for } v \in [2-u, 1] \\ D - vL_{14} - \frac{v}{2}E_4 - (u+v-2)E_1 - (v-1)L_{23} \text{ for } v \in [1, 4-2u] \\ D - vL_{14} - (u+v-2)(E_1 + E_4) - (v-1)L_{23} - (2u+v-4)E_5 \text{ for } v \in [4-2u, 3-u] \end{cases}$$

and

$$N(v) = \begin{cases} \frac{v}{2}E_4 \text{ for } v \in [0, 2-u] \\ \frac{v}{2}E_4 + (u+v-2)E_1 \text{ for } v \in [2-u, 1] \\ \frac{v}{2}E_4 + (u+v-2)E_1 + (v-1)L_{23} \text{ for } v \in [1, 4-2u] \\ (u+v-2)(E_1 + E_4) + (v-1)L_{23} + (2u+v-4)E_5 \text{ for } v \in [4-2u, 3-u] \end{cases}$$

Moreover

$$P(v)^2 = \begin{cases} 5 - 2v - v^2/2 - u^2 \text{ for } v \in [0, 2-u] \\ 9 - 4u - 6v + v^2/2 + 2uv \text{ for } v \in [2-u, 1] \\ \frac{(v-2)(3v+4u-10)}{2} \text{ for } v \in [1, 4-2u] \\ 2(u+v-3)^2 \text{ for } v \in [4-2u, 3-u] \end{cases}$$

and

$$P(v) \cdot \mathcal{C} = \begin{cases} v/2 + 1 \text{ for } v \in [0, 2-u] \\ 3 - u - v/2 \text{ for } v \in [2-u, 1] \\ 4 - u - 3v/2 \text{ for } v \in [1, 4-2u] \\ 2(3 - u - v) \text{ for } v \in [4-2u, 3-u] \end{cases}$$

Thus,

$$\begin{aligned} S_D(\mathcal{C}) &= \frac{1}{5-u^2} \left(\int_0^{2-u} 5 - 2v - v^2/2 - u^2 dv + \int_{2-u}^1 9 - 4u - 6v + v^2/2 + 2uv dv + \right. \\ &\quad \left. + \int_1^{4-2u} \frac{(v-2)(3v+4u-10)}{2} dv + \int_{4-2u}^{3-u} 2(u+v-3)^2 dv \right) = \frac{3u^3 - 12u^2 + 6u + 13}{15 - 3u^2} \end{aligned}$$

Thus, $\delta_P(T, D) \leq \frac{15-3u^2}{3u^3-12u^2+6u+13}$ for $P \in L_{14}$. Note that we have:

- if $P \in L_{14} \setminus (E_4 \cup E_1 \cup L_{23} \cup E_5)$ then

$$h_D(v) = \begin{cases} \frac{(v/2+1)^2}{2} & \text{for } v \in [0, 2-u] \\ \frac{(3-u-v/2)^2}{2} & \text{for } v \in [2-u, 1] \\ \frac{(4-u-3v/2)^2}{2} & \text{for } v \in [1, 4-2u] \\ 2(3-u-v)^2 & \text{for } v \in [4-2u, 3-u] \end{cases}$$

- if $P = L_{14} \cap E_1$ then

$$h_D(v) = \begin{cases} \frac{(v/2+1)^2}{2} & \text{for } v \in [0, 2-u] \\ \frac{(6-2u-v)(2u+3v-2)}{8} & \text{for } v \in [2-u, 1] \\ \frac{(8-2u-3v)(2u+v)}{8} & \text{for } v \in [1, 4-2u] \\ (3-u-v) & \text{for } v \in [4-2u, 3-u] \end{cases}$$

- if $P = L_{14} \cap L_{23}$ then

$$h_D(v) = \begin{cases} \frac{(v/2+1)^2}{2} & \text{for } v \in [0, 2-u] \\ \frac{(3-u-v/2)^2}{2} & \text{for } v \in [2-u, 1] \\ \frac{(8-2u-3v)(4-2u+v)}{8} & \text{for } v \in [1, 4-2u] \\ 2(2-u)(3-u-v) & \text{for } v \in [4-2u, 3-u] \end{cases}$$

So we have

- if $P \in L_{14} \setminus (E_4 \cup E_1 \cup L_{23} \cup E_5)$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^C; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} \frac{(v/2+1)^2}{2} dv + \int_{2-u}^1 \frac{(3-u-v/2)^2}{2} dv + \right. \\ &\quad \left. + \int_1^{4-2u} \frac{(4-u-3v/2)^2}{2} dv + \int_{4-2u}^{3-u} 2(3-u-v)^2 dv \right) = \\ &= \frac{21-u^3-6u}{2(15-3u^2)} \leq \frac{3u^3-12u^2+6u+13}{15-3u^2} \end{aligned}$$

- if $P = L_{14} \cap E_1$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^C; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} \frac{(v/2+1)^2}{2} dv + \int_{2-u}^1 \frac{(6-2u-v)(2u+3v-2)}{8} dv + \right. \\ &\quad \left. + \int_1^{4-2u} \frac{(8-2u-3v)(2u+v)}{8} dv + \int_{4-2u}^{3-u} (3-u-v) dv \right) = \\ &= \frac{19-2u^3}{2(15-3u^2)} \end{aligned}$$

- if $P = L_{14} \cap L_{23}$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^C; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} \frac{(v/2+1)^2}{2} dv + \int_{2-u}^1 \frac{(3-u-v/2)^2}{2} dv + \right. \\ &\quad \left. + \int_1^{4-2u} \frac{(8-2u-3v)(4-2u+v)}{8} dv + \int_{4-2u}^{3-u} 2(2-u)(3-u-v) dv \right) = \\ &= \frac{26-12u^2+3u^3}{2(15-3u^2)} \leq \frac{3u^3-12u^2+6u+13}{15-3u^2} \end{aligned}$$

We obtain that

$$\delta_P(T, D) = \frac{15-3u^2}{3u^3-12u^2+6u+13} \text{ for } P \in L_{14} \setminus (E_4 \cup E_5 \cup E_1) \text{ and } u \in [1, 3/2]$$

and

$$\delta_P(T, D) \geq \begin{cases} \frac{15 - 3u^2}{3u^3 - 12u^2 + 6u + 13} & \text{for } P = L_{14} \cap E_1 \text{ and } u \in [1, b] \\ \frac{2(15 - 3u^2)}{19 - 2u^3} & \text{for } P = L_{14} \cap E_1 \text{ and } u \in [b, 3/2] \end{cases}$$

where b is a root of $8u^3 - 24u^2 + 12u + 7$ on $[1, 3/2]$. Note that $b \in [1.261, 1.262]$.

Step 4.2. Suppose $P \in L_{12} \cup L_{14}$ and $u \in [3/2, 2]$. In this case we set $\mathcal{C} = L_{14}$. Then $\tau(\mathcal{C}) = 3 - u$. Without loss of generality, we can assume that $P \in L_{14}$. The Zariski Decomposition of the divisor $D - vL_{14}$ is given by:

$$P(v) = \begin{cases} D - vL_{14} - \frac{v}{2}E_4 & \text{for } v \in [0, 2 - u] \\ D - vL_{14} - \frac{v}{2}E_4 - (u + v - 2)E_1 & \text{for } v \in [2 - u, 4 - 2u] \\ D - vL_{14} - (u + v - 2)(E_1 + E_4) - (2u + v - 4)E_5 & \text{for } v \in [4 - 2u, 1] \\ D - vL_{14} - (u + v - 2)(E_1 + E_4) - (v - 1)L_{23} - (2u + v - 4)E_5 & \text{for } v \in [1, 3 - u] \end{cases}$$

and

$$N(v) = \begin{cases} \frac{v}{2}E_4 & \text{for } v \in [0, 2 - u] \\ \frac{v}{2}E_4 + (u + v - 2)E_1 & \text{for } v \in [2 - u, 4 - 2u] \\ (u + v - 2)(E_1 + E_4) + (2u + v - 4)E_5 & \text{for } v \in [4 - 2u, 1] \\ (u + v - 2)(E_1 + E_4) + (v - 1)L_{23} + (2u + v - 4)E_5 & \text{for } v \in [1, 3 - u] \end{cases}$$

Moreover

$$P(v)^2 = \begin{cases} 5 - 2v - v^2/2 - u^2 & \text{for } v \in [0, 2 - u] \\ 9 - 4u - 6v + v^2/2 + 2uv & \text{for } v \in [2 - u, 4 - 2u] \\ 2u^2 + 4uv + v^2 - 12u - 10v + 17 & \text{for } v \in [4 - 2u, 1] \\ 2(u + v - 3)^2 & \text{for } v \in [1, 3 - u] \end{cases}$$

and

$$P(v) \cdot \mathcal{C} = \begin{cases} 1 + v/2 & \text{for } v \in [0, 2 - u] \\ 3 - u - v/2 & \text{for } v \in [2 - u, 4 - 2u] \\ 5 - 2u - v & \text{for } v \in [4 - 2u, 1] \\ 2(3 - u - v) & \text{for } v \in [1, 3 - u] \end{cases}$$

Thus,

$$S_D(\mathcal{C}) = \frac{1}{5 - u^2} \left(\int_0^{2-u} 5 - 2v - v^2/2 - u^2 dv + \int_{2-u}^{4-2u} 9 - 4u - 6v + v^2/2 + 2uv dv + \int_{4-2u}^1 2u^2 + 4uv + v^2 - 12u - 10v + 17 dv + \int_1^{3-u} 2(u + v - 3)^2 dv \right) = \frac{3u^3 - 12u^2 + 6u + 13}{15 - 3u^2}$$

Thus, $\delta_P(T, D) \leq \frac{15 - 3u^2}{3u^3 - 12u^2 + 6u + 13}$ for $P \in L_{14}$. Note that we have:

- if $P \in L_{14} \setminus (E_4 \cup E_1 \cup L_{23} \cup E_5)$ then

$$h_D(v) = \begin{cases} \frac{(1+v/2)^2}{2} & \text{for } v \in [0, 2 - u] \\ \frac{(3-u-v/2)^2}{2} & \text{for } v \in [2 - u, 4 - 2u] \\ \frac{(5-2u-v)^2}{2} & \text{for } v \in [4 - 2u, 1] \\ 2(3 - u - v)^2 & \text{for } v \in [1, 3 - u] \end{cases}$$

- if $P = L_{14} \cap E_1$ then

$$h_D(v) = \begin{cases} \frac{(1+v/2)^2}{2} & \text{for } v \in [0, 2-u] \\ \frac{(6-2u-v)(2u+3v-2)}{8} & \text{for } v \in [2-u, 4-2u] \\ \frac{(v+1)(5-2u-v)}{2} & \text{for } v \in [4-2u, 1] \\ 2(3-u-v) & \text{for } v \in [1, 3-u] \end{cases}$$

- if $P = L_{14} \cap L_{23}$ then

$$h_D(v) = \begin{cases} \frac{(1+v/2)^2}{2} & \text{for } v \in [0, 2-u] \\ \frac{(3-u-v/2)^2}{2} & \text{for } v \in [2-u, 4-2u] \\ \frac{(5-2u-v)^2}{2} & \text{for } v \in [4-2u, 1] \\ 2(2-u)(3-u-v) & \text{for } v \in [1, 3-u] \end{cases}$$

- if $P \in L_{14} \setminus (E_4 \cup E_1 \cup L_{23} \cup E_5)$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^C; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} \frac{(v/2+1)^2}{2} dv + \int_{2-u}^{4-2u} \frac{(3-u-v/2)^2}{2} dv + \right. \\ &\quad \left. + \int_{4-2u}^1 \frac{(5-2u-v)^2}{2} dv + \int_1^{3-u} 2(3-u-v)^2 dv \right) = \\ &= \frac{7u^3 - 36u^2 + 48u - 6}{2(15-3u^2)} \leq \frac{3u^3 - 12u^2 + 6u + 13}{15-3u^2} \end{aligned}$$

- if $P = L_{14} \cap E_1$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^C; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} \frac{(v/2+1)^2}{2} dv + \int_{2-u}^{4-2u} \frac{(6-2u-v)(2u+3v-2)}{8} dv + \right. \\ &\quad \left. + \int_{4-2u}^1 \frac{(v+1)(5-2u-v)}{2} dv + \int_1^{3-u} 2(3-u-v) dv \right) = \\ &= \frac{3u^3 - 18u^2 + 27u - 4}{15-3u^2} \end{aligned}$$

- if $P = L_{14} \cap L_{23}$ then

$$\begin{aligned} S_D(W_{\bullet,\bullet}^C; P) &= \frac{2}{5-u^2} \left(\int_0^{2-u} \frac{(v/2+1)^2}{2} dv + \int_{2-u}^{4-2u} \frac{(3-u-v/2)^2}{2} dv + \right. \\ &\quad \left. + \int_{4-2u}^1 \frac{(5-2u-v)^2}{2} dv + \int_1^{3-u} 2(2-u)(3-u-v) dv \right) = \\ &= \frac{3u^3 - 12u^2 + 26}{2(15-3u^2)} \leq \frac{3u^3 - 12u^2 + 6u + 13}{15-3u^2} \end{aligned}$$

We obtain that

$$\delta_P(T, D) = \frac{15-3u^2}{3u^3 - 12u^2 + 6u + 13} \text{ for } P \in L_{14} \setminus (E_1 \cup E_4 \cup E_5) \text{ and } u \in [3/2, 2]$$

and

$$\delta_P(T, D) \geq \frac{15-3u^2}{3u^3 - 18u^2 + 27u - 4} \text{ for } P = L_{14} \cap E_1 \text{ and } u \in [3/2, 2]$$

□

Corollary A.3. Let P be a point in T that is contained in $L_{12} \cup L_{14} \cup L_{24} \cup E_2 \cup E_3 \cup E_4 \cup E_4$ then

$$\delta_P(T, D) \geq \begin{cases} \frac{15 - 3u^2}{16 + 3u - 9u^2 + 2u^3} \text{ for } u \in [1, a], \\ \frac{15 - 3u^2}{11 - u^3} \text{ for } u \in [a, 2] \end{cases}$$

Corollary A.4. Suppose O is a point on a del Pezzo surface \bar{T} with two \mathbb{A}_1 singularities and nine lines such that $\delta_O(T) \leq \frac{6}{5}$ then

$$\delta_O(\bar{T}, \bar{D}) \geq \begin{cases} \frac{15 - 3u^2}{16 + 3u - 9u^2 + 2u^3} \text{ for } u \in [1, a], \\ \frac{15 - 3u^2}{11 - u^3} \text{ for } u \in [a, 2] \end{cases}$$


A.3. Polarized δ -invariant on Del Pezzo surface of degree 4 with \mathbb{A}_2 singularity. Now, let us use the notations and assumptions of Section 2 with a minor difference: we assume that \bar{T} has a singular point of type \mathbb{A}_2 . Let us show that in the case when O is the singular point of the surface \bar{T} we have

$$\delta_O(\bar{T}, \bar{D}) = \frac{u^3 - 6u^2 + 19}{15 - 3u^2}$$

which immediately implies that $\delta_O(\bar{T}, W_{\bullet, \bullet}^{\bar{T}}) \leq \frac{80}{83}$. In this case, the morphism η is a blow up of \mathbb{P}^2 at points P_1, P_2 , and P_3 in general position; after that blowing up a point P_4 which belongs to the exceptional divisor corresponding to P_3 and no other negative curve and after that a point P_5 which belongs to the exceptional divisor corresponding to P_4 and no other negative curve. By [7, Section 6.5] we have:

$$\delta_P(T) = \begin{cases} 6/7 \text{ if } P \in E_3 \cup E_4, \\ 8/7 \text{ if } P \in (L_{13} \cup L_{23} \cup L_{34} \cup E_5) \setminus (E_3 \cup E_4), \\ 4/3 \text{ if } P \in (L_{12} \cup C_2) \cap (E_1 \cup E_2), \\ 18/13 \text{ if } P \in (L_{12} \cup C_2 \cup E_1 \cup E_2) \setminus ((L_{12} \cup C_2) \cap (E_1 \cup E_2)), \\ 3/2, \text{ otherwise} \end{cases}$$

where E_1, E_2, E_3, E_4, E_5 are exceptional divisors corresponding to P_1, P_2, P_3, P_4, P_5 respectively, C_2 is a strict transform of a (-1) -curve coming from the conic on \mathbb{P}^2 , L_{ij} are strict transforms of the lines passing through P_i and P_j for $(i, j) \in \{(1, 2), (1, 3), (2, 3)\}$ and L_{34} is a strict transform of a (-1) -curve coming from a line passing through P_3 on \mathbb{P}^2 . The dual graph of (-1) and (-2) -curves is given in the following picture:



Now let's prove that:

$$\delta_P(T, D) = \frac{u^3 - 6u^2 + 19}{15 - 3u^2} \text{ for } P \in E_4 \setminus (L_{34} \cup E_5)$$

Proof. Suppose $P \in E_4 \setminus (L_{34} \cup E_5)$. In this case we set $\mathcal{C} = E_4$. Then $\tau(\mathcal{C}) = 2$. The Zariski Decomposition of the divisor $D - vE_4$ is given by:

$$P(v) = \begin{cases} -K_T - (u-1)C_2 - vE_4 - \frac{v}{2}E_3 & \text{for } v \in [0, 2-u] \\ -K_T - (u-1)C_2 - vE_4 - \frac{v}{2}E_3 - (u+v-2)E_5 & \text{for } v \in [2-u, 1] \\ -K_T - (u-1)C_2 - vE_4 - \frac{v}{2}E_3 - (u+v-2)E_5 - (v-1)L_{34} & \text{for } v \in [1, 2] \end{cases}$$

and

$$N(v) = \begin{cases} \frac{v}{2}E_3 & \text{for } v \in [0, 2-u] \\ \frac{v}{2}E_3 + (u+v-2)E_5 & \text{for } v \in [2-u, 1] \\ \frac{v}{2}E_3 + (u+v-2)E_5 + (v-1)L_{34} & \text{for } v \in [1, 2] \end{cases}$$

Moreover,

$$P(v)^2 = \begin{cases} 5 - u^2 - 3v^2/2 & \text{for } v \in [0, 2-u] \\ 9 - 4u - 4v + 2uv - 1/2v^2 & \text{for } v \in [2-u, 1] \\ \frac{(v-2)(v+4u-10)}{2} & \text{for } v \in [1, 2] \end{cases}$$

and

$$P(v) \cdot \mathcal{C} = \begin{cases} 3v/2 & \text{for } v \in [0, 2-u] \\ 2-u+v/2 & \text{for } v \in [2-u, 1] \\ 3-u-v/2 & \text{for } v \in [1, 2] \end{cases}$$

Thus,

$$S_D(\mathcal{C}) = \frac{1}{5-u^2} \left(\int_0^{2-u} 5 - u^2 - 3v^2/2 dv + \int_{2-u}^1 9 - 4u - 4v + 2uv - 1/2v^2 dv + \int_1^2 \frac{(v-2)(v+4u-10)}{2} dv \right) = \frac{19 + u^3 - 6u^2}{15 - 3u^2}$$

Thus, $\delta_P(T, D) \leq \frac{15-3u^2}{19+u^3-6u^2}$ for $P \in E_4$. Note that for $P \in E_4 \setminus (E_5 \cup L_{34})$ we have:

$$h_D(v) = \begin{cases} \frac{9v^2}{8} & \text{for } v \in [0, 2-u] \\ \frac{(2-u+v/2)^2}{2} & \text{for } v \in [2-u, 1] \\ \frac{(3-u-v/2)^2}{2} & \text{for } v \in [1, 2] \end{cases}$$

So we have

$$S_D(W_{\bullet, \bullet}^{\mathcal{C}}; P) = \frac{2}{5-u^2} \left(\int_0^{2-u} \frac{9v^2}{8} dv + \int_{2-u}^1 \frac{(2-u+v/2)^2}{2} dv + \int_1^2 \frac{(3-u-v/2)^2}{2} dv \right) = \frac{21 + 6u - 18u^2 + 5u^3}{2(15 - 3u^2)} \leq \frac{19 + u^3 - 6u^2}{15 - 3u^2}$$

So we obtain that

$$\delta_P(T, D) = \frac{u^3 - 6u^2 + 19}{15 - 3u^2} \text{ for } P \in E_4 \setminus (L_{34} \cup E_5).$$

□

REFERENCES

- [1] H. Abban, Z. Zhuang, *K-stability of Fano varieties via admissible flags*, Forum of Mathematics, Pi **10** (2022), Paper No. e15, 43 p.
- [2] C. Araujo, A.-M. Castravet, I. Cheltsov, K. Fujita, A.-S. Kaloghiros, J. Martinez-Garcia, C. Shramov, H. Süß, N. Viswanathan, *The Calabi problem for Fano threefolds*, LMS Lecture Notes in Mathematics **485**, Cambridge University Press, 2023.
- [3] Grigory Belousov, Konstantin Loginov *K-stability of Fano threefolds of rank 4 and degree 24*, preprint, arXiv:2206.12208, 2022
- [4] H. Blum, Y. Liu, C. Xu, *Openness of K-semistability for Fano varieties*, Duke Math. J. **171** (2022), 2753–2797.
- [5] Ivan Cheltsov, *K-stability of Fano 3-folds of Picard rank 3 and degree 22* preprint, arXiv:2401.02818 , 2024
- [6] I. Cheltsov, V. Przyjalkowski, C. Shramov, *Fano threefolds with infinite automorphism groups*, Izv. Math. **83** (2019), 860–907.
- [7] E. Denisova, *δ -invariant of Du Val del Pezzo surfaces of degree ≥ 4* , preprint, arXiv:2304.11412 (2023).
- [8] K. Fujita, *A valuative criterion for uniform K-stability of \mathbb{Q} -Fano varieties*, J. Reine Angew. Math. **751** (2019), 309–338.
- [9] K. Fujita, *On K-stability and the volume functions of \mathbb{Q} -Fano varieties*, Proc. Lond. Math. Soc. **113** (2016), 541–582.
- [10] C. Li, *K-semistability is equivariant volume minimization*, Duke Math. Jour. **166** (2017), 3147–3218.
- [11] C. Xu, *K-stability of Fano varieties: an algebro-geometric approach*, EMS Surveys in Mathematical Sciences, 8 (2021), 265–354.

Elena Denisova

University of Edinburgh, Edinburgh, Scotland
 s2223072@ed.ac.uk