



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

# Missed opportunity for alcohol use disorder screening and management in primary health care facilities in northern rural Tanzania

a cross-sectional survey

### Citation for published version:

Mushi, D, Moshiro, C, Hanlon, C, Francis, JM & Teferra, S 2022, 'Missed opportunity for alcohol use disorder screening and management in primary health care facilities in northern rural Tanzania: a cross-sectional survey', *Substance Abuse Treatment, Prevention, and Policy*, vol. 17, no. 1. <https://doi.org/10.1186/s13011-022-00479-x>

### Digital Object Identifier (DOI):

[10.1186/s13011-022-00479-x](https://doi.org/10.1186/s13011-022-00479-x)

### Link:

[Link to publication record in Edinburgh Research Explorer](#)

### Document Version:

Publisher's PDF, also known as Version of record

### Published In:

Substance Abuse Treatment, Prevention, and Policy

### General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



RESEARCH

Open Access



# Missed opportunity for alcohol use disorder screening and management in primary health care facilities in northern rural Tanzania: a cross-sectional survey

Dorothy Mushi<sup>1,2,3\*</sup>, Candida Moshiro<sup>4</sup>, Charlotte Hanlon<sup>1,2,5</sup>, Joel M. Francis<sup>4,6</sup> and Solomon Teferra<sup>2</sup>

## Abstract

**Objective:** The study aimed to identify the missed opportunity for detection and management of alcohol use disorder by primary health care workers.

**Design:** A cross-sectional survey

**Setting:** Outpatient services in the six governmental primary health care facilities in Moshi district council in Tanzania.

**Participants:** A total of 1604 adults were screened for alcohol use disorder (AUD) using the Alcohol Use Disorder Identification Test (AUDIT). Participants scoring 8 or above then provided details about their help-seeking behavior and barriers to seeking care. Participants' records were reviewed to assess the screening and management of AUD.

**Results:** In the last 12 months, 60.7% reported alcohol use, and heavy episodic drinking (HED) was reported by 37.3%. AUD (AUDIT  $\geq 8$ ) was present in 23.9%. Males were more likely to have HED (aPR = 1.43; 95% CI: 1.3 to 1.4) or AUD (aPR = 2.9; 95% CI 1.9 to 4.2). Both HED and AUD increased with age. Only one participant (0.3%) had documented AUD screening and management. Only 5% of participants screening positive for AUD had sought help. Reasons for not seeking care were thinking that the problem would get better by itself (55.0%), wanting to handle the problem alone (42.0%), or not being bothered by the problem (40.0%).

**Conclusion:** While reported alcohol use, HED, and AUD are common among patients presenting to primary health-care facilities in northern Tanzania, help-seeking behavior and detection are very low. Not screening for AUD in primary health care is a missed opportunity for early detection and management. There is an urgent need to develop interventions to increase the detection of AUD by health care providers, while also addressing help-seeking behavior and barriers to seeking care.

**Keywords:** Alcohol use disorder, Primary health care, Screening, Detection, Alcohol use disorder management, Help-seeking, Barriers to seeking care

## Introduction

Alcohol use is socially accepted in most cultures worldwide. Excessive alcohol use accounts for substantial adverse consequences for health, social wellbeing, and economic status [1, 2]. Approximately one in 20 of all disability-adjusted life-years (5.1% DALYs) and 5.3% of

\*Correspondence: [dorrymush@yahoo.com](mailto:dorrymush@yahoo.com)

<sup>3</sup> Centre for Innovative Drug Development and Therapeutics Trial for Africa (CDT-Africa) College of Health Science, Addis Ababa University, Addis Ababa, Ethiopia  
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

all deaths globally are attributed to the harmful use of alcohol [1]. The mortality resulting from alcohol use is higher than that caused by other non-communicable or communicable diseases such as tuberculosis, HIV/AIDS, and diabetes [1]. Alcohol use disorder is responsible for around half of all alcohol-related harm [3].

Alcohol Use Disorder is defined as a cluster of cognitive, behavioral, and physiological symptoms indicating that the individual continues using alcohol despite significant alcohol-related problems [4]. In 2016, an estimated 283 million people aged 15 years and above had an alcohol use disorder (AUD) (5.1% of all adults). The 12-month prevalence of AUD was 3.7% in Africa in 2016 [1]. In Tanzania, the estimated prevalence of AUD varies across populations, ranging from 5.7% to 28% [5–7]. Heavy episodic drinking is also of public health concern. HED is a pattern of drinking whereby a person consumed at least 60 g or more of pure alcohol on at least one occasion in the past 30 days. Consumption of 60 g of pure alcohol corresponds approximately to six standard alcoholic drinks [1]. In 2016, the worldwide prevalence of HED was 18.2% and was associated with the different types of non-communicable as well as infectious diseases [1].

Studies conducted in healthcare facilities in sub-Saharan Africa (SSA) have shown that approximately one in five people attending healthcare facilities met the criteria for AUD [8–12]. This suggests the possibility that persons with AUD may frequently access health care services [13], though not specifically for their AUD. There is also evidence that most persons with AUD also have co-occurring acute or chronic physical or mental health problems [14–17]. Even with such a substantial magnitude of AUD and the clear adverse consequences, AUD appears to be rarely detected by health care providers [18, 19], leading to a wide treatment gap [18, 20, 21]. This gap is particularly marked in low-income and lower-middle-income countries [17, 18, 21].

Integration of evidence-based interventions for AUD into existing health care systems is recommended to address the AUD treatment gap [22–24]. Studies demonstrate that such integrated services can minimize cost and maximize the effectiveness of managing AUD and co-occurring health problems [22, 25–28]. An emerging and robust evidence base indicates the feasibility of delivering AUD interventions in primary health care settings [18, 29, 30]. However, studies have identified bottlenecks to the integration process and delivery of AUD interventions in health care systems [17, 31, 32]. These bottlenecks include low help-seeking behavior for people with AUD [17, 24, 33, 34] for example, due to stigma attached to AUD and lack of awareness. Further barriers to care exist for those who seek

help [17, 24], categorized into individual, structural, and contextual factors [31, 32].

To improve detection, help-seeking, and barriers to care for AUD and consequently narrow the treatment gap for AUD, more evidence from diverse contexts is needed. The purpose of this study was to identify the missed opportunity for detection and management of AUD by primary health care workers in Tanzania. The findings will inform the adaptation and piloting of a model for the integration of AUD interventions at the primary health care delivery level [35].

## Methods

### Study design and settings

A facility-based, cross-sectional survey was conducted from September to November 2019 in the six government health centers in Moshi district council, Kilimanjaro region, Tanzania. Kilimanjaro region is located around 530 km from the main economic city of Dar es Salaam (the former capital city of Tanzania). According to Tanzania's census [36], the Moshi district council had a population of 539,586 (262,897 males and 276,689 females) in 2012. Two faith-based hospitals have been designated by the Tanzanian Ministry of Health as the district hospitals, and there are also eight health centers and 88 dispensaries. Primary health care (PHC) services in Tanzania are provided at the district level through the health centers and dispensaries. Dispensaries are the first contact point for basic health services in the community.

The study area was selected based on the existence of previous studies of alcohol use disorder in Tanzania. In these previous studies, alcohol use (68.0% to 70.0%) and problematic alcohol use (20.0% to 47.0%) were found to be highly prevalent in the community [6, 7]. Based on previous findings, it was recommended that access to interventions for AUD should be expanded [6, 7, 37, 38]. Therefore, this study was conducted to inform the adaptation of a model for integrating services for AUD in PHC in the study area [35].

### Participants and sample size

Adult outpatients (aged 18 years and above) attending PHC during the study period and willing to give consent were invited into the study. The sample size was determined based on the assumption that the prevalence of AUD would range from 10.0% to 20.0% in the PHC setting [6, 7]. We assumed a detection rate of AUD of 10.0% from a previous study conducted in five low and middle-income countries [19]. The possibility of clustering was considered at the health facility level, the following were the reasons, (i) health care providers working in one PHC are more likely to share similarities such as clinical knowledge, practices, health care delivery infrastructure,

and system. (ii) Likewise, for the people who attend the same PHC, because they are living in the same area, they are more likely to share similarities in terms of their health beliefs, practices, and socioeconomic status. We could not find the published studies from Tanzania to guide the estimation of the intra-cluster correlation (ICC) for the detection of AUD in health centers. Therefore, we assumed the ICC to be 0.01, from a review of 31 studies [39], with a design effect of 1.95, which resulted in a sample size of 1600. The sample size was distributed over six PHC facilities based on the size of the catchment population. To achieve a total sample in each PHC facility, all men were recruited and a systematic sample of one in three women was recruited. Men were oversampled to ensure an adequate sample size that aimed for equal representation of men and women, as more women than men attend PHC, with an approximate ratio of men to women of 1:3.

#### Data collection and variables

The collection of data through the face-to-face interview was conducted electronically, through a mobile phone or tablet using Open Data Kit (ODK) software. Data collection tools were piloted, pre-tested, and modified accordingly. This helped the research team plan for the number of interviews to be conducted by one data collector per day while minimizing interference to the provision of clinical services and extra time spent by the participant at the health facility. Interviews were conducted after the participant's clinic appointment. This facilitated participant

engagement and cooperation because the participant had received the treatment that brought them to the health facility. Furthermore, healthcare professionals were not biased by the research interview findings. The questionnaire was administered in Swahili by ten final-year medical students supervised by one study coordinator and the principal investigator.

#### Measures

Socio-demographic variables collected in the study included age, sex, marital status, level of education, and occupation.

#### Alcohol use and alcohol use disorder

The World Health Organization's Alcohol Use Disorder Identification Test (AUDIT) was used to assess alcohol consumption during the past 12 months [40]. The AUDIT assessed alcohol consumption in terms of quantity, drinking pattern, and problems associated with alcohol use. Each item on the AUDIT was rated on a five-point Likert scale from 0 to 4, giving a total score ranging from zero to 40. The first question, which asks how often

does someone have a drink containing alcohol, informed alcohol use during the past 12 months. Heavy episodic drinking was defined based on the third question which asks 'how often do you have six or more drinks on one occasion?' A total score of eight or more indicates a tendency to problematic drinking as follows: a score of 8–15 indicates hazardous alcohol use, a score of 16–19 indicates harmful alcohol use, and an AUDIT score of 20 and above requires further evaluation for suspected alcohol dependence [41]. In this study, for participants who had an AUDIT score of  $\geq 8$ , data on help-seeking, and barriers to seeking care for AUD were collected. In addition, a clinical consultation checklist form that contained patient particulars, a description of the patient's symptoms/ complaint, diagnosis, and treatment plan was completed to assess whether the PHC provider had documented the AUD diagnosis and if the patient received any AUD intervention. The estimated mean consultation time in PHC facilities is 10 min [42]

The AUDIT has been validated across different populations in low and middle-income countries' settings [40, 43, 44]. The AUDIT has been adapted into the national language (Swahili) and used previously in Tanzania [5–7, 45]. The Swahili version of the AUDIT has been validated and the psychometric properties found to be acceptable [45]. For traditional brews, we applied the ethanol concentration data from the previous assessment of the locally available brews including the traditional liquor (spirit) and beers [46]. A pictorial chart demonstrating the standard drinks of the most commonly consumed traditional and industrial alcoholic beverages has been adapted [7] and was used in this study.

#### Barriers to seeking care

Barriers to accessing care based on various factors, including cost, service satisfaction, logistics, and stigma, were assessed using The Barriers to Accessing Care (BACE) questionnaire [47]. The BACE measures several barriers that may prevent individuals from seeking or continuing with care for a mental health problem. The barriers include individual and structural factors, such as individual knowledge and attitude about mental health care, social support, stigma, infrastructure, previous experience with mental health care, and treatment cost. The BACE has been adapted previously for use in low-income African countries [17, 48, 49].

#### Help-seeking behavior

Awareness of having an AUD and the individual's perspective that AUD is a problem that needs care was assessed using an adapted version of the PRIME help-seeking behavior questionnaire [50]. The questionnaire was administered to participants with an AUDIT score

of  $\geq 8$ . First, participants were asked if they had ever sought help for their alcohol use (Yes or No). Those who had sought help were then asked about the time, type, and perceived usefulness of the help they received.

#### Screening and management of AUD at PHC

Any documentation of detection and/or other interventions for problematic alcohol use (not just that the person drinks alcohol) in the clinical consultation form was considered as evidence for the screening and management of AUD at PHC.

#### Data management and analysis

Sociodemographic characteristics were summarized using proportions for categorical variables or median, and interquartile ranges for continuous variables. Screening and management of AUD were calculated as the proportion of participants with a probable AUD for whom there was documentation indicating detection and/or other interventions for problematic alcohol use. Odds ratios (OR) from a cross-sectional study might overestimate the effect when the outcome is common [51, 52], so prevalence ratios were obtained using a Poisson working model [53, 54]. A multivariable model was used to examine factors associated with alcohol use, HED, and AUD. All exposures (sex, age, marital status, employment, and education level) are known confounders [55, 56] and therefore were included in the multivariable models [57]. Crude and adjusted prevalence ratios with corresponding 95% confidence intervals and p-values were reported. We used weights to account for unequal selection probabilities, and adjusted for clustering and stratification (STATA *svy* commands) while calculating the prevalence of HED and AUD, and fitting regression models. [58]. A two-sided  $P < 0.05$  was considered statistically significant. Data were analyzed using STATA version 13 [59].

#### Ethics approval and consent to participant

This study was approved by the Addis Ababa University College of Health Science Institutional Review Board (protocol number: 023/19/psyc) and the Muhimbili University of Health and Allied Sciences Institutional Review Board (Ref.No.DA.282/298/01.C/). All participants gave written informed consent; if non-literate, participants gave a thumbprint to signify consent in presence of a data collector and another witness.

## Results

#### Sociodemographic characteristics

We recruited 1615 participants, 11 refused to participate in the study, and lack of time due to other responsibilities was the reason reported by almost all of the participants. Of 1604 individuals, the median age of study participants

was 41 years (interquartile range 29–56). Over half, 58.2% ( $n=933$ ) of participants were female. About two-thirds of participants had completed primary education, 65.0% ( $n=607$ ) of females and 63.0% ( $n=425$ ) of males. 63.7% ( $n=595$ ) of females and 70.5% ( $n=473$ ) of males were currently married. Details of sociodemographic characteristics are presented in Table 1.

#### Alcohol use and prevalence of AUD

Overall, 60.7% of the participants reported to use alcohol in the last 12 months ( $n=974$ ) (95% CI: 59.1 to 63.8%). Using the last 12 months timeframe, the weighted prevalence of HED was 37.3% ( $n=599$ ), and AUD (AUDIT  $\geq 8$ ) was 23.9% ( $n=378$ ), (95% CI: 18.6 to 30.2%). The prevalence of AUD was 3 times higher among males 38.7% ( $n=260$ ) than females 13.1% ( $n=118$ ),  $p$ -value  $< 0.001$ . Based on the AUDIT score among those who reported alcohol use in the last 12 months, 18.0% ( $n=281$ ) were classified as hazardous alcohol use (AUDIT score 8–15), 3.0% ( $n=42$ ) harmful alcohol use (AUDIT score 16–19) and 3.0% ( $n=55$ ) probable alcohol dependence (AUDIT score  $\geq 20$ ).

#### Factors associated with alcohol use and AUD

In the final multivariable analysis, reported alcohol use was independently associated with the male gender (aPR 1.42, 95% CI: 1.3 to 1.5) and older age (Table 2). Moreover, males were more likely to have HED (aPR = 1.43; 95% CI: 1.3 to 1.4) (Table 3) or AUD (aPR = 2.9; 95% CI 1.9 to 4.2) (Table 4). Both HED and AUD increased with age, participants aged 25 years and above had more prevalence of HED (Table 3) and AUD (Table 4) when compared to participants aged less than 25 years.

#### Screening and management of AUD

Out of 378 participants with an AUDIT score  $\geq 8$ , screening and management of AUD were recorded for only one participant.

#### Help-seeking behavior and barriers to seeking care among participants with AUD (AUDIT $\geq 8$ )

Only 5% (20/378) had reported seeking help from informal and non-informal sources for their alcohol problem at least once in their lifetime, clan leaders/heads ( $n=7$ ), friends ( $n=3$ ), medical providers ( $n=3$ ), religious or spiritual leaders ( $n=6$ ), and a traditional healer ( $n=1$ ). About 148 out of 378 participants (39.1%) reported at least one barrier to seeking care for their alcohol problem. The leading reported barriers were as follows: thought that the problem would get better by itself (55.0%), wanting to handle the problem alone (42.0%), and not being bothered by the problem (40.0%). However, being not satisfied with available services (9.0%), not thinking it

**Table 1** Socio-demographic characteristics of PHC attendees by sex, in Moshi district council, Tanzania (2019)

Characteristic	Categories	Total N (%) n = 1604	Sex	
			Female N (%) n = 933	Male N (%) n = 671
Age in years	< 25	225 (14.0)	157 (16.8)	68 (10.1)
	25–39	526 (32.8)	333 (35.7)	179 (28.8)
	40–49	292 (18.2)	163 (17.5)	129 (19.2)
	50–59	229 (14.3)	120 (12.9)	109 (16.2)
	≥ 60	332 (20.7)	160 (17.1)	172 (25.6)
Education level	Never went to School	84 (5.2)	44 (4.7)	40 (5.9)
	Incomplete primary school	100(6.2)	56 (6.0)	44 (6.6)
	Primary education	1032 (64.3)	607 (65.0)	425 (63.3)
	Secondary education	283 (17.6)	174 (18.7)	109 (16.2)
	University/ college	105 (6.6)	52 (5.6)	53 (7.9)
Marital status	Married	1068(66.5)	595 (63.7)	473 (70.5)
	Separated/divorced	78(4.8)	51 (5.5)	27 (4.0)
	Widowed	130(8.1)	85 (9.1)	45 (6.7)
	Cohabiting	54(3.4)	38 (4.1)	16 (2.4)
	Single	274(17.1)	164 (17.6)	110 (16.4)
Employment	Employed	183 (11.4)	84 (9.0)	99 (14.8)
	Self-employment	933 (58.2)	544 (58.8)	384 (57.2)
	Unemployed/	362 (22.6)	238 (25.5)	124 (18.5)
	Volunteered/ Student	59 (3.7)	39 (4.2)	20 (3.0)
	Retired	67 (4.2)	23 (2.5)	44 (6.6)

**Table 2** Factors associated with alcohol use among PHC attendees in Moshi district council, Tanzania (2019)

Factor	Categories	Total	Reported AU n (%)	Crude PR (95% CI)	Adjusted PR (95% CI)
Age in years	< 25	86	33 (38.2)	1	1
	25–39	302	173 (57.4)	1.52(1.3–1.8) **	1.43(1.2–1.7) **
	40–49	196	131(67.1)	1.76(1.5–2.1) **	1.58(1.3–1.9) **
	50–59	165	119 (72.0)	1.91(1.6–2.3) **	1.70(1.4–2.1) **
	≥ 60	225	152 (67.7)	1.78(1.5–2.1) **	1.54(1.3–1.9) **
Education level	Never went to School	49	28(58.3)	0.91(0.7–1.1)	0.87(0.7–1.0)
	Incomplete primary	66	43(66.0)	1.03(0.9–1.2)	0.99(0.9–1.2)
	Primary	648	406(62.7)	0.96(0.8–1.1)	
	Secondary	148	77(52.3)	0.83(0.7–0.9) *	0.89(0.8–1.0)
	University/ college	63	38 (60.0)	0.96(0.8–1.1)	0.96(0.8–1.1)
Sex	Female	507	383(75.5)	1	1
	Male	467	233 (50.0)	1.48(1.4–1.6) **	1.42(1.3–1.5) **
Marital status	Married	681	434 (63.7)	1	1
	Separated/divorced	48	30 (61.5)	0.96(0.8–1.2)	1.00(0.8–1.2)
	Widowed	86	57 (66.1)	1.02(0.9–1.2)	1.01(0.9–1.2)
	Cohabiting	28	14(51.8)	0.81(0.6–1.1)	0.98(0.8–1.2)
	Single	131	63 (47.8)	0.76(0.7–0.9) **	0.90(0.8–1.0)
Employment	Unemployed	296	189 (60.7)		1
	Self-employed	131	94 (71.5)	1.15(1.0–1.3) *	1.12(1.0–1.3)
	Employed	547	320 (58.6)	0.95(0.9–1.0)	0.89(0.8–1.0) *

AU = Alcohol use, PR prevalence ratio, (%) Row percentage

\*  $p < 0.05$ \*\*  $p < 0.001$

**Table 3** Factors associated with heavy episodic drinking among PHC attendees in Moshi district council, Tanzania (2019)

Factor	Categories	Total	Reported HED n (%)	Crude PR (95% CI)	Adjusted PR (95% CI)
Age in years	< 25	47	25(54.6)	1	1
	25–39	208	143(68.8)	1.29(1.0–1.6) *	1.42(1.2–1.7) **
	40–49	124	78 (63.2)	1.17(0.9–1.5)	1.58(1.3–1.9) **
	50–59	102	63 (61.8)	1.16(0.9–1.5)	1.70(1.4–2.1) **
	≥ 60	118	62 (52.4)	0.97(0.8–1.2)	1.54(1.3–1.9) **
Education level	Never went to School	30	18 (61.2)	1.03(0.8–1.3)	1.07(0.9–1.3)
	Incomplete primary	37	21 (56.0)	0.92(0.7–1.2)	0.99(0.8–1.2)
	Primary	391	236 (60.3)	1	1
	Secondary	96	62 (64.8)	1.09(1.0–1.2)	1.08(0.9–1.2)
	University/ college	45	32 (71.4)	1.17(1.0–1.4)	1.14(0.9–1.4)
Sex	Female			1	1
	Male	361	257(71.2)	1.38(1.2–1.5) **	1.43(1.3–1.5) **
Marital status	Married	424	264(62.2)	1	1
	Separated/divorced	33	23(68.7)	1.11(0.9–1.4)	1.00(0.8–1.2)
	Widowed	36	15 (41.8)	0.68(0.5–0.9)	1.01(0.9–1.2)
	Cohabiting	20	14 (71.4)	1.14(0.9–1.5)	0.98(0.8–1.2)
	Single	86	56 (65.6)	1.06(0.9–1.2)	0.90(0.8–1.0)
Employment	Unemployed	143	69(48.3)		1
	Self-employed	89	60(67.9)	1.39(1.2–1.7) **	1.12(1.0–1.3)
	Employed	367	246 (67.0)	1.38(1.2–1.6) **	0.89(0.8–1.0) *

PR prevalence ratio, (%) Row percentage, and HED heavy episodic drinking

\*  $p < 0.05$

\*\*  $p < 0.001$

was a treatable problem (9.0%), and being concerned that family members would not approve (7.0%) were the least frequently reported barriers (Fig. 1).

## Discussion

This study identified a missed opportunity for detection and management of alcohol use disorder by primary health care workers, with a detection rate of 0.3%. Our study found that reported alcohol use, HED, and AUD are common among people in need of health care services in PHC facilities, although reported help-seeking behavior was low. Therefore, it is a missed opportunity to not screen alcohol use and AUD routinely among patients attending PHC.

Reported alcohol use replicates the findings from previous studies conducted in community settings in Tanzania [6, 7] and from a systematic review and meta-analysis of alcohol use in Eastern Africa [55]. However, it was higher compared to the study conducted in health care settings in Ghana [10]. The reasons could be due to the use of different screening methods for the identification of alcohol use and duration of alcohol use. Our study used the AUDIT which is self-reported and assessed the use of alcohol during the past 12 months while the previous study used a breathalyzer or saliva alcohol test strip which could detect alcohol in the body within 24 h of intake. The

prevalence of HED in this study was similar to another study conducted in a clinical setting in four African countries [60]. In comparison to previous studies conducted in SSA health care settings, the prevalence of AUD in this study was akin to the study in South Africa [12] but lower than in other studies in the Eastern and Southern parts of SSA [8, 11, 61, 62]. A possible reason for the lower prevalence of AUD in the present study could be because this study was conducted in the outpatient clinics while the previous studies were conducted in HIV clinics among people living with HIV. The systematic review and meta-analysis study on the prevalence.

of AUD among people living with HIV found out that people living with HIV have a higher prevalence of AUD than the general population [56].

Alcohol use was associated with male gender and older age in line with other studies in Eastern Africa [49, 63]. Likewise, for HED [64]. In this study, the male gender was associated with an increased prevalence of AUD. The findings align with various studies in Tanzania [5, 7, 38] and in other parts of the world [17, 24, 65]. The findings are reflected in the World Health Organization's world mental health survey cross-national epidemiology study on AUD [65] which reported that the prevalence of AUD is much higher for men than

**Table 4** Factors associated with alcohol use disorder among PHC attendees in Moshi district council, Tanzania (2019)

Factor	Categories	Total	Reported AUD n (%)	Crude PR (95% CI)	Adjusted PR (95% CI)
Age in years	< 25	225	26 (11.6)	1	1
	25–39	526	135 (25.7)	2.4 (1.3–4.6) *	(1.1–4.0) *
	40–49	292	80 (27.4)	2.5 (1.4–4.4) *	(1.1–3.9) *
	50–59	229	79 (34.5)	3.2 (1.7–5.8) **	2.6 (1.4–4.9) *
	≥ 60	332	58 (17.5)	1.6 (0.8–3.2)	1.4 (0.6–3.1)
Education level	Never went to School	84	19 (22.6)	0.9 (0.5–1.7)	0.9 (0.6–1.3)
	Incomplete primary	100	18 (18.0)	0.7 (0.5–0.9) *	0.8 (0.6–1.1)
	Primary	1032	254 (4.6)	1	1
	Secondary	283	63 (22.3)	0.9 (0.7–1.3)	0.9 (0.7–1.4)
	University/ college	105	24 (22.9)	(0.6–1.5)	0.8 (0.6–1.2)
Sex	Female	933	118 (12.7)	1	1
	Male	671	260 (38.8)	2.9 (1.9–4.4) **	2.9 (1.9–4.2) **
Marital status	Married	1068	271 (25.4)	1	1
	Separated/divorced	78	23 (29.5)	1.2 (0.9–1.4)	1.2 (1.0–1.5) *
	Widowed	130	20 (15.4)	0.6 (0.4–0.9) *	0.8 (0.6–1.2)
	Cohabiting	54	9 (16.7)	0.7 (0.4–1.2)	0.9 (0.5–1.5)
	Single	274	55 (20.1)	0.8 (0.6–1.0)	0.9 (0.8–1.2)
Employment	Unemployed	488	94 (19.3)	1	1
	Self-employed	933	220 (23.6)	1.2 (0.7–1.9)	0.9 (0.8–1.2)
	Employed	183	64 (35.0)	1.7 (1.1–2.8) *	1.3 (0.9–1.6)

AUD Alcohol use disorder (AUDIT ≥ 8), PR prevalence ratio, (%) Row percentage

\*  $p < 0.05$

\*\*  $p < 0.001$

women cross-nationally. In keeping with other studies in health care facilities and community settings in SSA [17, 24, 49, 66, 67], an increase in age was positively and significantly associated with AUD.

In this study, almost no participants who screened positive for AUD were either offered an alcohol screening or any form of alcohol intervention by the health care providers. The possible reason could be that Tanzania's guidelines have not strictly indicated a routine screening and management of AUD [68]. Moreover, this is in line with large treatment gaps for mental health disorders (including AUD) in low and middle-income countries (LMIC) [17–19, 24, 69]. In addition to a lack of alcohol screening and brief interventions, reported help-seeking behavior for AUD services was low. Other studies in LMICs [17, 18, 24, 69], also reported low help-seeking behaviors among people attending health care facilities as well as in the community. But, the reported help-seeking behavior for the formal and informal care in this study is relatively low when compared to other studies conducted in high-income countries and that is possibly influenced by differences in socio-economic and cultural factors between high-income and low-income countries [70–72].

Several barriers to seeking care were reported by the participants in this study. The main barriers to care that were


reported included, thinking that problem would get better by itself, wanting to handle the problem alone, and not being bothered by the problem. These were similar to other studies conducted in Eastern Africa [17, 24]. In comparison to another study [17], the current study did not find other reported main barriers to care such as feeling unsure about where to go and being concerned about the cost of receiving professional help. A reason for that may be due to differences in the study setting. The previous study was conducted in the community while this study was conducted in health-care facilities such that participants were already in the healthcare facilities and receiving professional help.

This study's findings should be interpreted in light of some limitations. This study is based on self-reported information that may be predisposed to social desirability and recall bias. Therefore, participants could lead to underreporting or overreporting alcohol use, help-seeking behavior, and barriers to seeking care for AUD.

#### Implications of the study

The findings call for an urgent need to integrate interventions for alcohol use and AUD services. The call is in keeping with the World Health Organization (WHO) Mental Health Gap Action Program (MhGap) for mental neurological and substance abuse (MNS) [73]. The WHO





MhGap guideline has emphasized integrating evidence-based interventions for AUD (including screening) in the general health care services to address the AUD treatment gap. For this to be possible the integrated interventions should equip PHC workers to detect and intervene on AUD. In addition to that, these interventions have to include strategies that will improve help-seeking behavior and address barriers to care for AUD services.

## Conclusions

While reported alcohol use, HED, and AUD are common among patients presenting to primary healthcare facilities in northern Tanzania, help-seeking behavior and detection are very low. Not screening for AUD in primary health care is a missed opportunity for early detection and management. There is an urgent need to develop interventions to increase the detection of AUD by health care providers, while also addressing help-seeking behavior and barriers to seeking care.

## Acknowledgements

We gratefully acknowledge the support received from the administrative offices in the Kilimanjaro region, Moshi district council, and Rombo district. Our appreciation should also go to the in-charges and staff from the respective primary health care facilities in the Moshi district council. We also extend our thanks to data collectors as well as to study participants.

## Authors' contributions

DM, CM, CH, JF, and ST contributed to the design of the study. DM analyzed the data with support from CM. DM wrote the first draft. All authors contributed to the interpretation of the findings and reviewed the full draft of the paper. All authors approved the final manuscript.

## Funding

DM acknowledges support from the Centre for Innovative Drug Development and Therapeutics Trial for Africa (CDT-Africa), a World Bank Africa Centre of excellence at Addis Ababa University as part of her training fellowship. The opinions in this paper are those of the authors and do not reflect the views of CDT-Africa. CH is funded through the ASSET research programme, supported

by the UK's National Institute of Health Research (NIHR) (NIHR Global Health Research Unit on Health Systems Strengthening in Sub-Saharan Africa at King's College London (16/136/54)) using UK aid from the UK Government. CH also receives support from NIHR through grant NIHR200842. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research, or the Department of Health and Social Care, England. CH also receives support from the Wellcome Trust (Grant 222154/Z20/Z).

## Availability of data and materials

All data used to write this paper is summarized in tables, graphs, or within text in the paper.

## Declarations

### Ethics approval and consent participate

This study received ethical approval from the Addis Ababa University College of Health Science Institutional Review Board and Muhimbili University of Health and Allied Sciences.

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

## Author details

<sup>1</sup>Department of Psychiatry and Mental Health, Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania. <sup>2</sup>Department of Psychiatry, School of Medicine, College of Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia. <sup>3</sup>Centre for Innovative Drug Development and Therapeutics Trial for Africa (CDT-Africa) College of Health Science, Addis Ababa University, Addis Ababa, Ethiopia. <sup>4</sup>Department of Epidemiology and Biostatistics Muhimbili, University of Health and Allied Sciences, Dar es Salaam, Tanzania. <sup>5</sup>Centre for Global Mental Health, Health Service and Population Research Department, Institute of Psychiatry, Psychology, and Neuroscience, King's College London, London, UK. <sup>6</sup>Department of Family Medicine and Primary Care, Witwatersrand University, Faculty of Health Sciences, Johannesburg, South Africa.

Accepted: 13 June 2022

Published online: 06 July 2022

## References

- World Health Organization. Global Status Report on Alcohol and Health. World Health Organization; 2019.
- Rehm J, Mathers C, Popova S, Thavorncharoensap M, Teerawattananon Y, Patra J. Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. *The Lancet*. 2009;373(9682):2223–33. [https://doi.org/10.1016/S0140-6736\(09\)60746-7](https://doi.org/10.1016/S0140-6736(09)60746-7) (PMID: 19560604).
- Rehm J, Gmel GE Sr, Gmel G, Hasan OSM, Imtiaz S, Popova S, et al. The relationship between different dimensions of alcohol use and the burden of disease—An update. *Addiction*. 2017;112(6):968–1001. <https://doi.org/10.1111/add.13757> (PMID: 28220587).
- Association AP. Diagnostic and statistical manual of mental disorders (DSM-5®): American psychiatric pub; 2013.
- Mbatia J, Jenkins R, Singleton N, White B. Prevalence of Alcohol Consumption and Hazardous Drinking, Tobacco and Drug Use in Urban Tanzania, and Their Associated Risk Factors. *IJERPH*. 2009;6(7):1991–2006. <https://doi.org/10.3390/ijerph6071991> (PMID: 19742167).
- Mongi AS, Baisley K, Ao TTH, Chilongani J, Aguirre-Andreasen A, Francis SC, et al. Factors associated with problem drinking among women employed in food and recreational facilities in Northern Tanzania. *PLoS ONE*. 2013;8(12). DOI: <https://doi.org/10.1371/journal.pone.0084447>. PMID: 24391956.
- Francis JM, Weiss HA, Mshana G, Baisley K, Grosskurth H, Kapiga SH. The Epidemiology of Alcohol Use and Alcohol Use Disorders among Young People in Northern Tanzania. *PLoS ONE*. 2015;10(10): e0140041. <https://doi.org/10.1371/journal.pone.0140041> (PMID: 26444441).
- Pary CD, Kekwaletswe C, Shuper PA, Nkosi S, Myers BJ, Morojele NK. Heavy alcohol use in patients on highly active antiretroviral therapy: What responses are needed? *SAMJ: South African Medical Journal*. 2016 Jun;106(6):567–8. DOI: <https://doi.org/10.7196/SAMJ>. 2016.v106i6.10639. PMID: 27245717
- Hahn JA, Emenyonu NI, Fatch R, Muyindike WR, Kekiibina A, Carrico AW, et al. Declining and rebounding unhealthy alcohol consumption during the first year of HIV care in rural Uganda, using phosphatidylethanol to augment self-report. *Addiction*. 2016;111(2):272–9. <https://doi.org/10.1111/add.13173> (PMID: 26381193).
- Forson PK, Gardner A, Odoro G, Bonney J, Biney EA, Oppong C, et al. Frequency of alcohol use among injured adult patients presenting to a Ghanaian emergency department. *Ann Emerg Med*. 2016;68(4):492–500.e6. <https://doi.org/10.1016/j.annemergmed.2016.04.033> (PMID: 27241887).
- Kader R, Seedat S, Govender R, Koch JR, Parry CD. Hazardous and Harmful use of Alcohol and/or Other Drugs and Health Status Among South African Patients Attending HIV Clinics. *AIDS Behav*. 2014;18(3):525–34. <https://doi.org/10.1007/s10461-013-0587-9> (PMID: 23921585).
- Peltzer K, Matseke G, Azwihangwisi M, Babor T. Evaluation of Alcohol Screening and Brief Intervention in Routine Practice of Primary Care Nurses in Vhembe District. *South Africa Croat Med J*. 2008;49(3):392–401. <https://doi.org/10.3325/cmj.2008.3.392> (PMID: 18581618).
- Luitel NP, Baron EC, Kohrt BA, Komproue IH, Jordans MJD. Prevalence and correlates of depression and alcohol use disorder among adults attending primary health care services in Nepal: a cross-sectional study. *BMC Health Serv Res*. 2018;18(1):215. <https://doi.org/10.1186/s12913-018-3034-9> (PMID: 29587724).
- Pan C, Ju TR, Lee CC, Chen Y-P, Hsu C-Y, Hung D-Z, et al. Alcohol use disorder tied to the development of chronic kidney disease: A nationwide database analysis. *PLoS ONE*. 2018;13(9): e0203410. <https://doi.org/10.1371/journal.pone.0203410> (PMID: 30188943).
- Udo T, Vásquez E, Shaw BA. A lifetime history of alcohol use disorder increases the risk for chronic medical conditions after stable remission. *Drug Alcohol Depend*. 2015;157:68–74. <https://doi.org/10.1016/j.drugalcdep.2015.10.008> (PMID: 26482092).
- Shield KD, Parry C, Rehm J. Chronic Diseases and Conditions Related to Alcohol Use. *Alcohol Res*. 2014;35(2):155–71 (PMID: 24881324).
- Zewdu S, Hanlon C, Fekadu A, Medhin G, Teferra S. Treatment gap, help-seeking, stigma and magnitude of alcohol use disorder in rural Ethiopia. *Substance Abuse Treatment, Prevention, and Policy*. 2019;14(1):4. <https://doi.org/10.1186/s13011-019-0192-7> (PMID: 30658659).
- Luitel NP, Jordans MJD, Kohrt BA, Rathod SD, Komproue IH. Treatment gap and barriers for mental health care: A cross-sectional community survey in Nepal. *PLoS ONE*. 2017;12(8): e0183223. <https://doi.org/10.1371/journal.pone.0183223> (PMID: 28817734).
- Rathod SD, Roberts T, Medhin G, Murhar V, Samudre S, Luitel NP, et al. Detection and treatment initiation for depression and alcohol use disorders: facility-based cross-sectional studies in five low-income and middle-income country districts. *BMJ Open*. 2018;8(10): e023421. <https://doi.org/10.1136/BMJopen-2018-023421>.
- Kohn R, Saxena S, Levav I, Saraceno B. The treatment gap in mental health care. *Bull World Health Organ*. 2004;82(11):858–66.
- Rathod SD, Silva MJD, Ssebunnya J, Breuer E, Murhar V, Luitel NP, et al. Treatment Contact Coverage for Probable Depressive and Probable Alcohol Use Disorders in Four Low- and Middle-Income Country Districts: The PRIME Cross-Sectional Community Surveys. *PLoS ONE*. 2016;11(9): e0162038 (PMID: 30309992).
- Funk M, Saraceno B, Drew N, Faydi E. Integrating mental health into primary healthcare. *Ment Health Fam Med*. 2008;5(1):5–8 (PMID: 22477840).
- Ferreira-Borges C, Parry CDH, Babor TF. Harmful Use of Alcohol: A Shadow over Sub-Saharan Africa in Need of Workable Solutions. *Int J Environ Res Public Health*. 2017 Apr;14(4). DOI: <https://doi.org/10.3390/ijerph14040346>. PMID: 28346373.
- Nalwadda O, Rathod SD, Nakku J, Lund C, Prince M, Kigozi F. Alcohol use in a rural district in Uganda: findings from community-based and facility-based cross-sectional studies. *Int J Ment Heal Syst*. 2018;12(1):12. <https://doi.org/10.1186/s13033-018-0191-5> (PMID: 29632551).
- Benegal V, Chand PK, Obot IS. Packages of Care for Alcohol Use Disorders in Low- And Middle-Income Countries. *PLoS Med*. 2009;6(10): e1000170. <https://doi.org/10.1371/journal.pmed.1000170> (PMID: 19859536).
- Ward C, Mertens J, Bresick G, Little F, Weisner C. Screening and Brief Intervention for Substance Misuse: Does It Reduce Aggression and HIV-Related Risk Behaviours? *Alcohol Alcohol*. 2015;1:50. <https://doi.org/10.1093/alcac/agv007> (PMID: 25731180).
- O'Donnell A, Anderson P, Newbury-Birch D, Schulte B, Schmidt C, Reimer J, et al. The Impact of Brief Alcohol Interventions in Primary Healthcare: A Systematic Review of Reviews. *Alcohol Alcohol*. 2014;49(1):66–78. <https://doi.org/10.1093/alcac/agt170> (PMID: 24232177).
- Babor TF, Del Boca F, Bray JW. Screening, Brief Intervention and Referral to Treatment: implications of SAMHSA's SBIRT initiative for substance abuse policy and practice. *Addiction*. 2017;112(Suppl 2):110–7. <https://doi.org/10.1111/add.13675> (PMID: 28074569).
- Room R, Babor T, Rehm J. Alcohol and public health. 2005;365:12. [https://doi.org/10.1016/S0140-6736\(05\)17870-2](https://doi.org/10.1016/S0140-6736(05)17870-2) (PMID: 15705462).
- Hargraves D, White C, Frederick R, Cinibulk M, Peters M, Young A, et al. Implementing SBIRT (Screening, Brief Intervention, and Referral to Treatment) in primary care: lessons learned from a multi-practice evaluation portfolio. *Public Health Rev*. 2017;29:38. <https://doi.org/10.1186/s40985-017-0077-0> (PMID: 29450101).
- Johnson M, Jackson R, Guillaume L, Meier P, Goyder E. Barriers and facilitators to implementing screening and brief intervention for alcohol misuse: a systematic review of qualitative evidence. *J Public Health (Oxf)*. 2011;33(3):412–21. <https://doi.org/10.1093/PubMed/fdq095> (PMID: 21169370).
- Rahm AK, Boggs JM, Martin C, Price DW, Beck A, Backer TE, et al. Facilitators and Barriers to Implementing Screening, Brief Intervention, and Referral to Treatment (SBIRT) in Primary Care in Integrated Health Care Settings. *Subst Abuse*. 2015;36(3):281–8. <https://doi.org/10.1080/0889077.2014.951140> (PMID: 25127073).
- Wells JE, Horwood LJ, Fergusson DM. Reasons why young adults do or do not seek help for alcohol problems. *Aust N Z J Psychiatry*. 2007;41(12):1005–12. <https://doi.org/10.1080/00048670701691218> (PMID: 17999273).
- Probst C, Manthey J, Martinez A, Rehm J. Alcohol use disorder severity and reported reasons not to seek treatment: a cross-sectional study in European primary care practices. *Substance Abuse Treatment, Prevention, and Policy*. 2015;10(1):32. <https://doi.org/10.1186/s13011-015-0028-z> (PMID: 26264215).
- Mushi DP, Hanlon C, Francis JM, Teferra S. Adaptation and piloting of an integrated intervention model for alcohol use disorders in primary health-care in rural Tanzania: a study protocol. *BMJ Open*. 2020;10(10): e038615. <https://doi.org/10.1136/BMJopen-2020-038615> (PMID: 33130566).

36. The United Republic of Tanzania. Tanzania census report 2012. Tanzania 2012.
37. Ghebremichael M, Paintsil E, Larsen U. Alcohol abuse, sexual risk behaviors, and sexually transmitted infections in women in Moshi urban district, northern Tanzania. *Sex Transm Dis*. 2009;36(2):102–7. <https://doi.org/10.1097/OLQ.0b013e31818b20e6> (PMID: 19060779).
38. Staton CA, Vissoci JRN, Toomey N, Abdelgadir J, Chou P, Haglund M, et al. The impact of alcohol among injury patients in Moshi, Tanzania: a nested case-crossover study. *BMC Public Health*. 2018;18(1):275. <https://doi.org/10.1186/s12889-018-5144-z> (PMID: 29466975).
39. Adams G, Gulliford MC, Ukoumunne OC, Eldridge S, Chinn S, Campbell MJ. Patterns of intra-cluster correlation from primary care research to inform study design and analysis. *J Clin Epidemiol*. 2004;57(8):785–94. <https://doi.org/10.1016/j.jclinepi.2003.12.013> (PMID: 15485730).
40. Saunders JB, Aasland OG, Babor TF, Fuente JRDL, Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption-II. *Addiction*. 1993;88(6):791–804. <https://doi.org/10.1111/j.1360-0443.1993.tb02093.x> (PMID: 8329970).
41. Monteiro MG, Gomel M. World Health Organization project on brief interventions for alcohol-related problems in primary health care settings. *Journal of Substance Misuse*. 1998;3(1):5–9.
42. Siyam A, Ir P, York D, Antwi J, Amponsah F, Rambique O, Funzamo C, Azeez A, Mboera L, Kumalija CJ, Rumisha SF, Mremi I, Boerma T, O'Neill K. The burden of recording and reporting health data in primary health care facilities in five low- and lower-middle income countries. *BMC Health Serv Res*. 2021;13(21):691. <https://doi.org/10.1186/s12913-021-06652-5>.
43. Chishinga N, Kinyanda E, Weiss HA, Patel V, Ayles H, Seedat S. Validation of brief screening tools for depressive and alcohol use disorders among TB and HIV patients in primary care in Zambia. *BMC Psychiatry*. 2011;11(1):75. <https://doi.org/10.1186/1471-244X-11-75> (PMID: 21542929).
44. Myer L, Smit J, Roux LL, Parker S, Stein DJ, Seedat S. Common Mental Disorders among HIV-Infected Individuals in South Africa: Prevalence, Predictors, and Validation of Brief Psychiatric Rating Scales. *AIDS Patient Care STDS*. 2008;22(2):147–58. <https://doi.org/10.1089/apc.2007.0102> (PMID: 18260806).
45. Vissoci JRN, Hertz J, El-Gabri D, Andrade Do Nascimento JR, Pestillo De Oliveira L, Mmbaga BT, et al. Cross-Cultural Adaptation and Psychometric Properties of the AUDIT and CAGE Questionnaires in Tanzanian Swahili for a Traumatic Brain Injury Population. *Alcohol Alcohol*. 2018 Jan 1;53(1):112–20. DOI: <https://doi.org/10.1093/alcald/agx058>. PMID: 29281046.
46. Francis JM, Grosskurth H, Kapiga SH, Weiss HA, Mwasiyua J, Changalucha J. Ethanol Concentration of Traditional Alcoholic Beverages in Northern Tanzania. *J Stud Alcohol Drugs*. 2017;78(3):476–7. <https://doi.org/10.15288/jsad.2017.78.476> (PMID: 28499118).
47. Clement S, Brohan E, Jeffery D, Henderson C, Hatch SL, Thornicroft G. Development and psychometric properties the Barriers to Access to Care Evaluation scale (BACE) related to people with mental ill-health. *BMC Psychiatry*. 2012;12(1):36. <https://doi.org/10.1186/1471-244X-12-36> (PMID: 22546012).
48. Azale T, Fekadu A, Hanlon C. Treatment gap and help-seeking for postpartum depression in a rural African setting. *BMC Psychiatry*. 2016;16(1):196. <https://doi.org/10.1186/s12888-016-0892-8> (PMID: 27287387).
49. Tefera S, Medhin G, Selamu M, Bhana A, Hanlon C, Fekadu A. Hazardous alcohol use and associated factors in a rural Ethiopian district: a cross-sectional community survey. *BMC Public Health*. 2016;3(16):218. <https://doi.org/10.1186/s12889-016-2911-6> (PMID: 26940221).
50. Programme for Improving Mental Healthcare (PRIME) (<http://www.prime.uct.ac.za/prime-tools>)
51. Barr M, Clark R, Steel D. Examining associations in cross-sectional studies. National Institute for Applied Statistics Research Australia Working Paper Series. 2016 Jan 1
52. Wilber ST, Fu R. Risk Ratios and Odds Ratios for Common Events in Cross-sectional and Cohort Studies. *Acad Emerg Med*. 2010;17(6):649–51. <https://doi.org/10.1111/j.1553-2712.2010.00773.x> (PMID: 20624147).
53. Petersen MR, Deddens JA. A comparison of two methods for estimating prevalence ratios. *BMC Med Res Methodol*. 2008;8(1):9. <https://doi.org/10.1186/1471-2288-8-9> (PMID: 18307814).
54. Barros AJ, Hirakata VN. Alternatives for logistic regression in cross-sectional studies: an empirical comparison of models that directly estimate the prevalence ratio. *BMC Med Res Methodol*. 2003;3(1):21. <https://doi.org/10.1186/1471-2288-3-21> (PMID: 14567763).
55. Francis JM, Grosskurth H, Changalucha J, Kapiga SH, Weiss HA. Systematic review and meta-analysis: prevalence of alcohol use among young people in eastern Africa. *Trop Med Int Health*. 2014;19(4):476–88. <https://doi.org/10.1111/tmi.12267> (Epub 2014 Jan 31 PMID: 24479379).
56. Duko B, Ayalew M, Ayano G. The prevalence of alcohol use disorders among people living with HIV/AIDS: a systematic review and meta-analysis. *Substance Abuse Treatment, Prevention, and Policy*. 2019;14(1):52. <https://doi.org/10.1186/s13011-019-0240-3> (PMID: 31727086).
57. Sander Greenland and Neil Pearce Statistical Foundations for Model-Based Adjustments. The Annual Review of Public Health. doi:<https://doi.org/10.1146/annurev-publhealth-031914-122559>
58. Lavallée, P. Beaumont, J.-F. (2015). Why We Should Put Some Weight on Weights. Survey Insights: Methods from the Field, Weighting: Practical Issues and 'How to' Approach, Invited article, DOI: <https://doi.org/10.13094/SMIF-2015-00001>
59. StataCorp. Stata Statistical Software: Release 13. College Station, TX: StataCorp LP. 2013.
60. Nouaman MN, Vinikoor M, Seydi M, Ekouevi DK, Coffie PA, Mulenga L, Tanon A, Egger M, Dabis F, Jaquet A, Wandeler G, leDEA. High prevalence of binge drinking among people living with HIV in four African countries. *J Int AIDS Soc*. 2018 Dec;21(12):e25202. DOI: <https://doi.org/10.1002/jia2.25202>. PMID: 30549445.
61. Soboka M, Tesfaye M, Feyissa GT, Hanlon C. Alcohol use disorders and associated factors among people living with HIV who are attending services in southwest Ethiopia. *BMC Res Notes*. 2014;7(1):828. <https://doi.org/10.1186/1756-0500-7-828> (PMID: 25417542).
62. Coates T, Hahn J, Hahn JA, Fatch R, Wanyenze RK, Baveewo S, et al. Decreases in self-reported alcohol consumption following HIV counseling and testing at Mulago Hospital, Kampala. Uganda. 2014. <https://doi.org/10.1186/1471-2334-14-403> (PMID: 25038830).
63. Reda AA, Moges A, Wondmagegn BY, Biadgilign S. Alcohol drinking patterns among high school students in Ethiopia: a cross-sectional study. *BMC Public Health*. 2012;20(12):213. <https://doi.org/10.1186/1471-2458-12-213> (PMID: 22433230).
64. Krieger H, Young CM, Anthenien AM, Neighbors C. The Epidemiology of Binge Drinking Among College-Age Individuals in the United States. *Alcohol Res*. 2018;39(1):23–30. PMID: 30557145; PMCID: PMC6104967.
65. Glantz MD, Bharat C, Degenhardt L, Sampson NA, Scott KM, Lim CCW, et al. The epidemiology of alcohol use disorders cross-nationally: Findings from the World Mental Health Surveys. *Addict Behav*. 2020;1(102): 106128. <https://doi.org/10.1016/j.addbeh.2019.106128> (PMID: 31865172).
66. Kullgren G, Alibusa S, Birabwa-Oketcho H. Problem drinking among patients attending primary healthcare units in Kampala. Uganda African journal of psychiatry. 2009;12(1):52–8. <https://doi.org/10.4314/ajpsy.v12i1.30279> (PMID: 19517048).
67. Naamara W, Muhwezi WW. Factors associated with alcohol dependence among adult male clients in Butabika Hospital, Uganda. *J Soc Work Pract Addict*. 2014;14(3):322–6. <https://doi.org/10.1080/1533256X.2014.936251> (PMID: 25264430).
68. The United Republic of Tanzania, Ministry of health. National Guideline for Primary Mental healthcare; 2006.
69. Rathod SD, Nadkarni A, Bhana A, Shidhaye R. Epidemiological features of alcohol use in rural India: a population-based cross-sectional study. *BMJ Open*. 2015;5(12): e009802. <https://doi.org/10.1136/BMJOopen-2015-009802> (PMID: 26685035).
70. Brown JS, Evans-Lacko S, Aschan L, et al. Seeking informal and formal help for mental health problems in the community: a secondary analysis from a psychiatric morbidity survey in South London. *BMC Psychiatry*. 2014;14:275. <https://doi.org/10.1186/s12888-014-0275-y>.
71. Oliver MI, Pearson N, Coe N, Gunnell D. Help-seeking behaviour in men and women with common mental health problems: a cross-sectional study. *Br J Psychiatry*. 2005;186:297–301. <https://doi.org/10.1192/bjp.186.4.297>.

72. National Academies of Sciences, Engineering, and Medicine; Health and Medicine Division; Board on Health Care Services; Committee on Health Care Utilization and Adults with Disabilities. Health-Care Utilization as a Proxy in Disability Determination. Washington (DC): National Academies Press (US); 2018 Mar 1. 2, Factors That Affect Health-Care Utilization. <https://www.ncbi.nlm.nih.gov/books/NBK500097/>
73. World Health Organization. MhGAP intervention guide for mental, neurological, and substance use disorders in non-specialized health settings. Geneva: WHO; 2017.

### **Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

