



**University of Dundee**

## **Editorial**

Tatullo, Marco

*Published in:*  
Frontiers in Cell and Developmental Biology

*DOI:*  
[10.3389/fcell.2024.1391626](https://doi.org/10.3389/fcell.2024.1391626)

*Publication date:*  
2024

*Licence:*  
CC BY

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

*Citation for published version (APA):*  
Tatullo, M. (2024). Editorial: Coordinated regulation of the balance between stem cell self-renewal and differentiation. *Frontiers in Cell and Developmental Biology*, 12, Article 1391626.  
<https://doi.org/10.3389/fcell.2024.1391626>

### **General rights**

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



## OPEN ACCESS

## EDITED AND REVIEWED BY

Valerie Kouskoff,  
The University of Manchester, United Kingdom

## \*CORRESPONDENCE

Marco Tatullo,  
✉ marco.tatullo@uniba.it

RECEIVED 26 February 2024

ACCEPTED 01 March 2024

PUBLISHED 06 March 2024

## CITATION

Tatullo M (2024), Editorial: Coordinated regulation of the balance between stem cell self-renewal and differentiation.  
*Front. Cell Dev. Biol.* 12:1391626.  
doi: 10.3389/fcell.2024.1391626

## COPYRIGHT

© 2024 Tatullo. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](#). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# Editorial: Coordinated regulation of the balance between stem cell self-renewal and differentiation

Marco Tatullo<sup>1,2,3\*</sup>

<sup>1</sup>Department of Translational Biomedicine and Neuroscience (DIBRAIN), University of Bari Aldo Moro, Bari, Italy, <sup>2</sup>School of Dentistry, University of Dundee, Dundee, United Kingdom, <sup>3</sup>Medical Institute for Regeneration and Repairing and Organ Replacement (MIRROR), Interdepartmental Center, University of Bari Aldo Moro, Bari, Italy

## KEYWORDS

stem cells, cell signaling, cell regulation, regenerative medicine, aging

## Editorial on the Research Topic

### Coordinated regulation of the balance between stem cell self-renewal and differentiation

The balance between stem cell self-renewal and differentiation is a complex mechanism regulated by different factors and signals actively impacting on the biology of the surrounding microenvironment. Understanding this balance is crucial for advancing regenerative medicine and treating diseases directly or indirectly caused by stem cell dysfunction (Pan et al., 2014).

The increasing cellular imbalance during ageing is a complex and multifactorial process that impact on the overall regulation of stem cell metabolism, mainly based on glycolysis, working in quiescent stem cells, and oxidative phosphorylation, working in differentiating stem cells (Tatullo, 2024).

In this landscape, the scientific literature has clearly demonstrated the strategic role of stem cell niches in influencing the ability to balance self-renewal and differentiation towards specific lineages, ensuring tissues homeostasis. Nonetheless, stem cells observed outside their niche often behave differently, showing fate flexibility that presupposes the presence of super-enhancers somewhat able to induce the lineage commitment and plasticity of adult stem cells (Adam et al., 2015).

This Research Topic has investigated some interesting topics on the regulatory mechanisms governing stem cells mitotic division and differentiation in invertebrates and vertebrates, posing intriguing insights on some clinical effects, variously related to molecular, biochemical, genetic or epigenetic cofactors, closely linked to stem cell activity.

During ageing, stem cells develop metabolic changes that can dramatically modify several important pathways, such as cell autophagy and mitophagy. Nonetheless, adult stem cells (ASC) show limited differentiation capabilities, typically restricted to such cell types found in the tissue of origin.

A clinical use-case that well describes the effect of ageing on human tissues is the Hair follicle (HF) homeostasis and the conditions impacting during hair aging, such as oxidative stress, hormonal disorders, inflammation, and DNA damage and repair defects. Liang et al. reported how these factors pose threats to HF cells, especially hair follicle stem cells, and mesenchymal stem cells involved in hair regeneration and pigmentation (Liang et al.).

It is, therefore, evident that declines in stem cell function during ageing are based on a number of alterations in developmental pathways and genetic regulators.

The function of the main genetic regulators of embryogenic development, the *Hox* genes, a subset of homeobox genes encoding transcription factors that are generally repressed in undifferentiated pluripotent stem cells, has not been well understood in stem cell aging. Steens and Klein have described the HOX genes, and how stem cell-related signaling pathways establish and regulate ASC-specific HOX expression pattern with different temporal-spatial topography (Steens and Klein).

The role of HOX genes is particularly important, as they are involved in the response to epigenetic stresses, which seems to induce stem cell aging in several tissues and cells, such as muscle stem cells (Muscle-SCs). In fact, an altered epigenetic stress response in Muscle-SCs has been linked to limitation of their function and self-renewal, resulting in a limitation in muscle repair and regeneration (Schworer et al., 2016).

Muscle repair and regeneration is an interesting topic, also considering the recent interest towards the production of muscle fibers for cultured meat on a large scale. Katayama et al. have addressed the importance to expand myoblasts in a serum-free medium to avoid cost, ethical, and environmental issues. By using the C2C12 cells as cell model, the authors have demonstrated that such cells differentiate quickly into myotubes and lose their ability to proliferate when plated in a serum-reduced medium; moreover, the addition of methyl- $\beta$ -cyclodextrin (M $\beta$ CD) seems to block further differentiation of myoblasts into myotubes, ensuring the balancing towards the proliferative behaviour of myoblasts in a serum-free condition for cultured meat production (Katayama et al.).

These studies have taken into consideration different cell models and molecular patterns that have basically confirmed their applicability *in vitro*; nonetheless, it is strategic to investigate also the regulatory mechanisms underlying the stem cell behaviour during replication or differentiation. There are, in fact, a number of co-factors able to influence the stem cell fate, such as the transduction of mechanical stimuli (Marrelli et al., 2015), the presence of inflammatory conditions (Gardin et al., 2020), or the activity of local enhancers, typically working at nuclear level. In this landscape, functional analyses using intestinal-derived organoids

have deepened the knowledge on Angiogenin-4 (Ang4), a member of the ribonuclease A superfamily. Abo et al. have demonstrated that Ang4 influences the intestinal stem cells (ISCs) in a concentration-dependent manner, inducing apoptosis of intestinal epithelial cells (IECs) at high concentration, and promoting the growth of Lgr5<sup>+</sup> ISCs-based organoids at low concentration (Abo et al.).

In conclusion, the coordinated and multifactorial regulation of the stem cell behaviour is the primary mechanism to control the proper morphogenesis of organs and tissues. In this Research Topic, we have assessed how different regulators participate, by direct or mediate activities, in development of multiple organs, influencing the stem cell behaviour both under *in vitro* experimental conditions and in *in vivo* biomimetic models.

## Author contributions

MT: Writing—original draft, Writing—review and editing.

## Funding

The author(s) declare that no financial support was received for the research, authorship, and/or publication of this article.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## References

- Adam, R. C., Yang, H., Rockowitz, S., Larsen, S. B., Nikolova, M., Oristian, D. S., et al. (2015). Pioneer factors govern super-enhancer dynamics in stem cell plasticity and lineage choice. *Nature* 521 (7552), 366–370. doi:10.1038/nature14289
- Gardin, C., Bosco, G., Ferroni, L., Quartesan, S., Rizzato, A., Tatullo, M., et al. (2020). Hyperbaric oxygen therapy improves the osteogenic and vasculogenic properties of mesenchymal stem cells in the presence of inflammation *in vitro*. *Int. J. Mol. Sci.* 21 (4), 1452. doi:10.3390/ijms21041452
- Marrelli, M., Falisi, G., Apicella, A., Apicella, D., Amantea, M., Cielo, A., et al. (2015). Behaviour of dental pulp stem cells on different types of innovative mesoporous and nanoporous silicon scaffolds with different functionalizations of the surfaces. *J. Biol. Regul. Homeost. Agents* 29 (4), 991–997.
- Pan, L., Wang, S., Lu, T., Weng, C., Song, X., Park, J. K., et al. (2014). Protein competition switches the function of COP9 from self-renewal to differentiation. *Nature* 514 (7521), 233–236. doi:10.1038/nature13562
- Schwörer, S., Becker, F., Feller, C., Baig, A. H., Köber, U., Henze, H., et al. (2016). Epigenetic stress responses induce muscle stem-cell ageing by Hoxa9 developmental signals. *Nature* 540 (7633), 428–432. doi:10.1038/nature20603
- Tatullo, M. (2024). Entropy meets physiology: should we translate aging as disorder? *Stem Cells Dayt. Ohio* 42 (2), 91–97. doi:10.1093/stmcls/sxad084