

The association between frailty, care receipt and unmet need for care with the
risk of hospital admissions

Asri Maharani^{1¶*}, David R Sinclair^{2¶}, Andrew Clegg³, Barbara Hanratty², James Nazroo⁴,
Gindo Tampubolon⁵, Chris Todd¹, Raphael Wittenberg⁶, Terence W O'Neill^{1&}, Fiona E
Matthews^{2&}

¹ National Institute for Health and Care Research (NIHR) Policy Research Unit in Older
People and Frailty / Healthy Ageing, School of Health Sciences, Faculty of Biology,
Medicine and Health, The University of Manchester, Manchester, M13 9PL, UK.

² National Institute for Health and Care Research (NIHR) Policy Research Unit in Older
People and Frailty / Healthy Ageing, Population Health Sciences Institute, Newcastle
University, Newcastle-upon-Tyne, NE4 5PL, UK.

³ Academic Unit for Ageing and Stroke Research, Bradford Institute for Health Research,
School of Medicine, University of Leeds, UK.

⁴ Cathie Marsh Institute for Social Research, School of Social Sciences, Faculty of Humanities, University of Manchester, UK.

⁵ Global Development Institute, University of Manchester, UK.

⁶ National Institute for Health and Care Research (NIHR) Policy Research Unit in Older People and Frailty / Healthy Ageing, Care Policy and Evaluation Centre, London School of Economics and Political Science, London, WC2A 2AE, UK.

*Corresponding author

Email: asri.maharani@manchester.ac.uk

[¶] These authors contributed equally to this work.

[&] These authors also contributed equally to this work.

Abstracts

Background: Frailty is characterised by a decline in physical, cognitive, energy, and health reserves and is linked to greater functional dependency and higher social care utilisation.

However, the relationship between receiving care, or receiving insufficient care among older people with different frailty status and the risk of unplanned admission to hospital for any cause, or the risk of falls and fractures remains unclear.

Methods and findings: This study used information from 7,656 adults aged 60 and older participating in the English Longitudinal Study of Ageing (ELSA) waves 6-8. Care status was assessed through received care and self-reported unmet care needs, while frailty was measured using a frailty index. Competing-risk regression analysis was used (with death as a potential competing risk), adjusted for demographic and socioeconomic confounders. Around a quarter of the participants received care, of which approximately 60% received low levels of care, while the rest had high levels of care. Older people who received low and high levels of care had a higher risk of unplanned admission independent of frailty status. Unmet need for care was not significantly associated with an increased risk of unplanned admission compared to those receiving no care. Older people in receipt of care had an increased risk of hospitalisation due to falls but not fractures, compared to those who received no care after adjustment for covariates, including frailty status.

Conclusions: Care receipt increases the risk of hospitalisation substantially, suggesting this is a group worthy of prevention intervention focus.

Introduction

Demand for care services for older people is increasing as the global population continues to age [1]. The World Health Organisation (WHO) Global Strategy and Action Plan on Ageing and Health 2016–2020 highlighted the right of older people to receive care and support to maintain their best possible functional abilities [2]. Frailty, which describes how we gradually lose our in-built reserves with increasing age [3], is a framework for understanding health discrepancies among older adults and a significant predictor of care receipt [4]. Estimates of frailty prevalence worldwide vary between 12% to 24% [5]. Almost all older people with frailty (93%) experience mobility problems, and over half of them have difficulties with washing, dressing or housework [6]. Older people with frailty are thus more likely to be in need of social care services.

A prior study estimated that caring for frail people will cost between 4 and 9 times as much as caring for healthy people [7]. In the UK, social care is provided through paid care from public and private funding and unpaid care from friends and family. Despite this, a report estimated that 1.5 million people over 65 in England have unmet care needs [8]. Our prior work estimated that around 0.7 million and 1.6 million people aged 65+ in England were frail and prefrail in 2018, respectively. However, only 0.5 million adults in the same age group received government funding for care [9]. We also found that 82% (124 from 151) of the local authorities in the study had a greater number of persons with frailty aged 65+ than care recipients within the same age range, suggesting, given that frail individuals are more likely to require care, that there is a care deficit present in much of the country.

Frailty is associated with increased healthcare use, and hospital admissions represent a substantial proportion of the overall costs associated with the condition [10,11]. Frailty is associated with an annual additional 1.0 million emergency admissions and 1.1 million elective admissions in England [12]. Frail patients are also more likely to be attended by an

ambulance for incidents which do not require conveyance to a hospital [13]. In addition, severely frail older people have seven times longer lengths of stay in hospital following emergency hospitalisation than non-frail older people. The negative consequences of unmet care needs among older people on mental health problems [14,15] and higher mortality rates [16] have been documented in the literature. However, there is limited evidence on the effect of care receipt and unmet need for care among older people with different frailty status and their future healthcare utilisation.

This study aimed to understand how care receipt and unmet need for care among older people with different frailty status are associated with the risk of unplanned admission to the hospital for any cause and for conditions associated with frailty, specifically falls [17,18] and fractures [19,20].

Materials and methods

Participants and Setting

The analysis uses a dataset that combines the English Longitudinal Study of Ageing (ELSA) [21] with the census of public hospital records in England, the Hospital Episode Statistics (HES) [22], and mortality data from the Office for National Statistics (ONS) [23]. ELSA is a panel survey of a representative sample of the household population aged 50+ in England [21]. ELSA waves are performed every two years, collecting information on demographic, socioeconomic, and health characteristics. To date, it has conducted nine waves. Our analysis used data from ELSA waves 6 to 8, covering 2012-2017, as the information on the types of care received is available from wave 6, and HES and ONS data were available until 31 January 2018. All individuals included in the analysis had data linked to HES and ONS mortality (including those who dropped out of the study after the baseline survey). In this study, we included ELSA participants aged 60 and older.

Measures

Frailty

Frailty was assessed using a frailty index derived from data collected as part of ELSA. The frailty index included 60 variables ('deficits') representing conditions that accumulate with age and are associated with adverse outcomes, including disability, mobility, sensory impairments, cognitive function, and chronic diseases. The full list of variables used to create the frailty index is shown in **S1 Table**. An individual's frailty index is calculated as the proportion of possible deficits present in an individual. Frailty indices with at least 30-40 deficits can predict adverse outcomes accurately [24,25]. Frailty was measured at baseline (Wave 6). We categorised the frailty index into three groups: robust (frailty index ≤ 0.08), prefrail (frailty index $>0.08-0.25$) and frail (frailty index ≥ 0.25) [26].

Level of care and unmet need for care

Respondents in ELSA were asked to respond to questions about their care if they reported having at least one difficulty with mobility, an Activity of Daily Living (ADL) or an Instrumental Activity of Daily Living (IADL) [27]. Based on the level of care received, we categorised respondents into those in receipt of: (1) high levels of care, if the respondents received help in the last month for using the toilet, getting in and out, eating, bathing/showering, walking across a room, dressing, and having meals on wheels; (2) low levels of care, if the respondents received help in the last month for grocery shopping, house or garden work, managing money, climbing at least one flight of stairs without resting, taking medication, walking 100 yards and if they had attended a day centre; and (3) did not receive care.

Participants who have received care were also asked whether their care meets their needs. We classified the respondents into having: (1) unmet care needs, if they answered that the care

they sometimes had or hardly met their needs; and (2) met care needs, if they answered that the care they had met or usually met their needs; and (3) did not receive care.

Outcome measures

Unplanned admissions were derived from the HES data linked by NHS Digital to ELSA participants' NHS number, date of birth, gender and postcode. An unplanned admission was defined as admission to the hospital through (1) accident and emergency (A&E); (2) general practitioner (GP) after request of immediate admission; (3) bed bureau [28]; (4) consultant clinic; (5) Mental Health Crisis Resolution team; and (6) other A&E [29]. The full list of the HES method of admission codes is shown in **S2 Table** [29].

Hospitalisation due to falls was defined as the first hospitalisation where a diagnosis of fall was recorded since baseline (wave 6) based on the International Classification of Disease 10th version (ICD-10) of falls, i.e., W00 to W19 [30,31]. Hospitalisation due to fractures was the first hospitalisation where a fracture diagnosis was recorded since the baseline corresponded to the ICD-10 M, S and T codes (see **S3 Table**).

Covariates

Age was included in the principal analysis as a continuous variable and in sensitivity analysis after categorisation into 5-year age groups (60-64; 65-69; 70-74; 75-79; 80-84; 85+). Gender (male/female), ethnicity (white/non-white) and marital status (married/not married) were categorised as indicated. Educational attainment was categorised into lower than secondary school (reference), secondary school, and college or higher. Wealth was measured by the net total wealth of the respondent's benefit unit (defined as a single adult, or a married or cohabiting couple, and any dependent children [32]). Net total wealth comprised the sum of savings and investments after subtracting financial debt. We split wealth into quintiles to investigate the hierarchical effects of wealth.

Statistical analysis

To examine the effect of the mismatch between levels of frailty and receipt of care on each hospitalisation category in this study, we employed competing-risk regression analysis using a version of the Fine and Gray analysis [33]. This analysis allows a competing risk – an event that might occur during the follow-up instead of the event of interest – to be considered in the model. Death is the potential competing risk in this study when examining hospital admissions. Mortality status was ascertained from linked register data up to the end of January 2018. Frailty, level of care and need for care were defined in wave 6 (2012/2013) and the follow-up time up to 31 January 2018. We present the results as the subdistribution hazard ratios (SHRs) and 95% confidence intervals (95% CIs) [34]. The subdistribution hazard function is defined as the instantaneous rate of occurrence of hospitalisation in older people who have not yet experienced it during the study [34]. The SHR is the ratio of these functions in the presence of two different values of a covariate (e.g., a person who is frail relative to a person who is not frail).

For unplanned admissions as the outcome, we performed the analysis separately for the level of care and need for care. The first analysis included frailty status (robust as the reference, prefrail, and frail) and level of care (no care as the reference, low and high levels of care), while the second analysis included frailty status (robust as the reference, prefrail, and frail) and need for care (no care as the reference, met care needs, and unmet care needs). All analyses were adjusted for age, gender, ethnicity, marital status, wealth and education.

We further performed the analysis by gender and categorised the care receipt into: (1) received care; and (2) did not receive care. The same categorisation was used to analyse conditions associated with frailty: falls and fractures.

We checked for the presence of an interaction between frailty status, level of care, and need for care by creating a second model for each analysis. In Model 2, we created nine main dependent variables combining frailty status and level of care: (1) robust and received no care (reference group); (2) robust and received low levels of care; (3) robust and received high levels of care; (4) prefrail and received no care; (5) prefrail and received low levels of care; (6) prefrail and received high levels of care; (7) frail and received no care; (8) frail and received low levels of care; and (9) frail and received high levels of care. For analysis of the need for care, we created eight main dependent variables combining frailty status and need for care (there were no robust respondents reporting the unmet need for care needs): (1) robust and received no care (reference group); (2) robust and received care; (3) prefrail and received no care; (4) prefrail and reported having met care needs; (5) prefrail and reported having unmet care needs; (6) frail and received no care; (7) frail and reported having met care needs; and (8) frail and reported having unmet care needs. We looked for an interaction between frailty status, level of care and need for care on the risk of hospitalisation by plotting the SHRs and 95% CIs using both models. In order to compare Model 1 (without interaction) with Model 2 (with interaction), we calculated the SHRs of each category (i.e., robust and received no care as the reference; robust and received low levels of care; robust and received high levels of care; prefrail and received no care; prefrail and received low levels of care; prefrail and received high levels of care; frail and received no care; frail and received low levels of care; and frail and received high levels of care) by adding the log of each frailty status, level of care, and need for care and then taking its exponential. An interaction effect was considered to exist if the two plots showed different values of the association of the categories and the risk of hospitalisation. **S1 Fig** and **S2 Fig** show that the two plots have similar values, suggesting no interaction between frailty status and care receipt in their

relationships with the risk of hospitalisation. The model without an interaction was thus preferable. Survey data was weighted using ELSA cross-sectional survey weight at wave 6.

Sensitivity analysis

We performed three types of sensitivity analyses. Firstly, we used age categorised into groups (60-64; 65-69; 70-74; 75-79; 80-84; 85+) instead of age as a continuous variable.

Secondly, we performed two analyses on different sets of short epochs of time. The first set of epochs of time are: (1) wave 6 as the baseline with 6 months follow-up; (2) wave 7 baseline with 6 months follow-up; and (3) wave 8 baseline; 6 months follow-up. The second set of epochs of time are: (1) wave 6 baseline with 12 months follow-up; (2) wave 7 baseline with 12 months follow-up; and (3) wave 8 baseline; 6 months follow-up. We performed two meta-analyses using those two sets of epochs of time. In those analyses, frailty status, level of care and need for care were defined at each wave 6, 7, and 8. The start date was defined as the interview date. Age was defined as the age at each wave, and we had two different follow-up lengths for each wave, except for wave 8: 6 and 12 months. We could not have a similar follow-up length in wave 8 as the data were only available until 31 January 2018 (6 months after Wave 8 enrolled).

Finally, we performed the analysis by putting a censor date between two interview dates if there were any changes in frailty status, level of care or need for care between the two waves of ELSA. When a person's response changed between waves, we assumed the change occurred midway between the waves (censor date). The respondents were followed up until the censor date, death or end of the study if they did not change frailty status.

Results

Participant characteristics

Descriptive characteristics of the study sample are presented in **Table 1**. A total of 7,656 participants, 3,535 men and 4,121 women, were included in the analysis. The mean age was 71.1 years. The majority (97.2%) were white and 65.3% were married. Almost half (48.8%) of the respondents graduated from college or higher education level. After applying sample weighting, the proportion of participants who were frail and prefrail was estimated as 17.7% and 40.6%, respectively.

Table 1 Descriptive characteristics of the respondents at baseline. *Notes:* * unweighted; ** weighted

	Total*	Robust**	Prefrail**	Frail**
Frailty index, mean (SD)	0.1 (0.1)			
<i>Frailty status, n (%)</i>				
Robust	3,357 (43.9)	2,910 (41.7)		
Prefrail	3,026 (39.5)		2,833 (40.6)	
Frail	1,268 (16.6)			1,239 (17.7)
Age, mean (SD)	71.1 (8.2)	68.10 (6.5)	72.73 (8.4)	76.28 (9.6)
<i>Sex, n (%)</i>				
Males	3,535 (46.2)	1,574 (48.8)	1,182 (36.7)	468 (14.5)

Females	4,121 (53.8)	1,336 (35.6)	1,651 (43.9)	771 (20.5)
<i>Ethnicity, n (%)</i>				
White	7,442 (97.2)	2,819 (41.8)	2,758 (40.9)	1,169 (17.3)
Non-White	214 (2.8)	91 (38.6)	75 (31.9)	69 (29.5)
<i>Married, n (%)</i>				
No	2,653 (34.7)	696 (28.6)	1,065 (43.7)	677 (27.8)
Yes	5,001 (65.3)	2,213 (48.7)	1,767 (38.9)	562 (12.4)
<i>Education attainment, n (%)</i>				
Less than secondary school	2,507 (32.7)	706 (28.1)	1,108 (44.1)	699 (27.8)
Secondary school	1,414 (18.5)	570 (45.4)	528 (42.0)	159 (12.6)
College or higher	3,735 (48.8)	1,634 (50.9)	1,197 (37.3)	381 (11.9)
<i>Wealth, n (%)</i>				
5 th quintile (most wealthy)	1,500 (20.0)	859 (61.6)	460 (33.0)	75 (5.4)
4 th	1,500 (20.0)	686 (49.1)	608 (43.5)	103 (7.4)
3 rd	1,499 (20.0)	614 (44.1)	604 (43.3)	177 (12.7)
2 nd	1,504 (20.0)	498 (35.7)	617 (44.2)	281 (20.1)
1 st quintile (least wealthy)	1,495 (19.9)	334 (23.9)	661 (47.3)	401 (28.7)

<i>Level of care received, n (%)</i>				
No care	5,213 (74.0)	2,869 (55.0)	2,154 (41.3)	190 (3.7)
Receiving low levels of care	1,080 (15.3)	43 (4.0)	620 (57.4)	417 (38.6)
Receiving high levels of care	749 (10.6)	6 (0.8)	190 (25.4)	553 (73.8)
<i>Need for care, n (%)</i>				
No care	5,213 (74.0)	2,869 (55.0)	2154 (41.3)	190 (3.7)
Met care needs	1,167 (16.6)	27 (2.3)	559 (47.9)	581 (49.8)
Unmet care needs	539 (7.7)	7 (1.2)	170 (31.6)	362 (67.2)

The proportion of respondents with pre-frailty and frailty increased with age. Almost 10% of people aged 60-64 were frail, increasing to 44.4% among those aged 85+. Compared to men, women were more likely to be frail (20.5% vs 14.5%) and prefrail (43.9% vs 36.7%).

Compared to those who did not complete high school, people who graduated from high school and college or higher were less likely to be frail and prefrail. The proportion of respondents with frailty increased from 5.4% among the wealthiest quintile to 28.7% among the least wealthy quintile.

Around a quarter of adults aged 60+ in England received care, of which approximately 60% received low levels, while the rest had high levels of care. The level of care receipt is proportionally higher among frail and prefrail than robust older people: 6.4% and 47.6% of

the prefrail and frail respondents received high levels of care, respectively. Around a fifth (20.9%) of respondents with prefrailty received low levels of care, while 36.0% of those with frailty had low levels of care. Characteristics of respondents at baseline by level of care are shown in **S4 Table**. It shows that the proportions of individuals receiving either low or high levels of care (compared to no care) were higher among those who were older, female, non-White, not married, those who had lower educational attainment and who were less wealthy. Around 16.6% of respondents with prefrailty stated that their care needs were met, while 7.7% reported unmet needs for care. Half of the respondents with frailty stated that they had met care needs, while almost one-third (31.3%) reported unmet care needs. Characteristics of respondents at baseline categorised by the need for care are shown in **S5 Table**. The proportions of individuals reporting unmet need for care were higher among those who were older, female, non-White, not married, had lower education attainment and were less wealthy.

Frailty status, level of care and risk of unplanned hospital admission

During five years of follow-up, there were 2,663 unplanned admissions and 310 deaths (**S6 Table**). In an unadjusted competing risk model, compared to those who were robust, the subdistribution hazard ratios (SHRs) for unplanned hospital admission among people who were prefrail and frail were 1.80 (95%CI: 1.64; 1.97) and 2.74 (95%CI: 2.47; 3.03) respectively, see **S7 Table**. Compared to those who received no care, those who received either low or high levels of care were more likely to have an unplanned hospital admission: SHR 1.70 (95%CI:1.55; 1.87) and 1.82 (95%CI:1.64; 2.02) respectively.

After adjustment for covariates, the SHRs for unplanned hospital admission among those who were prefrail and frail were attenuated (see **Table 2**). Compared to those who were robust, the adjusted SHR for unplanned admission for prefrailty was 1.76 (95%CI: 1.59;

1.95) and for frailty 2.46 (95%CI:2.13; 2.84). After adjustment for covariates including frailty status, compared to those not receiving care, the adjusted SHR for unplanned admission for those with low levels of care was 1.19 (95%CI:1.06; 1.33) and for those with high levels of care was 1.29 (95%CI:1.12; 1.48).

Table 2 Subdistribution hazard ratio (95% CI) for the association between frailty status, level of care received, need for care and unplanned admissions.


	Level of care	Need for care
<i>Frailty status, reference: robust</i>		
Prefrail	1.76 (1.59; 1.95)	1.77 (1.60; 1.95)
Frail	2.46 (2.13; 2.84)	2.51 (2.18; 2.89)
<i>Level of care received, reference: no care</i>		
Receiving low levels of care	1.19 (1.06; 1.33)	
Receiving high levels of care	1.29 (1.12; 1.48)	
<i>Need for care, reference: no care</i>		
Met care needs		1.22 (1.09; 1.35)
Unmet care needs		1.21 (0.91; 1.61)

Note: Unplanned admissions N=2,662, competing event deaths N=310. All models were adjusted for age, gender, marital status, wealth in quintiles and education attainment.

Taking account of death as a competing risk, the cumulative incidence of unplanned hospital admissions increased over time for all frailty categories; the slope was greater among those who were frail and prefrail than those who were robust (see **Fig 1A**). The slope was also

greater within frailty categories for those who received care than those who did not. The cumulative incidence curve for frail people with high levels of care increased steeply with time, followed by frail people with low levels of care.

Fig 1 Estimates of the cumulative incidence of unplanned hospitalisation according to frailty status and (A) level of care received and (B) need for care. Death was the competing risk.



Frailty status, need for care and risk of unplanned hospital admission

In an unadjusted competing risk model, compared to those who were not in receipt of care, the SHRs for unplanned hospital admission among people who were in receipt of care and whose care needs were met was 1.82 (95%CI:1.64; 2.02), whilst for those with an unmet need of care the SHR was 2.07 (95%CI:1.61; 2.67), see **S7 Table**.

After adjustment for covariates, including frailty status, the strength of the SHRs was attenuated. Compared to those not receiving care, the adjusted SHR for unplanned admission for those in receipt of care and whose care needs were met was 1.22 (95%CI: 1.09; 1.35), with a similar SHR for unmet need for care 1.21 (95%CI: 0.91; 1.61), though with the confidence interval embracing unity, see **Table 2**.

Taking account of death as a competing risk, the cumulative incidence of unplanned hospital admissions was higher within frailty categories for those who were in receipt of care and whose care needs were met than those with an unmet need for care (**Fig 1B**).

For the first sensitivity analysis, we analysed the interaction between frailty with the level of care and need for care. **S1 Fig** and **S2 Fig** show that the analysis of the interaction between frailty with the level of care and need for care have similar values with those excluding the interaction, suggesting no interaction between frailty status and care receipt in their relationships with the risk of hospitalisation. The results of the sensitivity analyses using age group as the covariates (**S8 Table**), five different epochs of time (**S3 Fig**), and varying times of analysis (**S9 Table**) are similar to our principal results, suggesting the results are robust.

Frailty, level of care and risk of unplanned admission: Influence of gender

Among men, after adjustment for covariates including frailty status, compared to those who received no care, those who received care were associated with an increased risk of unplanned hospitalisation (SHRs 1.30; 95% CI 1.09, 1.54), see **S10 Table**. This was also true for women (SHRs 1.31; 95% CI 1.14, 1.50). **S4 Fig** shows that among men, those who were frail and received care had the steepest estimated cumulative incidence, followed by those who were frail and did not receive care. This order was similar for women, as being frail and receiving care had a steeper estimated cumulative incidence of frail.

Frailty status, receipt of care and the risk of admissions due to falls and fractures

During five years of follow-up, there were 586 admissions due to falls and 432 admissions due to fractures (**S6 Table**). **Table 3** reports the SHR for the association between frailty and care receipt levels and the risk of hospitalisation due to a fall estimated using competing risk analysis. The adjusted SHRs for hospitalisation due to a fall among older adults who were prefrail and frail were 2.18 (95%CI: 1.68; 2.83) and 2.73 (95%CI: 1.95; 3.80), respectively, compared with those who were robust. Receiving care was associated with a 1.30 (95% CI: 1.03; 1.63) higher risk of admissions due to falls.

Table 3 Subdistribution hazard ratio (95% CI) for the association between frailty status and care receipt with hospitalisation due to falls and fractures, England 2012-2018

	Hospitalisation due to falls ^a	Hospitalisation due to fractures ^a
--	---	---

Frailty status, reference: robust

Prefrail	2.18 (1.68; 2.83)	1.78 (1.35; 2.34)
Frail	2.73 (1.95; 3.80)	2.11 (1.45; 3.07)

Received care, reference: No


Yes	1.30 (1.03; 1.63)	1.25 (0.95; 1.63)
-----	-------------------	-------------------

Note: ^aAdjusted for age, gender, ethnicity, marital status, wealth and education.

The adjusted SHRs for hospitalisation due to a fracture among older adults who were prefrail and frail were 1.78 (95%CI: 1.35; 2.34) and 2.11 (95%CI: 1.45; 3.07), respectively, compared with those who were robust. Receiving care (SHR: 1.25; 95% CI: 0.95; 1.63) was not significantly associated with an increased risk of admissions due to fractures.

Fig 2A shows that frail older people had the steepest estimated cumulative incidence curves for hospitalisation due to falls, followed by those who were prefrail and robust. For fractures, prefrail older people with care had the steepest estimated cumulative incidence curve, followed by frail older people with no care (**Fig 2B**). In both cases (falls -2A and fractures -2B), the estimated cumulative incidence curves for hospitalisation were the steepest for frail older people regardless of whether or not they received care. It is also noticeable that each reason for admission (falls and fractures) for each group (frail, prefrail, robust) receiving care always fares worse than those not receiving care.

Fig 2. Estimates of the cumulative incidence curves of risk of hospitalisation due to falls and fractures according to frailty status and receipt of care. Death was the competing risk. (A) Hospitalisation due to falls; (B) Hospitalisation due to fractures.



Discussion

Using a large population-based survey (ELSA) linked to national hospitalisation and mortality records, we found that 15.2% and 10.4% of adults aged 60+ in England received low and high levels of care, respectively, with the proportion reporting care receipt higher among prefrail and frail than robust individuals. The data are consistent with previous findings [35-37]. For instance, a study based on primary care in Norwich found that the average number of care plans required per referral was higher among severely frail older patients (2.97) than fit patients (2.22), indicating more complex care needs in the community

[36]. In a cross-sectional study in the Netherlands, frail older adults with more ADL limitations and a higher frailty score were more likely to have higher care needs [38].

Our results suggest that compared to those receiving no care, receiving low or high levels of care was associated with a higher risk of unplanned admission and hospital admissions due to falls independent of frailty status. The finding may suggest the presence of other factors relating to falls were not captured by the frailty index, including Parkinson's disease [39], history of falls [40,41], and polypharmacy [42]. Future studies may include these factors in predicting medical care usage. Another factor which may affect hospitalisation is living status. The risk of falls might be higher among older people living alone [43] because of the amount of time between carer visits, no one around to help with the toilet, and concern that it is not a 'safe' environment to leave someone in post-fall.

In our analysis, the proportion of unmet care needs was highest among frail older people. An unmet need for care was associated with a small though non-significant risk of unplanned hospitalisation, with the magnitude of the risk similar to those whose care needs were reported as being met. However, caution is needed in interpreting these data as our definition of care needs focuses on the *adequacy* (met / unmet) of those who were already receiving care. There is a relative lack of data concerning the role of the unmet need for care as a contextual factor when examining frailty and adverse health outcomes in older adults, for which further research is needed. Supporting our finding, data from a Canadian study suggest that perceived unmet need for care among adults with chronic conditions was not associated with an increased risk of hospital admission [44], while two American studies did find an association [45,46].

We found that 40.6% and 17.7% of adults aged 60+ in England were prefrail and frail, respectively. Both frailty and prefrailty (compared to being robust) were associated with a higher risk of unplanned hospital admission and hospital admissions due to falls and fractures

after adjusting with care receipt and unmet need for care. These findings corroborate previous studies that report an association between frailty and an increase in emergency and elective hospital admissions [12,35,36,47]. The impact of frailty on healthcare utilisation is substantial: the length of inpatient stay for severely frail patients was seven times longer than for non-frail patients [12]. In relation to the influence of gender, our data suggest that after adjusting for covariates, receiving care (compared to receiving no care) was associated with a higher risk of unplanned admissions among men and women and with a magnitude of risk similar in men and women.

Strengths of our analysis include the nationally representative sample of non-institutionalised individuals, which is generalisable to the English population. Furthermore, the survey used in this study was linked to national hospitalisation and mortality data, which minimised loss at follow-up. Additionally, this study used a competing risk analysis strategy to consider mortality as a competing event rather than a survival analysis. Competing risk analysis accommodates the competing nature of multiple causes of the same event.

Several limitations need to be considered in interpreting the findings. First, care receipt and need for care were measured only at baseline, with no follow-up data. It was not possible, therefore, to address how changes in care receipt and care needs may have affected hospitalisation among older people. Second, questions about care were only asked when a respondent reported having difficulties in mobility, ADL or IADL in ELSA. Thus, information on care receipt and the need for care excluded those who did not report any functional difficulties; it is possible that more people would have reported care receipt and care needs if the entire sample had been asked. In addition, perceived unmet needs were measured using only one question in ELSA, which did not distinguish between different care needs. A cross-sectional study among frail older adults in the Netherlands examined different types of unmet care needs, i.e., environmental (accommodation, household activities, food,

and caring for another), physical needs (physical health, medication use, visual/hearing impairment, mobility/falls, and self-care), and psychosocial needs (memory, company, daytime activities, and information) [38]. The respondents reported the highest proportion of unmet care needs in the psychosocial domain. It is possible that different types of unmet needs may affect adverse health outcomes differently. Finally, the frailty index constructed in this study did not include the diagnosis of sarcopenia and nutritional status due to the unavailability of the information in ELSA. The Italian frailty index, for instance, includes the nutritional index and provides good reliability and validity in predicting mortality, disability and hospitalisation [48]. Future research may include sarcopenia, nutritional status, and other geriatric assessments in constructing a frailty index to allow for a more comprehensive assessment of an older adult's health.

Our findings have potential implications. In our analysis, frailty was associated with an increased risk of unplanned admission to the hospital. As frailty is a potentially reversible health state [49], early screening and intervention, good-quality and timely diagnosis of prefrailty and frailty in the community, and effective interventions at an early stage could be effective strategies for reducing or delaying the utilisation of secondary care services. Prior study shows that low social support is associated with long-term mortality among older people [50]. Our data suggest that older people with frailty or prefrailty who are *already in receipt of care* are at significantly greater risk of unplanned hospitalisation and, therefore, a group who may potentially benefit from more detailed assessment and targeted or personalised community-based interventions with the aim of reducing their risk.

In conclusion, older men and women who are in receipt of care are at increased risk of unplanned hospitalisation and other adverse outcomes. Those who are frail or prefrail are at greater risk of hospitalisation, providing opportunities for targeted community-based interventions to reduce the impact on already overstretched secondary care services.

Acknowledgements: We thank our academic and professional support colleagues from the National Institute for Health and Care Research Policy Research Unit in Older People and Frailty / Healthy Ageing, with whom we discussed the ideas presented in this paper during unit meetings.

Main text word counts: 4928

Number of data elements: 5

References

1. Government Office for Science. Future of an Ageing Population. London.
<https://www.gov.uk/government/publications/future-of-an-ageing-population>.
Published 7 July 2016.
2. World Health Organization. Global Strategy and Action Plan on Ageing and Health 2016–2020. Geneva.
<https://www.who.int/publications/i/item/9789241513500>. Published 2 January 2017.
3. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older adults: Evidence for a phenotype. *J Gerontol Ser A Biol Sci Med Sci*. 2001;56(3):M146-M57. doi: 10.1093/gerona/56.3.m146.
4. Kojima G. Frailty as a predictor of disabilities among community-dwelling older people: A systematic review and meta-analysis. *Disabil Rehabil*. 2017;39(19):1897-908. doi: 10.1080/09638288.2016.1212282.
5. O’Caoimh R, Sezgin D, O’Donovan MR, Molloy DW, Clegg A, Rockwood K, Liew A. Prevalence of frailty in 62 countries across the world: A systematic review and meta-analysis of population-level studies. *Age Ageing*. 2021;50(1):96-104. doi: 10.1093/ageing/afaa219.
6. Gale CR, Cooper C, Aihie Sayer A. Prevalence of frailty and disability: findings from the English Longitudinal Study of Ageing. *Age Ageing*. 2014;44(1):162-5. doi: 10.1093/ageing/afu148.
7. Nikolova S, Heaven A, Hulme C, West R, Pendleton N, Humphrey S, et al. Social care costs for community-dwelling older people living with frailty. *Health Soc Care Community*. 2022;30(3):e804-e11. doi: 10.1111/hsc.13450.

8. Age UK. Estimating need in older people: Findings for England. Available at: <https://www.ageuk.org.uk/globalassets/age-uk/documents/reports-and-publications/reports-and-briefings/active-communities/id204303-estimating-needs-report.pdf>. Published 24 December 2019.
9. Sinclair D, Maharani A, Chandola T, Bower P, Hanratty B, Nazroo J, et al. Using individual and neighbourhood profiles and trends to understand frailty with nationally representative population data. Part 2: Frailty and receipt for care in England. England: NIHR Policy Research Unit Older People and Frailty. <https://documents.manchester.ac.uk/display.aspx?DocID=58453>. Published June 2021.
10. Hoogendijk EO, Afilalo J, Ensrud KE, Kowal P, Onder G, Fried LP. Frailty: implications for clinical practice and public health. *Lancet*. 2019;394(10206):1365-75. doi: 10.1016/S0140-6736(19)31786-6.
11. García-Nogueras I, Aranda-Reneo I, Peña-Longobardo L, Oliva-Moreno J, Abizanda P. Use of health resources and healthcare costs associated with frailty: the FRADEA study. *J Nutr Health Aging*. 2017;21(2):207-14. doi: 10.1007/s12603-016-0727-9.
12. Han L, Clegg A, Doran T, Fraser L. The impact of frailty on healthcare resource use: A longitudinal analysis using the Clinical Practice Research Datalink in England. *Age Ageing*. 2019;48(5):665-71. doi: 10.1093/ageing/afz088.
13. Charlton K, Sinclair DR, Hanratty B, Burrow E, Stow D. Measuring frailty and its association with key outcomes in the ambulance setting: A cross sectional observational study. *BMC Geriatr*. 2022;22(1):1-10. doi: 10.1186/s12877-022-03633-z.

14. Hu B, Wang J. Unmet long-term care needs and depression: The double disadvantage of community-dwelling older people in rural China. *Health Soc Care Community*. 2019;27(1):126-38. doi: 10.1111/hsc.12630.
15. Song H, Sun H. Association of unmet long-term care needs with depression and anxiety among older adults in urban and rural China. *Geriatr Nurs*. 2023;49:115-21. doi: 10.1016/j.gerinurse.2022.11.017
16. He S, Craig BA, Xu H, Covinsky KE, Stallard E, Thomas III J, et al. Unmet need for ADL assistance is associated with mortality among older adults with mild disability. *J Gerontol A Biol Sci Med Sci*. 2015;70(9):1128-32. doi: 10.1093/gerona/glv028.
17. Cheng MH, Chang SF. Frailty as a risk factor for falls among community dwelling people: Evidence from a meta-analysis. *J Nurs Scholarsh*. 2017;49(5):529-36. doi: 10.1111/jnu.12322.
18. Kojima G. Frailty as a predictor of future falls among community-dwelling older people: a systematic review and meta-analysis. *J Am Med Dir Assoc*. 2015;16(12):1027-33. doi: 10.1016/j.jamda.2015.06.018.
19. Kojima G. Frailty as a predictor of fractures among community-dwelling older people: A systematic review and meta-analysis. *Bone*. 2016;90:116-22. doi: 10.1016/j.bone.2016.06.009.
20. Chen KW, Chang SF, Lin PL. Frailty as a predictor of future fracture in older adults: A systematic review and meta-analysis. *Worldviews Evid Based Nurs*. 2017;14(4):282-93. doi: 10.1111/wvn.12222.
21. Banks J, Batty G, Breedvelt J, Coughlin K, Crawford R, Marmot M, et al. English Longitudinal Study of Ageing: Waves 0-9, 1998-2019 [data collection]. 36th Edition. UK Data Service. SN: 5050. 2021.

22. NHS Digital. Hospital Episode Statistics (HES). 2022.
23. Office for National Statistics. Death Registration data. [Online]. 2019.
24. Rockwood K, Mitnitski A. Frailty defined by deficit accumulation and geriatric medicine defined by frailty. *Clin Geriatr Med*. 2011;27(1):17-26. doi: 10.1016/j.cger.2010.08.008.
25. Rockwood K, Mitnitski A. Frailty in relation to the accumulation of deficits. *J Gerontol A Biol Sci Med Sci*. 2007;62(7):722-7. doi: 10.1093/gerona/62.7.722.
26. Romero-Ortuno R. An alternative method for Frailty Index cut-off points to define frailty categories. *Eur Geriatr Med*. 2013;4(5):299-303. doi: 10.1016/j.eurger.2013.06.005.
27. Steptoe A, Breeze E, Banks J, Nazroo J. Cohort profile: the English Longitudinal Study of Ageing. *Int J Epidemiol*. 2013;42(6):1640-8. doi: 10.1093/ije/dys168.
28. White T, Carruthers I. NHS Jargon Explained: CRC Press; 2018.
29. NHS. NHS Data Model and Dictionary: Admission method. 2022. https://www.datadictionary.nhs.uk/attributes/admission_method.html.
30. World Health Organization. International Classification of Diseases, Tenth Revision (ICD-10). 2nd ed. 2004. <https://www.cdc.gov/nchs/icd/icd10.htm>.
31. Herbert A, Wijlaars L, Zylbersztejn A, Cromwell D, Hardelid P. Data resource profile: hospital episode statistics admitted patient care (HES APC). *Int J Epidemiol*. 2017;46(4):1093-i. doi: 10.1093/ije/dyx015.
32. Department for Work and Pensions. Benefit units in receipt of at least one income related benefit or tax credit, and the number of individuals within those benefit units, Great Britain. London. https://assets.publishing.service.gov.uk/media/5a7c0370e5274a7318b90802/IM17_22_adhoc.pdf. Published July 2012.

33. Fine JP, Gray RJ. A proportional hazards model for the subdistribution of a competing risk. *Journal of the American Statistical Association*. 1999;94(446):496-509. doi: 10.2307/2670170.
34. Austin PC, Fine JP. Practical recommendations for reporting fine-gray model analyses for competing risk data. *Stat Med*. 2017;36(27):4391-400. doi: 10.1002/sim.7501.
35. O'Halloran AM, Hartley P, Moloney D, McGarrigle C, Kenny RA, Romero-Ortuno R. Informing patterns of health and social care utilisation in Irish older people according to the Clinical Frailty Scale. *HRB Open Res*. 2021;4. doi: 10.12688/hrbopenres.13301.1.
36. Boyd PJ, Nevard M, Ford JA, Khondoker M, Cross JL, Fox C. The electronic frailty index as an indicator of community healthcare service utilisation in the older population. *Age Ageing*. 2019;48(2):273-7. doi: 10.1093/ageing/afy181.
37. Kelly S, O'Brien I, Smuts K, O'Sullivan M, Warters A. Prevalence of frailty among community dwelling older adults in receipt of low level home support: a cross-sectional analysis of the North Dublin Cohort. *BMC Geriatr*. 2017;17(1):1-11. doi: 10.1186/s12877-017-0508-2.
38. Hoogendijk EO, Muntinga ME, van Leeuwen KM, van der Horst HE, Deeg DJ, Frijters DH, et al. Self-perceived met and unmet care needs of frail older adults in primary care. *Arch Gerontol Geriatr*. 2014;58(1):37-42. doi: 10.1016/j.archger.2013.09.001.
39. Hoskovicová M, Dušek P, Sieger T, Brožová H, Zárubová K, Bezdíček O, et al. Predicting falls in Parkinson disease: What is the value of instrumented testing in off medication state? *PLoS One*. 2015;10(10):e0139849. doi: 10.1371/journal.pone.0139849.

40. Pellicer-García B, Antón-Solanas I, Ramón-Arbués E, García-Moyano L, Gea-Caballero V, Juárez-Vela R. Risk of falling and associated factors in older adults with a previous history of falls. *Int J Environ Res Public Health*. 2020;17(11):4085. doi: 10.3390/ijerph17114085.
41. Lee Y-Y, Chen C-L, Lee I-C, Lee I-C, Chen N-C. History of falls, dementia, lower education levels, mobility limitations, and aging are risk factors for falls among the community-dwelling elderly: a cohort study. *Int J Environ Res Public Health*. 2021;18(17):9356. doi: 10.3390/ijerph18179356.
42. Zaninotto P, Huang Y-T, Di Gessa G, Abell J, Lassale C, Steptoe A. Polypharmacy is a risk factor for hospital admission due to a fall: evidence from the English Longitudinal Study of Ageing. *BMC Public Health*. 2020;20:1-7. doi: 10.1186/s12889-020-09920-x.
43. Kojima G, Taniguchi Y, Kitamura A, Fujiwara Y. Is living alone a risk factor of frailty? A systematic review and meta-analysis. *Ageing Res Rev*. 2020;59:101048. doi: 10.1016/j.arr.2020.101048.
44. Ronksley PE, Sanmartin C, Quan H, Ravani P, Tonelli M, Manns B, Hemmelgarn BR. Association between perceived unmet health care needs and risk of adverse health outcomes among patients with chronic medical conditions. *Open Med*. 2013;7(1):e21. PMID: 23687534; PMCID: PMC3654502.
45. Hass Z, DePalma G, Craig BA, Xu H, Sands LP. Unmet need for help with activities of daily living disabilities and emergency department admissions among older medicare recipients. *Gerontologist*. 2017;57(2):206-10. doi: 10.1093/geront/gnv142.
46. Sands LP, Wang Y, McCabe GP, Jennings K, Eng C, Covinsky KE. Rates of acute care admissions for frail older people living with met versus unmet activity

- of daily living needs. *J Am Geriatr Soc*. 2006;54(2):339-44. doi: 10.1111/j.1532-5415.2005.00590.x.
47. Roe L, Normand C, Wren M-A, Browne J, O'Halloran AM. The impact of frailty on healthcare utilisation in Ireland: evidence from the Irish longitudinal study on ageing. *BMC Geriatrics*. 2017;17(1):1-12. doi: 10.1186/s12877-017-0579-0.
48. Abete P, Basile C, Bulli G, Curcio F, Liguori I, Della-Morte D, et al. The Italian version of the “frailty index” based on deficits in health: a validation study. *Aging Clinical And Experimental Research*. 2017;29:913-926. doi: 10.1007/s40520-017-0793-9.
49. Lang P-O, Michel J-P, Zekry D. Frailty syndrome: a transitional state in a dynamic process. *Gerontology*. 2009;55(5):539-49. doi: 10.1159/000211949.
50. Mazzella F, Cacciatore F, Galizia G, Della-Morte D, Rossetti M, Abbruzzese R, et al. Social support and long-term mortality in the elderly: role of comorbidity. *Archives of Gerontology and Geriatrics*. 2010;51(3):323-328. doi: 10.1016/j.archger.2010.01.011.

Supporting information

S1 Table. Deficit variables included in the ELSA Frailty Index.

S2 Table. Hospital Episode Statistics – Method of admission categories.

S3 Table. ICD-10 codes for fractures.

S1 Fig. Comparison between Model 1 and Model 2 in identifying the association between frailty status and level of care with unplanned admissions.

S2 Fig. Comparison between Model 1 and Model 2 identifying the association between frailty status and need for care with unplanned admissions.

S4 Table. Descriptive characteristics of the respondents (n=6,984) by level of care in ELSA wave 6 (2012/2013).

S5 Table. Descriptive characteristics of the respondents (N=6,984) by need for care in ELSA wave 6 (2012/2013).

S6 Table. The number of hospital admissions and death in each outcome. Presented are number (%)

S7 Table. Unadjusted subdistribution hazard ratio (95% CI) for the association between frailty status, level of care, need for care and each of the covariates with unplanned admissions. Unplanned admissions N=2,662, competing event deaths N=310.

S8 Table. Subdistribution hazard ratio (95% CI) for the association between frailty status, frequency for care and need for care with unplanned admissions with age group as the determinant.

S3 Fig. Subdistribution hazard ratio (95% CI) for the association between frailty status and level of care with unplanned admissions in as the determinant in each epoch of time (particular period of time)

S9 Table. Subdistribution hazard ratio (95% CI) for the association between frailty status, level of care, and need for care with unplanned admissions with varying time analysis. Adjusted for age group, gender, ethnicity, marital status, wealth and education.

S10 Table. Subdistribution hazard ratio (95% CI) for the association between frailty status and receiving care with unplanned admissions by gender. Unplanned admissions N=2,662, competing event deaths N=310. Adjusted for age group, gender, ethnicity, marital status, wealth and education.

S4 Fig. Estimates of the cumulative incidence curves of risk of unplanned hospitalisation according to frailty status and receipt of care by gender. Death was the competing risk.

The effect of levels of frailty and receipt of care on unplanned admissions and admissions due to falls and fractures

Supplementary Materials

Supplementary File 1: Additional information on statistical analysis

We checked for the presence of an interaction between frailty status, level of care, and need for care by creating a second model for each analysis. In Model 2, we created nine main dependent variables combining frailty status and level of care: [1] robust and received no care [reference group]; [2] robust and received low levels of care; [3] robust and received high levels of care; [4] prefrail and received no care; [5] prefrail and received low levels of care; [6] prefrail and received high levels of care; [7] frail and received no care; [8] frail and received low levels of care; and [9] frail and received high levels of care. For analysis of the need for care, we created eight main dependent variables combining frailty status and need for care (there were no robust respondents reporting unmet need for care needs): [1] robust and received no care [reference group]; [2] robust and received care; [3] prefrail and received no care; [4] prefrail and reported having met care needs; [5] prefrail and reported having unmet care needs; [6] frail and received no care; [7] frail and reported having met care needs; and [8] frail and reported having unmet care needs.

We looked for an interaction between frailty status, level of care and need for care on the risk of hospitalisation by plotting the SHRs and 95% CIs using both models. In order to compare with Model 2, we calculated the SHRs of each category (i.e., robust and received no care as the reference; robust and received low levels of care; robust and received high levels of care; prefrail and received no care; prefrail and received low levels of care; prefrail and received high levels of care; frail and received no care; frail and received low levels of care; and frail and received high levels of care) by adding the log of each frailty status, level of care, and need for care and then taking its exponential. An interaction effect was considered to exist if the two plots showed different values of the association of the categories and the risk of hospitalisation. **Supplementary Figures 1 and 2** show that the two plots have similar values, suggesting no interaction between frailty status and care receipt in their relationships with the risk of hospitalisation. The model without an interaction was thus preferable. Survey data was weighted using ELSA cross-sectional survey weight at wave 6.

Supplementary Table 1: Deficit variables included in the ELSA Frailty Index

Description	Assigned values (1 indicates a deficit, 0 no deficit)				
1. Difficulty with walking 100 yards	No=0	Yes=1			
2. Difficulty sitting for about two hours	No=0	Yes=1			
3. Difficulty getting up from a chair after sitting for long periods	No=0	Yes=1			
4. Difficulty climbing several flights of stairs without resting	No=0	Yes=1			
5. Difficulty climbing one flight of stairs without resting	No=0	Yes=1			
6. Difficulty stooping, kneeling, or crouching	No=0	Yes=1			
7. Difficulty reaching or extending arms above shoulder level	No=0	Yes=1			
8. Difficulty pulling or pushing large objects like a living room chair	No=0	Yes=1			
9. Difficulty lifting or carrying weights over 10 pounds, like a heavy bag	No=0	Yes=1			
10. Difficulty picking up a 5p coin from a table	No=0	Yes=1			
11. Difficulty dressing, including putting on shoes and socks	No=0	Yes=1			
12. Difficulty walking across a room	No=0	Yes=1			
13. Difficulty bathing or showering	No=0	Yes=1			
14. Difficulty eating, such as cutting up your food	No=0	Yes=1			
15. Difficulty getting in or out of bed	No=0	Yes=1			
16. Difficulty using the toilet, including getting up or down	No=0	Yes=1			
17. Difficulty using a map to figure out how to get around in a strange place	No=0	Yes=1			
18. Difficulty preparing a hot meal	No=0	Yes=1			
19. Difficulty shopping for groceries	No=0	Yes=1			
20. Difficulty making telephone calls	No=0	Yes=1			
21. Difficulty taking medications	No=0	Yes=1			
22. Difficulty managing money, (e.g. paying bills and keeping track of expenses)	No=0	Yes=1			
23. Difficulty doing work around the house or garden	No=0	Yes=1			
24. Self-reported general health	Excellent=0	V.good=0.25	Good=0.5	Fair=0.75	Poor=1
25. Whether respondent has felt depressed much of the time during the past week	No=0	Yes=1			
26. Whether respondent felt everything they did during the past week was an effort	No=0	Yes=1			
27. Whether respondent felt their sleep was restless much of the time during the past week	No=0	Yes=1			
28. Whether respondent was happy much of the time during the past week	Yes=1	No=0			
29. Whether respondent felt lonely much of the time during the past week	No=0	Yes=1			
30. Whether the respondent enjoyed life much of the time during the past week	Yes=1	No=0			
31. Whether respondent felt sad much of the time during the past week	No=0	Yes=1			

32. Whether respondent could not get going much of the time during the past week	No=0	Yes=1				
33. High blood pressure or hypertension (self-reported)	No=0	Yes=1				
34. Angina (self-reported)	No=0	Yes=1				
35. Heart attack (including MI or coronary thrombosis) (self-reported)	No=0	Yes=1				
36. Congestive heart failure (self-reported)	No=0	Yes=1				
37. An abnormal heart rhythm (self-reported)	No=0	Yes=1				
38. Diabetes or high blood sugar (self-reported)	No=0	Yes=1				
39. A stroke (cerebral vascular disease) (self-reported)	No=0	Yes=1				
40. Chronic lung disease such as chronic bronchitis or emphysema (self-reported)	No=0	Yes=1				
41. Asthma (self-reported)	No=0	Yes=1				
42. Arthritis (including osteoarthritis , or rheumatism) (self-reported)	No=0	Yes=1				
43. Osteoporosis, sometimes called thin or brittle bones (self-reported)	No=0	Yes=1				
44. Cancer or a malignant tumor (excluding minor skin cancers) (self-reported)	No=0	Yes=1				
45. Parkinson's disease (self-reported)	No=0	Yes=1				
46. Any emotional, nervous or psychiatric problems (self-reported)	No=0	Yes=1				
47. Alzheimer's disease (self-reported)	No=0	Yes=1				
48. Dementia, organic brain syndrome, senility or any other serious memory impairment (self-reported)	No=0	Yes=1				
49. Self-reported eyesight (while using lenses, if appropriate)	Excellent=0	V.good=0.2	Good=0.4	Fair=0.6	Poor=0.8	Blind=1
50. Self-reported hearing (while using hearing aid if appropriate)	Excellent=0	V.good=0.25	Good=0.5	Fair=0.75	Poor=1	
51. Whether respondent has fallen down at all /last year /last 2years	No=0	Yes=1				
52. Whether respondent has fractured hip ever /in last 2 years	No=0	Yes=1				
53. Whether respondent has had joint replacement ever	No=0	Yes=1				
54. Identify today's date: day of month	Yes=0	No=1				
55. Identify today's date: month	Yes=0	No=1				
56. Identify today's date: year	Yes=0	No=1				
57. Identify the day of the week?	Yes=0	No=1				
58. Immediate word recall (sample organized into quartiles)	1st quintile=0	2nd=0.3	3rd=0.6	4th quintile=1		
59. Delayed word recall (sample organized into quintiles)	1st quintile=0	2nd=0.25	3rd=0.5	4th=0.75	5th quintile=1	
60. Have pain while performing the walking test	Yes=0	No=1				

Supplementary Table 2: Hospital Episode Statistics – Method of admission categories

Code	Method of admission
11	Waiting list
12	Booked
13	Planned
21	Accident and Emergency
22	GP-after request of immediate admission
23	Bed bureau
24	Consultant clinic
25	Mental Health Crisis Resolution team
28	Other: A&E
31	Admitted ante partum
32	Admitted post-partum
81	Transfer of any admitted patient from other hospital provider other than in an emergency
82	Baby birth
83	Baby born outside
99	Not known

Supplementary Table 3: ICD-10 codes for fractures

Code	Method of admission
M 484	Fatigue fracture of vertebra
M 495	Collapsed vertebra in diseases classified elsewhere
M 80	Osteoporosis with pathological fracture
M 843	Stress fracture, not elsewhere classified
M 844	Pathological fracture, not elsewhere classified
M 907	Fracture of bone in neoplastic disease
M 966	Fracture of bone following insertion of orthopaedic implant, joint prosthesis or bone plate
S 02	Fracture of skull and facial bones
S 12	Fracture of neck
S 22	Fracture of rib(s), sternum and thoracic spine
S 32	Fracture of lumbar spine and pelvis
S 42	Fracture of shoulder and upper arm
S 52	Fracture of forearm
S 62	Fracture of wrist and hand level
S 72	Fracture of femur
S 82	Fracture of lower leg, including ankle
S 92	Fracture of upper limb, except ankle
T 02	Fractures involving multiple body regions
T 08	Fracture of spine, level unspecified
T 10	Fracture of upper limb, level unspecified
T 12	Fracture of lower limb, level unspecified
T 142	Fracture of unspecified body region

Supplementary Table 4: Descriptive characteristics of the respondents (n=6,984) by level of care in ELSA wave 6 (2012/2013).

	No care	Receiving low levels of care	Receiving high levels of care
<i>Age group, n (%)</i>			
60-64	1,540 (86.7)	145 (8.2)	89 (5.0)
65-69	1,372 (83.1)	159 (9.6)	118 (7.1)
70-74	962 (79.1)	142 (11.6)	112 (9.1)
75-79	707 (70.7)	181 (18.0)	112 (11.2)
80-84	407 (56.5)	190 (26.3)	123 (17.0)
85+	210 (33.8)	240 (38.7)	170 (27.4)
<i>Gender, n (%)</i>			
Men	2,617 (81.1)	306 (9.5)	301 (9.3)
Women	2,577 (68.5)	755 (20.0)	427 (11.3)
<i>Ethnicity, n (%)</i>			
White	5,039 (74.6)	1,018 (15.0)	690 (10.2)
Non-White	154 (65.4)	43 (18.3)	38 (16.1)
<i>Married, n (%)</i>			
No	1,517 (62.2)	629 (25.8)	292 (11.9)
Yes	3,675 (80.8)	423 (9.5)	436 (9.5)
<i>Education attainment, n (%)</i>			
Less than secondary school	1,598 (63.5)	532 (21.1)	384 (15.2)
Secondary school	995 (79.1)	165 (13.1)	97 (7.7)
College or higher	2,602 (80.9)	365 (11.3)	247 (7.6)
<i>Wealth, n (%)</i>			
1 st quintile (least wealthy)	749 (62.5)	250 (20.8)	199 (16.6)
2 nd	978 (63.7)	325 (21.1)	232 (15.1)
3 rd	1,073 (75.7)	212 (15.0)	131 (9.2)
4 th	1,173 (82.6)	155 (10.9)	92 (6.4)
5 th quintile (most wealthy)	1,147 (86.1)	116 (8.7)	69 (5.1)

Supplementary Table 5: Descriptive characteristics of the respondents (N=6,984) by need for care in ELSA wave 6 (2012/2013).

	No care	Met care needs	Unmet care needs
<i>Age group, n (%)</i>			
60-64	1,553 (87.5)	195 (11.0)	26 (1.4)
65-69	1,397 (84.6)	240 (14.5)	13 (0.7)
70-74	977 (80.3)	224 (18.4)	15 (1.2)
75-79	733 (73.3)	247 (24.7)	19 (1.9)
80-84	428 (59.3)	275 (38.1)	18 (2.4)
85+	228 (36.7)	360 (58.0)	32 (5.2)
<i>Gender, n (%)</i>			
Men	2,646 (82.0)	544 (16.8)	34 (1.0)
Women	2,666 (70.9)	1,005 (26.7)	89 (2.3)
<i>Ethnicity, n (%)</i>			
White	5,153 (76.3)	1,480 (21.9)	115 (1.7)
Non-White	159 (67.3)	68 (29.0)	9 (3.6)
<i>Married, n (%)</i>			
No	1,590 (81.8)	767 (31.4)	82 (3.3)
Yes	3,720 (65.1)	782 (17.2)	42 (0.9)
<i>Education attainment, n (%)</i>			
Less than secondary school	1,658 (65.9)	790 (31.4)	65 (2.6)
Secondary school	1,015 (80.7)	223 (17.7)	19 (1.5)
College or higher	2,638 (82.0)	536 (16.6)	39 (1.2)
<i>Wealth, n (%)</i>			
1 st quintile (least wealthy)	780 (65.0)	378 (31.5)	40 (3.3)
2 nd	1,011 (65.9)	479 (31.2)	44 (2.8)
3 rd	1,098 (77.5)	297 (20.9)	21 (1.4)
4 th	1,190 (83.8)	221 (15.5)	9 (0.6)
5 th quintile (most wealthy)	1,157 (86.8)	165 (12.4)	9 (0.7)

Supplementary Table 6 The number of hospital admissions and death in each outcome.
Presented are number (%)

Analysis outcomes	Number (%) of participants admitted to the hospital during the follow-up	Number (%) of participants died during the follow-up
Unplanned admissions	2663 (37.78)	310 (4.05)
Admissions due to fall	586 (7.65)	939 (12.26)
Admission due to fracture	432 (5.64)	1020 (13.32)

Supplementary Table 7: Unadjusted subdistribution hazard ratio (95% CI) for the association between frailty status, level of care, need for care and each of the covariates with unplanned admissions.

	Unadjusted SHRs (95% CIs)
<i>Frailty status, reference: robust</i>	
Prefrail	1.80 (1.64; 1.97)
Frail	2.74 (2.47; 3.03)
<i>Level of care, reference: no care</i>	
Received low levels of care	1.70 (1.55; 1.87)
Received high levels of care	1.82 (1.64; 2.02)
<i>Need for care, reference: no care</i>	
Met care needs	1.80 (1.66; 1.95)
Unmet care needs	2.07 (1.61; 2.67)
Age (years)	1.06 (1.06; 1.07)
Women (vs Men)	0.98 (0.92; 1.06)
Non White (vs White)	1.08 (0.86; 1.36)
Married (vs Non married)	0.73 (0.68; 0.79)
<i>Wealth, reference: 1st quintile (least wealthy)</i>	
2 nd	0.97 (0.87; 1.09)
3 rd	0.77 (0.68; 0.87)
4 th	0.71 (0.63; 0.80)
5 th quintile (most wealthy)	0.56 (0.50; 0.64)
<i>Education, reference: less than high school</i>	
High school	0.81 (0.73; 0.90)
College or higher	0.73 (0.67; 0.79)

Note: Unplanned admissions N=2,662, competing event deaths N=310.

Supplementary Table 8: Subdistribution hazard ratio (95% CI) for the association between frailty status, frequency for care and need for care with unplanned admissions with age group as the determinant

	Level of care	Need for care
<i>Frailty status, reference: robust</i>		
Prefrail	1.75 (1.58; 1.93)	
Frail	2.48 (2.15; 2.85)	
<i>Level of care, reference: no care</i>		
Received low levels of care	1.20 (1.07; 1.35)	
Received high levels of care	1.34 (1.17; 1.55)	
<i>Need for care, reference: no care</i>		
Met care needs		1.25 (1.13; 1.39)
Unmet care needs		1.26 (0.95; 1.66)
<i>Age group, reference: 60-64</i>		
65-69	1.16 (1.01; 1.33)	1.17 (1.02; 1.34)
70-74	1.75 (1.53; 2.01)	1.75 (1.53; 2.01)
75-79	2.23 (1.95; 2.55)	2.22 (1.94; 2.55)
80-84	2.58 (2.22; 3.00)	2.58 (2.22; 3.00)
85+	3.06 (2.60; 3.60)	3.05 (2.59; 3.59)
Women (vs Men)	0.76 (0.70; 0.83)	0.76 (0.70; 0.83)
Non White (vs White)	1.25 (0.96; 1.62)	1.26 (0.97; 1.63)
Married (vs Non Married)	0.90 (0.83; 0.99)	0.90 (0.83; 0.99)
<i>Wealth, reference: 1st quintile (least wealthy)</i>		
2 nd	0.90 (0.79; 1.01)	0.90 (0.80; 1.01)
3 rd	0.79 (0.69; 0.89)	0.78 (0.69; 0.89)
4 th	0.77 (0.67; 0.88)	0.77 (0.67; 0.87)
5 th quintile (most wealthy)	0.67 (0.59; 0.78)	0.68 (0.59; 0.78)
<i>Education, reference: less than high school</i>		
High school	1.04 (0.92; 1.16)	1.04 (0.92; 1.16)
College or higher	0.98 (0.90; 1.08)	0.98 (0.90; 1.08)

Supplementary Table 9: Subdistribution hazard ratio (95% CI) for the association between frailty status, level of care, and need for care with unplanned admissions with varying time analysis. Adjusted for age group, gender, ethnicity, marital status, wealth and education.


	Level of care	Need for care
<i>Frailty status, reference: robust</i>		
Prefrail	1.95 (1.73; 2.20)	
Frail	3.19 (2.72; 3.74)	
<i>Level of care, reference: no care</i>		
Received low levels of care	1.19 (1.04; 1.35)	
Received high levels of care	1.41 (1.21; 1.64)	
<i>Need for care, reference: no care</i>		
Met care needs		1.27 (1.13; 1.43)
Unmet care needs		1.34 (1.01; 1.78)
Age	1.04 (1.04; 1.05)	1.04 (1.04; 1.05)
Women (vs Men)	0.76 (0.69; 0.83)	0.76 (0.69; 0.83)
Non White (vs White)	1.33 (0.99; 1.79)	1.34 (0.99; 1.81)
Married (vs Non Married)	0.90 (0.81; 0.99)	0.90 (0.81; 0.99)
<i>Wealth, reference: 1st quintile (least wealthy)</i>		
2 nd	0.85 (0.74; 0.97)	0.85 (0.75; 0.97)
3 rd	0.80 (0.70; 0.92)	0.80 (0.70; 0.92)
4 th	0.77 (0.66; 0.89)	0.77 (0.66; 0.89)
5 th quintile (most wealthy)	0.68 (0.58; 0.80)	0.68 (0.58; 0.80)
<i>Education, reference: less than high school</i>		
High school	1.04 (0.92; 1.18)	1.04 (0.92; 1.19)
College or higher	0.98 (0.89; 1.09)	0.98 (0.89; 1.09)

Supplementary table 10 Subdistribution hazard ratio (95% CI) for the association between frailty status and receiving care with unplanned admissions by gender


	Male	Female
<i>Frailty status, reference: robust</i>		
Prefrail	1.73 (1.51; 1.98)	1.79 (1.54; 2.09)
Frail	2.39 (1.93; 2.94)	2.57 (2.12; 3.11)
<i>Receiving care, reference: no</i>		
Yes	1.30 (1.09; 1.54)	1.31 (1.14; 1.50)

Note: Unplanned admissions N=2,662, competing event deaths N=310. ^aAdjusted for age group, gender, ethnicity, marital status, wealth and education.


Supplementary Figure 1: Comparison between Model 1 and Model 2 in identifying the association between frailty status and level of care with unplanned admissions



Supplementary Figure 2: Comparison between Model 1 and Model 2 identifying the association between frailty status and need for care with unplanned admissions




Supplementary Figure 3: Subdistribution hazard ratio (95% CI) for the association between frailty status and level of care with unplanned admissions in as the determinant in each epoch of time (particular period of time)



Notes: Epochs: [1] wave 6 as the baseline with 6 months follow-up; [2] wave 6 baseline with 12 months follow-up; [3] wave 7 baseline with 6 months follow-up; [4] wave 7 baseline with 12 months follow-up; and [5] wave 8 baseline; 6 months follow-up.

Supplementary Figure 4 Estimates of the cumulative incidence curves of risk of unplanned hospitalisation according to frailty status and receipt of care by gender. Death was the competing risk.

A. Male



B. Female

