

Advancements in Predictive Microbiology

Taiwo, Oluseyi Rotimi; Onyeaka, Helen; Oladipo, Elijah K; Oloke, Julius Kola; Chukwugozie, Deborah C

DOI:

[10.1155/2024/6612162](https://doi.org/10.1155/2024/6612162)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Taiwo, OR, Onyeaka, H, Oladipo, EK, Oloke, JK & Chukwugozie, DC 2024, 'Advancements in Predictive Microbiology: Integrating New Technologies for Efficient Food Safety Models', *International journal of microbiology*, vol. 2024, 6612162. <https://doi.org/10.1155/2024/6612162>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Review Article

Advancements in Predictive Microbiology: Integrating New Technologies for Efficient Food Safety Models

Oluseyi Rotimi Taiwo,¹ Helen Onyeaka ,² Elijah K. Oladipo,^{1,3} Julius Kola Oloke,⁴ and Deborah C. Chukwugozie⁵

¹Genomics Unit, Helix Biogen Institute, Ogbomosho, Oyo, Nigeria

²School of Chemical Engineering, University of Birmingham, Edgbaston B15 2TT, Birmingham, UK

³Department of Microbiology, Laboratory of Molecular Biology, Immunology and Bioinformatics, Adeleke University, Ede, Osun, Nigeria

⁴Department of Natural Science, Microbiology Unit, Precious Cornerstone University, Ibadan, Oyo, Nigeria

⁵Department of Microbiology, Federal University Otuoke, Otuoke, Bayelsa, Nigeria

Correspondence should be addressed to Helen Onyeaka; h.onyeaka@bham.ac.uk

Received 13 July 2023; Revised 1 April 2024; Accepted 23 April 2024; Published 17 May 2024

Academic Editor: Todd R. Callaway

Copyright © 2024 Oluseyi Rotimi Taiwo et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Predictive microbiology is a rapidly evolving field that has gained significant interest over the years due to its diverse application in food safety. Predictive models are widely used in food microbiology to estimate the growth of microorganisms in food products. These models represent the dynamic interactions between intrinsic and extrinsic food factors as mathematical equations and then apply these data to predict shelf life, spoilage, and microbial risk assessment. Due to their ability to predict the microbial risk, these tools are also integrated into hazard analysis critical control point (HACCP) protocols. However, like most new technologies, several limitations have been linked to their use. Predictive models have been found incapable of modeling the intricate microbial interactions in food colonized by different bacteria populations under dynamic environmental conditions. To address this issue, researchers are integrating several new technologies into predictive models to improve efficiency and accuracy. Increasingly, newer technologies such as whole genome sequencing (WGS), metagenomics, artificial intelligence, and machine learning are being rapidly adopted into newer-generation models. This has facilitated the development of devices based on robotics, the Internet of Things, and time-temperature indicators that are being incorporated into food processing both domestically and industrially globally. This study reviewed current research on predictive models, limitations, challenges, and newer technologies being integrated into developing more efficient models. Machine learning algorithms commonly employed in predictive modeling are discussed with emphasis on their application in research and industry and their advantages over traditional models.

1. Introduction

Food safety effectively refers to all measures to guarantee that only food suitable for human consumption is passed on to the human population. Generally, this includes strategies, policies, and legislation to protect food from microbial, chemical, physical, and allergenic contamination during food production, distribution, and consumption [1]. Globally, food pollutants such as heavy metals, mycotoxins, molds, antibiotics, and pesticide residues are

becoming increasingly prevalent [2]. The burden of food-borne diseases remains significant with an estimate 600 million illnesses and 420,000 deaths annually [3]. Moreover, the landscape of food safety is evolving due to changing dietary patterns, the industrialization of food production, and the effects of climate change. These factors present new challenges in combating food-borne infections, leading to a rapid increase in cases of foodborne diseases that have adverse effects on public health and the global economy [4].

Moreover, the landscape of food safety is evolving due to changing dietary patterns, the industrialization of food production, and the effects of climate change. These factors present new challenges in combating foodborne infections, leading to a rapid increase in infectious diseases that have adverse effects on public health and the global economy [4].

Globally, food safety measures have evolved significantly to protect consumers on a large scale. In the 19th century, the recognition of the germ theory of disease prompted a shift toward prioritizing sanitation and animal health, leading to the establishment of enhanced safety standards for dairy products. These advancements quickly permeated other sectors of the food industry [5]. The rise of industrialization and mass food processing industries underscored the importance of public health measures aimed at guaranteeing the safety of food consumed by the public. Today, rigorous screening protocols are in place to mitigate microbial hazards (such as pathogenic microbes, metabolites, and mycotoxins), chemical hazards (including heavy metals, pesticides, and carcinogens), as well as allergens and physical hazards like sharp objects [6–9]. Biological hazards pose the most significant threat among these, often leading to devastating outbreaks [10]. According to a review by Lee and Yoon [11], norovirus is associated with the highest number of foodborne morbidities per year globally. It is followed by *Campylobacter*, *Salmonella*, and *Listeria monocytogenes*. Also, a WHO report showed that foodborne bacteria illnesses were seen to have more prevalence than viral and parasitic diseases [12].

Historically, classical food safety measures employed a hazard-based approach relying heavily on a routine inspection of processing facilities and end-product sampling [13, 14]. However, these techniques are highly fallible due to their subjective nature. Hazard-based meat inspection techniques, for example, are based on physical inspection of animal carcasses by incision and palpation to observe for signs of internal organ pathology and lymphadenopathy, which can indicate foodborne pathogens [15, 16]. However, this method may not be effective in detecting diseases such as colibacillosis, campylobacteriosis, or prion-related diseases (e.g., Bovine spongiform encephalopathy) or early infection [17, 18]. Furthermore, invasive practices such as palpation and incision of carcasses are currently being discouraged in most developing countries in favor of a risk-based approach due to their potential to cause cross-contamination of foodborne pathogens [19].

Another hazard-based approach to food safety is the end testing of food products [20]. However, this presents several challenges; first, it is a reactive rather than a proactive approach; in other words, this method is used to detect microbial contamination rather than prevent it from occurring in the first place. Second, negative results (indicating no microbial contamination) do not guarantee that the product is entirely free from contamination [21]. Moreover, this method, which is predominantly carried out through microbial culture techniques based has limited sensitivity and is limited by the microbial load present; consequently, the absence of a particular microbe during testing does not necessarily mean they were never present or will not appear

later. Due to their limited sensitivity, culture-based methods of end testing are also prone to false positives and false negatives and are currently being replaced, or in some cases, augmented by molecular detection methods (such as whole genome sequencing) which are more accurate [22]. Hazard-based end testing of food is also labor intensive and costly due to the expenses associated with the technique and the potential losses incurred upon detecting microbial contamination [23].

Moreover, relying solely on a hazard-based approach can be impractical and unsustainable. Strict enforcement of this approach may prohibit essential and nutritious foods if they contain naturally occurring toxins, such as genotoxic hydrazines in mushrooms, glycoalkaloids in tomatoes and potatoes, and hydrocyanic acid in cassava. Furthermore, traditional food processing methods like frying potatoes have been linked to producing carcinogens such as acrylamide. Banning these food items and cooking techniques outright may be unrealistic, particularly in developing countries. Such a ban could significantly disrupt food availability and consumption [24].

As part of measures towards overcoming these challenges with hazard-based approaches, the World Trade Organization (WTO) and the Codex Alimentarius Commission (CAC), in collaboration with different public health bodies and national governments, came up with a regulatory risk-based framework on food hygiene, processing, and practices in the food industry. The Codex Alimentarius Commission was founded in 1963 as an intergovernmental body under the Food and Agriculture Organization United Nations and the World Health [25]. During the 19th and 20th sessions of the CAC in 1991 and 1993, the decision was taken to base food safety decisions on risk assessment; the committee also accepted a recommendation to harmonize standard-setting methodologies. The drafting of the Codex Alimentarius document on “Guidelines for the application of the Hazard Analysis Critical Control Point (HACCP) system” (Codex, 1993) and the subsequent edition, “Hazard Analysis and Critical Control Point System and Guidelines for its Application” [26] played a key role in the adoption of HACCP protocols in food safety globally [27, 28].

Towards the end of protecting consumers, a risk analysis framework prescribing regulations such as the Appropriate Level of Protection (ALOP) and the Food Safety Objective (FSO) was developed. The ALOP, also known as ALR (Acceptable Level of Risk), is a concept used in risk analysis to establish the appropriate level of protection against hazards in food production and ensure the safety of food for human consumption. It is described as “the level of protection deemed appropriate by the member establishing a sanitary or phytosanitary measure to protect the human, animal, and plant life or health within its territory” [29]. The determination of the ALOP or ALR is based on a combination of scientific knowledge and expert judgment, taking into account factors such as the severity and likelihood of harm, the availability and effectiveness of risk mitigation measures, societal values, and expectations, as well as epidemiologic surveillance data on the occurrence and distribution of foodborne illnesses.

The Food Safety Objective (FSO) is a metric that describes the maximal threshold of a microbial hazard in a food at the point of consumption; it was designed to translate the ALOP into practical measures for stakeholders in food manufacturing and distribution [1, 30]. Maximal threshold hazard levels at other points on the food processing chain are called performance objectives (POs) [31]. Other regulatory standards developed in line with the risk-based safety approaches include GAP (good agricultural production practices), GHP (good hygiene practices), GMP (good manufacturing practices), and HACCP [32].

Adopting risk-based approaches in food safety has shifted objectives towards preventing hazards rather than relying solely on inspection-based methods [33]. This shift acknowledges the importance of proactively identifying and mitigating risks to food safety. While these approaches have contributed to improved food safety globally, there is a growing recognition of the need for increased efficiency, precision, and accuracy in food safety practices [34]. As a result, researchers are turning to mathematical microbial modeling to complement existing methods [35]. Predictive microbiology, in particular, has emerged as a valuable tool.

Microbial predictive modeling involves using mathematical models to predict the growth and behavior of microorganisms in food products under different environmental conditions [36]. This field of study is focused on developing models that can help food manufacturers and regulators understand how microorganisms respond to changes in factors such as temperature, pH, water activity, and other environmental conditions. The objective is to utilize this knowledge in forecasting the potential growth of harmful foodborne microorganisms and to predict the potential growth of harmful microorganisms that can cause foodborne illness [37, 38]. Predictive microbiology is established on the notion that the behavior of microorganisms, however complex, can be simulated and effectively reproduced in nonbiological models by applying mathematical and computational principles [35]. By utilizing mathematical models, predictive microbiology enables the identification of microorganisms of interest and provides insights into microbial ecology, sources of contamination, and potential contamination points. Predictive modeling plays a crucial role in ensuring food safety by allowing food manufacturers and regulatory agencies to anticipate and control the growth and behavior of microorganisms in food products. The goal of predictive microbiology is to provide tools that can be used to design food processing and preservation strategies that can help reduce the risk of foodborne illness and improve the safety and quality of food products.

The application of predictive microbiology has found its way into diverse areas of food microbiology, such as analysis of food formulation processes and their effect on shelf life, evaluation of processing operations, and hazard analysis critical control point (HACCP).

Currently, the advancement of modeling techniques in food safety is heavily influenced by the integration of artificial intelligence (AI) and other cutting-edge technologies, such as whole genome sequencing [39, 40]. Machine

learning algorithms such as random forest models and support vector machines (SVM) are becoming integral components of these evolving models [41, 42]. Whole genome sequencing (WGS) technologies, alongside genomics and other omics-based methods, generate vast amounts of high-throughput data. These data serve as invaluable resources for training models using various AI algorithms [43]. By harnessing the power of AI, researchers can effectively analyze and interpret complex datasets to derive meaningful insights. These sophisticated modeling tools find applications across diverse domains within food safety, notably in risk assessment and management.

The use of predictive models in food safety complements risk-based safety approaches; it also has several advantages over traditional microbiological testing, including the following.

Enhanced speed and accuracy: predictive models can quickly and accurately predict the growth and behavior of microorganisms in food products, allowing for faster and more accurate risk assessments and decision-making.

Improved risk assessment: predictive models can provide more detailed and accurate risk assessments of food products, helping to identify potential hazards and develop effective control measures.

Cost savings: using predictive models can reduce the need for expensive and time-consuming microbiological testing, saving both time and money.

Optimization of food processing and preservation methods: predictive models can optimize food processing and preservation methods, ensuring that food products are safe, high quality, and have a longer shelf life.

This review aims to provide a comprehensive overview of the field of predictive microbiology and its applications in improving food safety. It also addresses current limitations and challenges in this field and explores emerging trends and approaches, including the use of new technologies such as genomics and metagenomics.

1.1. Factors Affecting Microbial Growth in Food. During the preharvest stage, food crops are vulnerable to contamination in the field [44]. This can happen through contact with waterborne pathogens during irrigation, exposure to soil-borne pathogens present in manure, or contact with wild-life such as migratory birds [45]. The harvest stage is another critical point where contamination can occur. Food crops are susceptible to contamination as they are picked by workers, who may introduce pathogens through improper handling practices [46]. Postharvest processes, including transportation, processing, distribution, and preparation, also present opportunities for microbial contamination. Food can come into contact with soil, air, and waterborne microorganisms during these stages, increasing the risk of contamination [45]. In animal-based foods like beef, microbial contamination can originate from high microbial loads present in the cattle gut [47]. During carcass preparation, there is a risk of transferring these pathogens to skeletal muscle, further emphasizing the importance of stringent hygiene practices throughout the food processing chain [48].

Most foods provide an ideal nutrition source for microorganisms [49]. Generally, factors influencing microbial growth in food are classified into intrinsic and extrinsic factors. Intrinsic factors are physicochemical characteristics inherent to the food, such as composition, pH, water activity, oxidation-reduction potential, and biological structures. On the other hand, extrinsic factors encompass environmental conditions surrounding the food, including temperature, relative humidity, gaseous environments, and competitor microorganisms [49].

Intrinsic factors play a pivotal role in microbial growth dynamics. Foods of plant or animal origin possess biological components that act as barriers against microbial invasion and colonization. Examples include the tough testa and shells of certain seeds and nuts and the thin membrane found in eggs. In addition, readily available nutrient sources in food significantly influence microbial proliferation. Bacteria typically require simple sugars and free amino acids for metabolism, favoring their growth in nutrient-rich environments [50]. Conversely, microbial growth can be hindered by natural antimicrobial substances such as bacteriocins, allicins, and eugenol [51]. Extremes of pH, such as highly alkaline or acidic conditions, also deter bacterial growth as most bacteria thrive within a narrow pH range compared to molds and yeasts [52].

Considering the water activity requirements of microbes, it is notable that most bacteria are halophiles (salt loving). Gram-negative bacteria typically exhibit a greater affinity for water than Gram-positive bacteria, molds, and yeasts. Consequently, fast-growing bacteria are often the primary spoilers of fresh foods with high water content [50].

Moreover, additional factors influencing bacterial growth in food encompass implicit factors related to the microbes, including interactions between different microbial species colonizing the food. These interactions may involve differing nutrient utilization, stress thresholds, and the production of chemicals that modulate the growth of other microorganisms. Processing factors, such as heating, cooling, and drying treatments, also significantly affect food composition and microbial ecology. Ultimately, a complex interplay between all the factors above dictates microbial growth dynamics in food systems [53].

2. Principles of Predictive Microbiology

The basic principle of predictive microbiology is anchored on the knowledge that microbial growth and survival can be simulated in mathematical models.

The effectiveness of mathematical growth models is affected by different criteria. More excellent resolution and predictive relevance can be achieved by reducing the number of parameters under analysis. Furthermore, a mechanistic approach assessing few meaningful parameters allows for more phases of microbial growth to be covered and consequently increases the model's accuracy [54].

2.1. Microbial Growth Models. The growth curve of most foodborne bacteria typically exhibits the following four distinct phases: an initial lag phase characterized by minimal or no detectable growth, followed by an exponential (log) phase marked by rapid cell division, then a plateau phase where growth stabilizes, and finally, a mortality phase where conditions become unfavorable for further growth [55].

Microbial models are classified based on a number of different criteria. They can be grouped into kinetic and probabilistic models based on the expected microbial response; empirical and mechanistic models based on the method of analysis; and primary, secondary, and tertiary models based on the dependent variable assessed [29, 56, 57].

2.1.1. Primary Models. Primary models assess bacterial response over time to specific conditions. Typically, these models aim to describe microbial growth using as few parameters as possible [58]. Primary models may be empirical, rate growth models, inactivation or survival models, or a combination thereof. Examples include the Gompertz and Buchanan models [59]. These models evaluate a discrete number of intuitive parameters, such as the relative growth rate, initial population size, and asymptotic population size. However, they may be unable to assess all microbial growth phases or account for variables such as prolonged lag phases [54].

The Gompertz equation is commonly written mathematically as

$$L = A + C \exp \{-\exp [B(T - M)]\} \quad [60].$$

It describes an asymmetric sigmoidal curve.

$L = \log+$ of bacteria count in colony forming units(cfu); $t =$ time in hours; $A \sim$ asymptotic log count with a decrease in time indefinitely; C is asymptotic log count with an indefinite increase in time; M is the time at the maximal growth rate; and B is the relative growth rate at time M .

This model has been used by the Food MicroModel consortium and the Pathogen Modelling Program (PMP) group in the UK and the US, respectively [61]. The model, however, is limited in its ability to describe the exponential phase. It also overestimates parameter values [62–64].

The Buchanan model:

$$\log(N) = \begin{cases} \log(N) & (t \leq r), \\ \log(N_0) + \mu_{\max} (t - r) & (r < t < t_m), \\ \log(N_{\max}) & (t \geq t_m). \end{cases} \quad (1)$$

$$\tau = \tau_i + t_m \quad i = 1, 2, \dots, N, \quad (2)$$

$$\mu_{\max} = \frac{\log(N_{\max}) - \log(N_0)}{t_m - \tau}. \quad (3)$$

Equations (1)–(3) express assumptions by the model in simulating growth. The Buchanan model assumes that there is zero growth during the lag phase, and growth during the exponential phase increases linearly with time and specific growth in the plateau phase is zero.

The Buchanan model was developed based on the following three assumptions: (1) during the lag phase, the specific growth rate is zero; (2) during the exponential phase, the logarithm of the bacterial population increases linearly with time; and (3) during the stationary phase, the specific growth rate is zero [65]. It is also regarded as a very simple model, making it very popular for microbiological modeling [65].

2.1.2. Secondary Models. Secondary models analyze the factors influencing the kinetic parameters identified by primary models. They establish the relationship between primary model parameters and intrinsic and extrinsic factors such as temperature and pH [66]. While primary models focus on estimating changes in microbial population over time and observing specific responses such as growth rate and lag phase, secondary models assess the impact of intrinsic and extrinsic factors of the food on these responses [29]. Several secondary models have been developed to evaluate the lag phase and growth rate concerning one or more environmental or physicochemical factors [58, 67]. Examples of secondary models include the response surface (RS) models and Arrhenius models [68]. A response surface model in predictive microbiology is a mathematical model used to describe the relationship between multiple independent variables (factors) and a response variable, typically microbial growth or survival, in a complex environment [69]. Response surface models (RSMs) are crucial in predictive microbiology. They help us understand how different factors such as temperature, pH, and humidity affect microbial growth or survival [70, 71]. These models show how these variables work together, impacting the microorganisms' environment. RSM works by mapping out a multidimensional space to predict outcomes based on various factors. It involves conducting experiments where these factors are varied systematically and then using math and statistics to build a model representing the response across all the experiments [72]. This model's surface guides us to find the best conditions for controlling microbial growth or death. In predictive microbiology, RSM is used to fine-tune food storage and safety. It helps find the right mix of conditions, such as temperature and moisture, to prevent harmful microbes from growing in food. For example, RSM can help determine the exact temperature and humidity levels needed to prevent pathogen growth in food products [70].

2.1.3. Tertiary Models. Tertiary models are algorithm-run software packages combining primary and secondary models with a Graphic User Interface (GUI), making them userfriendly for amateur modelers. They are primarily used in the food industry and research to consolidate findings from primary and secondary models. The Unified Growth

Prediction Model (UGPM) software is an example of a tertiary model based on Baranyi Roberts's primary and temperature-dependent secondary models [73].

2.2. Empirical and Mechanistic Models

2.2.1. Empirical Models. Empirical models are pragmatic, rigid models in which relationships between different parameters are put forth as mathematical equations, usually in first or second-degree polynomial mathematical relationships [58, 68]. In an empirical model, predictions are made without considering variables such as physicochemical parameters capable of influencing the outcomes of predictions. An example is the quadratic response surface used by Gibson et al. [74].

2.2.2. Mechanistic or Deterministic Models. Mechanistic or deterministic models are established on theories and allow interpretation of the response in terms of known phenomena and processes. These models typically contain fewer parameters, fit the data better, and describe the response better. They are also known to extrapolate more effectively and are preferable to empirical models [66].

2.3. Kinetic and Probabilistic Models

2.3.1. Kinetic Models. Kinetic models determine the expected response rates (growth or death). They are used to predict concentration levels associated with a certain level of microbial strain and thus characterize risk (infection or intoxication-associated risk) [58]. Examples are the Gompertz and square root models, which describe rates of response, such as lag time, specific growth rate, and maximum population density, or inactivation/survival models that describe destruction or survival over time [75–78].

2.3.2. Probability Models. Probability models, on the contrary, are associated only with the probability of growth or toxin production and do not predict the rate at which this occurs [79, 80]. They are used to show the absolute limit of microbial growth within specific environments and demonstrate stress threshold levels, which may limit growth but ultimately permit it [79, 80].

2.4. Applications of Predictive Microbiology in Food Safety

2.4.1. Quality Control of Food Products. Predictive microbiology can be applied to validate the effectiveness of microbial inactivation processes such as drying, heat treatment, and refrigeration. Food processing industries producing yogurt, milk, wine, and sous-vide processed products require strict adherence to specific refrigeration temperature and heat treatment [81–83]. However, these treatment processes are often inadequate or improperly managed, leading to microbial growth. In the dairy industry, for instance, the insufficiency of heat treatments solely as the control measure against fungal spore germination has been

highlighted; this is due to the ability of some spore-forming fungal pathogens to survive pasteurization or thrive at low temperatures. Organisms such as *Bacillus sporothermourens* are known to survive the ultrahigh temperatures reached during heat treatments [84, 85]. Several research studies on cold chains have also revealed severe noncompliance with regulations at distribution, retail, and storage with the final consumer. This, therefore, necessitates the need for quality control points along the food chain.

A study by Gougouli et al. [86] demonstrated that microbial growth in yogurt, evidenced by signs like mycelial formation, is significantly influenced by variables such as warehousing duration, storage temperature, and microbial strain. Consequently, predictive models capable of anticipating fungal growth by analyzing these factors are highly valuable in the dairy industry. These models can enhance quality control measures by integrating them into the final testing of yogurt products for fungal growth before market release. In such applications, predictive models aid in determining optimal conditions for the growth of various microbial species in end-product challenge testing.

It is essential to recognize that different microbial contaminants have varying conditions permissive for growth. Therefore, the selection of environmental conditions during end-challenge tests cannot be made arbitrarily without a solid scientific basis, as these can lead to false negative tests and inadvertent marketing of contaminated products. Other predictive models in the dairy industry have modeled the influence of parameters such as temperature, pH, water activity, and inoculum size on the growth of *Listeria monocytogenes* in milk as well as the growth of *Yersinia enterocolitica* in Camembert-type cheese [87].

2.4.2. Risk Assessment and Management. Risk assessment is a systematic, scientific method designed to calculate human risk associated with exposure to foodborne risks [26]. It comprises the following four sequential stages: hazard identification, exposure assessment, hazard characterization, and risk characterization [88, 89]. Data from MRA are heavily utilized in developing policy-making and legislation for the most relevant foodborne pathogens [25].

Conventional methods for risk assessments of food products and additives are deterministic [90]. They are based on the assumption that estimated parameters are constants, while these parameters are variables, and their measurements are uncertain. Consequently, discrepancies between challenge tests and laboratory assays lead to compromised decisions when deterministic approaches are utilized alone in decision-making. Hence, more sensitive methods, such as predictive modeling, are necessary. By applying predictive models in the food industry, microbial hazards can be estimated from production to final consumption [91]. Information from these models can then be utilized in guiding decisions such as determining acceptable levels of microbial exposure and necessary actions to minimize risk to the final consumer [20].

A study was conducted by Pouillot and Lubran. [92] to assess the effectiveness of different established predictive models in quantitative risk assessment of listeriosis from contaminated cold-smoked salmon and the growth of *L. monocytogenes*. In this study, parameters such as the initial \log_{10} concentration of *L. monocytogenes* temperature fluctuation in a cooling device and storage duration were monitored using the dose-response model for invasive listeriosis developed by [89].

$$\Pr(\text{illness}|D) = 1 - \exp(-rD), \text{ and } D = C \times 10^{X_{\text{end}}}$$

With r as the probability of illness after ingesting one *L. monocytogenes* (number of organisms), the dose-response r was deemed $r = 1.06 \times 10^{-12}$ (based on data from an FAO report on population susceptibility).

D is the predicted dose of *L. monocytogenes* per serving.

X_{end} (\log_{10} cfu/g) = concentration of *L. monocytogenes* in the cold-smoked salmon after storage.

The final output of the model was calculated using Monte Carlo simulation as

$$(i) \bar{R} = 1/n \sum_{i=1}^n (1 - \exp(-r \times C \times 10^{X_{\text{end},i}})).$$

Different models, including the Buchanan model and the Baranyi model, were assessed. The result of the study showed the importance of accurately measuring bacterial growth in determining risk and the significance of selecting an appropriate primary growth model. The study also revealed the effect of different parameters on the estimated risk. Parameters such as the maximum achievable population density and lag phase were seen to be important in risk estimation.

The study also highlighted the gap between predictive microbiology models and current models being used in microbial risk assessment, as well as the difficulty in selecting appropriate models.

MicroHibro is an example of a predictive microbiology modeling software for evaluating spoilage and pathogenic microbes in the food industry. It provides estimates on the exposure level and associated risk. The software utilizes a Predictive Microbiology Model Database (PMDB), incorporating parameters such as growth, inactivation, transfer, and dose-response models. Estimating food microbial risk is done in MicroHibro by describing steps in the food chain using four microbial processes (growth, inactivation, transfer, and partitioning). It then estimates the microbial load and prevalence in the food of interest [93]. Other risk assessment tools that have been developed include the Dairy Products Safety Predictor, Food Spoilage and Safety Predictor, Listeria Meat Model, and FDA-iRISK [94].

2.4.3. HACCP (Hazard Analysis and Critical Control Point) Systems. Hazard Analysis and Critical Control Point (HACCP) is a food safety system designed to identify and prevent potential problems throughout food production, distribution, and consumption.

It employs a systematic approach to identify pathogens in raw materials and processing entry points, implement appropriate methods for their elimination, and detect

potential issues with the finished product resulting from improper handling. Predictive food microbiology plays a crucial role in implementing the HACCP concept.

The application of Quantitative Microbial Risk Assessment (QMRA) is essential in hazard analysis by evaluating potential microbial hazards in the food chain and determining critical control points (CCPs) and critical limits (CL). Predictive models can also be developed to evaluate systems established to monitor CCPs and verify the effectiveness of the HACCP system. These models can predict CCPs by determining levels for different parameters that permit microbial growth. In addition, they estimate various levels of microbial behavior to suggest permissible levels and thresholds for critical limits. Hence, integrating HACCPs with predictive models has excellent potential in decision-making [88].

In a study by Cassin et al. [95]; Monte Carlo simulation was used together with data from microbiological analysis in a dose-response analysis to predict the risk of hemolytic uremic syndrome (HUS) from *Escherichia coli* O157: H7 from beef hamburgers. The study modeled the pathogen behavior through processing, handling and final consumption and the determined the risk of the illness in different age groups. The authors then went on to assess the effectiveness of different mitigation strategies and critical control points in storage temperature control, preslaughter screening, and cooking temperature. From their results, the probability of illness was reduced through a strategy aimed at minimizing the bacteria growth during retail storage through accurate temperature monitoring. However, the authors also acknowledged the limitation of uncertainties in hygiene during individual operations during processing, production and handling. The authors were also unable to model certain steps in the production, processing, hamburger distribution, and consumption during the study. For instance, not all bacteria that survive the cooking process would be infective after. Also, the probability of HUS cannot be totally ascribed to dose.

2.4.4. Shelf-Life Determination. Traditional microbiological methods for determining shelf life are often time consuming, requiring substantial bacterial cell growth before overt spoilage reactions become apparent. Alternatively, newer methods may require advanced and expensive equipment. However, knowledge gained from predictive modeling of microbial behavior in food provides a robust foundation for developing devices capable of monitoring food shelf life throughout warehousing, transport, distribution, and retail sales.

Accurate shelf-life prediction models must fulfill several requirements, including identifying spoilage reactions (SRs) such as slime production, color changes, and offensive odors. They must also identify the specific microorganisms (SSOs) responsible for these spoilage reactions and analyze the spoilage domain (SD), which refers to the environmental conditions permitting the growth and function of a particular SSO. Developing and validating a shelf-life prediction model involve experiments demonstrating the spoilage

organisms, reactions, and domain. Subsequently, modeling microbial behavior within this domain is crucial for determining the “minimum spoilage level,” i.e., the concentration of SSO required to cause product rejection. Notably, spoilage organisms’ metabolic activities, rather than their numbers alone, play a significant role in determining spoilage [96].

Numerous predictive models have been developed for shelf-life studies in the food industry. Examples include models for predicting shelf life in yogurt by Mataragas et al. [97]; minced beef by Limbo et al. [98]; and Nutri cereal baby food by Rasane [99]. In addition, studies have focused on the impact of specific bacteria on spoilage in particular foods, such as *Pseudomonas* on pork and poultry by Bruckner et al. [100] and lactic acid bacteria on cooked ham by Kreyenschmidt [101]. These predictive models enhance food quality and safety by providing insights into the factors influencing shelf life and potential spoilage mechanisms.

2.5. Limitations of Predictive Models. Models provide many benefits in the decision-making process. However, it must be kept in mind that they are simplified representations of complex biological processes at best. Therefore, predictions based on model results should be made cautiously, considering previous experiences and other microbial ecological factors for which the models may not fully account. For instance, it is crucial to understand that models can only be extrapolated for values within the experimental ranges to which they were formulated, especially for parameters like temperature or water activity. This limitation arises because models, particularly empirical ones, are derived from fitting observed data and may only partially simulate actual microbial behavior. Research by Fakruddin et al. [68] also reports that specific models may predict faster growth rates compared to actual real-life scenarios. This discrepancy has been attributed to the development of most models using laboratory media, reducing their predictive relevance in the food industry even though they are validated in foods [102]. Practitioners in the food industry have also highlighted the incompatibility of models developed under stable environmental conditions in evaluating real-life products that experience fluctuating conditions of changing temperature, pH, and water activity. Moreover, predictive models cannot consider all the variables that affect food spoilage and microbial growth [103]. Most models evaluate just one or more parameters involved in food spoilage [104].

Researchers have also highlighted several more specific challenges, such as the prediction of the lag phase. The lag phase is a period of adjustment with little or no bacteria multiplication. Modeling this phase of bacterial growth has been challenging due to the variable length and the minimal level of microbial growth [103]. Population kinetics factors, like the lag phase and growth rate, are not easily predicted in rapidly evolving, complex ecosystems such as that of natural food. Modeling the shelf life of fish and seafood has also been difficult due to the scarcity of data on changes in fish microbial response after subjecting them to temperature changes during the lag and/or the exponential phase.

Therefore, applicable and accurate models for quantitative microbial risk assessment in seafood are unavailable [105].

Other limitations with the use of predictive models include the inability of models to model intraspecies diversity, the effect of natural food structure complex, as well as the effect of nutrient depletion and buildup of toxic bacteria waste in the medium [106, 107].

2.5.1. The Role of Industry Stakeholders in the Adoption of Predictive Models. Industry stakeholders applying modeling in food safety need to be informed of the predictive limits of models before making inferences. Thus, proposed models must be backed up with evidence showing a track record of valid predictions to aid adoption.

Table 1 outlines the role of AI, whole genome sequencing (WGS), and metagenomics in developing more efficient and accurate predictive models.

3. Challenges and Future Directions

Most microbial models are simplistic, primarily describing observable changes in response to environmental dynamics. While these models have proven effective in predicting parameters such as growth or inactivation rates within optimal temperature ranges, their ability to model complex phenomena like lag phases and adaptive responses remains questionable. Consequently, there is a growing need for more versatile and mechanistic next-generation models that integrate cellular information to unravel complex microbial behavior [109, 120].

Newer modeling approaches should prioritize understanding the behavior of individual cells to comprehend the dynamics of entire populations more accurately. Recent studies have emphasized the significance of individual cell heterogeneity as a significant source of variability [121, 122]. Given that food contamination can occur with very low numbers of pathogenic bacteria, understanding the mechanisms of individual cell behavior is crucial for robust microbial risk assessment [123, 124].

To address these challenges, research is underway on stochastic individual cell modeling based on systems biology approaches. These studies have revealed the variation of bacterial cells within a clone and the association between gene expression and phenotypic expression [125].

Next-generation models must also analyze complex interactions such as intraspecies diversity found in spore-forming bacteria and their relevance in risk assessment. This characteristic has not been adequately modeled in current approaches [126].

3.1. Integration of Predictive Microbiology with Other Technologies. Advances in whole genome sequencing (WGS) technologies, genomics, and metagenomics are revolutionizing pathogen detection, characterization, and identification in the field of food safety, giving rise to an emerging field known as foodomics [39, 127]. Furthermore, proteomics and metabolomics methods are being applied in detecting bacterial toxins and mycotoxins in foodstuffs,

making omics-based tools a vital aspect of risk assessment methods and food safety surveillance in the 21st century. The integration of WGS in surveillance programs has been applied in resolving outbreaks of listeriosis and Salmonellosis that previously would not have been detected [128–130]. By applying WGS technologies in predictive microbiology, researchers have access to details such as serotype, virulence factors, antimicrobial resistance genes, and genetic variations such as single nucleotide polymorphisms (SNPs) [39, 118]. This enables more precise risk assessment that better translates microbial genotype into differing clinical outcomes [131]. For instance, Shiga toxin-producing *Escherichia coli* is associated with different clinical outcomes ranging from diarrhea to hemolytic uremic syndrome (HUS) and other chronic sequelae. Applying WGS technologies in risk assessment makes it possible to link different hazards with particular clinical outcomes accurately [118, 132]. This was carried out in a study by Pielaat et al. [133] in which molecular data from WGS were applied to characterize hazards from *Escherichia coli* O157: H7 [134].

In addition, metagenomics can be used to study the interactions between microorganisms in a community [116]. Microbial communities are complex systems, and one microbe's behavior can influence others' behavior. By using metagenomics to study these interactions, it is possible to develop more sophisticated predictive models that take into account the dynamics of microbial communities.

Another tool with massive potential for integration into current predictive modeling systems is artificial intelligence and machine learning (ML) technology. The process of developing a predictive model generates large datasets that often cannot be analyzed using conventional statistical tools. Artificial intelligence and machine learning tools can be used to analyze large datasets and identify cryptic patterns and correlations. Artificial Intelligence, therefore, makes it possible to develop more sophisticated and accurate models that consider multiple factors that influence microbial growth and behavior. Several scientific reports have been written on the application of AI in this field [40, 43].

Machine learning is a subset of artificial intelligence focused on developing more efficient machines by integrating algorithms that enable them to learn from available datasets. Machine learning algorithms learn over time and are being employed in a holistic approach towards risk prediction due to their ability to consider complex genetic variations and interactions [135]. They allow for the evaluation of individual parameters and their interactions with other variables, which are vital in assessing strain diversity and the differences in observed phenotypic outcomes [136]. Microbial risk assessment incorporating these algorithms can capture changes in genetic constitution with time and consequently aid in faster identification of new strains with novel virulence characteristics [131]. It would also help in resolving limitations associated with predictive models. Machine learning algorithms are more flexible and dynamic than conventional predictive microbiology models as they do not require rigid formulas linking microbial parameters and observed features in food [137]. Algorithms employed in ML can discern complex relationships between input data

TABLE 1: The role of artificial intelligence (AI), whole genome sequencing (WGS), and metagenomics in developing more efficient and accurate predictive models.

Challenge associated with current predictive models	Advances made due to WGS, metagenomics, and artificial intelligence	References
Simplistic, unable to model complex microbial behavior	Metagenomics and WGS provide biological data on complex interactions in food ML and AI are used to visualize and model these interactions	[108, 109]
Errors in measurement and variability	Bayesian modelling and AI approaches provide reasonable estimates of uncertainty and variability	[110–112]
Understanding variability caused by individual cell heterogeneity	Stochastic individual cell modelling based on systems biology and artificial intelligence	[113, 114]
High intraspecific diversity in food	Metagenomic helps in characterising large microbial populations AI can be integrated into the development of sophisticated models that can take into consideration intraspecific differences and strain variability	[114–116]
Insufficient data available on microbial ecology	Whole genome sequencing and metagenomics helps to generate high quality data for predictive models Machine learning and artificial intelligence tools are applied in accurate, high quality data analysis	[117, 118]
Variability among different food materials	Bayesian networks incorporate prior knowledge about the properties of different foods	[37, 119]

and, from thence, make predictions [131]. Therefore, they do not rely on additional theories or mechanisms (as in mechanistic or empirical predictive models). AI is trained with large datasets, too many parameters to learn complex relationships, making it possible to predict even without clearly understood mechanisms accurately [137].

Machine learning models have also been shown to be more adept at exploring complex interactions among input features (e.g., growth kinetic parameters, type of pathogen, and sublethal injury rate) with fewer parameters [114].

A study by Lin et al. [114] applying machine learning for predicting the single-cell lag time of *Salmonella enteritidis* after heat and chlorine treatment on single-cell analysis demonstrates the following. Unlike traditional models, which need different sets of parameters to fit the data for varying microbes and processing methods, AI model requires only a set of hyperparameters when evaluating the impact of a sublethal injury on single-cell lag time, making it possible to build a single model capable of assessing single-cell variability among different pathogens to different treatments; an ideal model for all rather than a model per pathogen per treatment process [138].

These tools have also helped automate food quality monitoring systems through technologies such as robots, time-temperature indicators (TTIs), and Internet of Things (IoT) technologies. Through the integration of predictive models with artificial intelligence technologies, robots are being developed to optimize the handling and storage of products, reducing the risk of microbial contamination [139]. Figure 1 illustrates the integration of newer technologies into predictive modeling. IoT devices can be used to monitor and control the temperature and humidity of food and other products. Time-temperature indicators (TTIs) are devices that monitor the time-temperature history of a product and provide an indication of the degree of microbial growth. By integrating these technologies into current predictive models, it is possible to monitor and maintain the conditions required for the safe storage and distribution of products. Predictive models are developed to ensure that conditions are optimized for food safety, while artificial intelligence technologies help to add human-like awareness and decision-making in product design [140].

3.2. Machine Learning Algorithms Commonly Employed in Predictive Modeling. Many different ML algorithms, such as gradient boosting regression tree (GBRT), artificial neural network (ANN), random forest (RF), support vector regression (SVR), and logit boost (LB), are commonly used [114, 141]. Table 2 shows the application of different machine learning algorithms that have been used with varying levels of success.

3.2.1. Random forest. This algorithm derives its name from the approach towards data selection and decision-making. Generally, a random forest algorithm works by integrating multiple decision trees into a single ensemble of models based on the principle that the aggregation of multiple models should perform better than a single one. The




FIGURE 1: Integration of newer technologies into predictive modeling. WGS and metagenomics help to understand the abundance and diversity of microbial communities present in food samples. Data generated by these tools are then analyzed by artificial intelligence (AI) and machine learning (ML), leading to the development of accurate, sophisticated algorithms that are being utilised in the development of robots, time-temperature indicators, and internet of things (IoT) devices. Image sources: Arthur Shlain; dDara TH; rukanicon ID; icon scout IN; WiStudio TH.

algorithm randomly selects input variables and observations from provided data, from which it then generates individual decision trees. The final result, hence, is an aggregation of multiple trees for regression. Random forest algorithms have shown suitability in predictive microbiology due to their accuracy with classifications and regression analysis and suitability in assessing causality between parameters and effects; RF also considers the significance of variables in handling data. It is also rather easy to use and understand due to the small number of parameters to optimize (count of decision trees and independent variables per split) [154]. Random forest (RF) methods, particularly nonlinear variations, are highly effective in modeling parameters that lack a linear correlation, making them suitable for a wide range of applications in predictive microbiology and food safety. For example, RF models have been successfully applied in predicting heavy metal concentrations in Turbot muscle and liver tissues with an accuracy of up to 70% [155].

In many cases, microbial interactions in food involve numerous variables, posing challenges for specifying logistic regression models with optimal biologically relevant forms and interactions [156]. RF models offer a solution by effectively handling complex interactions and large datasets, such as those encountered in food spoilage studies. For instance, Chen et al. [157] utilized RF models to predict quality changes in Pacific white shrimp across a temperature range of -20° – 4° C, showcasing their utility in food preservation research. In this study, RF models and Arrhenius models were compared to determine specific markers of Pacific white shrimp quality. The RF models were seen to have lower error rates with values of 0.15%–1.24% (pH),

TABLE 2: The application of machine learning algorithms.

Application of machine learning	Algorithm	Reference
Risk prediction for highly pathogenic avian influenza at poultry farms in the Republic of Korea	Random forest Gradient boosting machine (GBM) Extreme gradient boosting	[142]
Effect of environmental factors on the abundance of <i>Vibrio parahaemolyticus</i> in oyster farms in Taiwan	Extreme gradient boosting	[143]
Prediction of skin flavonoid content from berry physical-mechanical characteristics in wine grapes	Regression trees Random forests Gradient boosting machine	[144]
Food source attribution of <i>Listeria monocytogenes</i>	Random forest Support vector machine Stochastic gradient boosting Logit boost	[145]
Microbial quality assessment of chicken liver inoculated or not with <i>Salmonella</i> using FTIR spectroscopy	Support vector machine	[146]
Microbial risk assessment using next generation sequencing data in predicting clinical outcomes in Shiga toxinogenic <i>Escherichia coli</i>	Random forest Support vector machine Logit boost	[131]
Single cell lag time of <i>Salmonella enteritidis</i> after heat and chlorine treatment	Gradient boosting regression tree Artificial neural network, random forest Support vector regression	[114]
Temperature prediction in apples	Artificial neural networks	[147]
Predicting perishable food temperatures along the supply chain	Artificial neural networks	[148]
Application of machine learning and RFID in the stability optimization of perishable foods	Kernel logistic regression	[149]
Application of whole genome sequences in determining food sources of HUMAN <i>Salmonella typhimurium</i> infection	Logit boost Random forest	[150]
Applying genomic data in determining disease severity in <i>Salmonella enterica</i> infection	Logistic regression Random forest Support vector machine, AdaBoost	[151]
Source of campylobacteriosis using whole genome data	K-nearest neighbour algorithm (KNN) eXGBoost Random forest	[152]
Prediction of <i>Escherichia coli</i> O157 growth by machine learning	Support vector regression, extremely randomized trees regression Gaussian process regression	[153]

0.21%–2.82% (thiobarbituric acid), 0.09%–2.51% (total sulfhydryl content), 0.60%–11.84% (Ca^{2+} -ATPase activity), and 0.01%–4.76% (total viable counts). The Arrhenius models were shown to have error rates of 0.33%–2.07%, 0.49%–19.84%, 1.73%–23.28%, 2.49%–36.32%, and 11.90%–37.94%, respectively.

Furthermore, RF models have been applied in predicting meat spoilage by integrating metabolome data from various analytical and imaging instruments, demonstrating their versatility in handling diverse types of data [41]. Similarly, Astuti et al. [158] utilized RF models trained on gas sensor array data from fresh chicken meat and *E. coli*-contaminated chicken meat to predict contamination and spoilage based on shelf time, achieving remarkable precision rates of up to 99.25% and 98.42%, respectively.

3.2.2. Support vector Machines (SVMs). SVMs are highly effective, flexible, holistic machine learning algorithms that can perform linear or nonlinear regression and classification tasks [159, 160]. Support vector regressions (SVRs) are an important branch of SVM applied specifically to regression problems [161]. SVM algorithms are based on a “kernel trick,” a technique that implicitly maps data inputs into high-dimensional spaces and draws a “hyperplane” between them. Basically, it draws a margin between differing data classes. Using the kernel trick technique, the hyperplane is drawn in a way that takes data points as close as possible to it while also maximizing the distance between the margin and the classes, hence minimizing the classification error [162, 163]. The hyperplane is a line (an interface) between two datasets. SVMs can find the most suitable line delineating the two datasets by using linear optimization [163]. Therefore, it can maximize the margin in the high-dimensional space. They are good at handling outliers and are most efficient with well-differentiated data parameters. They are specially used for classification and regression challenges. It is, however, limited by poor effectiveness in handling large and noisy datasets [164, 165]. Despite this, they have been successfully applied in various areas of food safety, including risk assessment in dairy production and prediction of contamination in food samples.

For example, Ma et al. [166] utilized SVMs for risk assessment in dairy production, demonstrating the model’s ability to process large datasets efficiently. Bonah et al. [167] employed SVM hyperparameter optimization in combination with data from an electric nose to develop a model capable of predicting *Salmonella typhimurium* contamination in fresh pork samples. Similarly, Zhang et al. [42] trained an SVM model on historical monitoring data to assess and predict the risk during the transportation and refrigeration of strawberries.

3.2.3. Artificial neural Networks. Artificial neural networks employs machine learning techniques that simulate the neural pathway of the brain. A basic neural network starts with an input layer of neurons (representing input such as temperature or water activity in a microbiological model), which activates neurons in hidden layers, which then activates neurons in the

output layer (depicting predictions from nonlinear activation functions in hidden layers). Each hidden layer contains a network of multiple neurons (also known as nodes or units), the basic unit of computation [168]. The neurons are a logistic regression unit that receives multiple numeric inputs (from other nodes or an external source) and computes one numeric output. The neuron output is computed by a nonlinear function called the activation function. If the weighted sum of the inputs exceeds an internal threshold value within the neuron, the output is activated; otherwise, it is inactivated.

The multilayer model makes it very efficient with nonlinear, complex, high-dimensional relationships between the input and output. However, ANNs require numerical data; hence, they require a transformation of categorical data before they can be used. Deep neural networks, which consist of multiple hidden layers, have been used to model the growth limits of *Bacillus* spp. spores with accuracy greater than 90% [137].

One advantage of ANNs is their parallel architecture, which enables fast computation and evaluation of complex biological processes in food processing operations. They have been successfully applied in modeling diverse food processes such as extraction, drying, fermentation, dairy processing, and quality evaluation. For example, Sun et al. [169] utilized ANNs to monitor the microwave vacuum drying process of carrots, demonstrating their effectiveness in food processing applications.

3.2.4. Gradient Boosting Regression Tree (GBRT). Models are based on the ensemble learning scheme like the random forest trees [170]. These algorithms work by creating multiple weak models that can be combined into one single model with strong learning capabilities. They are different from conventional boosting algorithms in that the new predictor of GBRT fits the residuals of the previous predictor. GBRT algorithms are designed to reduce loss function by current learner by ensuring that loss function reduces along with gradients direction; that way, final residuals approach zero with continuous iteration, adding up all the tree results to get the final prediction. Like random forests, Gb are tree based; they process interactions effectively, are robust to outliers, and can select variables automatically.

In a study by Sheng et al. [171], GBRT models were trained using multiwavelength data from multichannel infrared spectral sensors and broadband infrared (IR) sensors to predict milk fat and protein content under different conditions accurately. In addition, GBRT models have been developed to predict outbreaks of foodborne diseases by utilizing data from foodborne disease surveillance studies [166].

3.2.5. Logit boost (LB) and Stochastic Gradient Boosting. They are part of a boosting family of algorithms developed in the 1990s [172]. Boosting algorithms function by aggregating (boosting) several weak classifiers (a weak classifier predicts marginally better than random) into an ensemble with improved accuracy [173]. A logit boost model was developed

by Benefo et al. [174], making use of whole genome sequence data to identify genes associated with *Salmonella* stress response in poultry.

3.3. Applications of A.I in Predictive Modeling

3.3.1. Artificial Intelligence in Modeling Climatic Factors and Their Effect on Microbial Ecology. Pang et al. [175] utilized logistic regression (LR) and random forest (RF) to analyze the association between weather-related factors and *Listeria* spp. in a mixed produce and dairy farm. Climatic factors, including temperature, precipitation, and wind speed, were considered for analysis. In both LR and RF models, wind speed and precipitation were found to play a significant role in transmitting *Listeria* spp. These experiments demonstrate that both models have good predictive capabilities in analyzing the impact of risk factors, such as weather, on microbial distribution.

Similarly, a random forest-based predictive model was developed by Hwang et al. [176] to quantify the relationship between climatic factors and the presence of *Salmonella* on pastured poultry farms. According to their analysis, the soil model identified humidity as the most significant meteorological variable associated with *Salmonella* prevalence. In contrast, the feces model identified high wind gust speed and average temperature as the most significant. The developed models also showed high accuracy with ROC (area under the curve) values of 0.884 for the soil model and 0.872 for the feces model.

Xu et al. [177] also developed a random forest predictive model that evaluated farm practices and processing variables to identify factors that can reduce the prevalence of *Campylobacter*, *Salmonella*, and *Listeria* on pastured poultry farms. In the study, the effect of farm practices on the *E. coli* population was used to predict the presence of other foodborne pathogens within pastured poultry farm environment. From the results, the probabilities of identifying *Campylobacter* in poultry feces and fecal samples at a very low concentration of *E. coli* were 88% and 98%.

3.3.2. Machine Learning Development of Devices for Quality Control. Direct measurement of a product's biochemical quality is cost-ineffective and tedious. As an alternative, time-temperature integration (TTI) has been developed to evaluate food quality by integrating product temperature assessment over time. TTIs are portable devices attached to food products; they assess biochemical interactions in food and then indicate the remaining shelf life. TTIs assess food products for breach of temperature threshold and impact on food quality [178]. These tools are being automated to send warning signs of deterioration indicative of failing storage to food operators.

Currently, AI is integrated into the Internet of Things (IoT) and time-temperature indicators to improve the characterization and detection of cold chain breaches. It assists in defining significant temperature variations, optimizing the placement of sensors (to accurately monitor the load in refrigeration equipment), and developing methods for detecting and characterizing temperature fluctuations.

However, it is crucial to establish well-defined alert thresholds to ensure optimal food safety and minimize wastage. Alerts must strike a balance between being stringent enough to prevent food spoilage and lenient enough to avoid unnecessary disposal of safe food. Achieving this balance requires careful consideration of various factors, including the specific characteristics of the food product, potential health risks associated with spoilage, and logistical costs associated with food waste management [179].

do Nascimento Nunes et al. [180] applied neural networks to predict the temperature of berries in a pallet after training them with ambient temperature data. Subsequently, Badia-Melis. [147] extended this approach to assess fruit temperature dynamics during refrigeration failure using data from cold chain breaks in freezing systems. These two studies demonstrated the effectiveness of artificial neural networks in evaluating the cold chain (in comparison with Kriging algorithms and capacitive heat transfer methods). They also highlighted the effect of sensor number and positioning on results. Results from these studies compared favorably with results from thermal imaging and were also shown to be more effective than methods such as Kriging or capacitive heat transfer.

Other technologies are also being integrated into ML in food quality control. In a case study by Dourou et al. [146], machine learning was combined with Fourier-transform infrared spectroscopy (FTIR) in real-time monitoring of food microbiota during changing storage conditions. The ability of foodborne pathogens (*Salmonella* in this case) to thrive and proliferate during prolonged refrigeration was demonstrated.

3.3.3. Machine Learning in Predictive Modeling of Individual Cell Heterogeneity. A study by Lin et al. [114] compared the effectiveness of four ML models (ANN, RF, GBRT, and SVR) in predicting the single-cell lag time of *Salmonella Enteritidis* after heat and chlorine treatment. The machine learning models were trained with a dataset comprising the following selected variables: population lag times (λ), maximum specific growth rate (μ_{max}), turbidity detection time (T_d), sublethal injury rate, and log reduction. It was demonstrated from the study that emerging machine learning models have the potential to predict the single lag time of foodborne pathogens by directly learning from data without assuming underlying mechanisms. The study also demonstrated the importance of population lag time and sublethal injury rate as vital parameters in the single-cell lag time analysis. Increasing population lag time could be positively correlated with a high value of single-cell lag times. The results also provide a framework for food operators to understand the risk from single cells of foodborne pathogens as well as the growth characteristic of *S. Enteritidis* single cells after disinfection treatment [114].

Increases in the mean and variance of single-cell lag time and the population lag time of treated cells showed the effect of external stress on individual cell heterogeneity. Generally, external stress (in this case, from disinfection and heat treatments) is known to play a significant role in increasing

individual cell heterogeneity [115]. A comparison of the four models was also made, with ANN shown to have the most accurate prediction of the four models, followed closely by RF. In this study, GBRT and SVR had suboptimal predictions. SVR's unsuitability to handle noise may be responsible for its underperformance [181, 182].

3.3.4. Machine Learning in Predicting Clinical Outcomes from Foodborne Hazards. Machine learning analysis of data from whole genome sequencing of foodborne bacteria pathogens and epidemiological data are being used to characterize food safety risks and predict clinical outcomes. In a study by Njage et al. [131], whole genome sequencing of foodborne Shiga toxin-producing *Escherichia coli* was combined with four machine learning models, namely, random forests, support vector machines, and logit boost in hazard characterization and prediction of clinical outcomes from infection. Their results indicated the significance of genotypic data from WGS in determining clinical outcomes by analyzing certain predictor proteins. Genetic predictors of riskier clinical outcomes were shown to include proteins involved in bacterial attachment, genomic islands, sex-pili formation, and protein acetylation. Furthermore, the association between clinical outcome and strain characteristics such as MLST type, lineage-specific polymorphism assay (LSPA), stx-subtype, and sublineage type, and demographic factors such as age and travel were analyzed by ML models. Clinical outcomes were divided into categories ranging from diarrhea, bloody diarrhea, hospitalization, and hemolytic uremic syndrome (HUS).

ML models were trained with a rich dataset comprising sequences from MLST, specific polymorphisms (LSPA-6), and other strain characteristics obtained from STEC outbreaks.

The logit boost model was found to be the best-performing model after comparing its accuracy and similarity with other models.

The application of machine learning helps to overcome the challenge posed by high-dimensional data analysis, which is often required in predicting phenotypic clinical outcomes from microbial genotypes while also considering the intricate interactions between genetic factors that define disease outcomes.

3.3.5. Artificial Intelligence in Pathogen Detection and Growth. Machine learning algorithms have been successfully applied in the detection of pathogens (Table 2). A study by Amado et al. [183] employed a collection of ML algorithms, including random forest, support vector regression, KNN, NBC, and artificial neural networks, to detect the presence of *Escherichia coli* and *Staphylococcus aureus* in beef. Data obtained from emitted gases in meat were used as the input dataset. Results from this study indicated the random forest model to be the most accurate in predicting

bacterial contamination. Another study by Tanui et al. [145] utilized the random forest algorithm, the support vector machine, the stochastic gradient boosting algorithm, and the logistic boost algorithm in predicting food sources of listeriosis. The study made use of multi-locus sequence typing (MLST) data of *Listeria monocytogenes* isolates from dairy, fruits and vegetables, meat, poultry, and seafood [145]. Results linked the bulk of infections (32.5%) to fruits, followed by 18.8% in vegetables. This study further highlights the use of genomic data combined with machine learning based on foodborne-disease tracking. Table 2 shows how various machine learning algorithms have been applied in different fields to predict the risk of highly pathogenic organisms.

3.4. Impact of Climate Change on Predictive Microbiology. Climate change generally plays a key role in the burden of foodborne pathogens. At several points along the food chain, changes in weather patterns globally have a significant influence on the emergence of food safety hazards [184]. Elevated water temperatures caused by global warming have been linked with increased expression of virulence genes and faster die off in waterborne bacteria such as pathogenic *Vibrio* spp. [185]. Increased antimicrobial resistance has also been positively correlated with increased local temperature [186]. Globally, changes in microbial ecology due to climate change have been associated with changes in the epidemiology of foodborne infections. A study by Bandyopadhyay et al. [187] showed the association between an increased incidence of diarrhoea and increased temperature in sub-Saharan Africa. Specifically, studies in Ethiopia have demonstrated an increased incidence of diarrhea during the hotter periods of the year [188].

The impact of climate change on changing microbial ecology is readily seen in the dairy industry, where fluctuations in weather patterns are readily associated with changes in microbial ecology. The direct impact of global warming on weather parameters such as temperature and relative humidity plays a major role in increasing the biodiversity of pathogenic microbes in raw milk [189, 190]. Furthermore, warm weather and associated heat stress in milking cows predispose to mastitis and, consequently, higher somatic cell counts and altered physicochemical properties [191]. It is imperative, therefore, that predictive models be developed with consideration of the impact of climatic conditions, seasonal fluctuations in climatic variables, and global warming generally.

A few studies have been carried out on developing predictive models which factor in the impact of climate change. Predictive models that incorporate time-series analysis together with epidemiological data have been used to forecast the significance of environmental temperature in risk assessment of salmonellosis, campylobacteriosis and listeriosis in Belgium [192]. El-Fadel et al. [193] demonstrated the application of climatic forecasts in building

Poisson regression models that can measure the correlation between climatic factors and the incidence of morbidities due to water- and foodborne pathogens.

4. Conclusion

The review discussed how predictive models are widely used in food microbiology to deduce the growth of microorganisms in food products although these models have some limitations in modeling complex microbial interactions in food dominated by different bacteria populations. The integration of new technologies such as whole genome sequencing (WGS), metagenomics, artificial intelligence, and machine learning and the use of devices based on robotics, the Internet of Things, and time-temperature indicators have, however, been improving the efficiency and accuracy of these models. The integration of machine learning models, in particular, has been greatly beneficial in developing more advanced models. Hence, continued research and development of these technologies are essential for improving food safety and reducing the risk of microbial contamination in food products [194, 195].

Data Availability

The data used to support the findings of this study are available from the corresponding author upon reasonable request.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

Oluseyi Rotimi Taiwo performed conceptualization, writing the original draft, and editing. Helen Onyeaka performed conceptualization, supervision, writing the original draft, and editing. Elijah K. Oladipo, Julius Kola Oloke, and Deborah C. Chukwugozie performed writing the original draft and editing.

References

- [1] A. Paparella, "Food safety: definitions and aspects," *Food Safety and Hazards*, vol. 1, 2020.
- [2] A. Kalla, S. R. Kutcharlapati, S. B. Gannavarapu, and A. Layam, "Isolation and identification of specific pathogens, presence of antibiotics, aflatoxins, pesticide residues and industrial contaminants in supply chain of milk in selected coastal districts of Andhra Pradesh," *Advances in Bioscience and Biotechnology*, vol. 6, no. 4, pp. 330–344, 2015.
- [3] World Health Organization (Who), "Food safety," 2022, <https://www.who.int/news-room/fact-sheets/detail/food-safety>.
- [4] J. Fanzo, *Can Fixing Dinner Fix the Planet?*, JHU Press, Baltimore, MD, USA, 2021.
- [5] M. T. Osterholm and M. E. Potter, "Irradiation pasteurization of solid foods: taking food safety to the next level," *Emerging Infectious Diseases*, vol. 3, no. 4, p. 575, 1997.
- [6] T. Kobets, B. P. Smith, and G. M. Williams, "Food-borne chemical carcinogens and the evidence for human cancer risk," *Foods*, vol. 11, no. 18, p. 2828, 2022.
- [7] W. Messens, M. Hempen, and K. Koutsoumanis, "Use of predictive modelling in recent work of the panel on biological hazards of the European food safety authority," *Microbial Risk Analysis*, vol. 10, pp. 37–43, 2018.
- [8] P. K. Singh, R. P. Singh, P. Singh, and R. L. Singh, "Food hazards: physical, chemical, and biological," in *Food Safety and Human Health*, pp. 15–65, Academic Press, Cambridge, MA, USA, 2019.
- [9] E. D. Van Asselt, A. Arrizabalaga-Larrañaga, M. Focker, B. J. A. Berendsen, M. G. M. Van de Schans, and H. J. Van der Fels-Klerx, "Chemical food safety hazards in circular food systems: a review," *Critical Reviews in Food Science and Nutrition*, vol. 63, no. 30, pp. 10319–10331, 2023.
- [10] R. Lawley, L. Curtis, and J. Davis, *The Food Safety hazard Guidebook*, Royal Society of Chemistry, London, UK, 2012.
- [11] H. Lee and Y. Yoon, "Etiological agents implicated in foodborne illness world wide," *Food science of animal resources*, vol. 41, no. 1, p. 1, 2021.
- [12] World Health Organization [Who], "WHO estimates of the global burden of foodborne diseases," 2015, https://apps.who.int/iris/bitstream/handle/10665/199350/9789241565165_eng.pdf?sequence=1.
- [13] S. M. Barlow, A. R. Boobis, J. Bridges et al., "The role of hazard-and risk-based approaches in ensuring food safety," *Trends in Food Science & Technology*, vol. 46, no. 2, pp. 176–188, 2015.
- [14] P. D. Hamilton, P. Gale, and S. J. Pollard, "A commentary on recent water safety initiatives in the context of water utility risk management," *Environment International*, vol. 32, no. 8, pp. 958–966, 2006.
- [15] Z. T. Rani, A. Hugo, C. J. Hugo, P. Vimiso, and V. Muchenje, "Effect of post-slaughter handling during distribution on microbiological quality and safety of meat in the formal and informal sectors of South Africa: a review," *South African Journal of Animal Science*, vol. 47, no. 3, pp. 255–267, 2017.
- [16] J. J. Webber, B. Dobrenov, J. Lloyd, and D. Jordan, "Meat inspection in the Australian red-meat industries: past, present and future," *Australian Veterinary Journal*, vol. 90, no. 9, pp. 363–369, 2012.
- [17] B. Antunović, B. Blagojević, S. Jöhler et al., "Challenges and opportunities in the implementation of new meat inspection systems in Europe," *Trends in Food Science & Technology*, vol. 116, pp. 460–467, 2021.
- [18] K. D. C. Stärk, S. Alonso, N. Dadios et al., "Strengths and weaknesses of meat inspection as a contribution to animal health and welfare surveillance," *Food Control*, vol. 39, pp. 154–162, 2014.
- [19] R. Laukkanen-Ninios, R. Rahkila, L. Oivanen, E. R. Wirta, and M. Fredriksson-Ahomaa, "Views of veterinarians and meat inspectors concerning the practical application of visual meat inspection on domestic pigs in Finland," *Journal of Consumer protection and food safety*, vol. 15, no. 1, pp. 5–14, 2020.
- [20] S. C. González, A. Possas, E. Carrasco et al., "MicroHibro: A software tool for predictive microbiology and microbial risk

- assessment in foods," *International Journal of Food Microbiology*, vol. 290, pp. 226–236, 2019.
- [21] S. Ceuppens, D. Li, M. Uyttendaele et al., "Molecular methods in food safety microbiology: interpretation and implications of nucleic acid detection," *Comprehensive Reviews in Food Science and Food Safety*, vol. 13, no. 4, pp. 551–577, 2014.
- [22] A. Gill, "The importance of bacterial culture to food microbiology in the age of genomics," *Frontiers in Microbiology*, vol. 8, Article ID 256356, 2017.
- [23] J. C. Jokerst, J. A. Adkins, B. Bisha, M. M. Mentele, L. D. Goodridge, and C. S. Henry, "Development of a paper-based analytical device for colorimetric detection of select foodborne pathogens," *Analytical Chemistry*, vol. 84, no. 6, pp. 2900–2907, 2012.
- [24] F. Wu, J. D. Groopman, and J. J. Pestka, "Public health impacts of food-borne mycotoxins," *Annual Review of Food Science and Technology*, vol. 5, pp. 351–372, 2014.
- [25] World Health Organization/Food and Agriculture Organization, *Risk Assessment of Listeria Monocytogenes in Ready-To-Eat Foods*, World Health Organization, Rome, Italy, 2016, http://www.who.int/foodsafety/publications/mra_5/en/.
- [26] Codex Alimentarius Commission, *Principles and Guidelines for the Conduct of Microbiological Risk Assessment. CAC/GL 30-1999*, Food and Agriculture Organization of the United Nations/World Health Organization, Rome/Geneva, Italy, 2014.
- [27] T. Mayes, "Risk analysis in HACCP: burden or benefit?" *Food Control*, vol. 9, no. 2-3, pp. 171–176, 1998.
- [28] P. A. Wester, *Hazard Analysis and Risk Based Preventive Controls: Building a (Better) Food Safety Plan*, Academic Press, Cambridge, MA, USA, 2017.
- [29] F. Perez-Rodriguez, A. Valero, F. Pérez-Rodríguez, and A. Valero, *Predictive Microbiology in Foods*, Springer, New York, NY, USA, 2013.
- [30] M. Schirone and P. Visciano, "Trends of major food-borne outbreaks in the European Union during the years 2015–2019," *Hygie*, vol. 1, no. 3, pp. 106–119, 2021.
- [31] M. Van Schothorst, M. H. Zwietering, T. Ross, R. L. Buchanan, and M. B. Cole, "Relating microbiological criteria to food safety objectives and performance objectives," *Food Control*, vol. 20, no. 11, pp. 967–979, 2009.
- [32] A. Drewnowski, "The economics of food choice behavior: why poverty and obesity are linked," in *Obesity Treatment and Prevention New Direction*, vol. 73, pp. 95–112, Karger Publishers, Berlin, Germany, 2012.
- [33] S. M. Barlow, A. R. Boobis, J. Bridges et al., "The role of hazard-and risk-based approaches in ensuring food safety," *Trends in Food Science & Technology*, vol. 46, no. 2, pp. 176–188, 2015.
- [34] J. M. Mehta and I. M. Rietjens, "Pros and cons of hazard-versus risk-based approaches to food safety regulation," in *Present Knowledge in Food Safety*, pp. 1068–1087, Academic Press, Cambridge, MA, USA, 2023.
- [35] A. J. Lopatkin and J. J. Collins, "Predictive biology: modeling, understanding and harnessing microbial complexity," *Nature Reviews Microbiology*, vol. 18, no. 9, pp. 507–520, 2020.
- [36] F. Pérez-Rodríguez, A. Valero, F. Pérez-Rodríguez, and A. Valero, "Application of predictive models in quantitative risk assessment and risk management," *Predictive microbiology in foods*, pp. 87–97, 2013.
- [37] J. C. Augustin, H. Bergis, G. Midelet-Bourdin et al., "Design of challenge testing experiments to assess the variability of *Listeria monocytogenes* growth in foods," *Food Microbiology*, vol. 28, no. 4, pp. 746–754, 2011.
- [38] V. Martinez-Rios, E. Gkogka, and P. Dalgaard, "Predicting growth of *Listeria monocytogenes* at dynamic conditions during manufacturing, ripening and storage of cheeses—Evaluation and application of models," *Food Microbiology*, vol. 92, Article ID 103578, 2020.
- [39] J. Kovac, H. den Bakker, L. M. Carroll, and M. Wiedmann, "Precision food safety: a systems approach to food safety facilitated by genomics tools," *TrAC, Trends in Analytical Chemistry*, vol. 96, pp. 52–61, 2017.
- [40] V. Sahni, S. Srivastava, and R. Khan, "Modeling techniques to improve the quality of food using artificial intelligence," *Journal of Food Quality*, vol. 2021, Article ID 2140010, 10 pages, 2021.
- [41] L. Estelles-Lopez, "An automated ranking platform for machine learning regression models for meat spoilage prediction using multi-spectral imaging and metabolic profiling," *Food Research International*, vol. 99, pp. 206–215, 2017.
- [42] G. Zhang, G. Li, and J. Peng, "Risk assessment and monitoring of green logistics for fresh produce based on a support vector machine," *Sustainability*, vol. 12, no. 18, p. 7569, 2020.
- [43] K. Veselkov, G. Gonzalez, S. Aljifri et al., "HyperFoods: machine intelligent mapping of cancer-beating molecules in foods," *Scientific Reports*, vol. 9, no. 1, p. 9237, 2019.
- [44] A. Lenzi, M. Marvasi, and A. Baldi, "Agronomic practices to limit pre-and post-harvest contamination and proliferation of human pathogenic Enterobacteriaceae in vegetable produce," *Food Control*, vol. 119, Article ID 107486, 2021.
- [45] E. A. Alum, S. M. O. C. Urom, and C. M. A. Ben, "Microbiological contamination of food: the mechanisms, impacts and prevention," *International Journal of Scientific & Technology Research*, vol. 5, no. 3, pp. 65–78, 2016.
- [46] S. Sharma, A. Mishra, K. Shukla, T. Jindal, and S. Shukla, "Food contamination: it's stages and associated illness," *International Journal of Pharmaceutical, Chemical and Biological Sciences*, vol. 10, no. 4, 2020.
- [47] E. Abebe, G. Gugsu, and M. Ahmed, "Review on major food-borne zoonotic bacterial pathogens," *Journal of Tropical Medicine*, vol. 2020, Article ID 4674235, 19 pages, 2020.
- [48] N. G. Marriott, M. W. Schilling, R. B. Gravani, N. G. Marriott, M. W. Schilling, and R. B. Gravani, "Food contamination sources," *Principles of food sanitation*, pp. 83–91, 2018.
- [49] C. Rolfe and H. Daryaei, "Intrinsic and extrinsic factors affecting microbial growth in food systems," *Food Safety Engineering*, pp. 3–24, 2020.
- [50] S. H. Hamad, "Factors affecting the growth of microorganisms in food," *Progress in food preservation*, pp. 405–427, 2012.
- [51] N. S. Al-Zoreky, "Antimicrobial activity of pomegranate (*Punica granatum* L.) fruit peels," *International Journal of Food Microbiology*, vol. 134, no. 3, pp. 244–248, 2009.
- [52] T. A. Krulwich, G. Sachs, and E. Padan, "Molecular aspects of bacterial pH sensing and homeostasis," *Nature Reviews Microbiology*, vol. 9, no. 5, pp. 330–343, 2011.
- [53] R. Bhat and G. Paliyath, Eds., *Progress in Food Preservation*, John Wiley & Sons, Hoboken, NJ, USA, 2012.
- [54] D. S. Esser, J. H. Leveau, and K. M. Meyer, "Modeling microbial growth and dynamics," *Applied Microbiology and Biotechnology*, vol. 99, pp. 8831–8846, 2015.
- [55] M. Peleg and M. G. Corradini, "Microbial growth curves: what the models tell us and what they cannot," *Critical*

- Reviews in Food Science and Nutrition*, vol. 51, no. 10, pp. 917–945, 2011.
- [56] J. Baranyi and N. B. da Silva, “The use of predictive models to optimize risk of decisions,” *International Journal of Food Microbiology*, vol. 240, pp. 19–23, 2017.
- [57] M. P. Jiang, S. Y. Zheng, H. Wang et al., “Predictive model of aflatoxin contamination risk associated with granary-stored corn with versicolorin A monitoring and logistic regression,” *Food Additives & Contaminants: Part A*, vol. 36, no. 2, pp. 308–319, 2019.
- [58] E. Stavropoulou and E. Bezirtzoglou, “Predictive modeling of microbial behavior in food,” *Foods*, vol. 8, no. 12, p. 654, 2019.
- [59] L. Huang, “Dynamic identification of growth and survival kinetic parameters of microorganisms in foods,” *Current Opinion in Food Science*, vol. 14, pp. 85–92, 2017.
- [60] R. L. Buchanan, “Predictive food microbiology,” *Trends in Food Science & Technology*, vol. 4, no. 1, pp. 6–11, 1993.
- [61] P. J. McClure, J. Baranyi, E. Boogard, T. M. Kelly, and T. A. Roberts, “A predictive model for the combined effect of pH, sodium chloride and storage temperature on the growth of *Brochothrix thermosphacta*,” *International Journal of Food Microbiology*, vol. 19, pp. 161–178, 1993.
- [62] J. Baranyi, T. A. Roberts, and P. J. McClure, “A non-autonomous differential equation to model bacterial growth,” *Food Microbiology*, vol. 10, pp. 43–59, 1993.
- [63] H. Li, G. Xie, and A. Edmondson, “Evolution and limitations of primary mathematical models in predictive microbiology,” *British Food Journal*, vol. 109, no. 8, pp. 608–626, 2007.
- [64] J. M. Membre, T. Ross, and T. McMeekin, “Behaviour of *Listeria monocytogenes* under combined chilling processes,” *Letters in Applied Microbiology*, vol. 28, pp. 216–220, 1999.
- [65] R. L. Buchanan, R. C. Whiting, and W. C. Damert, “When is simple good enough: a comparison of the Gompertz, Baranyi, and three-phase linear models for fitting bacterial growth curves,” *Food Microbiology*, vol. 14, pp. 313–326, 1997.
- [66] H. Li, G. Xie, and A. S. Edmondson, “Review of secondary mathematical models of predictive microbiology,” *Journal of Food Products Marketing*, vol. 14, no. 2, pp. 57–74, 2008.
- [67] A. Valero, E. Carrasco, and R. M. García-Gimeno, “Principles and methodologies for the determination of shelf-life in foods,” *Trends in vital food and control engineering*, vol. 1, pp. 3–42, 2012.
- [68] M. D. Fakruddin, R. M. Mazumder, and K. S. B. Mannan, “Predictive microbiology: modeling microbial responses in food,” *Ceylon Journal of Science*, vol. 40, no. 2, pp. 121–131, 2011.
- [69] S. J. M. Breig and K. J. K. Luti, “Response surface methodology: a review on its applications and challenges in microbial cultures,” *Materials Today: Proceedings*, vol. 42, pp. 2277–2284, 2021.
- [70] B. A. D. Begoude, R. Lahlali, D. Friel, P. R. Tondje, and M. H. Jijakli, “Response surface methodology study of the combined effects of temperature, pH, and aw on the growth rate of *Trichoderma asperellum*,” *Journal of Applied Microbiology*, vol. 103, no. 4, pp. 845–854, 2007.
- [71] E. Mahdinia, S. Liu, A. Demirci, and V. M. Puri, “Microbial growth models,” *Food safety engineering*, pp. 357–398, 2020.
- [72] H. Zhuge and Y. Xing, “Probabilistic resource space model for managing resources in cyber-physical society,” *IEEE Transactions on Services Computing*, vol. 5, no. 3, pp. 404–421, 2011.
- [73] A. Psomas, “Development of mathematical models for spoilage prediction,” Agricultural University of Athens, Athens, Greece, 2012, Ph.D. thesis.
- [74] A. M. Gibson, N. Bratchell, and T. A. Roberts, “The effect of sodium chloride and temperature on the rate and extent of growth of *Clostridium botulinum* type A in pasteurized pork slurry,” *Journal of Applied Microbiology*, vol. 62, no. 6, pp. 479–490, 1987.
- [75] W. S. Choi, N. Son, J. I. Cho et al., “Predictive model of *Staphylococcus aureus* growth on egg products,” *Food Science and Biotechnology*, vol. 28, pp. 913–922, 2019.
- [76] L. Huang, “IPMP Global Fit—A one-step direct data analysis tool for predictive microbiology,” *International Journal of Food Microbiology*, vol. 262, pp. 38–48, 2017.
- [77] V. Valdramidis, “Predictive microbiology,” in *Modeling in Food Microbiology*, pp. 1–15, Elsevier, Amsterdam, Netherlands, 2016.
- [78] Z. Yi and J. Xie, “Prediction in the dynamics and spoilage of *Shewanella putrefaciens* in bigeye tuna (*Thunnus obesus*) by gas sensors stored at different refrigeration temperatures,” *Foods*, vol. 10, no. 9, p. 2132, 2021.
- [79] J. Horowitz, M. D. Normand, M. G. Corradini, and M. Peleg, “Probabilistic model of microbial cell growth, division, and mortality,” *Applied and Environmental Microbiology*, vol. 76, no. 1, pp. 230–242, 2010.
- [80] K. Koutsoumanis, A. Pavlis, G. J. E. Nychas, and K. Xanthiakos, “Probabilistic model for *Listeria monocytogenes* growth during distribution, retail storage, and domestic storage of pasteurized milk,” *Applied and Environmental Microbiology*, vol. 76, no. 7, pp. 2181–2191, 2010.
- [81] L. R. Granados, F. Pérez, and A. Valero, “Food quality management systems in the dairy industry: a case study on the application of predictive microbiology in the microbial quality of milks,” in *Technological Approaches for Novel Applications in Dairy Processing*, IntechOpen, London, UK, 2018.
- [82] L. Petruzzi, D. Campaniello, M. R. Corbo et al., “Wine microbiology and predictive microbiology: a short overview on application, and perspectives,” *Microorganisms*, vol. 10, no. 2, p. 421, 2022.
- [83] M. Ziyaina, B. N. Govindan, B. Rasco, T. Coffey, and S. S. Sablani, “Monitoring shelf life of pasteurized whole milk under refrigerated storage conditions: predictive models for quality loss,” *Journal of Food Science*, vol. 83, no. 2, pp. 409–418, 2018.
- [84] S. Flint, P. Bremer, J. Brooks et al., “Bacterial fouling in dairy processing,” *International Dairy Journal*, vol. 101, Article ID 104593, 2020.
- [85] S. Jindal, S. Anand, K. Huang, J. Goddard, L. Metzger, and J. Amamcharla, “Evaluation of modified stainless steel surfaces targeted to reduce biofilm formation by common milk spore formers,” *Journal of Dairy Science*, vol. 99, no. 12, pp. 9502–9513, 2016.
- [86] M. Gougouli, K. Kalantzi, E. Beletsiotis, and K. P. Koutsoumanis, “Development and application of predictive models for fungal growth as tools to improve quality control in yogurt production,” *Food Microbiology*, vol. 28, no. 8, pp. 1453–1462, 2011.
- [87] J. Kowalik and A. Lobacz, “Development of a predictive model describing the growth of *Yersinia enterocolitica* in Camembert-type cheese,” *International Journal of Food Science and Technology*, vol. 50, no. 3, pp. 811–818, 2015.

- [88] S. Benito, "The management of compounds that influence human health in modern winemaking from an HACCP point of view," *Fermentation*, vol. 5, no. 2, p. 33, 2019.
- [89] C. N. Haas, J. B. Rose, and C. P. Gerba, *Quantitative Microbial Risk Assessment*, John Wiley & Sons, Hoboken, NJ, USA, 2014.
- [90] A. García-Moraleja, G. Font, J. Mañes, and E. Ferrer, "Analysis of mycotoxins in coffee and risk assessment in Spanish adolescents and adults," *Food and Chemical Toxicology*, vol. 86, pp. 225–233, 2015.
- [91] J. M. Membre and S. Guillou, "Latest developments in foodborne pathogen risk assessment," *Current Opinion in Food Science*, vol. 8, pp. 120–126, 2016.
- [92] R. Pouillot and M. B. Lubran, "Predictive microbiology models vs. modeling microbial growth within *Listeria monocytogenes* risk assessment: what parameters matter and why," *Food Microbiology*, vol. 28, no. 4, pp. 720–726, 2011.
- [93] S. C. González, A. Possas, E. Carrasco et al., "MicroHibro: A software tool for predictive microbiology and microbial risk assessment in foods," *International Journal of Food Microbiology*, vol. 290, pp. 226–236, 2019.
- [94] F. Tenenhaus-Aziza and M. Ellouze, "Software for predictive microbiology and risk assessment: a description and comparison of tools presented at the ICPMF8 Software Fair," *Food Microbiology*, vol. 45, pp. 290–299, 2015.
- [95] M. H. Cassin, A. M. Lammerding, E. C. Todd, W. Ross, and R. S. McColl, "Quantitative risk assessment for *Escherichia coli* O157: H7 in ground beef hamburgers," *International Journal of Food Microbiology*, vol. 41, no. 1, pp. 21–44, 1998.
- [96] P. Dalgaard and H. Henrik Huss, "Mathematical modeling used for evaluation and prediction of microbial fish spoilage," in *Seafood Safety, Processing, and Biotechnology*, pp. 73–89, CRC Press, Boca Raton, FL, USA, 2020.
- [97] M. Mataragas, V. Dimitriou, P. N. Skandamis, and E. H. Drosinos, "Quantifying the spoilage and shelf-life of yoghurt with fruits," *Food Microbiology*, vol. 28, no. 3, pp. 611–616, 2011.
- [98] S. Limbo, L. Torri, N. Sinelli, L. Franzetti, and E. Casiraghi, "Evaluation and predictive modeling of shelf life of minced beef stored in high-oxygen modified atmosphere packaging at different temperatures," *Meat Science*, vol. 84, no. 1, pp. 129–136, 2010.
- [99] P. Rasane, A. Jha, and N. Sharma, "Predictive modeling for shelf-life determination of nutriceal-based fermented baby food," *Journal of Food Science and Technology*, vol. 52, pp. 5003–5011, 2015.
- [100] S. Bruckner, A. Albrecht, B. Petersen, and J. Kreyenschmidt, "A predictive shelf life model as a tool for the improvement of quality management in pork and poultry chains," *Food Control*, vol. 29, no. 2, pp. 451–460, 2013.
- [101] J. Kreyenschmidt, A. Hübner, E. Beierle, L. Chonsch, A. Scherer, and B. Petersen, "Determination of the shelf life of sliced cooked ham based on the growth of lactic acid bacteria in different steps of the chain," *Journal of Applied Microbiology*, vol. 108, no. 2, pp. 510–520, 2010.
- [102] T. Ross, J. Baranyi, and T. A. McMeekin, "Predictive microbiology and food safety," in *Encyclopaedia of Food Microbiology*, R. Robinson, C. A. Batt, and P. Patel, Eds., pp. 1699–1710, Academic Press, London, UK, 2000.
- [103] T. Ross and T. A. McMeekin, "Modeling microbial growth within food safety risk assessments," *Risk Analysis: International Journal*, vol. 23, no. 1, pp. 179–197, 2003.
- [104] P. Dalgaard, "Qualitative and quantitative characterization of spoilage bacteria from packed fish," *International Journal of Food Microbiology*, vol. 26, no. 3, pp. 319–333, 1995.
- [105] A. Hassoun and R. Karoui, "Quality evaluation of fish and other seafood by traditional and nondestructive instrumental methods: advantages and limitations," *Critical Reviews in Food Science and Nutrition*, vol. 57, no. 9, pp. 1976–1998, 2017.
- [106] L. Aldars-García, M. Berman, J. Ortiz, A. J. Ramos, and S. Marín, "Probability models for growth and aflatoxin B1 production as affected by intraspecies variability in *Aspergillus flavus*," *Food Microbiology*, vol. 72, pp. 166–175, 2018.
- [107] J. Van Impe, F. Poschet, B. M. Nicolai, and A. Geeraerd, "S & P-type models: a novel class of predictive microbial growth models," in *13th World Congress of Food Science & Technology 2006*, p. 1134, IUFOST World Congress, Leuven, Belgium, 2006.
- [108] K. P. Koutsoumanis and Z. Aspidou, "Moving towards a risk-based food safety management," *Current Opinion in Food Science*, vol. 12, pp. 36–41, 2016.
- [109] C. Pin, A. Metris, and J. Baranyi, "Next generation of predictive models," in *Advances in Microbial Food Safety*, J. Sofos, Ed., pp. 498–515, Woodhead Publ, Cambridge, UK, 2013.
- [110] R. L. Buchanan and M. H. Golden, "Critical factors for predictive microbiology," in *Compendium of Methods for the Microbiological Examination of Foods*, pp. 1–14, American Public Health Association, Washington, DC, USA, 5th edition, 2013.
- [111] S. S. T. Karamcheti, "Exploring the opportunities for incorporating dynamic Bayesian modeling in real-world food microbiology applications," University of Otago, Otago, New Zealand, 2022, Doctoral dissertation.
- [112] T. Ross, "Limitations to the predictive value of microbial challenge testing," *International Journal of Food Microbiology*, vol. 210, pp. 16–21, 2015.
- [113] K. P. Koutsoumanis and Z. Aspidou, "Individual cell heterogeneity in predictive food microbiology: challenges in predicting a "noisy" world," *International Journal of Food Microbiology*, vol. 240, pp. 3–10, 2017.
- [114] Z. Lin, X. Qin, J. Li et al., "Machine learning approach for predicting single cell lag time of *Salmonella* Enteritidis after heat and chlorine treatment," *Food Research International*, vol. 156, Article ID 111132, 2022.
- [115] J. S. Aguirre, M. R. Rodriguez, and G. D. G. de Fernando, "Effects of electron beam irradiation on the variability in survivor number and duration of lag phase of four foodborne organisms," *International Journal of Food Microbiology*, vol. 149, no. 3, pp. 236–246, 2011.
- [116] K. Faust, "Metagenomics meets time series analysis: unravelling microbial community dynamics," *Current Opinion in Microbiology*, vol. 25, pp. 56–66, 2015.
- [117] S. Athmaja, M. Hanumanthappa, and V. Kavitha, "A survey of machine learning algorithms for big data analytics," in *Proceedings of the 2017 International conference on innovations in information, embedded and communication systems (ICIIECS)*, pp. 1–4, IEEE, Coimbatore, India, March, 2017.
- [118] E. Franz, P. Delaquis, S. Morabito et al., "Exploiting the explosion of information associated with whole genome sequencing to tackle Shiga toxin-producing *Escherichia coli* (STEC) in global food production systems," *International Journal of Food Microbiology*, vol. 187, pp. 57–72, 2014.
- [119] C. Baudrit, M. Sicard, P. H. Wuillemin, and N. Perrot, "Towards a global modelling of the Camembert-type cheese ripening process by coupling heterogeneous knowledge with dynamic Bayesian networks," *Journal of Food Engineering*, vol. 98, no. 3, pp. 283–293, 2010.

- [120] S. Brul, F. I. Mensonides, K. J. Hellingwerf, and M. J. T. de Mattos, "Microbial systems biology: new frontiers open to predictive microbiology," *International Journal of Food Microbiology*, vol. 128, no. 1, pp. 16–21, 2008.
- [121] Z. Aspidou and K. P. Koutsoumanis, "Individual cell heterogeneity as variability source in population dynamics of microbial inactivation," *Food Microbiology*, vol. 45, pp. 216–221, 2015.
- [122] S. D. Papagianeli, A. Lianou, Z. Aspidou, L. Stathas, and K. Koutsoumanis, "The magnitude of heterogeneity in individual-cell growth dynamics is an inherent characteristic of *Salmonella enterica* ser. Typhimurium strains," *Food Research International*, vol. 162, Article ID 111991, 2022.
- [123] Y. Hara-Kudo and K. Takatori, "Contamination level and ingestion dose of foodborne pathogens associated with infections," *Epidemiology and Infection*, vol. 139, no. 10, pp. 1505–1510, 2011.
- [124] N. Tegtmeier, M. Rohde, and S. Backert, "Clinical presentations and pathogenicity mechanisms of bacterial foodborne infections," in *Microbial Food Safety: An Introduction*, pp. 13–31, Springer New York, New York, NY, USA, 2011.
- [125] M. Ackermann, "A functional perspective on phenotypic heterogeneity in microorganisms," *Nature Reviews Microbiology*, vol. 13, no. 8, pp. 497–508, 2015.
- [126] J. C. Augustin, "Challenges in risk assessment and predictive microbiology of food-borne spore-forming bacteria," *Food Microbiology*, vol. 28, no. 2, pp. 209–213, 2011.
- [127] J. Giacometti and D. Josic, "Foodomics in microbial safety," *TrAC, Trends in Analytical Chemistry*, vol. 52, pp. 16–22, 2013.
- [128] A. B. Hassena, J. Haendiges, S. Zormati et al., "Virulence and resistance genes profiles and clonal relationships of non-typhoidal food-borne *Salmonella* strains isolated in Tunisia by whole genome sequencing," *International Journal of Food Microbiology*, vol. 337, Article ID 108941, 2021.
- [129] O. A. Olowe, O. J. Adefioye, T. A. Ajayeoba et al., "Phylogenetic grouping and biofilm formation of multidrug resistant *Escherichia coli* isolates from humans, animals and food products in South-West Nigeria," *Science Africa*, vol. 6, Article ID e00158, 2019.
- [130] Z. Zakaria, L. Hassan, N. Ahmad et al., "Discerning the antimicrobial resistance, virulence, and phylogenetic relatedness of *Salmonella* isolates across the human, poultry, and food materials sources in Malaysia," *Frontiers in Microbiol.*, vol. 12, Article ID 652642, 2021.
- [131] P. M. K. Njage, P. Leekitcharoenphon, and T. Hald, "Improving hazard characterization in microbial risk assessment using next generation sequencing data and machine learning: predicting clinical outcomes in shigatoxigenic *Escherichia coli*," *International Journal of Food Microbiology*, vol. 292, pp. 72–82, 2019.
- [132] J. M. Spinale, R. L. Ruebner, L. Copelovitch, and B. S. Kaplan, "Long-term outcomes of Shiga toxin hemolytic uremic syndrome," *Pediatric Nephrology*, vol. 28, pp. 2097–2105, 2013.
- [133] A. Pielat, M. P. Boer, L. M. Wijnands et al., "First step in using molecular data for microbial food safety risk assessment; hazard identification of *Escherichia coli* O157: H7 by coupling genomic data with in vitro adherence to human epithelial cells," *International Journal of Food Microbiology*, vol. 213, pp. 130–138, 2015.
- [134] J. A. Carriço, A. J. Sabat, A. W. Friedrich, M. Ramirez, and on behalf of the Escmid Study Group for Epidemiological Markers C, "Bioinformatics in bacterial molecular epidemiology and public health: databases, tools and the next-generation sequencing revolution," *Euro Surveillance*, vol. 18, no. 4, 2013.
- [135] M. W. Libbrecht and W. S. Noble, "Machine learning application in genetics and genomics," *Nature Reviews Genetics*, vol. 16, no. 6, pp. 321–332, 2015.
- [136] S. Okser, T. Pahikkala, and T. Aittokallio, "Genetic variants and their interactions in disease risk prediction—machine learning and network perspectives," *BioData Mining*, vol. 6, no. 1, pp. 1–16, 2013.
- [137] S. Kuroda, H. Okuda, W. Ishida, and S. Koseki, "Modeling growth limits of *Bacillus* spp. spores by using deep-learning algorithm," *Food Microbiology*, vol. 78, pp. 38–45, 2019.
- [138] A. Métris, S. M. George, B. M. Mackey, and J. Baranyi, "Modeling the variability of single-cell lag times for *Listeria innocua* populations after sublethal and lethal heat treatments," *Applied and Environmental Microbiology*, vol. 74, no. 22, pp. 6949–6955, 2008.
- [139] I. Lvovich, Y. Lvovich, A. Preobrazhenskiy, Y. Preobrazhenskiy, and O. Choporov, "Manag. dev. internet of things systems based on classification and predictive models and algorithms," in *Proceedings of the 2021 International Seminar on Electrical Devices Design and Production (SED)*, pp. 1–5, IEEE, Prague, Czech Republic, April, 2021.
- [140] L. Lu, W. Zheng, Z. Lv, and Y. Tang, "Dev. and appl. of time-temp. indicators used on food during the cold chain logistics," *Packaging Technological Science*, vol. 26, pp. 80–90, 2013.
- [141] S. J. Goodswen, J. L. Barratt, P. J. Kennedy, A. Kaufer, L. Calarco, and J. T. Ellis, "Machine learning and applications in microbiology," *FEMS Microbiology Reviews*, vol. 45, no. 5, Article ID fuab015, 2021.
- [142] D. S. Yoo, Y. H. Song, D. W. Choi, J. S. Lim, K. Lee, and T. Kang, "Machine learning-driven dynamic risk prediction for highly pathogenic avian influenza at poultry farms in Republic of Korea: daily risk estimation for individual premises," *Transboundary and Emerging Diseases*, vol. 69, no. 5, pp. 2667–2681, 2022.
- [143] N. Ndraha, H. I. Hsiao, Y. Z. Hsieh, and A. K. Pradhan, "Predictive models for the effect of environmental factors on the abundance of *Vibrio parahaemolyticus* in oyster farms in Taiwan using extreme gradient boosting," *Food Control*, vol. 130, Article ID 108353, 2021.
- [144] L. Brillante, F. Gaiotti, L. Lovat et al., "Investigating the use of gradient boosting machine, random forest and their ensemble to predict skin flavonoid content from berry physical-mechanical characteristics in wine grapes," *Computers and Electronics in Agriculture*, vol. 117, pp. 186–193, 2015.
- [145] C. K. Tanui, E. O. Benefo, S. Karanth, and A. K. Pradhan, "A machine learning model for food source attribution of *Listeria monocytogenes*," *Pathogens*, vol. 11, no. 6, p. 691, 2022.
- [146] D. Dourou, "Rapid microbial quality assessment of chicken liver inoculated or not with *Salmonella* Using FTIR spectroscopy and machine learning," *Frontiers in Microbiology*, vol. 11, Article ID 623788, 2021.
- [147] R. Badia-Melis, U. Mc Carthy, and I. Uysal, "Data estimation methods for predicting temperatures of fruit in refrigerated containers," *Biosystems Engineering*, vol. 151, pp. 261–272, 2016.
- [148] S. Mercier and I. Uysal, "Neural network models for predicting perishable food temperatures along the supply chain," *Biosystems Engineering*, vol. 171, pp. 91–100, 2018.

- [149] E. Mohebi and L. Marquez, "Application of machine learning and RFID in the stability optimization of perishable foods," 2014, https://www.academia.edu/28109063/Application_of_Machine_Learning_and_RFID_in_the_Stability_Optimization_of_Perishable_Foods.
- [150] N. Munck, P. M. K. Njage, P. Leekitcharoenphon, E. Litrup, and T. Hald, "Application of whole-genome sequences and machine learning in source attribution of *Salmonella Typhimurium*," *Risk Analysis*, vol. 40, no. 9, pp. 1693–1705, 2020.
- [151] S. Karanth, C. K. Tanui, J. Meng, and A. K. Pradhan, "Exploring the predictive capability of advanced machine learning in identifying severe disease phenotype in *Salmonella enterica*," *Food Research International*, vol. 151, Article ID 110817, 2022.
- [152] N. Arning, S. K. Sheppard, S. Bayliss, D. A. Clifton, and D. J. Wilson, "Machine learning to predict the source of campylobacteriosis using whole genome data," *PLoS Genetics*, vol. 17, no. 10, Article ID e1009436, 2021.
- [153] K. Koyama, K. Kubo, S. Hiura, and S. Koseki, "Is skipping the definition of primary and secondary models possible? Prediction of *Escherichia coli* O157 growth by machine learning," *Journal of Microbiological Methods*, vol. 192, Article ID 106366, 2022.
- [154] J. O. Ogutu, H. P. Piepho, and T. Schulz-Streeck, "A comparison of random forests, boosting and support vector machines for genomic selection," in *BMC Proc*, vol. 5, no. 3, pp. 1–5, BioMed Central, London, UK, 2011.
- [155] Ş. M. Petrea, M. Costache, D. Cristea et al., "A machine learning approach in analyzing bioaccumulation of heavy metals in turbot tissues," *Molecules*, vol. 25, no. 20, p. 4696, 2020.
- [156] W. Gu, A. R. Vieira, R. M. Hoekstra, P. M. Griffin, and D. Cole, "Use of random forest to estimate population attributable fractions from a case-control study of *Salmonella enterica* serotype Enteritidis infections," *Epidemiology and Infection*, vol. 143, no. 13, pp. 2786–2794, 2015.
- [157] S. Chen, F. Tao, C. Pan et al., "Modeling quality changes in Pacific white shrimp (*Litopenaeus vannamei*) during storage: comparison of the Arrhenius model and Random Forest model," *Journal of Food Processing and Preservation*, vol. 45, no. 1, Article ID e14999, 2021.
- [158] S. D. Astuti, M. H. Tamimi, A. A. Pradhana et al., "Gas sensor array to classify the chicken meat with *E. coli* contaminant by using random forest and support vector machine," *Biosensors and Bioelectronics X*, vol. 9, Article ID 100083, 2021.
- [159] M. Kuhn, "Building predictive models in R using the caret package," *Journal of Statistical Software*, vol. 28, pp. 1–26, 2008.
- [160] A. Pradhan, "Support vector machine-a survey," *International Journal of Emerging Technology and Advanced Engineering*, vol. 2, no. 8, pp. 82–85, 2012.
- [161] A. García-Florian, C. López-Martín, C. Yáñez-Márquez, and A. Abran, "Support vector regression for predicting software enhancement effort," *Information and Software Technology*, vol. 97, pp. 99–109, 2018.
- [162] B. Mahesh, "Machine learning algorithms-a review," *International Journal of Science and Research*, vol. 9, no. 1, pp. 381–386, 2020.
- [163] B. Scholkopf and A. J. Smola, *Learning with Kernels: Support Vector Machines, Regularization, Optimization, and beyond*, MIT press, Cambridge, MA, USA, 2018.
- [164] S. Amarappa and S. V. Sathyanarayana, "Data classification using Support vector Machine (SVM), a simplified approach," *International Journal of Electron Computing Science and Engineering*, vol. 3, pp. 435–445, 2014.
- [165] E. Naranjo Bejarano, "Kernels Methods in Machine Learning," 2022, <https://arxiv.org/pdf/math/0701907>.
- [166] Y. Ma, Y. Hou, Y. Liu, and Y. Xue, "Research of food safety risk assessment methods based on big data," in *Proceedings of the 2016 IEEE International Conference on Big Data Analysis (ICBDA)*, pp. 1–5, IEEE, Hangzhou, China, March, 2016.
- [167] E. Bonah, X. Huang, Y. Hongying et al., "Detection of *Salmonella Typhimurium* contamination levels in fresh pork samples using electronic nose smellprints in tandem with support vector machine regression and metaheuristic optimization algorithms," *Journal of Food Science and Technology*, vol. 58, pp. 3861–3870, 2021.
- [168] S. Karsoliya, "Approximating number of hidden layer neurons in multiple hidden layer BPNN architecture," *International Journal of Engineering Trends and Technology*, vol. 3, no. 6, pp. 714–717, 2012.
- [169] Q. Sun, M. Zhang, A. S. Mujumdar, and P. Yang, "Combined LF-NMR and artificial intelligence for continuous real-time monitoring of carrot in microwave vacuum drying," *Food and Bioprocess Technology*, vol. 12, pp. 551–562, 2019.
- [170] A. Torres-Barrán, Á. Alonso, and J. R. Dorronsoro, "Regression tree ensembles for wind energy and solar radiation prediction," *Neurocomputing*, vol. 326, pp. 151–160, 2019.
- [171] T. Sheng, S. Shi, Y. Zhu, D. Chen, and S. Liu, "Analysis of protein and fat in milk using multiwavelength gradient-boosted regression tree," *IEEE Transactions on Instrumentation and Measurement*, vol. 71, pp. 1–10, 2022.
- [172] K. H. Chu, "Fitting the Gompertz equation to asymmetric breakthrough curves," *Journal of Environmental Chemical Engineering*, vol. 8, no. 3, Article ID 103713, 2020.
- [173] M. Kuhn and K. Johnson, *Applied Predictive Modeling*, Springer, New York, NY, USA, 2013.
- [174] E. O. Benefo, S. Karanth, and A. K. Pradhan, "A machine learning approach to identifying *Salmonella* stress response genes in isolates from poultry processing," *Food Research International*, vol. 175, Article ID 113635, 2024.
- [175] H. Pang, R. McEgan, A. Mishra, S. A. Micallef, and A. K. Pradhan, "Identifying and modeling meteorological risk factors associated with pre-harvest contamination of *Listeria* species in a mixed produce and dairy farm," *Food Research International*, vol. 102, pp. 355–363, 2017.
- [176] D. Hwang, M. J. Rothrock Jr, H. Pang, M. Guo, and A. Mishra, "Predicting *Salmonella* prevalence associated with meteorological factors in pastured poultry farms in southeastern United States," *Science of the Total Environment*, vol. 713, Article ID 136359, 2020.
- [177] X. Xu, M. J. Rothrock Jr, J. Reeves, G. D. Kumar, and A. Mishra, "Using *E. coli* population to predict foodborne pathogens in pastured poultry farms," *Food Microbiology*, vol. 108, Article ID 104092, 2022.
- [178] K. B. Biji, C. N. Ravishankar, C. O. Mohan, and T. K. Srinivasa Gopal, "Smart packaging systems for food applications: a review," *Journal of Food Science and Technology*, vol. 52, pp. 6125–6135, 2015.
- [179] T. O. Hafliðason, G. Ólafsdóttir, S. Bogason, and G. Stefánsson, "Criteria for temperature alerts in cod supply chains," *International Journal of Physical Distribution & Logistics Management*, vol. 42, no. 4, pp. 355–371, 2012.
- [180] M. C. do Nascimento Nunes, M. Nicometo, J. P. Emond, R. B. Melis, and I. Uysal, "Improvement in fresh fruit and vegetable logistics quality: berry logistics field studies," *Philosophical Transactions of the Royal Society A:*

- Mathematical, Physical & Engineering Sciences*, vol. 372, no. 2017, Article ID 20130307, 2014.
- [181] J. Huo, T. Shi, and J. Chang, "Comparison of Random Forest and SVM for electrical short-term load forecast with different data sources," in *Proceedings of the 2016 7th IEEE International Conference on Software Engineering and Service Science (ICSESS)*, pp. 1077–1080, IEEE, Beijing, China, August, 2016.
- [182] H.-xia Luo, D. A. I. Sheng-pei, L. I. Mao-fen et al., "Comparison of machine learning algorithms for mapping mango plantations based on Gaofen-1 imagery," *Journal of Integrative Agriculture*, vol. 19, no. 11, pp. 2815–2828, 2020.
- [183] T. M. Amado, M. R. Bunuan, R. F. Chicote et al., "Development of predictive models using machine learning algorithms for food adulterants bacteria detection," in *Proceedings of the 2019 IEEE 11th International Conference on Humanoid, Nanotechnology, Information Technology, Communication and Control, Environment, and Management (HNICEM)*, pp. 1–6, IEEE, Laoag, Philippines, November, 2019.
- [184] M. C. Tirado, R. Clarke, L. A. Jaykus, A. McQuatters-Gollop, and J. M. Frank, "Climate change and food safety: a review," *Food Research International*, vol. 43, no. 7, pp. 1745–1765, 2010.
- [185] FAO (food and agriculture organization of the united Nations), *Climate Change: Implications for Food Safety*, FAO, Rome, Italy, 2008, <http://www.fao.org/docrep/010/i0195e/i0195e00.htm>.
- [186] D. A. Hutchins, J. K. Jansson, J. V. Remais, V. I. Rich, B. K. Singh, and P. Trivedi, "Climate change microbiology problems and perspectives," *Nature Reviews Microbiology*, vol. 17, no. 6, pp. 391–396, 2019.
- [187] S. Bandyopadhyay, S. Kanji, and L. Wang, "The impact of rainfall and temperature variation on diarrheal prevalence in Sub-Saharan Africa," *Applied Geography*, vol. 33, pp. 63–72, 2012.
- [188] M. Azage, A. Kumie, A. Worku, A. Bagtzoglou, and E. Anagnostou, "Effect of climatic variability on childhood diarrhea and its high risk periods in northwestern parts of Ethiopia," *PLoS One*, vol. 12, no. 10, Article ID e0186933, 2017.
- [189] N. Li, Y. Wang, C. You et al., "Variation in raw milk microbiota throughout 12 months and the impact of weather conditions," *Scientific Reports*, vol. 8, no. 1, p. 2371, 2018.
- [190] M. Van der Spiegel, H. J. Van der Fels-Klerx, and H. J. P. Marvin, "Effects of climate change on food safety hazards in the dairy production chain," *Food Research International*, vol. 46, no. 1, pp. 201–208, 2012.
- [191] C. A. Becker, R. J. Collier, and A. E. Stone, "Invited review: physiological and behavioral effects of heat stress in dairy cows," *Journal of Dairy Science*, vol. 103, no. 8, pp. 6751–6770, 2020.
- [192] N. de, C. Maertens, B. Devleeschauwer et al., "Burden of salmonellosis, campylobacteriosis and listeriosis: a time series analysis, Belgium, 2012 to 2020," *Euro Surveillance*, vol. 22, no. 38, Article ID 30615, 2017.
- [193] M. El-Fadel, S. Ghanimeh, R. Maroun, and I. Alameddine, "Climate change and temperature rise: implications on food-and water-borne diseases," *Science of the Total Environment*, vol. 437, pp. 15–21, 2012.
- [194] Centers for Disease Control and Prevention (Cdc), "Outbreaks of *Escherichia coli* O157: H7 infection and cryptosporidiosis associated with drinking unpasteurized apple cider--Connecticut and New York, October 1996," *MMWR. Morbidity and mortality weekly report*, vol. 46, no. 1, pp. 4–8, 1997.
- [195] C. N. Okafo, V. J. Umoh, and M. Galadima, "Occurrence of pathogens on vegetables harvested from soils irrigated with contaminated streams," *Science of the Total Environment*, vol. 311, no. 1-3, pp. 49–56, 2003.