Correction: Multiarm multistage (MAMS) randomised selection designs: impact of treatment selection rules on the operating characteristics

ChoodariOskooei, Babak; Blenkinsop, Alexandra; Handley, Kelly; Pinkney, Thomas; Parmar, Mahesh K. B.

[Link to publication on Research at Birmingham portal](#)

CORRECTION

Correction: Multi-arm multi-stage (MAMS) randomised selection designs: impact of treatment selection rules on the operating characteristics

Babak Choodari-Oskooei[1*†], Alexandra Blenkinsop[2†], Kelly Handley[3], Thomas Pinkney[4] and
Mahesh K. B. Parmar[1]

**Correction: BMC Med Res Methodol 24, 124 (2024)
https://doi.org/10.1186/s12874-024-02247-w**

Following publication of the original article [1], the authors reported an error found on page 4 and replacement of Fig. 2.

1) On page 4, the following text "\sigma _{\hat{\theta}_ {jk}}" should be replaced with $\sigma_{\widehat{\theta}_{jk}}$.
2) On page 4, the "ψ_jk=1" should be replaced with $\psi_{jk} = 1$.
3) In Fig. 2, the title of the two bottom graphs, "a-1" and "b-1" should be replaced with "a-2" and "b-2", respectively. The revised version of Fig. 2 is shown below.

The original article [1] has been updated.

Published online: 02 July 2024

Reference
1. Choodari-Oskooei B, Blenkinsop A, Handley K, et al. Multi-arm multi-stage (MAMS) randomised selection designs: impact of treatment selection rules on the operating characteristics. BMC Med Res Methodol. 2024;24:124. https://doi.org/10.1186/s12874-024-02247-w.

†Babak Choodari-Oskooei and Alexandra Blenkinsop contributed equally to this work.

The original article can be found online at https://doi.org/10.1186/s12874-024-02247-w.

*Correspondence:
Babak Choodari-Oskooei
b.choodari-oskooei@ucl.ac.uk
1 MRC Clinical Trials Unit at UCL, Institute of Clinical Trials and Methodology, UCL, 90 High Holborn, London WC1V 6LJ, UK
2 Department of Mathematics, Imperial College London, London, UK
3 Birmingham Clinical Trials Unit, University of Birmingham, Birmingham, UK
4 Institute of Applied Health Research, University of Birmingham, Birmingham, UK

Fig. 2 FWER (left) and overall power (right) by the timing of the treatment selection at stage 1 (top) and stage 2 (bottom) and subset selection rule for a three-stage design. The overall power is calculated when one research arm is effective with the target effect size. The X-axis is control arm information time in all graphs