



Assessment of Academic Resilience and its associated factors among Pharmacy Students in Twelve Countries

Elnaem, M. H., Asikin bt Wan Salam, W. N. A., Thabit, A. K., Mubarak, N., AbouKhatwa, M. M., Ramatillah, D. L., Isah, A., Barakat, M., Al-Jumaili, A. A., Mansour, N. O., Fathelrahman, A., Adam, M. F., Jamil, S., Baraka, M., Rabbani, S. A., Abdelaziz, D. H., Elrggal, M. E., Okuyan, B., & Elcioglu, H. K. (2024). Assessment of Academic Resilience and its associated factors among Pharmacy Students in Twelve Countries. *American Journal of Pharmaceutical Education*, 88(5), 1-7. Article 100693. Advance online publication. <https://doi.org/10.1016/j.ajpe.2024.100693>

[Link to publication record in Ulster University Research Portal](#)

Published in:

American Journal of Pharmaceutical Education

Publication Status:

Published online: 02/04/2024

DOI:

[10.1016/j.ajpe.2024.100693](https://doi.org/10.1016/j.ajpe.2024.100693)

Document Version

Publisher's PDF, also known as Version of record

General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

Research

Assessment of Academic Resilience and Its Associated Factors Among Pharmacy Students in Twelve Countries



Mohamed Hassan Elnaem^{a,*}, Wan Nur Ain Asikin Wan Salam^b, Abrar K. Thabit^c, Naeem Mubarak^d, Merna Mahmoud Abou Khatwa^e, Diana Laila Ramatillah^f, AbdulMuminu Isah^g, Muna Barakat^h, Ali Azeez Al-Jumailiⁱ, Noha O. Mansour^j, Ahmed Ibrahim Fathelrahman^k, Mohammed Fathelrahman Adam^l, Safayat Jamil^m, Mohamed Barakaⁿ, Syed Arman Rabbani^o, Doaa H. Abdelaziz^{p,t}, Mahmoud E. Elrggal^q, Betul Okuyan^r, Hatice Kübra Elcioglu^s

^a Ulster University, School of Pharmacy and Pharmaceutical Sciences, Coleraine, United Kingdom

^b International Islamic University Malaysia, Faculty of Pharmacy, Department of Pharmacy Practice, Kuantan, Pahang, Malaysia

^c King Abdulaziz University, Faculty of Pharmacy, Pharmacy Practice Department, Jeddah, Saudi Arabia

^d University of Health Sciences, Lahore Medical & Dental College, Department of Pharmacy Practice, Lahore, Pakistan

^e Alexandria University, Faculty of Pharmacy, Department of Clinical Pharmacy and Pharmacy Practice, Alexandria, Egypt

^f Universitas 17 Agustus 1945 Jakarta, Faculty of Pharmacy, Jakarta, Indonesia

^g University of Nigeria, Department of Clinical Pharmacy and Pharmacy Management, Nsukka, Nigeria

^h Applied Science Private University, Faculty of Pharmacy, Department of Clinical Pharmacy and Therapeutics, Amman, Jordan

ⁱ University of Baghdad College of Pharmacy, College of Pharmacy, Baghdad, Iraq

^j Mansoura University, Faculty of Pharmacy, Clinical Pharmacy and Pharmacy Practice Department, Mansoura, Egypt

^k Taif University, College of Pharmacy, Department of Clinical Pharmacy, Taif, Saudi Arabia

^l University of Science and Technology, Faculty of Pharmacy, Omdurman, Sudan

^m Khwaja Yunus Ali University, Department of Pharmacy, Sirajganj, Bangladesh

ⁿ Fatima College of Health Sciences, Department of Pharmacy, Abu Dhabi, United Arab Emirates

^o RAK Medical and Health Sciences University, RAK College of Pharmacy, Department of Clinical Pharmacy and Pharmacology, Ras Al-Khaimah, United Arab Emirates

^p Future University in Egypt, Faculty of Pharmacy, Pharmacy Practice & Clinical Pharmacy Department, Cairo, Egypt

^q Umm Al-Qura University, College of Medicine, Al-Qunfudah, Saudi Arabia

^r Marmara University, Faculty of Pharmacy, Department of Clinical Pharmacy, Istanbul, Türkiye

^s Marmara University, Faculty of Pharmacy, Department of Pharmacology, Istanbul, Türkiye

^t Al-Baha University, Faculty of Pharmacy, Department of Clinical Pharmacy, Al-Baha, Saudi Arabia.

ARTICLE INFO

Keywords:

Academic resilience
Pharmacy
Students
Education
International

ABSTRACT

Objective: Academic resilience, a critical determinant of academic achievement, is affected by various factors. There is a paucity of large-scale international assessments of academic resilience among pharmacy students. Therefore, this study aimed to assess academic resilience among pharmacy students in 12 countries and to evaluate factors associated with their academic resilience levels.

Methods: A cross-sectional online survey-based study was conducted among randomly selected pharmacy students in 12 countries: Egypt, Türkiye, Indonesia, Pakistan, Bangladesh, Iraq, Jordan, Nigeria, Malaysia, Saudi Arabia, Sudan, and the United Arab Emirates. After pilot testing, the validated 30-item academic resilience scale (ARS) was used for the assessment. The data were collected between November 1, 2022 and April 15, 2023. Descriptive and inferential statistics were performed, as appropriate.

Results: A total of 3950 were received from the 12 participating countries. The mean age was 21.68 ± 2.62 years. About two-thirds of the responses were from female participants and those studying for Bachelor of Pharmacy degrees. Overall, the findings show moderate academic resilience, which varied across countries. The median (IQR) of the total ARS-30 was 114 (103–124). Females exhibited lower negative affective and emotional response subscale levels than males. There were significant cross-country variations in the ARS-30 and all subscales. The highest overall levels were reported for Sudan, Pakistan, and Nigeria and the lowest were

* Corresponding author.

E-mail address: m.elnaem@ulster.ac.uk (M.H. Elnaem).

reported for Indonesia and Türkiye. Students in private universities tended to have higher overall ARS levels than public university students. Higher academic performance was significantly associated with ARS levels, whereas those with excellent performance exhibited the highest ARS levels. Students with exercise routines had higher ARS levels than those without exercise routines. Finally, students who were engaged in extracurricular activities had higher ARS levels than those who did not participate in these activities.

Conclusion: The study offers insights into the factors affecting academic resilience in pharmacy students across several countries. The findings could guide interventions and support activities to improve resilience and academic outcomes.

1. Introduction

Undergraduate pharmacy programs, similar to other health professional education programs, involve an integrated mix of multiple didactic courses, practical and laboratory sessions, numerous projects, and several months of virtual and on-site training before graduation. Such extensive educational programs have been shown in previous studies to be associated with stress, burnout, and negative impacts on mental health.^{1,2} To succeed academically, the ability to respond to difficulties and overcome challenges, setbacks, and adversity during the academic program (eg, low grades, difficult assignments, meeting deadlines, and examination pressure) defines academic resilience.³ Academic resilience is a positive indicator of mental well-being and personal growth.⁴

Previous studies have used different methods to evaluate academic resilience among undergraduate pharmacy students. However, the most common method is the academic resilience scale (ARS), a self-reported measure that assesses 4 dimensions of academic resilience.⁵ These include negative affect and emotional response (how students manage stress and negative emotions), adaptive help-seeking (how well students reflect on their experiences and seek help when needed), adaptive thought processes (how well students maintain a positive attitude and focus on their goals), and perseverance (how well students persist in the face of academic challenges).⁵

Academic resilience is a complex construct influenced by various factors, including personality, coping skills, social support, and life experiences.^{1,3} Previous studies found that factors such as younger age, higher grade point average, self-directed learning capacity, social support from family or friends, having time for fun, and engaging in mindfulness-based stress reduction therapy were associated with academic resilience.⁶⁻¹⁰ Among pharmacy students, academic resilience declined as the students advanced in the program, which was also associated with negative academic performance and a higher attrition rate.¹ Another study found that academic resilience positively correlated with academic achievement and success because it acts as a way to cope with academic burnout.¹¹ Therefore, achieving and maintaining academic resilience among pharmacy students is crucial to ensure successful academic outcomes.¹² Moreover, recent research on resilience among early career pharmacists has identified the beneficial role of activities and interventions that help build resilience during undergraduate studies.¹³

Few studies have evaluated academic resilience among undergraduate pharmacy students, given the unique structure of pharmacy programs and training compared with other health professional education programs, such as medicine and nursing. Therefore, this study aimed to assess academic resilience levels among undergraduate pharmacy students in 12 countries and to evaluate factors associated with their academic resilience levels (including their coping strategies and social support).

2. Methods

2.1. Study Design

A cross-sectional online survey-based study was conducted among pharmacy students in 12 countries with relatively large pharmacy

student populations in Asia, the Middle East, Africa, and Europe: Egypt, Türkiye, Indonesia, Pakistan, Bangladesh, Iraq, Jordan, Nigeria, Malaysia, Saudi Arabia, Sudan, and the United Arab Emirates. The study used a validated, self-administered survey prepared in English, Turkish, and Arabic on Google Forms. The study coinvestigators in each country disseminated all forms via private social media and other educational platforms, such as Microsoft Teams. Participants selected the language of the form and responded to only 1 version to avoid duplicate responses. The form settings were adjusted to limit only 1 response per participant. The data were collected between November 1, 2022 and April 15, 2023.

2.2. Inclusion and Exclusion Criteria

The study involved undergraduate pharmacy students who studied for at least 1 full semester at an accredited university in 1 participating country. Students from different pharmacy programs across all years of the study and training were eligible to participate in the study. All received responses were checked for completeness, and incomplete forms were excluded from the final data analysis.

2.3. Sample Size

This study was directed toward cross-country comparisons. Because there were considerable variations among countries regarding the number of pharmacy schools, target student population, and accessible sampling frames, the target sample varied across countries. Using the Raosoft sample size calculator, assuming an estimated proportion of 50% and a 95% CI and confirming that at least 1 principal pharmacy school will be participating in each country, the minimum required sample size was estimated to be 180 students at least in every setting. All countries met this successfully, with larger sample frames exceeding these minimum requirements. The data collection duration was extended in some countries compared with others to satisfy the sample size requirement.

2.4. Instrument Structure and Translation

The survey consisted of 3 sections: demographic data, items on common coping strategies and social support, and the validated ARS-30. After the consent form in the first section, questions on sex, age, country of residence, ethnicity, living arrangement, type of pharmacy program, type of institution, average exercise, sleep frequency, engaging in extracurricular activities (eg, arts, sports, personal skills, debate competitions, and community services), and cumulative grade point average were asked. The second section asked about coping strategies (eg, exercising and seeking religious advice) and social support (eg, their family, classmates, and institution). The third section required participants to answer the ARS-30. Formal permission to use the ARS's original and translated forms was given before the study initiation.^{14,15}

2.5. Validity and Pilot Testing

A panel of 7 academicians who are pharmacy practice experts evaluated the content validity for the first 2 sections, which are the

demographic part and questions regarding students' perception of coping strategies and social support. The questionnaire content validity was analyzed by estimating the content validity index for each item involving a minimum of 3 experts with at least a 0.78 content validity index score.¹⁶ The items that were < 0.78 for clarity were improved for better clarity according to the experts' suggestions, and no items were deleted. For the Turkish form, an expert panel ($n = 3$) for translation, cultural adaptation, and pilot testing for face validity among pharmacy students were performed.

The third section, ARS-30, was developed by Simon Cassidy and designated to evaluate students' reactions (cognitive-affective and behavioral) toward academic challenges.⁵ On a 5-point Likert scale, participants answered the questions based on their perception of a scenario or vignette because they will be asked to visualize themselves encountering an academic situation. The items were constructed using positive and negative wordings to perceive the reaction to distress. A 5-point Likert scale ranging from 1 (strongly agree) to 5 (strongly disagree) resides on each item. The overall score will be calculated by summation, where the theoretical range is 30 to 150, reflecting the level of academic resilience.⁵ The items represented 3 factors: perseverance, reflecting and adaptive help-seeking, and interpreted negative affect and emotional response. Perseverance is the first factor that highlights the perception of hardship as a chance to confront challenges and make improvements, strive, hold onto goals, embrace and use feedback, and creative problem-solving. The second factor is reflecting and the adaptive help-seeking attribute to assessing strengths and weaknesses, adjusting the study method and looking for support, help, and inspiration. The third factor is interpreted as the negative affect and emotional response that comprises anxiety, overthinking, optimism, vulnerability, and suppressing emotion. Previous research emphasized the role of these factors in shaping academic resilience and predicting academic achievement.^{17,18} All the reverse coding was done as per the developer's guide. Finally, the total ARS scores and individual factor scores were calculated for each participant.

2.6. Data Collection

The final survey was distributed to potential participants through the study coinvestigators via private social media and educational platforms (eg, WhatsApp and Microsoft Teams). In addition, a weekly reminder was sent throughout the period allocated for data collection until the target sample size was reached.

2.7. Statistical Analysis

The current study used the IBM SPSS Statistics for Windows, Version 28.0 (IBM Corp Released 2021. IBM SPSS Statistics for Windows, Version 28.0; IBM Corp) to analyze the data. The responses were subjected to descriptive statistics in terms of frequencies and percentages. Initially, the total ARS scores and individual factors' scores were computed. Internal consistency was measured by calculating Cronbach α . Subsequently, the Mann-Whitney was used to assess the significance of the differences in these scores across 2-group variables of participants' demographics, academic-related information, and coping and support strategies, given that normal distribution assumptions were not met. Similarly, the Kruskal-Wallis test was used to examine the score differences between variables with more than 2 groups. We adjusted the P values for multiple comparisons using the Bonferroni adjustment. The resulting adjusted P values were reported for each pairwise comparison. A P value of < .05 was considered statistically significant for all other comparisons.

2.8. Ethical Approval

Ethical approval for this study was initially provided by the International Islamic University Malaysia (IIUM) Research Ethics

Committee (2022–081). Further ethical approval and administrative agreements were obtained from coinvestigator institutions when required (Marmara University Faculty of Medicine Clinical Trials Ethical Committee - 06.01.2023/09.2023.109). The introductory page of the online form included the participation information sheet and informed consent that required approval before accessing the main survey. By

Table 1
General Information of Study Participants ($N = 3950$).

Characteristics	Frequency	Percent
Sex		
Female	2600	65.8
Male	1350	34.2
Country of residence		
Egypt	681	17.2
Pakistan	469	11.9
Nigeria	411	10.4
Indonesia	378	9.6
Bangladesh	355	9.0
Türkiye	293	7.4
Saudi Arabia	251	6.4
Sudan	249	6.3
Iraq	242	6.1
Malaysia	233	5.9
Jordan	221	5.6
United Arab Emirates	167	4.2
Race		
African (non-Arab)	424	10.7
Arab	1779	45.0
Asian (non-Arab)	1469	37.2
Caucasian	278	7.0
Living arrangement		
On campus (sharing hostel)	777	19.7
Living alone	317	8.0
Living with family	2362	59.8
Living with friend(s) outside campus	494	12.5
Type of institution		
Private (not funded by government)	1593	40.3
Public (funded by the government)	2357	59.7
Pharmacy program		
Bachelor's degree (BS Pharm)/with honors	2604	65.9
PharmD (Doctor of Pharmacy)	1346	34.1
Year of study		
Year 1	670	17.0
Year 2	517	13.1
Year 3	649	16.4
Year 4	1308	33.1
Year 5	712	18.0
Year 6	94	2.4
Academic performance up to the previous semester/annual examination		
Excellent	899	22.8
Very good	1420	35.9
Good	1056	26.7
Average	362	9.2
Below average	164	4.2
Poor	49	1.2
Exercise status		
No	1428	36.2
Yes. Irregular exercise	2170	54.9
Yes. Regular exercise	352	8.9
Participation in extracurricular activities (arts, sports, personal skills, debate competitions, and community service)		
Maybe (participate occasionally in some activities)	1790	45.3
No	1550	39.2
Yes (regularly participate in almost all activities)	610	15.4
The average number of hours sleep a day		
4 to 6 h/d	1990	50.4
7 to 10 h/d	1864	47.2
More than 10 h/d	96	2.4
Total	3950	100.0

Table 2
Assessment of Coping Strategies and Social Support (N = 3950).

Coping strategy/social support	1 (strongly agree)	2 (agree)	3 (neutral)	4 (disagree)	5 (strongly disagree)
Coping strategy					
Doing activities such as exercise or recreational activities helps you in dealing with problems or tough times.	1468 (37.2)	1637 (41.4)	645 (16.3)	142 (3.6)	58 (1.5)
Spiritual or religious support helps you to get through the problems or tough time.	2351 (59.5)	1157 (29.3)	321 (8.1)	69 (1.7)	52 (1.3)
Acting like nothing has happened helps you go through the problems.	675 (17.1)	1103 (27.9)	1087 (27.5)	723 (18.3)	362 (9.2)
Social support					
Your family supported you when you chose the pharmacy course.	2279 (57.7)	1119 (28.3)	392 (9.9)	96 (2.4)	64 (1.6)
You get the help and support you need from your family.	2139 (54.2)	1167 (29.5)	460 (11.6)	104 (2.6)	80 (2)
You get the help and support you need from your classmates.	967 (24.5)	1753 (44.4)	894 (22.6)	227 (5.7)	109 (2.8)
You get the help and support you need from your institution.	710 (18)	1250 (31.6)	1159 (29.3)	502 (12.7)	329 (8.3)

All values are given in n (%).

Table 3
Median and IQR of Total Academic Resilience (ARS-30) Scale and Subscales (N = 3950).

Score definition	Median	IQR	Max and min theoretical values
Total ARS score	114	103-124	30-150
Subscale 1 (perseverance)	56	51-61	14-70
Subscale 2 (reflecting and adaptive help-seeking):	36	33-41	9-45
Subscale 3 (negative affective and emotional response)	22	17-26	7-35

Abbreviations: ARS, academic resilience scale; max, maximum; min, minimum.

approving the consent form, the participants were deemed to have consented to participate in this research. They were also free to withdraw their consent during the study.

3. Results

3.1. Overall Characteristics of the Study Participants

A total of 4013 responses were initially received. Sixty-three forms were not complete and were excluded from the final analysis. Therefore, 3950 responses from the 12 participating countries were finally considered for formal analysis. The mean age was 21.68 ± 2.62 years. About two-thirds of the responses were from female participants and those studying for Bachelor of Pharmacy degrees. Approximately 60% belong to public universities. More than 50% of the students were in years 4 and 5 of their studies. One-third of the sample did not report exercise routines, and about 40% did not participate in extracurricular activities. Table 1 shows the overall participants' characteristics.

3.2. Coping and Support Strategies Among the Study Participants

The participants were asked how they had been supported or tried to cope with their academic-related demands. The highest agreements were for family support and help started with choosing the pharmacy program. Approximately 80% reported agreement with the role of exercise and activities in helping them overcome tough times. About 90% agreed with the positive impact of religious and spiritual support on their ability to handle problems. Table 2 shows an overview of the study participants' responses concerning key coping and support strategies.

3.3. Academic Resilience Scales and Subscales Reliability Analysis and Assessment Findings

A reliability analysis of the total scale and subscales was conducted to confirm the internal consistency of using the ASR-30 among our study participants. In this study, the results indicated that the ARS-30 has good internal consistency, with $\alpha = 0.874$ for the overall scale, $\alpha = 0.786$ for the perseverance subscale, $\alpha = 0.876$ for the reflecting

and adaptive help-seeking subscale, and $\alpha = 0.743$ for the negative affect and emotional response subscale. Overall, the findings show moderate academic resilience, which varied across countries. The median (IQR) of total ARS-30 was 114 (103–124). The perseverance subscale was 56 (51–61), whereas the reflecting and adaptive help-seeking subscale was 36 (33–41). Finally, the negative affective and emotional response subscale was 22 (17–26). Table 3 presents the median and IQRs of the overall ARS-30 and its 3 subscales.

3.4. Cross-tabulation of Mean Ranks of ARS-30 and Its 3 Subscales Across Various Participants' Characteristics

Overall, the ARS-30 and its subscales exhibited variations across different characteristics of participants. Females had lower levels in the third negative affective and emotional response subscale than males ($P = .003$). There were significant cross-country variations in the ARS-30 and all subscales. The highest overall levels were reported for Sudan, Pakistan, and Nigeria and the lowest were reported for Indonesia and Türkiye. Students in private universities tended to have higher overall ARS levels than public university students ($P = .035$). Higher academic performance was significantly associated with ARS levels, whereas those with excellent performance exhibited the highest ARS levels ($P < .001$). Students with exercise routines had higher ARS levels ($P < .001$) than those without exercise routines. Finally, students who were engaged in extracurricular activities had higher ARS levels than those who did not participate in these activities ($P < .001$). Table 4 presents the cross-tabulation of the overall ARS and its subscales against the participants' characteristics.

4. Discussion

To the best of our knowledge, the present study possibly reports the largest-scale academic resilience assessment in pharmacy education. We reported moderate levels of academic resilience that varied considerably across countries. The findings also reported variations in academic resilience levels and subscales based on sex, institution, year of study, academic performance, and participation in extracurricular activities. The study also described the status of coping and support strategies among participants, where exercise, religion, and family

Table 4
Associations With the Total Academic Resilience (ARS-30) Scale and Subscales Expressed in Mean Rank ($N = 3950$).

Characteristics	ARS-30 score	Subscale 1 (perseverance)	Subscale 2 (reflecting and adaptive help-seeking):	Subscale 3 (negative affective and emotional response)
Sex				
Female ($n = 2600$)	1971.16	1988.46	1977	1936.36
Male ($n = 1350$)	1983.86	1950.53	1972.60	2050.87
P value ^a	.740	.321	.908	.003
Country of residence				
Egypt ($n = 681$)	1969.04	1996.09	1940.95	1932.37
Pakistan ($n = 469$)	2205.78	2043.88	2237.76	2239.35
Nigeria ($n = 411$)	2339.85	2419.38	2135.85	2163.56
Indonesia ($n = 378$)	1544.78	1373.99	1752.42	1943.67
Bangladesh ($n = 355$)	1968.50	1841.18	2175.31	1970.83
Türkiye ($n = 293$)	1382.90	1569.67	1407.79	1438.80
Saudi Arabia ($n = 251$)	1826.81	2030.08	1805.22	1670.58
Sudan ($n = 249$)	2501.92	2450.40	2339.80	2408.06
Iraq ($n = 242$)	1917.14	1964.35	1747.81	2039.71
Malaysia ($n = 233$)	1954.74	2023.97	1928.62	1891.04
Jordan ($n = 221$)	1982.63	2045.41	2024.72	1877.47
United Arab Emirates ($n = 167$)	2030.62	2032.05	2104.42	1939.04
P value ^b	< .001	< .001	< .001	< .001
Race				
African (non-Arab) ($n = 424$)	2330.43	2406.03	2136.72	2155.70
Arab ($n = 1779$)	2021.67	2064.34	1976.56	1972.61
Asian (non-Arab) ($n = 1469$)	1930.05	1819	2038.93	2028.28
Caucasian ($n = 278$)	1378.84	1577.33	1387.63	1440.30
P value ^b	< .001	< .001	< .001	< .001
Living arrangement				
On campus (sharing hostel) ($n = 777$)	1996.56	2030.02	1994.90	1923.95
Living alone ($n = 317$)	1881.41	1867.64	1886.18	1985.82
Living with family ($n = 2362$)	1991.08	1990.35	1980.32	1990.31
Living with friend(s) outside campus ($n = 494$)	1928.27	1887.94	1979.27	1979.16
P value ^b	.294	.047	.527	.568
Type of institution				
Private (not funded by government) ($n = 1593$)	2021.99	1944.66	2097.33	2049.53
Public (funded by the government) ($n = 2357$)	1944.08	1996.34	1893.16	1925.47
P value ^a	.035	.162	< .001	< .001
Pharmacy program				
Bachelor's degree (BS Pharm)/with honors ($n = 2604$)	1967.73	1965.50	1965.98	1975.26
PharmD (Doctor of Pharmacy) ($n = 1346$)	1990.53	1994.84	1993.91	1975.97
P value ^a	.551	.443	.465	.985
Year of study				
Year 1 ($n = 670$)	2132.95	2040.95	2146.71	2188.57
Year 2 ($n = 517$)	1962.21	1977.67	1962.05	1953.68
Year 3 ($n = 649$)	1912.22	1972.96	1923.30	1839.35
Year 4 ($n = 1308$)	1940.40	1938.21	1957.08	1946.57
Year 5 ($n = 712$)	1981.67	1983.29	1928.99	1999.74
Year 6 ($n = 94$)	1804.86	1974.48	1798.06	1735.77
P value ^b	.003	.601	.001	< .001
Academic Performance up to the previous semester/annual exam				
Excellent ($n = 899$)	2202.70	2207.50	2153	2096.65
Very good ($n = 1420$)	1973.76	1971.03	1969.81	1981.95
Good ($n = 1056$)	1904.26	1884.23	1926.57	1951.95
Average ($n = 362$)	1895.27	1900.76	1879.30	1963.81
Below average ($n = 164$)	1572.16	1619.30	1726.27	1670.05
Poor ($n = 49$)	1335.53	1559.76	1483.29	1182.23
P value ^b	< .001	< .001	< .001	< .001
Exercise status				
No ($n = 1428$)	1859.95	1909.21	1872.45	1841.08
Yes. Irregular exercise ($n = 2170$)	2039.77	2020.60	2031.62	2032.07
Yes. Regular exercise ($n = 352$)	2048.09	1966.37	2047.56	2172.08
P value ^b	< .001	.016	< .001	< .001
Participation in extracurricular activities (arts, sports, personal skills, debate competitions, and community service)				
Maybe (participate occasionally in some activities) ($n = 1790$)	2044.69	2055.92	1983.25	2036.60
No ($n = 1550$)	1888.47	1903.42	1883.87	1914.34
Yes (regularly participate in almost all activities) ($n = 610$)	1993.60	1922.66	2185.60	1951.63
P value ^b	< .001	.016	< .001	.007
The average number of hours sleep a day				
4 to 6 h/d ($n = 1990$)	2033.27	2028.87	2036.23	2005.04
7 to 10 h/d ($n = 1864$)	1925.81	1933.13	1916.15	1949.81
More than 10 h/d ($n = 96$)	1742.88	1962	1869.12	1861.86
P value ^b	.002	.002	.003	.198

Abbreviation: ARS-30, 30-item academic resilience scale.

The significance level is 0.050.

^a Mann-Whitney U test.

^b Kruskal-Wallis test.

appeared to have a significant role in shaping how students cope with tough times throughout their studies. These findings provide valuable insights into the factors that influence academic resilience among pharmacy students and can be used to develop interventions to support students and improve their outcomes.

Our findings showed significant variations in overall academic resilience across countries, with the highest levels recorded in Sudan, Pakistan, and Nigeria. In contrast, relatively lower levels were reported in Indonesia and Türkiye. Other countries, such as Malaysia, Jordan, Saudi Arabia, and the United Arab Emirates, tended to have relatively moderate levels. This variation was anticipated in the context of differences across countries and institutions in the offered framework of student integration into the academic system and the range of support activities, contributing directly to students' level of well-being and academic resilience.^{19,20} This was further amplified by the unique demands of pharmacy programs concerning practical and experiential learning activities that exhibited significant differences across countries in the design, demands, and institutional support reported in 2 recent reports in Arab and Asia-Pacific countries.^{21,22}

Although previous large-scale studies highlighted differences in the overall academic resilience based on sex,¹ the current study found only sex, students' mental well-being and academic resilience. Given its multifaceted nature, further research is imperative-based differences in the third subscale 3 of negative affective and emotional responses but not in the overall academic resilience measures. This might suggest a more refined impact of gender on academic resilience and a guide for providing additional support. Comparable to previous studies, our study found a significant difference in academic resilience across years of study, where first-year students tended to have higher academic resilience than other years. Unsurprisingly, our findings support that academic resilience directly correlates with academic performance, whereas students with higher academic resilience levels have higher academic grades. This is consistent with a previous Malaysian study that showed a positive correlation between academic resilience and cumulative grade points.¹²

Earlier large-scale assessments of mental well-being and grit, essential traits for academic achievement, identified an impact of institution type on the overall assessment.^{20,23} This impact was also evident in our findings because students from private universities had higher academic resilience levels than their counterparts from public universities. This, again, can be attributed initially to student support activities that might be more likely to be available in a well-structured format in private schools. One of these support modalities might be linked to sports engagement because exercise seems to be associated with students' resilience, possibly through the positive link between exercise routine and mental well-being among pharmacy students.²⁰

On the other hand, our findings did not show any significant difference in academic resilience levels based on living arrangements, such as staying alone, sharing a hostel, or living with family. However, an earlier study in Thailand reported significantly higher resilience levels among those with friends and family surrounding them during their studies.²⁴

The implications of this study hold significant weight in the realm of pharmacy education. Initially, the findings indicate the necessity of cultivating and implementing interventions designed to uphold academic resilience among pharmacy students. These interventions may center around the cultivation of coping mechanisms, the provision of accessible support services, and the creation of an environment that fosters support.² Moreover, the findings propose that pharmacy programs should be customized to cater to the unique needs of individual students. For instance, programs may need to extend additional support to female students, those from certain ethnic backgrounds, penultimate and final-year students, and those encountering academic challenges.²⁵ Finally, the findings suggest that pharmacy programs should actively encourage student participation in extracurricular activities. Such activities are linked to academic achievement, character, social

development, and community involvement.²⁶ All these elements collectively contribute to an enhanced state of academic resilience.

The main strengths of the current study were its large sample and that it is a multinational survey involving representative samples of pharmacy students from 12 countries representing varied geographic regions extending from Asia, the Middle East, Africa, and Europe. Another strength was the use of a validated ARS-30.

There are some limitations of this study. There is a risk of selection bias; pharmacy students with higher academic resilience might be willing to participate in this online survey. In addition, the self-reported nature of the survey may limit the overall assessment accuracy among participants. In addition, points related to family obligations and detailed program requirements in every country might have benefited from a thorough, detailed assessment to ascertain its impact on academic resilience.

5. Conclusion

The present study provides valuable insights into the factors that could impact the academic resilience of pharmacy students. These findings could inform the development of interventions and structured support activities to boost academic resilience and enhance overall academic outcomes. Pharmacy colleges should activate academic advising services to support the students' mental well-being and academic resilience. Given its multifaceted nature, it is imperative to conduct further research to gain in-depth insights into the specific traits or characteristics associated with academic resilience.

Author Contributions

Conceptualization: M.H.E. *Data curation:* M.H.E., W.N.A.A.B.W.S. *Formal analysis:* M.H.E., W.N.A.A.B.W.S. *Funding acquisition:* M.H.E., M.E.E. *Investigation:* All authors. *Methodology:* M.H.E., W.N.A.A.B.W.S., B.O., H.K.E. *Project administration:* All authors. *Resources:* M.H.E. *Software:* M.H.E. *Supervision:* M.H.E., A.K.T. *Validation:* M.H.E., W.N.A.A.B.W.S. *Visualization:* M.H.E. *Writing – original draft:* M.H.E., W.N.A.A.B.W.S., A.K.T. *Writing – review & editing:* All authors.

Declaration of Competing Interest

None declared.

Acknowledgments

The authors would like to thank Prof Mesut Sancar for his valuable input and support for study protocol approval in Türkiye.

Funding/Support

The authors extend their appreciation to the Deputyship for Research and Innovation of the Ministry of Education in Saudi Arabia for funding this research work through project number IFP22UQU4320605DSR246.

References

- Cassidy S, Mawdsley A, Langran C, Hughes L, Willis SC. A large-scale multicenter study of academic resilience and well-being in pharmacy education. *Am J Pharm Educ.* 2023;87(2):ajpe8998. <https://doi.org/10.5688/ajpe8998>
- Mawdsley A, Willis SC. Academic resilience in UK pharmacy education – a pilot study applying love and break up letters methodology. *BMC Med Educ.* 2023;23(1):441. <https://doi.org/10.1186/s12909-023-04380-4>
- Martin AJ, Marsh HW. Academic buoyancy: towards an understanding of students' everyday academic resilience. *J Sch Psychol.* 2008;46(1):53–83. <https://doi.org/10.1016/j.jsp.2007.01.002>
- Sanderson B, Brewer M. What do we know about student resilience in health professional education? A scoping review of the literature. *Nurse Educ Today.* 2017;58:65–71. <https://doi.org/10.1016/j.nedt.2017.07.018>
- Cassidy S. The Academic Resilience Scale (ARS-30): a new multidimensional construct measure. *Front Psychol.* 2016;7:1787. <https://doi.org/10.3389/fpsyg.2016.01787>

6. Furtado M. Assessment of academic resilience as a non-cognitive variable in entry-level doctor of physical therapy students. *J Allied Health*. 2022;51(3):189–197.
7. Shahidi Delshad E, Nobahar M, Raiesdana N, Yarahmadi S, Saberian M. Academic resilience, moral perfectionism, and self-compassion among undergraduate nursing students: a cross-sectional, multi-center study. *J Prof Nurs*. 2023;46:39–44. <https://doi.org/10.1016/j.profnurs.2023.02.006>
8. Baumgartner, Baumgartner JN, Schneider TR. A randomized controlled trial of mindfulness-based stress reduction on academic resilience and performance in college students. *J Am Coll Health*. 2023;71(6):1916–1925. <https://doi.org/10.1080/07448481.2021.1950728>
9. Hwang EH, Kim KH. Relationship between optimism, emotional intelligence, and academic resilience of nursing students: the mediating effect of self-directed learning competency. *Front Public Health*. 2023;11:1182689. <https://doi.org/10.3389/fpubh.2023.1182689>
10. Ye Y, Huang X, Liu Y. Social support and academic burnout among university students: a moderated mediation model. *Psychol Res Behav Manag*. 2021;14:335–344. <https://doi.org/10.2147/PRBM.S300797>
11. Romano L, Consiglio P, Angelini G, Fiorilli C. Between academic resilience and burnout: the moderating role of satisfaction on school context relationships. *Eur J Invest Health Psychol Educ*. 2021;11(3):770–780. <https://doi.org/10.3390/ejihpe11030055>
12. Abubakar U, Azli NASM, Hashim IA, et al. Research article: the relationship between academic resilience and academic performance among pharmacy students. *Pharm Educ*. 2021;21(1):705–712.
13. Halimi SN, Rowett D, Whitfield K, Luetsch K. How early career pharmacists understand resilience – a qualitative study of experiences, challenges and strategies. *Res Social Adm Pharm*. 2023;19(3):486–494. <https://doi.org/10.1016/j.sapharm.2022.10.012>
14. Aliyev R, Akbaş U, Özbay Y. Mediating role of internal factors in predicting academic resilience. *Int J Sch Educ Psychol*. 2021;9(3):236–251. <https://doi.org/10.1080/21683603.2021.1904068>
15. Cengiz S, Peker A. Adaptation of the Academic Resilience Scale (ARS-30) Turkish version validity and reliability study. *Turk Psychol Couns Guid J*. 2022;12(65):215–228.
16. Almasreh E, Moles R, Chen TF. Evaluation of methods used for estimating content validity. *Res Social Adm Pharm*. 2019;15(2):214–221. <https://doi.org/10.1016/j.sapharm.2018.03.066>
17. Thorsen C, Yang Hansen K, Johansson S. The mechanisms of interest and perseverance in predicting achievement among academically resilient and non-resilient students: evidence from Swedish longitudinal data. *Br J Educ Psychol*. 2021;91(4):1481–1497. <https://doi.org/10.1111/bjep.12431>
18. Yang S, Wang W. The role of academic resilience, motivational intensity and their relationship in EFL learners' academic achievement. *Front Psychol*. 2021;12:823537. <https://doi.org/10.3389/fpsyg.2021.823537>
19. Prebble T, Hargraves H, Leach L, Naidoo K, Suddaby G, Zepke N. Impact of student support services and academic development programmes on student outcomes in undergraduate tertiary study: a synthesis of the research report to the Ministry of Education. <www.minedu.govt.nz>. Accessed October 7, 2023.
20. Elnaem MH, Mubarak N, Salim KTM, et al. Assessment of mental wellbeing of undergraduate pharmacy students from mm countries: the role of gender, lifestyle, health-related, and academic-related factors. *Front Public Health*. 2022;10:1011376. <https://doi.org/10.3389/fpubh.2022.1011376>
21. Al-Ghananeem AM, Malcom DR, Shammass S, Aburjai T. A call to action to transform pharmacy education and practice in the Arab world. *Am J Pharm Educ*. 2018;82(9):7014. <https://doi.org/10.5668/ajpe7014>
22. Mohamed MHN, Mak V, Sumalatha G, et al. Research article: pharmacy education during and beyond COVID-19 in six Asia-Pacific countries: changes, challenges, and experiences. *Pharm Educ*. 2020;20(2):183–195.
23. Elnaem MH, Barakat M, Mubarak N, et al. Evaluation of grit and its associated factors among undergraduate pharmacy students from 14 Asian and Middle Eastern countries amid the COVID-19 pandemic. *Saudi Pharm J*. 2023;31(3):410–416. <https://doi.org/10.1016/j.jsps.2023.01.008>
24. Auttama N, Seangpraw K, Ong-Artborirak POA, Tonchay P. Factors associated with self-esteem, resilience, mental health, and psychological self-care among university students in Northern Thailand. *J Multidiscip Healthc*. 2021;14:1213–1221. <https://doi.org/10.2147/JMDH.S308076>
25. Rhoney DH, Singleton S, Nelson NR, Anderson SM, Hubal R. Forces driving change in pharmacy education: opportunities to take academic, social, technological, economic, and political into the future. *J Am Coll Clin Pharm*. 2021;4(5):639–651. <https://doi.org/10.1002/jac5.1407>
26. Christison C. The benefits of participating in extracurricular activities. *Bu J Grad Stud Educ*. 2013;5(2):17–20.