













## NATURE NOTES

# Long-distance dispersal in the short-distance dispersing house sparrow (*Passer domesticus*)

Peter S. Ranke<sup>1,2</sup>  | Michael L. Pepke<sup>1,3</sup>  | Jørgen S. Søraker<sup>1,4</sup>  | Gabriel David<sup>1,5</sup>  |  
Yimen G. Araya-Ajoy<sup>1</sup>  | Jonathan Wright<sup>1</sup>  | Ådne M. Nafstad<sup>1</sup>  |  
Bernt Rønning<sup>1,6</sup>  | Henrik Pärn<sup>1,7</sup>  | Thor Harald Ringsby<sup>1</sup>  | Henrik Jensen<sup>1</sup>  |  
Bernt-Erik Sæther<sup>1,8</sup> 

<sup>1</sup>Centre for Biodiversity Dynamics (CBD), Department of Biology, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

<sup>2</sup>BirdLife Norway, Trondheim, Norway

<sup>3</sup>Center for Evolutionary Hologenomics, Globe Institute, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

<sup>4</sup>Edward Grey Institute, Department of Biology, University of Oxford, Oxford, UK

<sup>5</sup>Animal Ecology, Department of Ecology and Genetics, Uppsala University, Uppsala, Sweden

<sup>6</sup>Department of Teacher Education, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

<sup>7</sup>Department of Aquatic Resources (SLU Aqua), Swedish University of Agricultural Sciences, Lysekil, Sweden

<sup>8</sup>The Gjørevoll Centre, Department of Biology, Norwegian University of Science and Technology (NTNU), Trondheim, Norway

## Correspondence

Peter S. Ranke, Centre for Biodiversity Dynamics (CBD), Department of Biology, Norwegian University of Science and Technology (NTNU), Trondheim, Norway. Email: [peter.s.ranke@ntnu.no](mailto:peter.s.ranke@ntnu.no)

## Funding information

Norges Forskningsråd, Grant/Award Number: 223257, 274930 and 302619; Norwegian Directorate for Nature Management; European Commission; European Research Council, Grant/Award Number: 101095997 and 268562

## Abstract

The house sparrow (*Passer domesticus*) is a small passerine known to be highly sedentary. Throughout a 30-year capture–mark–recapture study, we have obtained occasional reports of recoveries far outside our main metapopulation study system, documenting unusually long dispersal distances. Our records constitute the highest occurrence of long-distance dispersal events recorded for this species in Scandinavia. Such long-distance dispersals radically change the predicted distribution of dispersal distances and connectedness for our study metapopulation. Moreover, it reveals a much greater potential for colonization than formerly recorded for the house sparrow, which is an invasive species across four continents. These rare and occasional long-distance dispersal events are challenging to document but may have important implications for the genetic composition of small and isolated populations and for our understanding of dispersal ecology and evolution.

## KEYWORDS

capture–mark–recapture, dispersal distance, dispersal distribution, dispersal scale, genetic composition, inbreeding

## TAXONOMY CLASSIFICATION

Biodiversity ecology, Biogeography, Conservation ecology, Conservation genetics, Demography, Evolutionary ecology, Population ecology, Population genetics, Spatial ecology

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd.

## 1 | INTRODUCTION

Knowledge of the extent of long-distance dispersal is important for predicting scale-dependent dispersal rates within study systems, which enables predictions of population viability for small and isolated populations (Sutherland et al., 2000; Trakhtenbrot et al., 2005). It is also necessary for a better understanding of both the rate of spread of expanding and invasive populations (Neubert & Caswell, 2000) and local adaptation processes (Garant et al., 2007). The house sparrow (*Passer domesticus*) is known to be highly sedentary (Anderson, 2006; Summers-Smith, 1988), and bird ringing records show that most house sparrows generally disperse over very short distances from their natal site (in Great Britain on average below 2 km, Paradis et al., 1998). In Finland, 90% of natal dispersal distances were less than 16 km (Kekkonen et al., 2011), in Denmark 98% were within 10 km (Bønløkke et al., 2006), and on the Faroe Islands, 99% were within 1 km (Hammer et al., 2014). However, in Denmark two records exceeded 100 km, one individual from Helgoland, Germany, to the Danish west coast (108 km), and one individual from Skåne, Sweden, to Bornholm (133 km, Bønløkke et al., 2006). In Sweden, seven records exceeded 50 km (three of which exceeded 100 km, Fransson & Hall-Karlsson, 2008).

In Norway, the house sparrow is an abundant bird species in human-populated areas and in agricultural landscapes across the country. It is even found in the northernmost regions (e.g., Kirkenes, Finnmark), where populations have been known to exist since the early 1900s (Collett, 1921). In our main house sparrow capture-mark-recapture study system on the Helgeland coast in northern Norway (Araya-Ajoy et al., 2021; Ranke et al., 2021), most individuals remain sedentary (ca. 83%), and the remaining ca. 17% disperse over relatively short distances (most often 10–15 km; Pärn et al., 2009; Ranke et al., 2021; Tufto et al., 2005). However, recent advances in genetic assignment methods have revealed that in some years and on some islands as many as around 22% of recruits may have dispersed during their first year of life in our study metapopulation (Saatoglu et al., 2021). Our study metapopulation covers an area of around 1600 km<sup>2</sup>, spanning 18 small islands in a coastal archipelago in northern Norway. Although this study system is large compared to the usual scale of movement distances in house sparrows (Kekkonen et al., 2011; Tufto et al., 2005), it is likely that we fail to document occasional dispersal distances exceeding 50 km.

During the years 1992–2022, we have in our main study metapopulation ringed a total of 25,425 individuals (16,349 nestlings and an additional 6932 fledged juveniles and 2144 recruits with unknown age). Recaptures and observations of these individuals throughout their lifetime enable us to record inter-island movements (Ranke et al., 2021). However, additional sampling along the Norwegian coast (Jensen et al., 2013) and research we have carried out in house sparrow populations between ca. 15 and 300 km south of the main study metapopulation during the years 2001–2022 (in total ca. 11,750 house sparrows ringed; Kvalnes et al., 2017; Nafstad et al., 2023; Ranke et al., 2017, 2020; Stubberud et al., 2017), combined with regular ringing recoveries by local inhabitants


(Bakken et al., 2006), allow us to explore patterns in a high number of individual house sparrow movements across long periods of time and over large geographical distances, which exceed those of our main study metapopulation (see Figure 1). Here, we aim to describe and document such long-distance dispersal events in reference to the dispersal dynamics within a metapopulation of house sparrows. The extent of long-distance dispersal events may have implications for dispersal distance distributions or kernels (Tufto et al., 2005), the estimation of fitness of local and dispersing individuals within spatially constrained study systems (Doligez & Pärt, 2008), the rate of immigration of novel genotypes affecting levels of inbreeding (Green & Hatchwell, 2018) and thus long-term population viability (Sutherland et al., 2000), the transmission of infectious diseases (Ferraguti et al., 2019), and for unraveling drivers of biogeographical colonization dynamics (Pedersen et al., 2018).

## 2 | METHODS

The house sparrow has been a well-suited study species for several research fields in biology due to its sedentary and synanthropic behavior and because it naturally breeds close to human settlements (Anderson, 2006; Summers-Smith, 1988). House sparrows have been studied in the Helgeland archipelago (66.30°–66.80° N, 12.00°–13.10° E; see black square in Figure 1) off the coast of northern Norway since 1992 (Baalsrud et al., 2014; Sæther et al., 1999). This involves research on small, fragmented populations within a metapopulation setting. The study system thus consists of small human-inhabited islands that have historically been sustained through fishing with some subsistence farming. Today, some islands are more densely populated with village centers (situated farther off the coast; hereafter referred to as “non-farm islands”), while others have remained scarcely populated with dairy farms (closer to the mainland; hereafter referred to as “farm islands”). Within our study metapopulation, the outer non-farm islands show proportionally more and longer distances of natal dispersal than the inner farm islands (Ranke et al., 2021).

Fieldwork is carried out at irregular intervals throughout the year, but almost continuously from May to mid-August and from mid-September to the end of October. All newly recorded individuals are given a unique combination of rings, consisting of a metal ring and three additional colored plastic rings (Figure 2). Captures using mist nets, and re-sightings of ringed individuals, enable us to closely monitor each population throughout the season. During the breeding season, nests are searched for thoroughly on each study island, and natural nests as well as nest boxes are checked at least every 10 days. Nestlings are ringed at 11 days (5–13 days) of age. A small blood sample is collected from adults and nestlings using venipuncture from the wing, allowing us to perform genetic analyses and confirm individual sex and identity (described in Jensen et al., 2013; Saatoglu et al., 2021).

Dispersal in house sparrows occurs mainly in the autumn by juveniles (i.e., natal dispersal, Altwegg et al., 2000). In the period



**FIGURE 1** Long-distance dispersal events of house sparrows (*Passer domesticus*) from and to the Helgeland archipelago in northern Norway during the years 1992–2022. Black dots represent ringing locations, and red dots represent the recovery sites. The black square in the main map indicates the main island archipelago metapopulation study site. Numbered identities of recoveries refer to [Table 1](#).

1992–2014, we have recorded average dispersal rates across years and islands of around 17% from ecological data (Ranke et al., 2021) and approximately 22% based on genetic analyses (Saatoglu et al., 2021). Here, we document individual dispersal events exceeding 50km (i.e., outside our main metapopulation study system; [Table 1](#), [Figure 1](#)), which we would have been unable to record without a wider Ringing Scheme system of ringing recoveries.

Dispersal distances were recorded for each individual as the shortest direct route from its natal site, either determined as the exact location of the nest where it was hatched, or measured as the geographic center of the population of house sparrows within the natal island ( $\pm 500\text{m}$ ) to the first breeding site. For our own records, the exact location of the recovery was obtained from our database, but in other cases we received records from the Ringing Scheme (Bakken et al., 2006) with negligible lower resolution (i.e., within  $\pm 1\text{km}$ ). Thus, minimum distance traveled could be calculated with acceptable certainty for all dispersal events.

### 3 | RESULTS AND DISCUSSION

The present study demonstrates that house sparrows may disperse over much longer distances than previously anticipated for this highly sedentary bird species, and at least up to 159 km in northern Norway. Our 12 long-distance (>50km) dispersal records represent a large proportion of long-distance dispersal events recorded for house sparrows in Scandinavia (seven in Sweden, four in Denmark (excluding one Danish record of exactly 50km), and five additional records elsewhere in Norway, see [Table A1](#)). Thus, out of a total 28 long-distance dispersal events in Scandinavia in the past 100 years, nearly half (43%;  $n=12$ ) of these are recorded in the Helgeland archipelago. The number of ringed house sparrows is similar across the three countries, and our metapopulation under study constitutes about one-third of the ringed house sparrows in Norway.

The proportion of undetected successful dispersal events cannot currently be distinguished from mortality in our study area.



**FIGURE 2** A house sparrow (*Passer domesticus*) male with a unique combination of color rings and a numbered metal ring at one of our study islands, Selv er, in the Helgeland archipelago in northern Norway. The above individual is shown carrying nest-building materials, en route to the nest site (photo by P. S. Ranke).

Moreover, the detectability outside our study area is lower than inside, and thus, many long-distance dispersal events might remain undetected. Previous examination of dispersal within our study system revealed that 17%–22% of the recruits had dispersed (Ranke et al., 2021; Saatoglu et al., 2021), but recruitment probabilities vary between 15% and 20% (Ringsby et al., 2002; S ether et al., 1999); thus, most juveniles are probably lost due to mortality by predation, parasite infection (Holand et al., 2014), starvation or at sea during dispersal. Out of 12 dispersal records exceeding 50 km, 10 were emigrating from and one immigrating to the outer non-farm islands in our study system (Table 1; Figure 1), which is consistent with the generally higher levels and longer dispersal distances on average for this set of islands (Ranke et al., 2021; Saatoglu et al., 2021).

Average dispersal distances recorded in this metapopulation exceed normal dispersal distances for house sparrows in other countries (Kekkonen et al., 2011; Paradis et al., 1998) and other sites in Norway (Bakken et al., 2006). This may partly be due to the geographical structure of our study system, where individuals are forced to move across larger distances over open sea compared with more contiguous non-coastal landscapes (Kekkonen et al., 2011; Ranke et al., 2021). This is analogous to Eurasian nuthatch (*Sitta europaea*) populations in patchy habitats in Belgium showing longer dispersal distances than populations in contiguous forest habitats (Matthysen et al., 1995). However, the number of long-distance house sparrow

**TABLE 1** Overview of long-distance dispersal events (>50 km;  $n = 12$ ) in house sparrows (*Passer domesticus*) from or to the Helgeland archipelago, northern Norway, during the years 1992–2022.

No.	ID	Sex	Age (cy)	Date	Distance	Journey	Circumstances
1	8M31147	M	1–2	2009-07-01 2010-02-11	159	Tr�ena Leka	Nestling Captured
2	8N13455	M	1–2	2014-06-29 2015-02-04	159	Tr�ena Leka	Nestling Captured
3	8P14515	F	1–2	2019-06-15 2020-06-23	124	Selv�er Bod�	Nestling Found dead
4	8P14582	X	1–2	2019-06-29 2020-05-18	120	Lovund Giv�er	Juvenile Captured
5	8N13408	M	4–5	2014-06-22 2018-06-09	113	Tr�ena R�st	Juvenile Photographed
6	8M31109	M	1–2	2009-06-25 2010-02-19	94	Tr�ena Vega	Nestling Captured
7	8L57742	M	1–2	2007-05-23 2008-02-24	92	Tr�ena Vega	Nestling Captured
8	8L89915	M	1–4	2009-05-21 2012-03-07	90	Tr�ena Vega	Nestling Captured
9	8866658	F	1–2	2002-06-19 2003-05-23	77	Vega Lovund	Nestling Captured
10	8N13261	F	1–2	2014-05-29 2015-02-23	75	Selv�er-Tr�ena Austb�	Nestling Captured
11	8N42048	F	1–2	2017-06-30 2018-04-16	54	Selv�er-Tr�ena Nesna	Nestling Captured
12	8M72794	F	1–2	2010-07-16 2011-10-10	52	Indre Kvar�y Sandnessj�en	Nestling Found dead

Note: Disperser identity (No.) corresponding to the map in Figure 1, ID (ring number, from the Ringing Scheme at Stavanger Museum), Sex (male 'M', female 'F' or unknown 'X'), Age range in calendar years (cy; note that 1 is equal to the year of birth) at the time when dispersal occurred, Date of ringing, Distance in km, Journey from and to (separated with line shift), and Circumstances at ringing and recovery (all recovered being either captured [ $n = 9$ ], photographed [ $n = 1$ ] or found dead [ $n = 2$ ]).

dispersal events recorded here is disproportionately high, as there have been only five other house sparrow dispersal records exceeding 50 km in Norway during 1914–2022 (Bakken et al., 2006; Norwegian Ringing Scheme *pers. comm.*; see Table A1 in Appendix). The extent and duration of our fieldwork campaign, resulting in high recapture probabilities across a large geographic area (Jensen et al., 2013; Kvalnes et al., 2017; Nafstad et al., 2023; Ranke et al., 2017, 2020), have additionally contributed to documenting rare dispersal events, and our intensive field campaign contributes to about half of our documented long-distance dispersal events. Moreover, color-ringed individuals would be more conspicuous and easier to detect also for local inhabitants, which might additionally increase detection probabilities of individuals from our study populations.

Furthermore, we suggest that the extreme northern location and marginal environmental conditions of our study site – at the edge of the global range distribution for house sparrows – and the low density of human settlements and farms, mean that there are large distances between adjacent viable habitats. For example, house sparrows in Kenya display greater genetic variation toward the edges of their range expansion, supporting the suggestion of a higher rate of long-distance dispersal in such areas (Schrey et al., 2014). However, similar to Schrey et al. (2014), we cannot rule out potential effects of human-mediated dispersal (in our area by ferries, fishing boats, etc.) potentially facilitating some of these long-distance dispersal events. We have observed sparrows on the ferries, but this is extremely rare, and so it is unclear whether ship-assisted dispersal has any general impact on dispersal rates within our study area. Moreover, similar long-distance dispersal events in North American house sparrows may have taken place during recent range expansion (Gibson, 2012; Schrey et al., 2011), and in some places this has included geographical landscapes similar to those of our study populations situated at high northern latitudes. In northern Norway, harsh weather conditions, including autumn storms during the period of natal dispersal, may affect wind-assisted geographic displacement over water and across large distances (e.g., Darlington, 1938).

These types of long-distance dispersal events highlight the potential for colonization of new habitats by house sparrows, which may be an important and hitherto overlooked part of this species success as a global colonizer and invasive species (Anderson, 2006; Hanson et al., 2020). Future studies may utilize these observations to investigate if individual phenotypic differences in, for example, morphology (Skjelseth et al., 2007), physiology (Nafstad et al., 2023; Pepke et al., 2022), or life-history characteristics (Pärn et al., 2009; Saatoglu et al., 2024) underlie longer-than-expected dispersal distances (Tufto et al., 2005) in a species with very low flight efficiency (Claramunt, 2021). If dispersing individuals can successfully recruit into a breeding population, then they can offer a valuable genetic contribution to small, isolated, and often inbred populations (Dickel et al., 2021), such as these house sparrow populations habituating to island life on the northern Norwegian coast (Niskanen et al., 2020; Ranke et al., 2020). Thus, although long-distance dispersal happens infrequently in house sparrows, long-distance dispersal can have major implications for gene flow across large geographic

areas (Garant et al., 2007), which in turn may influence the rate of inbreeding (Keller et al., 2001), population (re)colonization and spatio-temporal population dynamics (Baalsrud et al., 2014; Billing et al., 2012; Ranke et al., 2021) and adaptive and non-adaptive population differentiation (Aase et al., 2022; Araya-Ajoy et al., 2019; Holand et al., 2011; Jensen et al., 2013). Furthermore, long-distance dispersal can affect estimates of fitness of dispersers compared to non-dispersers, because of the bias in the number of recorded offspring, especially if dispersal behavior has a heritable component (Doligez & Pärt, 2008; Saatoglu et al., 2024). This study therefore reveals the biological and methodological implications of a long tail of upper values in the distribution of dispersal distances, even in an otherwise sedentary species, and therefore the need for large-scale longitudinal field studies quantifying these rare but important dispersal events.

#### AUTHOR CONTRIBUTIONS

**Peter S. Ranke:** Conceptualization (lead); data curation (lead); visualization (lead); writing – original draft (lead); writing – review and editing (equal). **Michael L. Pepke:** Writing – review and editing (equal). **Jørgen S. Søraker:** Writing – review and editing (equal). **Gabriel David:** Writing – review and editing (equal). **Yimen G. Araya-Ajoy:** Writing – review and editing (equal). **Jonathan Wright:** Writing – review and editing (equal). **Ådne M. Nafstad:** Writing – review and editing (equal). **Bernt Rønning:** Writing – review and editing (equal). **Henrik Pärn:** Writing – review and editing (equal). **Thor Harald Ringsby:** Writing – review and editing (equal). **Henrik Jensen:** Writing – review and editing (equal). **Bernt-Erik Sæther:** Funding acquisition (lead); writing – review and editing (equal).

#### ACKNOWLEDGEMENTS

We would like to thank the inhabitants of our study area for their great hospitality over more than 30 years of study. Moreover, we thank the large number of scientific staff, students, and volunteers that have contributed to the fieldwork. We thank the Ringing Schemes in Scandinavia for providing updated information on their dispersal records for the house sparrow: Håvard Husebø at Stavanger Museum, Norway; Thord Fransson at Naturhistoriska riksmuseet, Sweden; and Amanda Johannisson at Statens Naturhistoriske Museum, Københavns Universitet, Denmark. This research was carried out with permissions from the Norwegian Environment Agency and the Ringing Centre at Stavanger Museum, Norway. This study was partly funded by the EU Commission METABIRD, Norwegian Directorate for Nature Management, the Research Council of Norway (Centres of Excellence scheme to CBD project no. 223257, and project nos. 274930 and 302619), and Advanced Grants from the European Research Council (project nos. 268562 and 101095997).

#### CONFLICT OF INTEREST STATEMENT

None declared.

#### DATA AVAILABILITY STATEMENT

All data are provided in Table 1, Table A1, and Figure 1.

## ORCID

- Peter S. Ranke  <https://orcid.org/0000-0003-3757-8626>
- Michael L. Pepke  <https://orcid.org/0000-0002-6280-1829>
- Jørgen S. Søraker  <https://orcid.org/0000-0001-8146-4576>
- Gabriel David  <https://orcid.org/0009-0007-2935-3335>
- Yimen G. Araya-Ajoy  <https://orcid.org/0000-0001-7844-0477>
- Jonathan Wright  <https://orcid.org/0000-0002-5848-4736>
- Ådne M. Nafstad  <https://orcid.org/0000-0002-2577-5414>
- Bernt Rønning  <https://orcid.org/0000-0002-4237-5893>
- Henrik Pärn  <https://orcid.org/0000-0001-7745-9950>
- Thor Harald Ringsby  <https://orcid.org/0000-0001-9089-7592>
- Henrik Jensen  <https://orcid.org/0000-0001-7804-1564>
- Bernt-Erik Sæther  <https://orcid.org/0000-0002-0049-9767>

## REFERENCES

- Aase, K., Jensen, H., & Muff, S. (2022). Genomic estimation of quantitative genetic parameters in wild admixed populations. *Methods in Ecology and Evolution*, 13, 1014–1026. <https://doi.org/10.1111/2041-210X.13810>
- Altwegg, R., Ringsby, T. H., & Sæther, B.-E. (2000). Phenotypic correlates and consequences of dispersal in a metapopulation of house sparrows *Passer domesticus*. *Journal of Animal Ecology*, 69, 762–770. <https://doi.org/10.1046/j.1365-2656.2000.00431.x>
- Anderson, T. R. (2006). *Biology of the ubiquitous house sparrow: From genes to populations*. Oxford University Press.
- Araya-Ajoy, Y., Niskanen, A. K., Froy, H., Ranke, P. S., Kvalnes, T., Rønning, B., Pepke, M. L., Jensen, H., Ringsby, T. H., Sæther, B.-E., & Wright, J. (2021). Variation in generation time reveals density regulation as an important driver of pace of life in a bird metapopulation. *Ecology Letters*, 24, 2077–2087. <https://doi.org/10.1111/ele.13835>
- Araya-Ajoy, Y. G., Ranke, P. S., Kvalnes, T., Rønning, B., Holand, H., Myhre, A. M., Pärn, H., Jensen, H., Ringsby, T. H., Sæther, B.-E., & Wright, J. (2019). Characterizing morphological (co)variation using structural equation models: Body size, allometric relationships and evolvability in a house sparrow metapopulation. *Evolution*, 73, 452–466. <https://doi.org/10.1111/evo.13668>
- Baalsrud, H. T., Sæther, B.-E., Hagen, I. J., Myhre, A. M., Ringsby, T. H., Pärn, H., & Jensen, H. (2014). Effects of population characteristics and structure on estimates of effective population size in a house sparrow metapopulation. *Molecular Ecology*, 23, 2653–2668. <https://doi.org/10.1111/mec.12770>
- Bakken, V., Runde, O., & Tjørve, E. (2006). *Norsk Ringmerkjingsatlas* (Vol. 2). Stavanger Museum.
- Billing, A. M., Lee, A. M., Skjelseth, S., Borg, A. A., Hale, M. C., Slate, J., Pärn, H., Ringsby, T. H., Sæther, B.-E., & Jensen, H. (2012). Evidence of inbreeding depression but not inbreeding avoidance in a natural house sparrow population. *Molecular Ecology*, 21(6), 1487–1499. <https://doi.org/10.1111/j.1365-294X.2012.05490.x>
- Bønløkke, J., Madsen, J. J., Thorup, K., Pedersen, K. T., Bjerrum, M., & Rahbek, C. (2006). *The Danish bird migration atlas*. Københavns Universitet, Forlaget Rhodos A/S & Zoologisk Museum, Trykt af Narayana Press.
- Claramunt, S. (2021). Flight efficiency explains differences in natal dispersal distances in birds. *Ecology*, 102(9), e03442. <https://doi.org/10.1002/ecy.3442>
- Collett, R. (1921). *Norges fugle* (by Ørjan Olsen) (Vol. I). H. Aschehoug & Co.
- Darlington, P. J. (1938). The origin of the fauna of the greater Antilles, with discussion of dispersal of animals over water and through the air. *The Quarterly Review of Biology*, 13, 274–300. <https://doi.org/10.1086/394561>
- Dickel, L., Arcese, P., Nietlisbach, P., Keller, L. F., Jensen, H., & Reid, J. M. (2021). Are immigrants outbred and unrelated? Testing standard assumptions in a wild metapopulation. *Molecular Ecology*, 30, 5674–5686. <https://doi.org/10.1111/mec.16173>
- Doligez, B., & Pärt, T. (2008). Estimating fitness consequences of dispersal: A road to 'know-where'? Non-random dispersal and the underestimation of dispersers' fitness. *Journal of Animal Ecology*, 77, 1199–1211. <https://doi.org/10.1111/j.1365-2656.2008.01446.x>
- Ferraguti, M., Martínez-de la Puente, J., García-Longoria Soriguer, R., Figuerola, J., & Marzal, A. (2019). From Africa to Europe: Evidence of transmission of a tropical plasmodium lineage in Spanish populations of house sparrows. *Parasites and Vectors*, 12, 548. <https://doi.org/10.1186/s13071-019-3804-1>
- Fransson, T., & Hall-Karlsson, S. (2008). *Svensk ringmärkningsatlas, volym 3. Tättingar*. Svensk Ornitologisk Förings Förlag.
- Garant, D., Forde, S. E., & Hendry, A. P. (2007). The multifarious effects of dispersal and gene flow on contemporary adaptation. *Functional Ecology*, 21, 434–443. <https://doi.org/10.1111/j.1365-2435.2006.01228.x>
- Gibson, D. D. (2012). On two fronts: Occurrence of the house sparrow in Alaska. *Western Birds*, 43, 248–254.
- Green, J. P., & Hatchwell, B. J. (2018). Inclusive fitness consequences of dispersal decisions in a cooperatively breeding bird, the long-tailed tit (*Aegithalos caudatus*). *Proceedings of the National Academy of Sciences of the United States of America*, 115, 12011–12016. <https://doi.org/10.1073/pnas.1815873115>
- Hammer, S., Madsen, J., Jensen, J.-K., Pedersen, K., Bloch, D., & Thorup, K. (2014). *Færøsk Trækfugleatlas - The Faroese bird migration atlas*. Fróðskapur - Faroe University Press.
- Hanson, H. E., Koussayer, B., Kilvitis, H. J., Schrey, A. W., Maddox, J. D., & Martin, L. B. (2020). Epigenetic potential in native and introduced populations of house sparrows (*Passer domesticus*). *Integrative and Comparative Biology*, 60, 1458–1468. <https://doi.org/10.1093/icb/icaa060>
- Holand, A. M., Jensen, H., Tufto, J., & Moe, R. (2011). Does selection or genetic drift explain geographic differentiation of morphological characters in house sparrows *Passer domesticus*? *Genetics Research*, 93, 367–379. <https://doi.org/10.1017/S0016672311000267>
- Holand, H., Jensen, H., Tufto, J., Soliman, M. M., Pärn, H., Sæther, B.-E., & Ringsby, T. H. (2014). Lower survival probability of house sparrows severely infected by the gapeworm parasite. *Journal of Avian Biology*, 45(4), 365–373. <https://doi.org/10.1111/jav.00354>
- Jensen, H., Moe, R., Hagen, I. J., Holand, A. M., Kekkonen, J., Tufto, J., & Sæther, B.-E. (2013). Genetic variation and structure of house sparrow populations: Is there an Island effect? *Molecular Ecology*, 22, 1792–1805. <https://doi.org/10.1111/mec.12226>
- Kekkonen, J., Seppä, P., Hanski, I. K., Jensen, H., Väisänen, R. A., & Brommer, J. E. (2011). Low genetic differentiation in a sedentary bird: House sparrow population genetics in a contiguous landscape. *Heredity*, 106, 183–190. <https://doi.org/10.1038/hdy.2010.32>
- Keller, L. F., Jeffery, K. J., Arcese, P., Beaumont, M. A., Hochachka, W. M., Smith, J. N., & Bruford, M. W. (2001). Immigration and the ephemerality of a natural population bottleneck: Evidence from molecular markers. *Proceedings of the Royal Society B: Biological Sciences*, 268(1474), 1387–1394. <https://doi.org/10.1098/rspb.2001.1607>
- Kvalnes, T., Ringsby, T. H., Jensen, H., Hagen, I. J., Rønning, B., Pärn, H., Holand, H., Engen, S., & Sæther, B.-E. (2017). Reversal of response to artificial selection on body size in a wild passerine bird. *Evolution*, 71, 2062–2079. <https://doi.org/10.1111/evo.13277>
- Matthysen, E., Adriaansen, F., & Dhondt, A. A. (1995). Dispersal distances of nuthatches, *Sitta europaea*, in a highly fragmented forest habitat. *Oikos*, 72, 375–381. <https://doi.org/10.2307/3546123>
- Nafstad, Å. M., Rønning, B., Aase, K., Ringsby, T. H., Hagen, I. J., Ranke, P. S., Kvalnes, T., Stawski, C., Räsänen, K., Sæther, B.-E., Muff, S., & Jensen, H. (2023). Spatial variation in the evolutionary potential

- and constraints of basal metabolic rate and body mass in a wild bird. *Journal of Evolutionary Biology*, 36, 650–662. <https://doi.org/10.1111/jeb.14164>
- Neubert, M. G., & Caswell, H. (2000). Demography and dispersal: Calculation and sensitivity analysis of invasion speed for structured populations. *Ecology*, 81, 1613–1628. [https://doi.org/10.1890/0012-9658\(2000\)081\[1613:DADCAS\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2000)081[1613:DADCAS]2.0.CO;2)
- Niskanen, A. K., Billing, A. M., Holand, H., Hagen, I. J., Huisman, J., Araya-Ajoy, Y. G., Husby, A., Rønning, B., Myhre, A. M., Ranke, P. S., Kvalnes, T., Pärn, H., Ringsby, T. H., Lien, S., Sæther, B.-E., Muff, S., & Jensen, H. (2020). Consistent scaling of inbreeding depression in space and time. *Proceedings of the National Academy of Sciences of the United States of America*, 117(25), 14584–14592. <https://doi.org/10.1073/pnas.1909599117>
- Paradis, E., Baillie, S. R., Sutherland, W. J., & Gregory, R. D. (1998). Patterns of natal and breeding dispersal in birds. *Journal of Animal Ecology*, 67, 518–536. <https://doi.org/10.1046/j.1365-2656.1998.00215.x>
- Pärn, H., Jensen, H., Ringsby, T. H., & Sæther, B.-E. (2009). Sex-specific fitness correlates of dispersal in a house sparrow metapopulation. *Journal of Animal Ecology*, 78, 1216–1225. <https://doi.org/10.1111/j.1365-2656.2009.01597.x>
- Pedersen, M. P., Irestedt, M., Joseph, L., Rahbek, C., & Jönsson, K. A. (2018). Phylogeography of a 'great speciator' (Aves: *Edolisoma tenuirostre*) reveals complex dispersal and diversification dynamics across the Indo-Pacific. *Journal of Biogeography*, 45, 826–837. <https://doi.org/10.1111/jbi.13182>
- Pepke, M. L., Kvalnes, T., Ranke, P. S., Araya-Ajoy, Y. G., Wright, J., Sæther, B.-E., Jensen, H., & Ringsby, T. H. (2022). Causes and consequences of variation in early-life telomere length in a bird metapopulation. *Ecology and Evolution*, 12, e9144. <https://doi.org/10.1002/ece3.9144>
- Ranke, P. S., Araya-Ajoy, Y., Ringsby, T. H., Pärn, H., Rønning, B., Jensen, H., Wright, J., & Sæther, B.-E. (2021). Spatial structure and dispersal dynamics in a house sparrow metapopulation. *Journal of Animal Ecology*, 90, 2767–2781. <https://doi.org/10.1111/1365-2656.13580>
- Ranke, P. S., Skjelseth, S., Hagen, I. J., Billing, A. M., Pedersen, Å. A. B., Pärn, H., Ringsby, T. H., Sæther, B.-E., & Jensen, H. (2020). Multi-generational genetic consequences of reinforcement in a bird metapopulation. *Conservation Genetics*, 21, 603–612. <https://doi.org/10.1007/s10592-020-01273-7>
- Ranke, P. S., Skjelseth, S., Pärn, H., Herfindal, I., Pedersen, Å. A. B., Stokke, B. G., Kvalnes, T., Ringsby, T. H., Sæther, B.-E., & Jensen, H. (2017). Demographic influences of translocated individuals on a resident population of house sparrows. *Oikos*, 126, 1410–1418. <https://doi.org/10.1111/oik.04065>
- Ringsby, T. H., Sæther, B.-E., Tufto, J., Jensen, H., & Solberg, E. J. (2002). Asynchronous spatiotemporal demography of a house sparrow metapopulation in a correlated environment. *Ecology*, 83, 561–569. [https://doi.org/10.1890/0012-9658\(2002\)083\[0561:ASDOAH\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2002)083[0561:ASDOAH]2.0.CO;2)
- Saatoglu, D., Lundregan, S., Fetterplace, E., Goedert, D., Husby, A., Niskanen, A. K., Muff, S., & Jensen, H. (2024). The genetic basis of dispersal in a vertebrate metapopulation. *Molecular Ecology*, 33, e17295. <https://doi.org/10.1111/mec.17295>
- Saatoglu, D., Niskanen, A. K., Kuismin, M., Ranke, P. S., Hagen, I. J., Araya-Ajoy, Y. G., Kvalnes, T., Pärn, H., Rønning, B., Ringsby, T. H., Sæther, B.-E., Husby, A., Sillanpää, M. J., & Jensen, H. (2021). Dispersal in a house sparrow metapopulation: An integrative case study of genetic assignment calibrated with ecological data and pedigree information. *Molecular Ecology*, 30(19), 4740–4756. <https://doi.org/10.1111/mec.16083>
- Sæther, B.-E., Ringsby, T. H., Bakke, O., & Solberg, E. J. (1999). Spatial and temporal variation in demography of a house sparrow metapopulation. *Journal of Animal Ecology*, 68, 628–637. <https://doi.org/10.1046/j.1365-2656.1999.00314.x>
- Schrey, A. W., Grispo, M., Awad, M., Cook, M. B., McCoy, E. D., Mushinsky, H. R., Albayrak, T., Bensch, S., Burke, T., Butler, L. K., Dor, R., Fokidis, H. B., Jensen, H., Imboma, T., Kessler-Rios, M. M., Marzal, A., Stewart, I. R. K., Westerdahl, H., Westneat, D. F., ... Martin, L. B. (2011). Broad-scale latitudinal patterns of genetic diversity among native European and introduced house sparrow (*Passer domesticus*) populations. *Molecular Ecology*, 20, 1133–1143. <https://doi.org/10.1111/j.1365-294X.2011.05001.x>
- Schrey, A. W., Liebl, A. L., Richards, C. L., & Martin, L. B. (2014). Range expansion of house sparrows (*Passer domesticus*) in Kenya: Evidence of genetic admixture and human-mediated dispersal. *Journal of Heredity*, 105, 60–69. <https://doi.org/10.1093/jhered/est085>
- Skjelseth, S., Ringsby, T. H., Tufto, J., Jensen, H., & Sæther, B.-E. (2007). Dispersal of introduced house sparrows *Passer domesticus*: An experiment. *Proceedings of the Royal Society B: Biological Sciences*, 274, 1763–1771. <https://doi.org/10.1098/rspb.2007.0338>
- Stubberud, M. W., Myhre, A. M., Holand, H., Kvalnes, T., Ringsby, T. H., Sæther, B.-E., & Jensen, H. (2017). Sensitivity analysis of effective population size to demographic parameters in house sparrows. *Molecular Ecology*, 26, 2449–2465. <https://doi.org/10.1111/mec.14057>
- Summers-Smith, J. D. (1988). *The sparrows: A study of the genus passer*. T. and A. D. Poyser.
- Sutherland, G. D., Harestad, A. S., Price, K., & Lertzman, K. P. (2000). Scaling of natal dispersal distances in terrestrial birds and mammals. *Conservation Ecology*, 4(1), 16. <http://www.consecol.org/vol4/iss1/art16/>
- Trakhtenbrot, A., Nathan, R., Perry, G., & Richardson, D. M. (2005). The importance of long-distance dispersal in biodiversity conservation. *Diversity and Distributions*, 11, 173–181. <https://doi.org/10.1111/j.1366-9516.2005.00156.x>
- Tufto, J., Ringsby, T. H., Dhondt, A. A., Adriaensen, F., & Matthysen, E. (2005). A parametric model for estimation of dispersal patterns applied to five passerine spatially structured populations. *The American Naturalist*, 165, E13–E26. <https://doi.org/10.1086/426698>

**How to cite this article:** Ranke, P. S., Pepke, M. L., Søraker, J. S., David, G., Araya-Ajoy, Y. G., Wright, J., Nafstad, Å. M., Rønning, B., Pärn, H., Ringsby, T. H., Jensen, H., & Sæther, B.-E. (2024). Long-distance dispersal in the short-distance dispersing house sparrow (*Passer domesticus*). *Ecology and Evolution*, 14, e11356. <https://doi.org/10.1002/ece3.11356>

## APPENDIX

TABLE A1 Overview of long-distance dispersal events (>50km;  $n = 5$ ) for house sparrows (*Passer domesticus*) in Norway outside the Helgeland archipelago during years 1914–2022.

No.	ID	Sex	Age (cy)	Date	Distance	Journey	Circumstances
1	8N05951	M	1–2	2012-05-26 2013-01-06	102	Lauvøya Støren	Nestling Photographed
2	849776	X	1–2	1955-07-08 1956-05-21	87	Nordfjordeid Tomrefjord	Nestling Found dead
3	72412	X	1–5	1954-08-03 1958-05-09	60	Nordfjordeid Florø	Nestling Found dead
4	032459	F	1–2	1948-06-03 1949-06-18	56	Steinkjer Overhalla	Nestling Killed
5	DB84457	X	1–2	1990-07-26 1991-05-13	55	Lista Kleven, Mandal	Juvenile Killed/window

Note: Disperser identity (No.), ID (ring number, from a Norwegian Ringing Scheme [Stavanger Museum, Zoological Museum or Stat. Vilt. Ås]), Sex (male 'M', female 'F' or unknown 'X'), Age range in calendar years (cy; note that 1 is equal to the year of birth) at the time when dispersal occurred, Date of ringing, Distance in km, Journey from and to (separated with line shift), and Circumstances at ringing and recovery (all recovered being either found dead [ $n = 2$ ], killed [ $n = 2$ ], or photographed [ $n = 1$ ]).