

Study of CP violation in $B^0 \rightarrow DK^*(892)^0$ decays with $D \rightarrow K\pi(\pi\pi)$, $\pi\pi(\pi\pi)$, and KK final states



The LHCb collaboration

E-mail: alex.gilman@physics.ox.ac.uk

ABSTRACT: A measurement of CP -violating observables associated with the interference of $B^0 \rightarrow D^0 K^*(892)^0$ and $B^0 \rightarrow \bar{D}^0 K^*(892)^0$ decay amplitudes is performed in the $D^0 \rightarrow K^\mp \pi^\pm (\pi^+ \pi^-)$, $D^0 \rightarrow \pi^+ \pi^- (\pi^+ \pi^-)$, and $D^0 \rightarrow K^+ K^-$ final states using data collected by the LHCb experiment corresponding to an integrated luminosity of 9 fb^{-1} . CP -violating observables related to the interference of $B_s^0 \rightarrow D^0 \bar{K}^*(892)^0$ and $B_s^0 \rightarrow \bar{D}^0 \bar{K}^*(892)^0$ are also measured, but no evidence for interference is found. The B^0 observables are used to constrain the parameter space of the CKM angle γ and the hadronic parameters $r_{B^0}^{DK^*}$ and $\delta_{B^0}^{DK^*}$ with inputs from other measurements. In a combined analysis, these measurements allow for four solutions in the parameter space, only one of which is consistent with the world average.

KEYWORDS: B Physics, CKM Angle Gamma, CP Violation, Hadron-Hadron Scattering

ARXIV EPRINT: [2401.17934](https://arxiv.org/abs/2401.17934)

Contents

1	Introduction	1
2	LHCb detector and simulation	3
3	Selection, efficiencies, and production and detection asymmetries	4
4	Measurement of CP-violating observables	6
4.1	Modelling the reconstructed mass distribution	6
4.2	Parameterisation of the reconstructed mass distribution	8
4.3	Systematic uncertainties	9
4.4	Results of the reconstructed mass fit	11
5	Interpretation and conclusions	13
A	Correlation matrices	17
	The LHCb collaboration	23

1 Introduction

The only observed phenomena of CP violation to date are attributed to complex phases in the Cabibbo-Kobayashi-Maskawa (CKM) matrix elements [1, 2], which describe the interactions mediated by W^\pm bosons between quarks of different flavour and form a three-by-three unitary matrix. The unitarity of the CKM matrix results in a set of constraints between the matrix elements. One such relation, particularly relevant to decays of B^0 mesons, is commonly parameterised in terms of three angles α , β , and γ which sum to 180° . The angle γ can be measured from the interference of $b \rightarrow c\bar{u}s$ and $b \rightarrow \bar{c}us$ amplitudes with negligible theoretical uncertainties [3]. Precise measurements of γ provide stringent tests of the Standard Model's requirement of CKM unitarity and thus its description of CP violation.

The determination of γ through direct measurements of $b \rightarrow c\bar{u}s$ and $b \rightarrow \bar{c}us$ interference is currently driven by the LHCb experiment. The world average of direct measurements from the HFLAV collaboration is $\gamma = (66.2_{-3.6}^{+3.4})^\circ$ [4], and is dominated by $B^+ \rightarrow DK^+$ decays that have been analysed with the full proton-proton (pp) collision datasets collected by the LHCb detector in 2011–2012 (Run 1) and 2015–2018 (Run 2) [5]. Here and throughout the article, D is used to represent a superposition of D^0 and \bar{D}^0 decays, with a similar convention for D^* . The value of γ can be determined indirectly by assuming CKM unitarity and performing a global fit to all measurements relating to the CKM matrix. With such techniques, the UTFit collaboration determined $\gamma = (64.9 \pm 1.4)^\circ$ [6] and the CKMFitter collaboration determined $\gamma = (65.5_{-1.2}^{+1.3})^\circ$ [7]. While the results of the direct and indirect determinations of γ are in agreement, the comparison is limited by the precision of direct measurements. As such, additional direct measurements of γ are required to provide more

stringent tests of CKM unitarity. Despite the smaller branching fraction of $B^0 \rightarrow DK^*(892)^0$ decays compared to $B^+ \rightarrow DK^+$ decays, a competitive precision on γ can be achieved due to the larger interference effects in these decays compared to B^+ decays [8]. The CP violation in these decays only depends on the flavour at decay, and thus can be studied independent of decay time.

This article presents the measurement of CP -violating observables in the decays $B^0 \rightarrow DK^*(892)^0$ and $\bar{B}^0 \rightarrow D\bar{K}^*(892)^0$ where the D meson is reconstructed through the $K^\mp\pi^\pm$, K^+K^- , $\pi^+\pi^-$, $K^\mp\pi^\pm\pi^+\pi^-$, and $\pi^+\pi^-\pi^+\pi^-$ decay modes with Run 1 and Run 2 pp collision data corresponding to an integrated luminosity of 9fb^{-1} collected at centre-of-mass energies between 7 and 13 TeV. The $K^*(892)^0$ meson is referred to throughout as K^{*0} and is implied to decay to a $K^+\pi^-$ final state. The charge of the kaon child from the K^{*0} candidate is used to determine the flavour of the parent B meson. The interpretation of the measured CP -violating observables in terms of γ and hadronic parameters for $B^0 \rightarrow DK^{*0}$ decays is also presented. This work examines the so-called Atwood-Dunietz-Soni (ADS) final states [9], which experience interference in the admixture of Cabibbo-favoured $D^0 \rightarrow K^-\pi^+(\pi^+\pi^-)$ and doubly Cabibbo-suppressed $\bar{D}^0 \rightarrow K^-\pi^+(\pi^+\pi^-)$ decays. The notation $K\pi(\pi\pi)$ is used to refer to the final state where the kaon child of the D candidate has the same charge as the kaon child of the K^{*0} candidate, while $\pi K(\pi\pi)$ refers to the final state where the two kaons have opposite charge. The CP -eigenstate final states $D \rightarrow \pi^+\pi^-$ and $D \rightarrow K^+K^-$ are also studied and referred to as Gronau-London-Wilder (GLW) decay modes [10, 11]. An extension to the GLW method [12] allows the inclusion of the $D \rightarrow \pi^+\pi^-\pi^+\pi^-$ decay mode which is predominantly CP -even.

These two classes of final states provide complementary sensitivity to the fundamental parameters of interest: the weak phase, γ , the ratio of amplitudes between the $B^0 \rightarrow \bar{D}^0 K^{*0}$ and $B^0 \rightarrow D^0 K^{*0}$ decays, $r_{B^0}^{DK^*}$, and the strong phase difference between the two amplitudes, $\delta_{B^0}^{DK^*}$. The interference can introduce asymmetries in the rates between $\bar{B}^0 \rightarrow D\bar{K}^{*0}$ and $B^0 \rightarrow DK^{*0}$ decays, and modulate the flavour-integrated decay rates of $B^0 \rightarrow DK^{*0}$ for each D meson decay. The effects of interference in the $B^0 \rightarrow D[K\pi(\pi\pi)]K^{*0}$ final state are expected to be small compared to the predicted experimental sensitivity, and so this final state is used as a normalisation channel to probe interference effects in the other final states. This strategy benefits from the cancellation of a large number of systematic uncertainties related to the reconstruction and selection of signal candidates. For the $B^0 \rightarrow D[\pi K(\pi\pi)]K^{*0}$ final state, the parameters

$$\mathcal{R}_{\pi K(\pi\pi)}^+ \equiv \frac{\Gamma(B^0 \rightarrow D[\pi K(\pi\pi)]K^{*0})}{\Gamma(B^0 \rightarrow D[K\pi(\pi\pi)]K^{*0})} \quad \text{and} \quad \mathcal{R}_{\pi K(\pi\pi)}^- \equiv \frac{\Gamma(\bar{B}^0 \rightarrow D[\pi K(\pi\pi)]\bar{K}^{*0})}{\Gamma(\bar{B}^0 \rightarrow D[K\pi(\pi\pi)]\bar{K}^{*0})} \quad (1.1)$$

are measured. The asymmetry of the $B^0 \rightarrow D[K\pi(\pi\pi)]K^{*0}$ final state,

$$\mathcal{A}_{K\pi} \equiv \frac{\Gamma(\bar{B}^0 \rightarrow D[K\pi(\pi\pi)]\bar{K}^{*0}) - \Gamma(B^0 \rightarrow D[K\pi(\pi\pi)]K^{*0})}{\Gamma(\bar{B}^0 \rightarrow D[K\pi(\pi\pi)]\bar{K}^{*0}) + \Gamma(B^0 \rightarrow D[K\pi(\pi\pi)]K^{*0})}, \quad (1.2)$$

is also measured.

For the GLW final states, the flavour-integrated decay rates relative to the $B^0 \rightarrow D[K\pi(\pi\pi)]K^{*0}$ decay are measured, leading to the cancellation of most systematic

uncertainties related to the estimation of selection efficiencies. The asymmetries do not require a normalisation channel for the cancellation of systematic uncertainties, so the observables of interest are

$$\mathcal{R}_{CP}^{hh(\pi\pi)} \equiv \frac{\Gamma(\bar{B}^0 \rightarrow D[hh(\pi\pi)]\bar{K}^{*0}) + \Gamma(B^0 \rightarrow D[hh(\pi\pi)]K^{*0})}{\Gamma(\bar{B}^0 \rightarrow D[K\pi(\pi\pi)]\bar{K}^{*0}) + \Gamma(B^0 \rightarrow D[K\pi(\pi\pi)]K^{*0})} \times \frac{\mathcal{B}(D^0 \rightarrow K\pi(\pi\pi))}{\mathcal{B}(D^0 \rightarrow hh(\pi\pi))} \quad (1.3)$$

and

$$\mathcal{A}_{CP}^{hh(\pi\pi)} \equiv \frac{\Gamma(\bar{B}^0 \rightarrow D[hh(\pi\pi)]\bar{K}^{*0}) - \Gamma(B^0 \rightarrow D[hh(\pi\pi)]K^{*0})}{\Gamma(\bar{B}^0 \rightarrow D[hh(\pi\pi)]\bar{K}^{*0}) + \Gamma(B^0 \rightarrow D[hh(\pi\pi)]K^{*0})}, \quad (1.4)$$

where $hh(\pi\pi)$ refers to KK , $\pi\pi$, or $\pi\pi\pi\pi$.

The CP -violating observables in the decay $B_s^0 \rightarrow D\bar{K}^{*0}$ are also measured in this analysis, and can be parameterised in an identical fashion. However, the effect of the interference in this process is expected to be negligible in all of the observed decay modes, as the parameter directly proportional to the interference effects, $r_{B_s^0}^{DK^*}$, is estimated to be roughly an order of magnitude smaller than the corresponding B^0 parameter based on the known CKM elements [4]. Similar notation is used to refer to the B_s^0 observables throughout the article, with an additional s in the subscript.

All of these observables have been measured previously by the LHCb collaboration using the data collected in Run 1 and 2015–2016, except the ones associated with the $D \rightarrow \pi\pi\pi\pi$ decay channel, for which only the 2015–2016 data was analysed [13]. The measurements presented in this article supersede the results of the previous analyses.

The rest of the article is organised as follows. The LHCb detector and simulation is described in section 2. The selection requirements placed on signal candidates are discussed in section 3. The determination of CP -violating observables from the signal candidates passing selection requirements is discussed in section 4. The interpretation of the determined CP -violating observables in terms of the fundamental physics parameters and concluding remarks are presented in section 5.

2 LHCb detector and simulation

The LHCb detector [14, 15] is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the pp interaction region [16], a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes [17] placed downstream of the magnet. The tracking system provides a measurement of the momentum, p , of charged particles with a relative uncertainty that varies from 0.5% at low momentum to 1.0% at 200 GeV/ c . The minimum distance of a track to a primary pp collision vertex (PV), the impact parameter (IP), is measured with a resolution of $(15 + 29/p_T) \mu\text{m}$, where p_T is the component of the momentum transverse to the beam, in GeV/ c . Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov (RICH) detectors [18, 19]. Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower detectors,

an electromagnetic and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers [20].

The events considered in the analysis are triggered at the hardware level either when one of the final-state tracks of the signal decay deposits enough energy in the calorimeter system, or when one of the other particles in the event, not reconstructed as part of the signal candidate, fulfils any trigger requirement. At the software level, it is required that at least one particle has high p_T and high χ_{IP}^2 , where χ_{IP}^2 is defined as the difference in the PV fit χ^2 with and without the inclusion of that particle. A multivariate algorithm [21] is used to identify displaced vertices consistent with being a two-, three-, or four-track b -hadron decay. The PVs are fitted with and without the B candidate tracks, and the PV that gives the smallest χ_{IP}^2 is associated with the B candidate.

Simulation is used to model the reconstructed mass distributions of the signal and background contributions and determine their selection efficiencies. In the simulation, pp collisions are generated using PYTHIA [22] with a specific LHCb configuration [23]. Decays of unstable particles are described by EVTGEN [24], in which final-state radiation is generated using PHOTOS [25]. The simulation of interactions of the generated particles with the detector and its response is implemented using the GEANT4 toolkit [26, 27] as described in ref. [28]. Some subdominant sources of background are generated with RapidSim, a fast simulation [29] that mimics the geometric acceptance and tracking resolution of the LHCb detector as well as the dynamics of the decay via EVTGEN.

3 Selection, efficiencies, and production and detection asymmetries

Signal B -meson candidates are built from a $K^{*0} \rightarrow K^+\pi^-$ candidate and a D -meson candidate. Selection requirements identical to those from ref. [30] are placed on the $K^{*0} \rightarrow K^+\pi^-$ candidate to select the K^{*0} resonance: the reconstructed mass of the K^{*0} candidate must be within $50 \text{ MeV}/c^2$ of the known K^{*0} meson mass [31] and a requirement is made on the angle θ^* between the K^{*0} kaon child momentum and the B^0 candidate in the K^{*0} rest frame of $|\cos \theta^*| > 0.4$. The reconstructed mass of the D candidate is required to be within $25 \text{ MeV}/c^2$ of the known D^0 meson mass [31].

With the above selections in place, boosted decision tree classifiers with gradient boost (BDT) [32] are trained to discriminate between signal candidates and combinatorial background. Two BDT classifiers are trained for each final state, one for each run period. The same BDT classifier is shared for each run period between the $B^0 \rightarrow [K\pi(\pi\pi)]K^{*0}$ and $B^0 \rightarrow [\pi K(\pi\pi)]K^{*0}$ final states. This gives ten separate BDT classifiers in total, each trained using simulated signal samples and background samples from data with a B^0 candidate reconstructed mass within the range $5800\text{--}5960 \text{ MeV}/c^2$. The classifiers are trained using the angle between the direction of the reconstructed B momentum and the direction defined by the primary and secondary vertices, the χ_{IP}^2 of the B^0 and D candidates, the reduced χ^2 of the B^0 vertex fit, the χ_{IP}^2 and the transverse momenta of the K^{*0} candidate children, the χ_{IP}^2 and the transverse momenta of the D candidate children (only for two-body D final states), and the transverse momentum imbalance of the B^0 candidate, defined as

$$I_{p_T} \equiv \frac{p_T(B^0) - \sum_X p_T(X)}{p_T(B^0) + \sum_X p_T(X)}, \quad (3.1)$$

where the summation is over all charged tracks X inconsistent with originating from the primary vertex within a cone around the B^0 candidate, excluding those used in the B^0 reconstruction [33]. Variables related to the D children are omitted for the four-body D final states to avoid any dependence on the modelling of the D decay dynamics. Requirements on the output of each BDT classifier are chosen based on an optimisation of the estimated sensitivity to CP -violating observables. Each BDT classifier retains roughly 85–90% of the signal candidates and removes over 90% of the combinatorial background.

Stringent particle identification (PID) requirements are placed on the kaon child of the K^{*0} candidate to reduce misidentification of the K^{*0} flavour to negligible levels, as the charge of this particle is used to determine the flavour of the parent B meson. More relaxed particle identification requirements are also placed on the pion from the K^{*0} candidate and all kaons and pions from the two-body D -meson decay candidates. For $D \rightarrow K\pi\pi\pi$ and $D \rightarrow \pi K\pi\pi$ decay candidates, the kaon and both pions with charge opposite to the D candidate kaon child are subject to PID requirements. For $D \rightarrow \pi\pi\pi\pi$ candidates, only the two pions with the same charge as the K^{*0} candidate kaon child are subject to PID requirements. All particles with PID requirements are also required to have track momentum between 3–200 GeV/ c to ensure suitable discrimination between kaons and pions in the RICH detectors.

Requirements are also placed on the displacement of the D candidate decay vertex from the B^0 candidate decay vertex to suppress charmless B^0 decays to the same final states, which proceed without an intermediate D meson. Different requirements are placed on the ADS and GLW final states due to different relative contributions of charmless backgrounds. For the GLW (ADS) final states, the D candidate decay vertex displacement is required to be at least three (two) times its uncertainty. A discussion of the remaining contributions from these charmless backgrounds is included in section 4.1.

Finally, vetoes are applied to each D final state to remove backgrounds or maintain consistency with the measurements of D hadronic parameters used in section 5. In the ADS final states, $D \rightarrow K\pi$ decays misidentified as $D \rightarrow \pi K$ due to two incorrect PID assignments are removed with the requirement that the reconstructed D mass where the mass hypotheses of the kaon and pion children have been swapped differs from the known D^0 meson mass by more than $15 \text{ MeV}/c^2$. Backgrounds from $B^\pm \rightarrow DK^\pm$ decays paired with a random pion are removed with a requirement that the DK^\pm reconstructed mass differ by more than $25 \text{ MeV}/c^2$ from the known B^\pm meson mass. Decays of the B^0 meson to the same final state proceeding through different charmed intermediate states, e.g. $B^0 \rightarrow D^- [K^+ \pi^- \pi^-] \pi^+$ misidentified as $B^0 \rightarrow D [\pi^+ \pi^-] K^+ \pi^-$, are removed with requirements that the reconstructed mass of a D child combined with the K^{*0} candidate to form a Cabibbo-favoured decay of another charmed meson does not fall within $15 \text{ MeV}/c^2$ of the known meson mass. Candidate $D \rightarrow K_S^0 \pi^+ \pi^-$ decays are removed from the $D \rightarrow \pi^+ \pi^- \pi^+ \pi^-$ sample with the requirement that the reconstructed $\pi^+ \pi^-$ masses are not within $480\text{--}505 \text{ MeV}/c^2$.

The CP -violating observables of interest are all proportional to the ratios of decay rates of different D final states, and thus much of the reconstruction and selection efficiency cancels in these ratios. However, due to the different kinematic distributions of each D final state, and due to different selection requirements placed on the different D decay modes, these efficiencies must be accounted for. Additionally, corrections for effects introducing asymmetries aside from those due to the interference discussed in section 1 must be applied. The selection

requirements are applied to simulated samples to estimate the selection efficiencies of each final state integrated across the two B flavours, with the exception of PID efficiencies. Charge-dependent PID efficiencies are estimated with calibration data samples of kaons and pions collected from D^* decays weighted to match the kinematics of signal decays [34]. The resulting efficiency of reconstructing and selecting signal candidates is approximately 2×10^{-3} .

Three other possible sources of flavour asymmetry must be accounted for, aside from that resulting from the interference discussed in section 1. First, an asymmetry can be introduced in the production of $B_{(s)}^0$ mesons and their antiparticles from the initial pp collisions, referred to as A_{prod} . Second, tracking and reconstruction can depend on the charge of final-state particles due to the difference of interactions between particles and anti-particles and the detector material, which is accounted for through the difference in detection asymmetry between kaons and pions $A_{K\pi} = A_K - A_\pi$. Lastly, similar to above, asymmetries can be introduced in PID which is accounted for implicitly in the PID efficiency corrections discussed previously. All discussed asymmetries are defined to match the flavour conventions introduced in section 1.

The quantity A_{prod} is determined from the measurement of B^0 and B_s^0 asymmetries in ref. [35]. The measured production asymmetries are $A_{\text{prod}} = (-8 \pm 5) \times 10^{-3}$ and $A_{s,\text{prod}} = (6 \pm 10) \times 10^{-3}$. Since these results are only determined using data from pp centre-of-mass energies of 7 and 8 TeV, and not at 13 TeV, the same central values are assumed for the full dataset, but with inflated uncertainty.

The value of $A_{K\pi}$ is estimated using calibration data samples of $D^+ \rightarrow K^- \pi^+ \pi^+$ and $D^+ \rightarrow K_S^0 \pi^+$ decays. The resulting predicted asymmetry weighted to match the signal kinematic distributions is $A_{K\pi} = (-9.8 \pm 5.5) \times 10^{-3}$. The total detection asymmetry for a $B^0 \rightarrow DK^{*0}$ candidate depends on the difference between the number of kaons and the number of pions of the same sign in the final state, e.g. the $B^0 \rightarrow D[K\pi]K^{*0}$ final state is corrected by $2A_{K\pi}$, the $B^0 \rightarrow D[\pi\pi]K^{*0}$ final state is corrected by $A_{K\pi}$, while the $B^0 \rightarrow D[\pi K]K^{*0}$ final state requires no such correction.

4 Measurement of CP -violating observables

The CP -violating observables discussed in section 1 are determined from the selected data through a simultaneous unbinned extended maximum-likelihood fit of the B^0 reconstructed mass for each flavour of each D final state, where constraints are placed on the D candidate mass and the direction of the B candidate momentum using the DECAYTREEFITTER package [36]. The reconstructed mass is calculated with these constraints unless otherwise specified. Only candidates with a reconstructed mass within the interval 5000–5800 MeV/ c^2 are considered.

4.1 Modelling the reconstructed mass distribution

The reconstructed mass distributions of both $B^0 \rightarrow DK^{*0}$ and $B_s^0 \rightarrow D\bar{K}^{*0}$ decays are modelled by modified Cruijff functions [37] parameterised as

$$f(M) = \begin{cases} e^{-\frac{(M-\mu)^2(1+\beta(M-\mu)^2)}{2\sigma^2+\alpha_L(M-\mu)^2}}, & M < \mu \\ e^{-\frac{(M-\mu)^2(1+\beta(M-\mu)^2)}{2(\rho\sigma)^2+\alpha_R(M-\mu)^2}}, & M > \mu \end{cases}. \quad (4.1)$$

For each decay mode, all the parameters of the $B_{(s)}^0 \rightarrow DK^{*0}$ modified Cruijff functions are fixed from fits to the corresponding simulation samples of $B^0 \rightarrow D[K\pi]K^{*0}$ and $B_s^0 \rightarrow D[\pi K]K^{*0}$ decays, with the exception of the mean, μ , and width, σ , which are left as free parameters. The constraints on the D meson mass minimise the dependence on the D decay final state, but the four-body D final states have a slightly larger width. The ratio of the two-body width and four-body width is determined from simulation and fixed in the fit to data.

The selection requirements accept $B_{(s)}^0 \rightarrow D^*K^{*0}$ decays with a similar efficiency to signal decays, as their topology is identical to signal decays with one additional missing neutral particle from the $D^* \rightarrow \gamma/\pi^0 D$ decay. The reconstructed masses of these decays are mainly located below the B^0 mass peak, but the right-hand tail of the $B_s^0 \rightarrow D^*\bar{K}^{*0}$ distribution extends to the B^0 meson mass. As this is the decay of a pseudoscalar particle into two vector particles, the final state is characterised by three independent helicity amplitudes $(0, \pm 1)$. However, two of these amplitudes (± 1) have indistinguishable reconstructed mass distributions. This leaves four different configurations of $B_{(s)}^0 \rightarrow D^*K^{*0}$ decays: two possible D^* decay chains each with two reconstructed mass distributions. Each configuration is modelled with either a single broad peaking structure or a double-peaked structure, as described in ref. [33]. The relative size of the $D^*[\gamma D]$ and $D^*[\pi^0 D]$ decays for each helicity state are fixed based on efficiencies predicted from simulation and the measured branching fractions. The interference effects of interest can affect the helicity amplitudes differently. Therefore, four free parameters are included in the fit to account for this: one for the ratio of helicity states in ADS $B^0 \rightarrow D^*K^{*0}$ decays, one ratio for each flavour of the GLW $B^0 \rightarrow D^*K^{*0}$ decays, and one ratio shared across all $B_s^0 \rightarrow D^*\bar{K}^{*0}$ final states.

Due to particle misidentification, $B^0 \rightarrow D\pi^+\pi^-$ and $B^0 \rightarrow D^*\pi^+\pi^-$ decays can also be mistaken for signal decays when one pion is misidentified as a kaon. The relatively stringent PID requirement on the K^+ child of the K^{*0} candidate suppresses these contributions significantly, but not to negligible levels. The reconstructed mass distribution of these decays is taken from simulated samples with PID weights determined from calibration data samples. The simulation samples only consider the contribution of $B^0 \rightarrow D\rho^0[\pi^+\pi^-]$ and $B^0 \rightarrow D^*\rho^0[\pi^+\pi^-]$, as the selection requirements on the K^{*0} candidate significantly suppress other contributions. The reconstructed mass distribution of $B^0 \rightarrow D\rho^0[\pi^+\pi^-]$ decays is modelled with a double-sided Crystal Ball function [38]. The probability distribution function (PDF) for $B^0 \rightarrow D^*\rho^0[\pi^+\pi^-]$ decays is parameterised in a similar fashion to the $B^0 \rightarrow D^*K^{*0}$ decays, and the PDF shares the ratio of helicity states with ADS $B^0 \rightarrow D^*K^{*0}$ decays. The peak of the $B^0 \rightarrow D\rho^0[\pi^+\pi^-]$ distribution is located between the B^0 and B_s^0 peaks, while the $B^0 \rightarrow D^*\rho^0[\pi^+\pi^-]$ distribution has minimal contribution near or above the B^0 mass peak.

Partially reconstructed $B^+ \rightarrow DK^+\pi^+\pi^-$ decays, where the π^+ is not included in the reconstruction, also pass final selection requirements. The reconstructed mass distribution of these decays is modelled with a smoothed kernel density estimation [39] of simulated $B^+ \rightarrow DK_1(1270)^+$ decays, and contributes negligibly near the B^0 mass peak, being distributed primarily around a reconstructed mass of $5050 \text{ MeV}/c^2$.

Requirements placed on the D candidate flight significance do not fully suppress the contribution of charmless B^0 and B_s^0 decays to the same final states as the observed signal channels. The size of these contributions and possible flavour asymmetries in each final

Branching Fraction	Value	
$\mathcal{B}(D^0 \rightarrow K^- \pi^+)$	$(3.999 \pm 0.045) \%$	[4]
$\mathcal{B}(D^0 \rightarrow \pi^+ \pi^-)$	$(1.490 \pm 0.027) \times 10^{-3}$	[4]
$\mathcal{B}(D^0 \rightarrow K^+ K^-)$	$(4.113 \pm 0.051) \times 10^{-3}$	[4]
$\mathcal{B}(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)$	$(8.22 \pm 0.14) \%$	[31]
$\mathcal{B}(D^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^-)$	$(8.06 \pm 0.20) \times 10^{-3}$	[31]

Table 1. Summary of branching fractions used as inputs in the measurement.

state are estimated using data in the sidebands of the D -candidate reconstructed mass distribution. However, as the default selection and reconstruction algorithm applies a mass constraint on the D candidate, the B^0 -candidate reconstructed-mass distribution in those sidebands is biased. To address this, the number of charmless decays in our selection sample is estimated with samples from the D -candidate sideband regions for each final state by fitting to the B^0 candidate reconstructed-mass distribution without any constraint applied on the reconstructed D -candidate mass. The number of events found in the sidebands are used to estimate the number of charmless decays that pass the selection requirements. The magnitude of the flavour asymmetry in these decays is estimated with a similar procedure, however with a significantly less stringent requirement on the D flight significance to increase statistical precision. Significant charmless contributions from B_s^0 decays are only observed in the $D \rightarrow \pi K$ and $D \rightarrow \pi K \pi \pi$ final states. The largest contribution from charmless decays is found in the $B^0 \rightarrow D[\pi\pi]K^{*0}$ and $\bar{B}^0 \rightarrow D[\pi\pi]\bar{K}^{*0}$ samples with a yield that is roughly 10% of the determined signal yield.

The combinatorial background is modelled by an exponential PDF. The parameters of the exponential PDF associated with each D decay mode vary freely in the fit, but are shared across B flavour for each D decay mode.

4.2 Parameterisation of the reconstructed mass distribution

The fit to determine the number of signal decays in each of the fourteen samples (two flavours for each of the seven D final states) is parameterised in terms of the CP -violating observables introduced in section 1, the flavour-integrated number of observed $B^0 \rightarrow D[K\pi]K^{*0}$ decays, and the flavour-integrated number of observed $B^0 \rightarrow D[K\pi\pi\pi]K^{*0}$ decays. These observables are related to the number of observed decays in each signal mode and the efficiency and asymmetry corrections discussed in section 4.1. The $\mathcal{R}_{CP}^{hh(\pi\pi)}$ parameters are dependent on measured D^0 branching fractions, reported in table 1.

The $B_s^0 \rightarrow D\bar{K}^{*0}$, $B^0 \rightarrow D^*K^{*0}$, $B_s^0 \rightarrow D^*\bar{K}^{*0}$, and $B^+ \rightarrow DK^+\pi^+\pi^-$ background components are parameterised in a similar fashion. The CP -violating observables of $B_s^0 \rightarrow D\bar{K}^{*0}$ decays are left as free parameters in the fit, as are the CP -violating observables of $B^0 \rightarrow D^*K^{*0}$ decays. The observables of $B_s^0 \rightarrow D^*\bar{K}^{*0}$ decays are fixed to have no effects from interference, and the CP -violating observables of $B^+ \rightarrow DK^+\pi^+\pi^-$ decays are fixed to the measurements for the two-body D final states from ref. [40] and predicted values based on the results of ref. [5] for the four-body D final states. The $B^0 \rightarrow D\pi^+\pi^-$ and $B^0 \rightarrow D^*\pi^+\pi^-$ yields are constrained to be equal for $B^0 \rightarrow D[K\pi]K^{*0}$ and $B^0 \rightarrow D[\pi K]K^{*0}$ decays, due to

an equal probability of misidentifying the π^+ meson and the π^- meson. For all other decay modes, the yields are constrained relative to the yields in $B^0 \rightarrow D[K\pi]K^{*0}$, and scaled by the ratio of D branching fractions and selection efficiencies. The number of charmless decays is subject to Gaussian constraints according to the determined values. The combinatorial backgrounds are parameterised in terms of the observed number of candidates, which is required to be equal for each flavour of a given D decay mode. The possibility of biases introduced in the fitting procedure is investigated by fitting toy data samples generated from the result of the total PDF fit to data. No evidence is seen for biases in the central values or uncertainties of any of the CP -violating observables.

4.3 Systematic uncertainties

The following sources of systematic uncertainties are considered: uncertainty in the determination of the asymmetry and efficiency corrections, uncertainty from the measured branching fractions used as inputs in the fit, and uncertainty in the modelling of the PDFs or in the determination of the fixed yields in the fit. Any systematic uncertainties that are estimated at less than 5% of the measured statistical uncertainty from the fit are considered negligible, and are excluded in the final determination of systematic uncertainties. A summary of the assigned systematic uncertainties is provided in tables 2 and 3. The total systematic uncertainty is calculated as the sum in quadrature of the individual sources. The dominant systematic uncertainties are due to the measurement of input branching fractions, asymmetry corrections, and estimation of particle identification efficiencies, but are generally small compared to statistical uncertainties. The assigned systematic uncertainties are slightly larger than those in the previous analysis from ref. [13] as more detailed study is possible with the larger dataset.

The uncertainties of A_{prod} and $A_{K\pi}$ are propagated to the parameters of interest based on the quoted uncertainties in section 3. The measured uncertainty of A_{prod} [35] is doubled for the data samples collected at 13 TeV to account for possible energy dependence in the asymmetry. Additionally, the possibility of asymmetries introduced by the hardware trigger is estimated to be at the level of 10^{-3} based on the results of ref. [33]. The uncertainty due to the selection efficiencies largely cancels, with the primary uncertainty arising from different PID requirements on different final states. Uncertainties in estimated PID efficiencies arise from the limited statistics of simulated samples for kinematic weighting and the finite size of bins used in the weighting of the PID efficiency. The propagated uncertainties of the input branching fractions listed in table 1 are also determined.

The uncertainty due to the modelling of the B^0 candidate reconstructed mass for each component in the fit is also investigated. The uncertainty due to the modelling of $B_s^0 \rightarrow D^*K^{*0}$ decays is assessed by varying the determined efficiencies and input branching fractions of the $D^*[\gamma D]$ and $D^*[\pi^0 D]$ components within their respective uncertainties. Two alternative PDFs are examined to assess uncertainty in the modelling of $B^+ \rightarrow DK^+\pi^+\pi^-$ decays: one based on simulated samples of $B^+ \rightarrow DK^*(1400)^0[K^+\pi^+\pi^-]$ decays, and another based on $B^+ \rightarrow DK^+\pi^+\pi^-$ decays collected from data. Performing the fit with either distribution results in negligible variations in the CP -violating observables. An alternative $B^0 \rightarrow D\pi^+\pi^-$ reconstructed mass distribution composed of an equal admixture of $B^0 \rightarrow D\rho^0[\pi^+\pi^-]$ and non-resonant $B^0 \rightarrow D(\pi\pi)_{S\text{-wave}}$ decays based on results in ref. [41] is also considered. These uncertainties are negligible for all CP -violating observables except $R_{\pi K(\pi\pi)}^\pm$.

	$\mathcal{A}_{K\pi}$	$\mathcal{R}_{\pi K}^+$	$\mathcal{R}_{\pi K}^-$	$\mathcal{R}_{CP}^{\pi\pi}$	$\mathcal{A}_{CP}^{\pi\pi}$	\mathcal{R}_{CP}^{KK}	\mathcal{A}_{CP}^{KK}	$\mathcal{A}_{K\pi\pi\pi}$	$R_{\pi K\pi\pi}^+$	$R_{\pi K\pi\pi}^-$	$\mathcal{R}_{CP}^{4\pi}$	$\mathcal{A}_{CP}^{4\pi}$
A_{prod}	0.009	0.001	0.001	0.002	0.009	-	0.009	0.009	0.001	0.001	-	0.009
$A_{K\pi}$	0.009	0.001	0.001	0.001	0.006	0.001	0.006	0.009	0.001	0.001	0.001	0.006
PID	0.008	0.004	0.004	0.012	0.010	0.010	0.010	0.010	0.004	0.004	0.017	0.011
Fit PDFs	-	0.003	0.003	-	-	-	-	-	0.003	0.003	-	-
Charmless Asymmetries	-	-	-	0.011	0.005	-	-	-	0.001	0.001	-	-
Combinatorial Asymmetries	-	0.002	0.002	-	0.002	-	0.001	-	0.004	0.004	-	0.001
Input Branching Fractions	-	-	-	0.020	-	0.013	-	-	-	-	0.028	-
Total systematic	0.015	0.005	0.005	0.026	0.016	0.017	0.015	0.016	0.006	0.006	0.033	0.016
Statistical	0.017	0.013	0.013	0.110	0.103	0.057	0.064	0.018	0.014	0.014	0.084	0.088

Table 2. Summary of uncertainties on the physics observables measured in $B^0 \rightarrow DK^{*0}$ decays. Uncertainties smaller than 5% of the statistical uncertainty are indicated with dashed lines.

	$\mathcal{A}_{s,\pi K}$	$\mathcal{R}_{s,K\pi}^+$	$\mathcal{R}_{s,K\pi}^-$	$\mathcal{R}_{CP}^{s,\pi\pi}$	$\mathcal{A}_{CP}^{s,\pi\pi}$	$\mathcal{R}_{CP}^{s,KK}$	$\mathcal{A}_{CP}^{s,KK}$	$\mathcal{A}_{s,\pi K\pi\pi}$	$R_{s,K\pi\pi}^+$	$R_{s,K\pi\pi}^-$	$\mathcal{R}_{CP}^{s,4\pi}$	$\mathcal{A}_{CP}^{s,4\pi}$
$A_{s,\text{prod}}$	0.018	0.003	0.003	-	0.018	-	0.018	0.018	0.003	0.003	-	0.018
$A_{K\pi}$	-	0.002	0.002	0.001	0.006	0.001	0.006	-	0.002	0.002	0.001	0.006
PID	0.008	0.004	0.004	0.012	0.010	0.010	0.010	0.010	0.004	0.004	0.017	0.011
Fit PDFs	-	0.002	0.002	-	-	-	-	-	0.002	0.002	-	-
Charmless Asymmetries	-	-	-	-	-	-	-	-	-	-	-	-
Combinatorial Asymmetries	-	0.002	0.002	-	0.002	-	0.001	-	0.004	0.004	-	0.001
Input Branching Fractions	-	-	-	0.020	-	0.013	-	-	-	-	0.028	-
Total Systematic	0.020	0.006	0.006	0.023	0.021	0.016	0.021	0.021	0.007	0.007	0.033	0.022
Statistical	0.011	0.002	0.002	0.056	0.057	0.032	0.034	0.012	0.004	0.004	0.044	0.048

Table 3. Summary of uncertainties on the physics observables measured in $B_s^0 \rightarrow DK^{*0}$ decays. Uncertainties smaller than 5% of the statistical uncertainty are indicated with dashed lines.




Figure 1. Reconstructed mass distributions for selected candidates in the (left) \bar{B}^0 and (right) B^0 samples for the (top) $D \rightarrow K\pi$ and (bottom) $D \rightarrow \pi K$ decay channels. The fit projection is overlaid.

The statistical uncertainty of the flavour-integrated number of charmless decays is accounted for through Gaussian constraints in the fit, but the asymmetry of the charmless yields is fixed based on the central values determined in data. The resulting uncertainty on the parameters of interest is determined through variations of the charmless asymmetries within their measured uncertainties.

The assumption of equal combinatorial background in B^0 and \bar{B}^0 candidate samples is studied using pseudoexperiments, where an ensemble of datasets is generated with asymmetric background yields. The magnitude of the asymmetry generated is determined from an alternative fit to the data. The pseudoexperiments are fit with the default model and biases in the fitted observables are taken as the systematic uncertainty.

4.4 Results of the reconstructed mass fit

The reconstructed B candidate mass distributions for different D decay modes are shown with the results of the fit overlaid in figures 1–3. The yields of the favoured decays, $N_{K\pi}$, $N_{K\pi\pi\pi}$, $N_{s,\pi K}$, and $N_{s,\pi K\pi\pi}$ are found to be roughly 3800, 3600, 8800, and 8200, respectively. The CP -violating observables determined in the fit for $B^0 \rightarrow DK^{*0}$ decays are given in table 4 and the results of $B_s^0 \rightarrow D\bar{K}^{*0}$ are given in table 5. The results from the B_s^0 decays conform to the expectation of unobservable effects of interference: all \mathcal{R}_{CP}^s parameters are consistent with unity within one standard deviation, and no statistically significant asymmetries are observed. However, statistically significant evidence of interference is seen in the $B^0 \rightarrow DK^{*0}$ results, most notably in \mathcal{R}_{CP}^{KK} at the 3σ level and $\mathcal{R}_{CP}^{4\pi}$ at the 2σ level.




Figure 2. Reconstructed mass distributions for selected candidates in the (left) \bar{B}^0 and (right) B^0 samples for the (top) $D \rightarrow K\pi\pi\pi$ and (bottom) $D \rightarrow \pi K\pi\pi$ decay modes. The fit projections are overlaid.

Parameter	Value
$\mathcal{A}_{K\pi}$	$0.031 \pm 0.017 \pm 0.015$
$\mathcal{R}_{\pi K}^+$	$0.069 \pm 0.013 \pm 0.005$
$\mathcal{R}_{\pi K}^-$	$0.093 \pm 0.013 \pm 0.005$
$\mathcal{A}_{K\pi\pi\pi}$	$-0.012 \pm 0.018 \pm 0.016$
$\mathcal{R}_{\pi K\pi\pi}^+$	$0.060 \pm 0.014 \pm 0.006$
$\mathcal{R}_{\pi K\pi\pi}^-$	$0.038 \pm 0.014 \pm 0.006$
\mathcal{R}_{CP}^{KK}	$0.811 \pm 0.057 \pm 0.017$
\mathcal{A}_{CP}^{KK}	$-0.047 \pm 0.063 \pm 0.015$
$\mathcal{R}_{CP}^{\pi\pi}$	$1.104 \pm 0.111 \pm 0.026$
$\mathcal{A}_{CP}^{\pi\pi}$	$-0.034 \pm 0.094 \pm 0.016$
$\mathcal{R}_{CP}^{4\pi}$	$0.882 \pm 0.086 \pm 0.033$
$\mathcal{A}_{CP}^{4\pi}$	$0.021 \pm 0.087 \pm 0.016$

Table 4. Fitted CP -violating observables related to $B^0 \rightarrow DK^{*0}$ decays. The first stated uncertainties are statistical and the second are systematic.

An approximately 2σ tension is observed between the values of \mathcal{R}_{CP}^{KK} and $\mathcal{R}_{CP}^{\pi\pi}$, which are expected to be consistent, but this difference is attributed to a statistical fluctuation. The asymmetries \mathcal{A}_{CP}^{KK} and $\mathcal{A}_{CP}^{\pi\pi}$ are consistent with each other. In aggregate, the results are self-consistent, as evidenced by our interpretation of the parameters in section 5.




Figure 3. Reconstructed mass distributions for selected candidates in the (left) \bar{B}^0 and (right) B^0 samples for the (top) $D \rightarrow \pi\pi$, (centre) $D \rightarrow KK$, and (bottom) $D \rightarrow \pi\pi\pi\pi$ decay modes. The fit projections are overlaid.

Statistical precision on the CP -violating observables has improved by around 60% in comparison to the previous results in ref. [13], an improvement which is consistent with the increase in signal yield. Central values are also consistent taking into account correlations and the more accurate determination of the contribution of charmless decays with the larger dataset.

5 Interpretation and conclusions

The measured parameters from table 4 can be expressed in terms of the fundamental physics parameters of interest, γ , $r_{B^0}^{DK^*}$, and $\delta_{B^0}^{DK^*}$, and some additional inputs. These additional inputs include the coherence factor κ_{B^0} , which quantifies the dilution of the interference effects of interest due to the selected $B^0 \rightarrow DK^+\pi^-$ decays that do not proceed through

Parameter	Value
$\mathcal{A}_{s,\pi K}$	$-0.009 \pm 0.011 \pm 0.020$
$\mathcal{R}_{s,K\pi}^+$	$0.004 \pm 0.002 \pm 0.006$
$\mathcal{R}_{s,K\pi}^-$	$0.004 \pm 0.002 \pm 0.006$
$\mathcal{A}_{s,\pi K\pi\pi}$	$-0.029 \pm 0.012 \pm 0.021$
$\mathcal{R}_{s,K\pi\pi\pi}^+$	$0.019 \pm 0.004 \pm 0.007$
$\mathcal{R}_{s,K\pi\pi\pi}^-$	$0.015 \pm 0.004 \pm 0.007$
$\mathcal{R}_{CP}^{s,KK}$	$1.000 \pm 0.034 \pm 0.016$
$\mathcal{A}_{CP}^{s,KK}$	$0.062 \pm 0.032 \pm 0.021$
$\mathcal{R}_{CP}^{s,\pi\pi}$	$0.996 \pm 0.057 \pm 0.023$
$\mathcal{A}_{CP}^{s,\pi\pi}$	$-0.001 \pm 0.056 \pm 0.021$
$\mathcal{R}_{CP}^{s,4\pi}$	$1.010 \pm 0.048 \pm 0.033$
$\mathcal{A}_{CP}^{s,4\pi}$	$0.017 \pm 0.044 \pm 0.022$

Table 5. Fitted CP -violating observables related to $B_s^0 \rightarrow D\bar{K}^{*0}$ decays. The first stated uncertainties are statistical and the second are systematic.

an intermediate K^{*0} resonance. The value of this parameter measured by ref. [30] is used, which placed identical requirements on the K^{*0} candidate to those presented in this article. There is additional dependence on the hadronic parameters associated with the D meson decays to a final state X : r_D^X , δ_D^X , and κ_D^X for ADS decays, and the CP -even fraction F_+^X for GLW decays. The values of the ADS parameters are taken from ref. [5] for the $D \rightarrow K\pi$ decay (except for $\kappa_D^{K\pi}$, which is exactly one) and from ref. [42] for the $D \rightarrow K\pi\pi\pi$ decay. For the two-body GLW modes, F_+^{hh} is exactly one, and the value of $F_+^{4\pi} = 0.746 \pm 0.013$ is taken from an average of the results from ref. [43] and ref. [44]. The GLW observables $\mathcal{A}_{CP}^{hh(\pi\pi)}$ and $\mathcal{R}_{CP}^{hh(\pi\pi)}$ relate to these parameters through

$$\begin{aligned}
 \mathcal{A}_{CP}^{hh(\pi\pi)} &= \frac{2\kappa_{B^0} r_{B^0}^{DK^*} (2F_+^{hh(\pi\pi)} - 1) \sin(\delta_{B^0}^{DK^*}) \sin(\gamma)}{1 + (r_{B^0}^{DK^*})^2 + 2\kappa_{B^0} r_{B^0}^{DK^*} (2F_+^{hh(\pi\pi)} - 1) \cos(\delta_{B^0}^{DK^*}) \cos(\gamma)}, \\
 \mathcal{R}_{CP}^{hh(\pi\pi)} &= \frac{1 + (r_{B^0}^{DK^*})^2 + 2\kappa_{B^0} r_{B^0}^{DK^*} (2F_+^{hh(\pi\pi)} - 1) \cos(\delta_{B^0}^{DK^*}) \cos(\gamma)}{1 + (r_{B^0}^{DK^*})^2 (r_D^{K\pi\pi\pi})^2 + 2\kappa_{B^0} r_{B^0}^{DK^*} r_D^{K\pi\pi\pi} \kappa_D^{K\pi\pi\pi} \cos(\delta_{B^0}^{DK^*} - \delta_D^{K\pi\pi\pi}) \cos(\gamma)}.
 \end{aligned}
 \tag{5.1}$$

The $\mathcal{R}_{\pi K(\pi\pi)}^+$ and $\mathcal{R}_{\pi K(\pi\pi)}^-$ observables can be expressed, neglecting effects of charm mixing, as

$$\mathcal{R}_{\pi K(\pi\pi)}^\pm = \frac{(r_{B^0}^{DK^*})^2 + (r_D^{K\pi(\pi\pi)})^2 + 2\kappa_{B^0} r_{B^0}^{DK^*} r_D^{K\pi(\pi\pi)} \kappa_D^{K\pi(\pi\pi)} \cos(\delta_{B^0}^{DK^*} + \delta_D^{K\pi(\pi\pi)} \pm \gamma)}{1 + (r_{B^0}^{DK^*})^2 (r_D^{K\pi(\pi\pi)})^2 + 2\kappa_{B^0} r_{B^0}^{DK^*} r_D^{K\pi(\pi\pi)} \kappa_D^{K\pi(\pi\pi)} \cos(\delta_{B^0}^{DK^*} - \delta_D^{K\pi(\pi\pi)} \pm \gamma)}.
 \tag{5.2}$$




Figure 4. Confidence levels from a frequentist analysis of the B^0 -related CP -violating observables from (left) just this analysis and (right) this analysis combined with the results from ref. [47] on the CKM angle γ .

Additional corrections are made due to the effects of charm mixing, as described in ref. [45], which are of the order of 1%. Finally, the parameters $\mathcal{A}_{K\pi}$ and $\mathcal{A}_{K\pi\pi\pi}$ can be expressed as

$$\mathcal{A}_{K\pi(\pi\pi)} = \frac{2\kappa_{B^0}\kappa_D^{K\pi(\pi\pi)}r_D^{K\pi(\pi\pi)}\sin\left(\delta_{B^0}^{DK^*} - \delta_D^{K\pi(\pi\pi)}\right)\sin(\gamma)}{1 + \left(r_{B^0}^{DK^*}\right)^2\left(r_D^{K\pi(\pi\pi)}\right)^2 + 2\kappa_{B^0}\kappa_D^{K\pi(\pi\pi)}r_D^{K\pi(\pi\pi)}\cos\left(\delta_{B^0}^{DK^*} - \delta_D^{K\pi(\pi\pi)}\right)\cos(\gamma)}. \quad (5.3)$$

With eqs. (5.1)–(5.3), the measured CP -violating observables from table 4, and the aforementioned inputs, limits are set on the $(\gamma, r_{B^0}^{DK^*}, \delta_{B^0}^{DK^*})$ parameter space using the GammaCombo package [46], which can implement both profile-likelihood and frequentist methods. The correlations of statistical and systematic uncertainties for the measured CP -violating observables are accounted for with correlation matrices shown in appendix A. Four solutions are found, and constraints on the parameter space are shown in figures 4–6. It should be noted that the probability contours shown in the two-dimensional figures do not correspond to 1σ , 2σ , 3σ , etc. when projected to one dimension. The preferred solution determines γ to be 172° with an approximate Gaussian uncertainty of 6° and $\delta_{B^0}^{DK^*}$ to be 296° with an approximate Gaussian uncertainty of 8° . However, the second-preferred solution is consistent with the world-average of direct measurements of $\gamma = \left(66.2_{-3.6}^{+3.4}\right)^\circ$ [4], finding γ to be 62° with an approximate Gaussian uncertainty of 8° and $\delta_{B^0}^{DK^*}$ to be 187° with an approximate Gaussian uncertainty of 6° . As can be seen in figure 5, all solutions give a consistent determination of $r_{B^0}^{DK^*}$, with the preferred solution of this analysis finding $r_{B^0}^{DK^*} = 0.235 \pm 0.017$.

A combined analysis of these results with the measurement of CP -violating observables from $B^0 \rightarrow DK^{*0}$ with $D \rightarrow K_S^0 h^+ h^-$ from ref. [47] is also performed. As the $D \rightarrow K_S^0 h^+ h^-$ decay probes D strong phases other than those close to 0 and π , this breaks the 90° degeneracy of the solutions found for γ in this analysis and provides additional precision. Projections of the allowed parameter space from the combined analysis are shown in figures 4–6. We estimate the central values and uncertainties of the three primary parameters of interest




Figure 5. Confidence level contours from a profile-likelihood analysis of the B^0 -related CP -violating observables from (left) just this analysis and (right) this analysis combined with the results from ref. [47] projected to the γ - $r_{B^0}^{DK^*0}$ plane. Contours contain 68.3%, 95.4%, and 99.7% of the distribution.




Figure 6. Confidence level contours from a profile-likelihood analysis of the CP -violating observables from (left) just this analysis and (right) this analysis combined with the results from ref. [47] projected to the γ - $\delta_{B^0}^{DK^*0}$ plane. Contours contain 68.3%, 95.4%, and 99.7% of the distribution.

with the PLUGIN method of GammaCombo, which implements a frequentist approach. The determinations of the PLUGIN method including all of the measured D decay modes in $B^0 \rightarrow DK^{*0}$ are $\gamma = (63.2^{+6.9}_{-8.1})^\circ$, $r_{B^0}^{DK^*} = 0.233 \pm 0.016$, and $\delta_{B^0}^{DK^*} = (192.1^{+6.7}_{-6.1})^\circ$.

In summary, measurements of CP -violating observables in $B_{(s)}^0 \rightarrow DK^{*0}$ decays are presented in this article. These are the most precise measurements to date and provide the most stringent limits to date on γ from B^0 decays. The CP -violating observables in B_s^0 decays are found to be consistent with no CP violation. A combined analysis of the CP -violating observables from B^0 decays finds a solution for γ that is consistent with the measurement from B^+ decays reported in the HFLAV average [4].

Acknowledgments

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); MOST and NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); NWO (Netherlands); MNiSW and NCN (Poland); MCID/IFA (Romania); MICINN (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); DOE NP and NSF (U.S.A.). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), and Polish WLCG (Poland). We are indebted to the communities behind the multiple open-source software packages on which we depend. Individual groups or members have received support from ARC and ARDC (Australia); Key Research Program of Frontier Sciences of CAS, CAS PIFI, CAS CCEPP, Fundamental Research Funds for the Central Universities, and Sci. & Tech. Program of Guangzhou (China); Minciencias (Colombia); EPLANET, Marie Skłodowska-Curie Actions, ERC and NextGenerationEU (European Union); A*MIDEX, ANR, IPhU and Labex P2IO, and Région Auvergne-Rhône-Alpes (France); AvH Foundation (Germany); ICSC (Italy); GVA, XuntaGal, GENCAT, Inditex, InTalent and Prog. Atracción Talento, CM (Spain); SRC (Sweden); the Leverhulme Trust, the Royal Society and UKRI (United Kingdom).

A Correlation matrices

The statistical correlation matrices of the determined CP -violating observables are shown in table 6. The systematic correlations are shown in table 7.

	$\mathcal{A}_{K\pi}$	$\mathcal{R}_{\pi K}^+$	$\mathcal{R}_{\pi K}^-$	$\mathcal{A}_{K\pi\pi\pi}$	$\mathcal{R}_{\pi K\pi\pi}^+$	$\mathcal{R}_{\pi K\pi\pi}^-$	\mathcal{A}_{CP}^{KK}	\mathcal{R}_{CP}^{KK}	$\mathcal{A}_{CP}^{\pi\pi}$	$\mathcal{R}_{CP}^{\pi\pi}$	$\mathcal{A}_{CP}^{\pi\pi\pi\pi}$	$\mathcal{R}_{CP}^{\pi\pi\pi\pi}$
$\mathcal{A}_{K\pi}$	100	9	-12	0	0	0	0	-1	0	0	0	0
$\mathcal{R}_{\pi K}^+$	100	100	6	0	0	0	0	3	0	2	0	0
$\mathcal{R}_{\pi K}^-$	100	100	100	0	0	0	0	3	0	2	0	0
$\mathcal{A}_{K\pi\pi\pi}$	100	100	100	100	7	-5	0	0	0	0	0	-1
$\mathcal{R}_{\pi K\pi\pi}^+$	100	100	100	100	100	8	0	0	0	0	0	2
$\mathcal{R}_{\pi K\pi\pi}^-$	100	100	100	100	100	100	0	0	0	0	0	1
\mathcal{A}_{CP}^{KK}	100	100	100	100	100	100	100	4	0	0	0	0
\mathcal{R}_{CP}^{KK}	100	100	100	100	100	100	100	100	0	5	0	0
$\mathcal{A}_{CP}^{\pi\pi}$	100	100	100	100	100	100	100	100	4	4	0	0
$\mathcal{R}_{CP}^{\pi\pi}$	100	100	100	100	100	100	100	100	100	100	0	0
$\mathcal{A}_{CP}^{\pi\pi\pi\pi}$	100	100	100	100	100	100	100	100	100	100	100	0
$\mathcal{R}_{CP}^{\pi\pi\pi\pi}$	100	100	100	100	100	100	100	100	100	100	100	100

Table 6. Statistical correlation matrix of B^0 observables, in percent.

	$\mathcal{A}_{K\pi}$	$\mathcal{R}_{\pi K}^+$	$\mathcal{R}_{\pi K}^-$	$\mathcal{A}_{K\pi\pi\pi}$	$\mathcal{R}_{\pi K\pi\pi}^+$	$\mathcal{R}_{\pi K\pi\pi}^-$	\mathcal{A}_{CP}^{KK}	\mathcal{R}_{CP}^{KK}	$\mathcal{A}_{CP}^{\pi\pi}$	$\mathcal{R}_{CP}^{\pi\pi}$	$\mathcal{A}_{CP}^{\pi\pi\pi\pi}$	$\mathcal{R}_{CP}^{\pi\pi\pi\pi}$
$\mathcal{A}_{K\pi}$	100	0	-5	81	15	-19	78	-3	79	-3	78	-3
$\mathcal{R}_{\pi K}^+$		100	15	14	7	6	12	5	11	3	5	-1
$\mathcal{R}_{\pi K}^-$			100	-19	7	4	-26	14	-25	4	-26	-1
$\mathcal{A}_{K\pi\pi\pi}$				100	-1	-9	77	0	77	-1	76	-30
$\mathcal{R}_{\pi K\pi\pi}^+$					100	15	18	0	17	1	19	4
$\mathcal{R}_{\pi K\pi\pi}^-$						100	-24	0	-24	1	-23	4
\mathcal{A}_{CP}^{KK}							100	-3	94	-2	93	0
\mathcal{R}_{CP}^{KK}								100	-2	15	-1	-2
$\mathcal{A}_{CP}^{\pi\pi}$									100	-3	95	0
$\mathcal{R}_{CP}^{\pi\pi}$										100	-2	0
$\mathcal{A}_{CP}^{\pi\pi\pi\pi}$											100	-3
$\mathcal{R}_{CP}^{\pi\pi\pi\pi}$												100

Table 7. Systematic correlation matrix of B^0 observables, in percent.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] N. Cabibbo, *Unitary symmetry and leptonic decays*, *Phys. Rev. Lett.* **10** (1963) 531 [[INSPIRE](#)].
- [2] M. Kobayashi and T. Maskawa, *CP-violation in the renormalizable theory of weak interaction*, *Prog. Theor. Phys.* **49** (1973) 652 [[INSPIRE](#)].
- [3] J. Brod and J. Zupan, *The ultimate theoretical error on γ from $B \rightarrow DK$ decays*, *JHEP* **01** (2014) 051 [[arXiv:1308.5663](#)] [[INSPIRE](#)].
- [4] HFLAV collaboration, *Averages of b -hadron, c -hadron, and τ -lepton properties as of 2021*, *Phys. Rev. D* **107** (2023) 052008 [[arXiv:2206.07501](#)] [[INSPIRE](#)].
- [5] LHCb collaboration, *Simultaneous determination of the CKM angle γ and parameters related to mixing and CP violation in the charm sector*, LHCb-CONF-2022-003, CERN, Geneva (2022).
- [6] UTFIT collaboration, *New UTfit analysis of the unitarity triangle in the Cabibbo-Kobayashi-Maskawa scheme*, *Rend. Lincei Sci. Fis. Nat.* **34** (2023) 37 [[arXiv:2212.03894](#)] [[INSPIRE](#)].
- [7] CKMFITTER GROUP collaboration, *CP violation and the CKM matrix: Assessing the impact of the asymmetric B factories*, *Eur. Phys. J. C* **41** (2005) 1 [[hep-ph/0406184](#)] [[INSPIRE](#)].
- [8] T. Gershon, *On the measurement of the unitarity triangle angle γ from $B^0 \rightarrow DK^{*0}$ decays*, *Phys. Rev. D* **79** (2009) 051301 [[arXiv:0810.2706](#)] [[INSPIRE](#)].
- [9] D. Atwood, I. Dunietz and A. Soni, *Improved methods for observing CP violation in $B^\pm \rightarrow KD$ and measuring the CKM phase γ* , *Phys. Rev. D* **63** (2001) 036005 [[hep-ph/0008090](#)] [[INSPIRE](#)].
- [10] M. Gronau and D. Wyler, *On determining a weak phase from CP asymmetries in charged B decays*, *Phys. Lett. B* **265** (1991) 172 [[INSPIRE](#)].
- [11] M. Gronau and D. London, *How to determine all the angles of the unitarity triangle from $B_d^0 \rightarrow DK_S^0$ and $B_s^0 \rightarrow D\phi$* , *Phys. Lett. B* **253** (1991) 483 [[INSPIRE](#)].
- [12] M. Nayak et al., *First determination of the CP content of $D \rightarrow \pi^+\pi^-\pi^0$ and $D \rightarrow K^+K^-\pi^0$* , *Phys. Lett. B* **740** (2015) 1 [[arXiv:1410.3964](#)] [[INSPIRE](#)].
- [13] LHCb collaboration, *Measurement of CP observables in the process $B^0 \rightarrow DK^{*0}$ with two- and four-body D decays*, *JHEP* **08** (2019) 041 [[arXiv:1906.08297](#)] [[INSPIRE](#)].
- [14] LHCb collaboration, *LHCb detector performance*, *Int. J. Mod. Phys. A* **30** (2015) 1530022 [[arXiv:1412.6352](#)] [[INSPIRE](#)].
- [15] LHCb collaboration, *The LHCb detector at the LHC*, *2008 JINST* **3** S08005 [[INSPIRE](#)].
- [16] R. Aaij et al., *Performance of the LHCb vertex locator*, *2014 JINST* **9** P09007 [[arXiv:1405.7808](#)] [[INSPIRE](#)].
- [17] LHCb OUTER TRACKER GROUP collaboration, *Performance of the LHCb outer tracker*, *2014 JINST* **9** P01002 [[arXiv:1311.3893](#)] [[INSPIRE](#)].
- [18] LHCb RICH GROUP collaboration, *Performance of the LHCb RICH detector at the LHC*, *Eur. Phys. J. C* **73** (2013) 2431 [[arXiv:1211.6759](#)] [[INSPIRE](#)].

- [19] R. Aaij et al., *Selection and processing of calibration samples to measure the particle identification performance of the LHCb experiment in Run 2*, *EPJ Tech. Instrum.* **6** (2019) 1 [[arXiv:1803.00824](#)] [[INSPIRE](#)].
- [20] A.A. Alves Jr. et al., *Performance of the LHCb muon system*, *2013 JINST* **8** P02022 [[arXiv:1211.1346](#)] [[INSPIRE](#)].
- [21] V.V. Gligorov and M. Williams, *Efficient, reliable and fast high-level triggering using a bonsai boosted decision tree*, *2013 JINST* **8** P02013 [[arXiv:1210.6861](#)] [[INSPIRE](#)].
- [22] T. Sjostrand, S. Mrenna and P.Z. Skands, *PYTHIA 6.4 physics and manual*, *JHEP* **05** (2006) 026 [[hep-ph/0603175](#)] [[INSPIRE](#)].
- [23] LHCb collaboration, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, *J. Phys. Conf. Ser.* **331** (2011) 032047 [[INSPIRE](#)].
- [24] D.J. Lange, *The EvtGen particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001) 152 [[INSPIRE](#)].
- [25] N. Davidson, T. Przedzinski and Z. Was, *PHOTOS interface in C++: Technical and physics documentation*, *Comput. Phys. Commun.* **199** (2016) 86 [[arXiv:1011.0937](#)] [[INSPIRE](#)].
- [26] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [[INSPIRE](#)].
- [27] J. Allison et al., *Geant4 developments and applications*, *IEEE Trans. Nucl. Sci.* **53** (2006) 270 [[INSPIRE](#)].
- [28] LHCb collaboration, *The LHCb simulation application, Gauss: Design, evolution and experience*, *J. Phys. Conf. Ser.* **331** (2011) 032023 [[INSPIRE](#)].
- [29] G.A. Cowan, D.C. Craik and M.D. Needham, *RapidSim: an application for the fast simulation of heavy-quark hadron decays*, *Comput. Phys. Commun.* **214** (2017) 239 [[arXiv:1612.07489](#)] [[INSPIRE](#)].
- [30] LHCb collaboration, *Constraints on the unitarity triangle angle γ from Dalitz plot analysis of $B^0 \rightarrow DK^+\pi^-$ decays*, *Phys. Rev. D* **93** (2016) 112018 [*Erratum ibid.* **94** (2016) 079902] [[arXiv:1602.03455](#)] [[INSPIRE](#)].
- [31] PARTICLE DATA GROUP collaboration, *Review of particle physics*, *PTEP* **2022** (2022) 083C01 [[INSPIRE](#)].
- [32] J. Therhaag, *TMVA Toolkit for multivariate data analysis in ROOT*, *PoS ICHEP2010* (2010) 510 [[INSPIRE](#)].
- [33] LHCb collaboration, *Measurement of CP observables in $B^\pm \rightarrow D^{(*)}K^\pm$ and $B^\pm \rightarrow D^{(*)}\pi^\pm$ decays using two-body D final states*, *JHEP* **04** (2021) 081 [[arXiv:2012.09903](#)] [[INSPIRE](#)].
- [34] L. Anderlini et al., *The PIDCalib package*, *LHCb-PUB-2016-021*, CERN, Geneva (2016).
- [35] LHCb collaboration, *Measurement of B^0 , B_s^0 , B^+ and Λ_b^0 production asymmetries in 7 and 8 TeV proton-proton collisions*, *Phys. Lett. B* **774** (2017) 139 [[arXiv:1703.08464](#)] [[INSPIRE](#)].
- [36] W.D. Hulsbergen, *Decay chain fitting with a Kalman filter*, *Nucl. Instrum. Meth. A* **552** (2005) 566 [[physics/0503191](#)] [[INSPIRE](#)].
- [37] LHCb collaboration, *Measurement of the CKM angle γ in $B^\pm \rightarrow DK^\pm$ and $B^\pm \rightarrow D\pi^\pm$ decays with $D \rightarrow K_S^0 h^+ h^-$* , *JHEP* **02** (2021) 169 [[arXiv:2010.08483](#)] [[INSPIRE](#)].
- [38] T. Skwarnicki, *A study of the radiative CASCADE transitions between the Upsilon-Prime and Upsilon resonances*, Ph.D. thesis, Institute of Nuclear Physics, Krakow, Poland (1986) [[INSPIRE](#)].

- [39] K.S. Cranmer, *Kernel estimation in high-energy physics*, *Comput. Phys. Commun.* **136** (2001) 198 [[hep-ex/0011057](#)] [[INSPIRE](#)].
- [40] LHCb collaboration, *Study of $B^- \rightarrow DK^- \pi^+ \pi^-$ and $B^- \rightarrow D \pi^- \pi^+ \pi^-$ decays and determination of the CKM angle γ* , *Phys. Rev. D* **92** (2015) 112005 [[arXiv:1505.07044](#)] [[INSPIRE](#)].
- [41] LHCb collaboration, *Dalitz plot analysis of $B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-$ decays*, *Phys. Rev. D* **92** (2015) 032002 [[arXiv:1505.01710](#)] [[INSPIRE](#)].
- [42] BESIII collaboration, *Measurement of the $D \rightarrow K^- \pi^+ \pi^+ \pi^-$ and $D \rightarrow K^- \pi^+ \pi^0$ coherence factors and average strong-phase differences in quantum-correlated $D\bar{D}$ decays*, *JHEP* **05** (2021) 164 [[arXiv:2103.05988](#)] [[INSPIRE](#)].
- [43] BESIII collaboration, *Measurement of the CP-even fraction of $D^0 \rightarrow \pi^+ \pi^- \pi^+ \pi^-$* , *Phys. Rev. D* **106** (2022) 092004 [[arXiv:2208.10098](#)] [[INSPIRE](#)].
- [44] S. Malde et al., *First determination of the CP content of $D \rightarrow \pi^+ \pi^- \pi^+ \pi^-$ and updated determination of the CP contents of $D \rightarrow \pi^+ \pi^- \pi^0$ and $D \rightarrow K^+ K^- \pi^0$* , *Phys. Lett. B* **747** (2015) 9 [[arXiv:1504.05878](#)] [[INSPIRE](#)].
- [45] M. Rama, *Effect of D-Dbar mixing in the extraction of gamma with $B^- \rightarrow D^0 K^-$ and $B^- \rightarrow D^0 \pi^-$* , *Phys. Rev. D* **89** (2014) 014021 [[arXiv:1307.4384](#)] [[INSPIRE](#)].
- [46] LHCb collaboration, *Measurement of the CKM angle γ from a combination of LHCb results*, *JHEP* **12** (2016) 087 [[arXiv:1611.03076](#)] [[INSPIRE](#)].
- [47] LHCb collaboration, *Measurement of the CKM angle γ in the $B^0 \rightarrow DK^{*0}$ channel using self-conjugate $D \rightarrow K_S^0 h^+ h^-$ decays*, *Eur. Phys. J. C* **84** (2024) 206 [[arXiv:2309.05514](#)] [[INSPIRE](#)].

The LHCb collaboration

R. Aaij³⁵, A.S.W. Abdelmotteleb⁵⁴, C. Abellan Beteta⁴⁸, F. Abudinén⁵⁴, T. Ackernley⁵⁸, B. Adeva⁴⁴, M. Adinolfi⁵², P. Adlarson⁷⁸, C. Agapopoulou⁴⁶, C.A. Aidala⁷⁹, Z. Ajaltouni¹¹, S. Akar⁶³, K. Akiba³⁵, P. Albicocco²⁵, J. Albrecht¹⁷, F. Alessio⁴⁶, M. Alexander⁵⁷, A. Alfonso Albero⁴³, Z. Aliouche⁶⁰, P. Alvarez Cartelle⁵³, R. Amalric¹⁵, S. Amato³, J.L. Amey⁵², Y. Amhis^{13,46}, L. An⁶, L. Anderlini²⁴, M. Andersson⁴⁸, A. Andreianov⁴¹, P. Andreola⁴⁸, M. Andreotti²³, D. Andreou⁶⁶, A. Anelli^{28,o}, D. Ao⁷, F. Archilli^{34,u}, M. Argenton²³, S. Arguedas Cuendis⁹, A. Artamonov⁴¹, M. Artuso⁶⁶, E. Aslanides¹², M. Atzeni⁶², B. Audurier¹⁴, D. Bacher⁶¹, I. Bachiller Perea¹⁰, S. Bachmann¹⁹, M. Bachmayer⁴⁷, J.J. Back⁵⁴, P. Baladron Rodriguez⁴⁴, V. Balagura¹⁴, W. Baldini²³, J. Baptista de Souza Leite², M. Barbetti^{24,l}, I. R. Barbosa⁶⁷, R.J. Barlow⁶⁰, S. Barsuk¹³, W. Barter⁵⁶, M. Bartolini⁵³, J. Bartz⁶⁶, F. Baryshnikov⁴¹, J.M. Basels¹⁶, G. Bassi^{32,r}, B. Batsukh⁵, A. Battig¹⁷, A. Bay⁴⁷, A. Beck⁵⁴, M. Becker¹⁷, F. Bedeschi³², I.B. Bediaga², A. Beiter⁶⁶, S. Belin⁴⁴, V. Bellee⁴⁸, K. Belous⁴¹, I. Belov²⁶, I. Belyaev⁴¹, G. Benane¹², G. Bencivenni²⁵, E. Ben-Haim¹⁵, A. Berezhnoy⁴¹, R. Bernet⁴⁸, S. Bernet Andres⁴², H.C. Bernstein⁶⁶, C. Bertella⁶⁰, A. Bertolin³⁰, C. Betancourt⁴⁸, F. Betti⁵⁶, J. Bex⁵³, Ia. Bezshyiko⁴⁸, J. Bhom³⁸, M.S. Bieker¹⁷, N.V. Biesuz²³, P. Billoir¹⁵, A. Biolchini³⁵, M. Birch⁵⁹, F.C.R. Bishop¹⁰, A. Bitadze⁶⁰, A. Bizzeti¹², M.P. Blago⁵³, T. Blake⁵⁴, F. Blanc⁴⁷, J.E. Blank¹⁷, S. Blusk⁶⁶, D. Bobulska⁵⁷, V. Bocharnikov⁴¹, J.A. Boelhauve¹⁷, O. Boente Garcia¹⁴, T. Boettcher⁶³, A. Bohare⁵⁶, A. Boldyrev⁴¹, C.S. Bolognani⁷⁶, R. Bolzonella^{23,k}, N. Bondar⁴¹, F. Borgato^{30,46}, S. Borghi⁶⁰, M. Borsato^{28,o}, J.T. Borsuk³⁸, S.A. Bouchiba⁴⁷, T.J.V. Bowcock⁵⁸, A. Boyer⁴⁶, C. Bozzi²³, M.J. Bradley⁵⁹, A. Brea Rodriguez⁴⁴, N. Breer¹⁷, J. Brodzicka³⁸, A. Brossa Gonzalo⁴⁴, J. Brown⁵⁸, D. Brundu²⁹, A. Buonauro⁴⁸, L. Buonincontri³⁰, A.T. Burke⁶⁰, C. Burr⁴⁶, A. Bursche⁶⁹, A. Butkevich⁴¹, J.S. Butter⁵³, J. Buytaert⁴⁶, W. Byczynski⁴⁶, S. Cadeddu²⁹, H. Cai⁷¹, R. Calabrese^{23,k}, L. Calefice¹⁷, S. Cali²⁵, M. Calvi^{28,o}, M. Calvo Gomez⁴², J. Cambon Bouzas⁴⁴, P. Campana²⁵, D.H. Campora Perez⁷⁶, A.F. Campoverde Quezada⁷, S. Capelli^{28,o}, L. Capriotti²³, R. Caravaca-Mora⁹, A. Carbone^{22,i}, L. Carcedo Salgado⁴⁴, R. Cardinale^{26,m}, A. Cardini²⁹, P. Carniti^{28,o}, L. Carus¹⁹, A. Casais Vidal⁶², R. Caspary¹⁹, G. Casse⁵⁸, J. Castro Godinez⁹, M. Cattaneo⁴⁶, G. Cavallero²³, V. Cavallini^{23,k}, S. Celani⁴⁷, J. Cerasoli¹², D. Cervenkov⁶¹, S. Cesare^{27,n}, A.J. Chadwick⁵⁸, I. Chahrour⁷⁹, M. Charles¹⁵, Ph. Charpentier⁴⁶, C.A. Chavez Barajas⁵⁸, M. Chefdeville¹⁰, C. Chen¹², S. Chen⁵, Z. Chen⁷, A. Chernov³⁸, S. Chernyshenko⁵⁰, V. Chobanova^{44,y}, S. Cholak⁴⁷, M. Chrzaszcz³⁸, A. Chubykin⁴¹, V. Chulikov⁴¹, P. Ciambone²⁵, M.F. Cicala⁵⁴, X. Cid Vidal⁴⁴, G. Ciezarek⁴⁶, P. Cifra⁴⁶, P.E.L. Clarke⁵⁶, M. Clemencic⁴⁶, H.V. Cliff⁵³, J. Closier⁴⁶, J.L. Cobbedick⁶⁰, C. Cocha Toapaxi¹⁹, V. Coco⁴⁶, J. Cogan¹², E. Cogneras¹¹, L. Cojocariu⁴⁰, P. Collins⁴⁶, T. Colombo⁴⁶, A. Comerma-Montells⁴³, L. Congedo²¹, A. Contu²⁹, N. Cooke⁵⁷, I. Corredoira⁴⁴, A. Correia¹⁵, G. Corti⁴⁶, J.J. Cottee Meldrum⁵², B. Couturier⁴⁶, D.C. Craik⁴⁸, M. Cruz Torres^{2,g}, E. Curras Rivera⁴⁷, R. Currie⁵⁶, C.L. Da Silva⁶⁵, S. Dadabaev⁴¹, L. Dai⁶⁸, X. Dai⁶, E. Dall’Occo¹⁷, J. Dalseno⁴⁴, C. D’Ambrosio⁴⁶, J. Daniel¹¹, A. Danilina⁴¹, P. d’Argent²¹, A. Davidson⁵⁴, J.E. Davies⁶⁰, A. Davis⁶⁰, O. De Aguiar Francisco⁶⁰, C. De Angelis^{29,j},

J. de Boer [ID](#)³⁵, K. De Bruyn [ID](#)⁷⁵, S. De Capua [ID](#)⁶⁰, M. De Cian [ID](#)^{19,46},
U. De Freitas Carneiro Da Graca [ID](#)^{2,b}, E. De Lucia [ID](#)²⁵, J.M. De Miranda [ID](#)², L. De Paula [ID](#)³,
M. De Serio [ID](#)^{21,h}, D. De Simone [ID](#)⁴⁸, P. De Simone [ID](#)²⁵, F. De Vellis [ID](#)¹⁷, J.A. de Vries [ID](#)⁷⁶,
F. Debernardis [ID](#)^{21,h}, D. Decamp [ID](#)¹⁰, V. Dedu [ID](#)¹², L. Del Buono [ID](#)¹⁵, B. Delaney [ID](#)⁶²,
H.-P. Dembinski [ID](#)¹⁷, J. Deng [ID](#)⁸, V. Denysenko [ID](#)⁴⁸, O. Deschamps [ID](#)¹¹, F. Dettori [ID](#)^{29,j}, B. Dey [ID](#)⁷⁴,
P. Di Nezza [ID](#)²⁵, I. Diachkov [ID](#)⁴¹, S. Didenko [ID](#)⁴¹, S. Ding [ID](#)⁶⁶, V. Dobishuk [ID](#)⁵⁰, A. D. Docheva [ID](#)⁵⁷,
A. Dolmatov [ID](#)⁴¹, C. Dong [ID](#)⁴, A.M. Donohoe [ID](#)²⁰, F. Dordei [ID](#)²⁹, A.C. dos Reis [ID](#)², L. Douglas [ID](#)⁵⁷,
A.G. Downes [ID](#)¹⁰, W. Duan [ID](#)⁶⁹, P. Duda [ID](#)⁷⁷, M.W. Dudek [ID](#)³⁸, L. Dufour [ID](#)⁴⁶, V. Duk [ID](#)³¹,
P. Durante [ID](#)⁴⁶, M. M. Duras [ID](#)⁷⁷, J.M. Durham [ID](#)⁶⁵, A. Dziurda [ID](#)³⁸, A. Dzyuba [ID](#)⁴¹, S. Easo [ID](#)^{55,46},
E. Eckstein [ID](#)⁷³, U. Egede [ID](#)¹, A. Egorychev [ID](#)⁴¹, V. Egorychev [ID](#)⁴¹, C. Eirea Orro [ID](#)⁴⁴, S. Eisenhardt [ID](#)⁵⁶,
E. Ejopu [ID](#)⁶⁰, S. Ek-In [ID](#)⁴⁷, L. Eklund [ID](#)⁷⁸, M. Elashri [ID](#)⁶³, J. Ellbracht [ID](#)¹⁷, S. Ely [ID](#)⁵⁹, A. Ene [ID](#)⁴⁰,
E. Epple [ID](#)⁶³, S. Escher [ID](#)¹⁶, J. Eschle [ID](#)⁴⁸, S. Esen [ID](#)¹⁹, T. Evans [ID](#)⁶⁰, F. Fabiano [ID](#)^{29,j,46},
L.N. Falcao [ID](#)², Y. Fan [ID](#)⁷, B. Fang [ID](#)^{71,13}, L. Fantini [ID](#)^{31,q}, M. Faria [ID](#)⁴⁷, K. Farmer [ID](#)⁵⁶,
D. Fazzini [ID](#)^{28,o}, L. Felkowski [ID](#)⁷⁷, M. Feng [ID](#)^{5,7}, M. Feo [ID](#)⁴⁶, M. Fernandez Gomez [ID](#)⁴⁴,
A.D. Fernez [ID](#)⁶⁴, F. Ferrari [ID](#)²², F. Ferreira Rodrigues [ID](#)³, S. Ferreres Sole [ID](#)³⁵, M. Ferrillo [ID](#)⁴⁸,
M. Ferro-Luzzi [ID](#)⁴⁶, S. Filippov [ID](#)⁴¹, R.A. Fini [ID](#)²¹, M. Fiorini [ID](#)^{23,k}, K.M. Fischer [ID](#)⁶¹,
D.S. Fitzgerald [ID](#)⁷⁹, C. Fitzpatrick [ID](#)⁶⁰, F. Fleuret [ID](#)¹⁴, M. Fontana [ID](#)²², L. F. Foreman [ID](#)⁶⁰,
R. Forty [ID](#)⁴⁶, D. Foulds-Holt [ID](#)⁵³, M. Franco Sevilla [ID](#)⁶⁴, M. Frank [ID](#)⁴⁶, E. Franzoso [ID](#)^{23,k},
G. Frau [ID](#)¹⁹, C. Frei [ID](#)⁴⁶, D.A. Friday [ID](#)⁶⁰, L. Frontini [ID](#)^{27,n}, J. Fu [ID](#)⁷, Q. Fuehring [ID](#)¹⁷, Y. Fujii [ID](#)¹,
T. Fulghesu [ID](#)¹⁵, E. Gabriel [ID](#)³⁵, G. Galati [ID](#)^{21,h}, M.D. Galati [ID](#)³⁵, A. Gallas Torreira [ID](#)⁴⁴,
D. Galli [ID](#)^{22,i}, S. Gambetta [ID](#)⁵⁶, M. Gandelman [ID](#)³, P. Gandini [ID](#)²⁷, H. Gao [ID](#)⁷, R. Gao [ID](#)⁶¹,
Y. Gao [ID](#)⁸, Y. Gao [ID](#)⁶, Y. Gao [ID](#)⁸, M. Garau [ID](#)^{29,j}, L.M. Garcia Martin [ID](#)⁴⁷, P. Garcia Moreno [ID](#)⁴³,
J. García Pardiñas [ID](#)⁴⁶, B. Garcia Plana [ID](#)⁴⁴, K. G. Garg [ID](#)⁸, L. Garrido [ID](#)⁴³, C. Gaspar [ID](#)⁴⁶,
R.E. Geertsema [ID](#)³⁵, L.L. Gerken [ID](#)¹⁷, E. Gersabeck [ID](#)⁶⁰, M. Gersabeck [ID](#)⁶⁰, T. Gershon [ID](#)⁵⁴,
Z. Ghorbanimoghaddam [ID](#)⁵², L. Giambastiani [ID](#)³⁰, F. I. Giasemis [ID](#)^{15,e}, V. Gibson [ID](#)⁵³,
H.K. Giemza [ID](#)³⁹, A.L. Gilman [ID](#)⁶¹, M. Giovannetti [ID](#)²⁵, A. Gioventù [ID](#)⁴³, P. Gironella Gironell [ID](#)⁴³,
C. Giugliano [ID](#)^{23,k}, M.A. Giza [ID](#)³⁸, E.L. Gkougkousis [ID](#)⁵⁹, F.C. Glaser [ID](#)^{13,19}, V.V. Gligorov [ID](#)¹⁵,
C. Göbel [ID](#)⁶⁷, E. Golobardes [ID](#)⁴², D. Golubkov [ID](#)⁴¹, A. Golutvin [ID](#)^{59,41,46}, A. Gomes [ID](#)^{2,a,†},
S. Gomez Fernandez [ID](#)⁴³, F. Goncalves Abrantes [ID](#)⁶¹, M. Goncerz [ID](#)³⁸, G. Gong [ID](#)⁴, J.
A. Gooding [ID](#)¹⁷, I.V. Gorelov [ID](#)⁴¹, C. Gotti [ID](#)²⁸, J.P. Grabowski [ID](#)⁷³, L.A. Granado Cardoso [ID](#)⁴⁶,
E. Graugés [ID](#)⁴³, E. Graverini [ID](#)^{47,s}, L. Grazette [ID](#)⁵⁴, G. Graziani [ID](#), A. T. Grecu [ID](#)⁴⁰,
L.M. Greeven [ID](#)³⁵, N.A. Grieser [ID](#)⁶³, L. Grillo [ID](#)⁵⁷, S. Gromov [ID](#)⁴¹, C. Gu [ID](#)¹⁴, M. Guarise [ID](#)²³,
M. Guittiere [ID](#)¹³, V. Guliaeva [ID](#)⁴¹, P. A. Günther [ID](#)¹⁹, A.-K. Guseinov [ID](#)⁴¹, E. Gushchin [ID](#)⁴¹,
Y. Guz [ID](#)^{6,41,46}, T. Gys [ID](#)⁴⁶, T. Hadavizadeh [ID](#)¹, C. Hadjivasiliou [ID](#)⁶⁴, G. Haefeli [ID](#)⁴⁷, C. Haen [ID](#)⁴⁶,
J. Haimberger [ID](#)⁴⁶, M. Hajheidari [ID](#)⁴⁶, M.M. Halvorsen [ID](#)⁴⁶, P.M. Hamilton [ID](#)⁶⁴, J. Hammerich [ID](#)⁵⁸,
Q. Han [ID](#)⁸, X. Han [ID](#)¹⁹, S. Hansmann-Menzemer [ID](#)¹⁹, L. Hao [ID](#)⁷, N. Harnew [ID](#)⁶¹, T. Harrison [ID](#)⁵⁸,
M. Hartmann [ID](#)¹³, J. He [ID](#)^{7,c}, K. Heijhoff [ID](#)³⁵, F. Hemmer [ID](#)⁴⁶, C. Henderson [ID](#)⁶³,
R.D.L. Henderson [ID](#)^{1,54}, A.M. Hennequin [ID](#)⁴⁶, K. Hennessy [ID](#)⁵⁸, L. Henry [ID](#)⁴⁷, J. Herd [ID](#)⁵⁹,
P. Herrero Gascon [ID](#)¹⁹, J. Heuel [ID](#)¹⁶, A. Hicheur [ID](#)³, G. Hijano Mendizabal [ID](#)⁴⁸, D. Hill [ID](#)⁴⁷,
S.E. Hollitt [ID](#)¹⁷, J. Horswill [ID](#)⁶⁰, R. Hou [ID](#)⁸, Y. Hou [ID](#)¹⁰, N. Howarth [ID](#)⁵⁸, J. Hu [ID](#)¹⁹, J. Hu [ID](#)⁶⁹,
W. Hu [ID](#)⁶, X. Hu [ID](#)⁴, W. Huang [ID](#)⁷, W. Hulsbergen [ID](#)³⁵, R.J. Hunter [ID](#)⁵⁴, M. Hushchyn [ID](#)⁴¹,
D. Hutchcroft [ID](#)⁵⁸, D. Ilin [ID](#)⁴¹, P. Ilten [ID](#)⁶³, A. Inglessi [ID](#)⁴¹, A. Iniukhin [ID](#)⁴¹, A. Ishteev [ID](#)⁴¹,
K. Ivshin [ID](#)⁴¹, R. Jacobsson [ID](#)⁴⁶, H. Jage [ID](#)¹⁶, S.J. Jaimes Elles [ID](#)^{45,72}, S. Jakobsen [ID](#)⁴⁶, E. Jans [ID](#)³⁵,

B.K. Jashal [ID](#)⁴⁵, A. Jawahery [ID](#)^{64,46}, V. Jevtic [ID](#)¹⁷, E. Jiang [ID](#)⁶⁴, X. Jiang [ID](#)^{5,7}, Y. Jiang [ID](#)⁷, Y. J. Jiang [ID](#)⁶, M. John [ID](#)⁶¹, D. Johnson [ID](#)⁵¹, C.R. Jones [ID](#)⁵³, T.P. Jones [ID](#)⁵⁴, S. Joshi [ID](#)³⁹, B. Jost [ID](#)⁴⁶, N. Jurik [ID](#)⁴⁶, I. Juszczak [ID](#)³⁸, D. Kaminaris [ID](#)⁴⁷, S. Kandybei [ID](#)⁴⁹, Y. Kang [ID](#)⁴, M. Karacson [ID](#)⁴⁶, D. Karpenkov [ID](#)⁴¹, M. Karpov [ID](#)⁴¹, A. M. Kauniskangas [ID](#)⁴⁷, J.W. Kautz [ID](#)⁶³, F. Keizer [ID](#)⁴⁶, D.M. Keller [ID](#)⁶⁶, M. Kenzie [ID](#)⁵³, T. Ketel [ID](#)³⁵, B. Khanji [ID](#)⁶⁶, A. Kharisova [ID](#)⁴¹, S. Kholodenko [ID](#)³², G. Khreich [ID](#)¹³, T. Kirn [ID](#)¹⁶, V.S. Kirsebom [ID](#)⁴⁷, O. Kitouni [ID](#)⁶², S. Klaver [ID](#)³⁶, N. Kleijne [ID](#)^{32,r}, K. Klimaszewski [ID](#)³⁹, M.R. Kmiec [ID](#)³⁹, S. Koliiev [ID](#)⁵⁰, L. Kolk [ID](#)¹⁷, A. Konoplyannikov [ID](#)⁴¹, P. Kopciwicz [ID](#)^{37,46}, P. Koppenburg [ID](#)³⁵, M. Korolev [ID](#)⁴¹, I. Kostiuik [ID](#)³⁵, O. Kot [ID](#)⁵⁰, S. Kotriakhova [ID](#), A. Kozachuk [ID](#)⁴¹, P. Kravchenko [ID](#)⁴¹, L. Kravchuk [ID](#)⁴¹, M. Kreps [ID](#)⁵⁴, S. Kretzschmar [ID](#)¹⁶, P. Krokovny [ID](#)⁴¹, W. Krupa [ID](#)⁶⁶, W. Krzemien [ID](#)³⁹, J. Kubat [ID](#)¹⁹, S. Kubis [ID](#)⁷⁷, W. Kucewicz [ID](#)³⁸, M. Kucharczyk [ID](#)³⁸, V. Kudryavtsev [ID](#)⁴¹, E. Kulikova [ID](#)⁴¹, A. Kupsc [ID](#)⁷⁸, B. K. Kutsenko [ID](#)¹², D. Lacarrere [ID](#)⁴⁶, A. Lai [ID](#)²⁹, A. Lampis [ID](#)²⁹, D. Lancierini [ID](#)⁴⁸, C. Landesa Gomez [ID](#)⁴⁴, J.J. Lane [ID](#)¹, R. Lane [ID](#)⁵², C. Langenbruch [ID](#)¹⁹, J. Langer [ID](#)¹⁷, O. Lantwin [ID](#)⁴¹, T. Latham [ID](#)⁵⁴, F. Lazzari [ID](#)^{32,s}, C. Lazzeroni [ID](#)⁵¹, R. Le Gac [ID](#)¹², S.H. Lee [ID](#)⁷⁹, R. Lefèvre [ID](#)¹¹, A. Leflat [ID](#)⁴¹, S. Legotin [ID](#)⁴¹, M. Lehurax [ID](#)⁵⁴, O. Leroy [ID](#)¹², T. Lesiak [ID](#)³⁸, B. Leverington [ID](#)¹⁹, A. Li [ID](#)⁴, H. Li [ID](#)⁶⁹, K. Li [ID](#)⁸, L. Li [ID](#)⁶⁰, P. Li [ID](#)⁴⁶, P.-R. Li [ID](#)⁷⁰, S. Li [ID](#)⁸, T. Li [ID](#)^{5,d}, T. Li [ID](#)⁶⁹, Y. Li [ID](#)⁸, Y. Li [ID](#)⁵, Z. Li [ID](#)⁶⁶, Z. Lian [ID](#)⁴, X. Liang [ID](#)⁶⁶, C. Lin [ID](#)⁷, T. Lin [ID](#)⁵⁵, R. Lindner [ID](#)⁴⁶, V. Lisovskyi [ID](#)⁴⁷, R. Litvinov [ID](#)^{29,j}, F. L. Liu [ID](#)¹, G. Liu [ID](#)⁶⁹, K. Liu [ID](#)⁷⁰, Q. Liu [ID](#)⁷, S. Liu [ID](#)^{5,7}, Y. Liu [ID](#)⁵⁶, Y. Liu [ID](#)⁷⁰, Y. L. Liu [ID](#)⁵⁹, A. Lobo Salvia [ID](#)⁴³, A. Loi [ID](#)²⁹, J. Lomba Castro [ID](#)⁴⁴, T. Long [ID](#)⁵³, J.H. Lopes [ID](#)³, A. Lopez Huertas [ID](#)⁴³, S. López Soliño [ID](#)⁴⁴, G.H. Lovell [ID](#)⁵³, C. Lucarelli [ID](#)^{24,l}, D. Lucchesi [ID](#)^{30,p}, S. Luchuk [ID](#)⁴¹, M. Lucio Martinez [ID](#)⁷⁶, V. Lukashenko [ID](#)^{35,50}, Y. Luo [ID](#)⁶, A. Lupato [ID](#)³⁰, E. Luppi [ID](#)^{23,k}, K. Lynch [ID](#)²⁰, X.-R. Lyu [ID](#)⁷, G. M. Ma [ID](#)⁴, R. Ma [ID](#)⁷, S. Maccolini [ID](#)¹⁷, F. Machefer [ID](#)¹³, F. Maciuc [ID](#)⁴⁰, B. M. Mack [ID](#)⁶⁶, I. Mackay [ID](#)⁶¹, L. M. Mackey [ID](#)⁶⁶, L.R. Madhan Mohan [ID](#)⁵³, M. M. Madurai [ID](#)⁵¹, A. Maevskiy [ID](#)⁴¹, D. Magdalinski [ID](#)³⁵, D. Maisuzenko [ID](#)⁴¹, M.W. Majewski [ID](#)³⁷, J.J. Malczewski [ID](#)³⁸, S. Malde [ID](#)⁶¹, B. Malecki [ID](#)^{38,46}, L. Malentacca [ID](#)⁴⁶, A. Malinin [ID](#)⁴¹, T. Maltsev [ID](#)⁴¹, G. Manca [ID](#)^{29,j}, G. Mancinelli [ID](#)¹², C. Mancuso [ID](#)^{27,13,n}, R. Manera Escalero [ID](#)⁴³, D. Manuzzi [ID](#)²², D. Marangotto [ID](#)^{27,n}, J.F. Marchand [ID](#)¹⁰, R. Marchevski [ID](#)⁴⁷, U. Marconi [ID](#)²², S. Mariani [ID](#)⁴⁶, C. Marin Benito [ID](#)⁴³, J. Marks [ID](#)¹⁹, A.M. Marshall [ID](#)⁵², P.J. Marshall [ID](#)⁵⁸, G. Martelli [ID](#)^{31,q}, G. Martellotti [ID](#)³³, L. Martinazzoli [ID](#)⁴⁶, M. Martinelli [ID](#)^{28,o}, D. Martinez Santos [ID](#)⁴⁴, F. Martinez Vidal [ID](#)⁴⁵, A. Massafferri [ID](#)², M. Materok [ID](#)¹⁶, R. Matev [ID](#)⁴⁶, A. Mathad [ID](#)⁴⁸, V. Matiunin [ID](#)⁴¹, C. Matteuzzi [ID](#)⁶⁶, K.R. Mattioli [ID](#)¹⁴, A. Mauri [ID](#)⁵⁹, E. Maurice [ID](#)¹⁴, J. Mauricio [ID](#)⁴³, P. Mayencourt [ID](#)⁴⁷, M. Mazurek [ID](#)⁴⁶, M. McCann [ID](#)⁵⁹, L. McConnell [ID](#)²⁰, T.H. McGrath [ID](#)⁶⁰, N.T. McHugh [ID](#)⁵⁷, A. McNab [ID](#)⁶⁰, R. McNulty [ID](#)²⁰, B. Meadows [ID](#)⁶³, G. Meier [ID](#)¹⁷, D. Melnychuk [ID](#)³⁹, M. Merk [ID](#)^{35,76}, A. Merli [ID](#)^{27,n}, L. Meyer Garcia [ID](#)³, D. Miao [ID](#)^{5,7}, H. Miao [ID](#)⁷, M. Mikhasenko [ID](#)^{73,f}, D.A. Milanes [ID](#)⁷², A. Minotti [ID](#)^{28,o}, E. Minucci [ID](#)⁶⁶, T. Miralles [ID](#)¹¹, S.E. Mitchell [ID](#)⁵⁶, B. Mitreska [ID](#)¹⁷, D.S. Mitzel [ID](#)¹⁷, A. Modak [ID](#)⁵⁵, A. Mödden [ID](#)¹⁷, R.A. Mohammed [ID](#)⁶¹, R.D. Moise [ID](#)¹⁶, S. Mokhnenko [ID](#)⁴¹, T. Mombächer [ID](#)⁴⁶, M. Monk [ID](#)^{54,1}, I.A. Monroy [ID](#)⁷², S. Monteil [ID](#)¹¹, A. Morcillo Gomez [ID](#)⁴⁴, G. Morello [ID](#)²⁵, M.J. Morello [ID](#)^{32,r}, M.P. Morgenthaler [ID](#)¹⁹, A.B. Morris [ID](#)⁴⁶, A.G. Morris [ID](#)¹², R. Mountain [ID](#)⁶⁶, H. Mu [ID](#)⁴, Z. M. Mu [ID](#)⁶, E. Muhammad [ID](#)⁵⁴, F. Muheim [ID](#)⁵⁶, M. Mulder [ID](#)⁷⁵, K. Müller [ID](#)⁴⁸, F. Muñoz-Rojas [ID](#)⁹, R. Murta [ID](#)⁵⁹, P. Naik [ID](#)⁵⁸, T. Nakada [ID](#)⁴⁷, R. Nandakumar [ID](#)⁵⁵, T. Nanut [ID](#)⁴⁶, I. Nasteva [ID](#)³, M. Needham [ID](#)⁵⁶, N. Neri [ID](#)^{27,n}, S. Neubert [ID](#)⁷³, N. Neufeld [ID](#)⁴⁶, P. Neustroev [ID](#)⁴¹, R. Newcombe [ID](#)⁵⁹,

J. Nicolini [ID](#)^{17,13}, D. Nicotra [ID](#)⁷⁶, E.M. Niel [ID](#)⁴⁷, N. Nikitin [ID](#)⁴¹, P. Nogga⁷³, N.S. Nolte [ID](#)⁶²,
 C. Normand [ID](#)^{10,29}, J. Novoa Fernandez [ID](#)⁴⁴, G. Nowak [ID](#)⁶³, C. Nunez [ID](#)⁷⁹, H. N. Nur [ID](#)⁵⁷,
 A. Oblakowska-Mucha [ID](#)³⁷, V. Obraztsov [ID](#)⁴¹, T. Oeser [ID](#)¹⁶, S. Okamura [ID](#)^{23,k,46}, R. Oldeman [ID](#)^{29,j},
 F. Oliva [ID](#)⁵⁶, M. Olocco [ID](#)¹⁷, C.J.G. Onderwater [ID](#)⁷⁶, R.H. O’Neil [ID](#)⁵⁶, J.M. Otalora Goicochea [ID](#)³,
 T. Ovsiannikova [ID](#)⁴¹, P. Owen [ID](#)⁴⁸, A. Oyanguren [ID](#)⁴⁵, O. Ozcelik [ID](#)⁵⁶, K.O. Padeken [ID](#)⁷³,
 B. Pagare [ID](#)⁵⁴, P.R. Pais [ID](#)¹⁹, T. Pajero [ID](#)⁶¹, A. Palano [ID](#)²¹, M. Palutan [ID](#)²⁵, G. Panshin [ID](#)⁴¹,
 L. Paolucci [ID](#)⁵⁴, A. Papanestis [ID](#)⁵⁵, M. Pappagallo [ID](#)^{21,h}, L.L. Pappalardo [ID](#)^{23,k},
 C. Pappenheimer [ID](#)⁶³, C. Parkes [ID](#)⁶⁰, B. Passalacqua [ID](#)^{23,k}, G. Passaleva [ID](#)²⁴, D. Passaro [ID](#)^{32,r},
 A. Pastore [ID](#)²¹, M. Patel [ID](#)⁵⁹, J. Patoc [ID](#)⁶¹, C. Patrignani [ID](#)^{22,i}, C.J. Pawley [ID](#)⁷⁶, A. Pellegrino [ID](#)³⁵,
 M. Pepe Altarelli [ID](#)²⁵, S. Perazzini [ID](#)²², D. Pereima [ID](#)⁴¹, A. Pereiro Castro [ID](#)⁴⁴, P. Perret [ID](#)¹¹,
 A. Perro [ID](#)⁴⁶, K. Petridis [ID](#)⁵², A. Petrolini [ID](#)^{26,m}, S. Petrucci [ID](#)⁵⁶, J. P. Pfaller [ID](#)⁶³, H. Pham [ID](#)⁶⁶,
 L. Pica [ID](#)^{32,r}, M. Piccini [ID](#)³¹, B. Pietrzyk [ID](#)¹⁰, G. Pietrzyk [ID](#)¹³, D. Pinci [ID](#)³³, F. Pisani [ID](#)⁴⁶,
 M. Pizzichemi [ID](#)^{28,o}, V. Placinta [ID](#)⁴⁰, M. Plo Casasus [ID](#)⁴⁴, F. Polci [ID](#)^{15,46}, M. Poli Lener [ID](#)²⁵,
 A. Poluektov [ID](#)¹², N. Polukhina [ID](#)⁴¹, I. Polyakov [ID](#)⁴⁶, E. Polycarpo [ID](#)³, S. Ponce [ID](#)⁴⁶, D. Popov [ID](#)⁷,
 S. Poslavskii [ID](#)⁴¹, K. Prasanth [ID](#)³⁸, C. Prouve [ID](#)⁴⁴, V. Pugatch [ID](#)⁵⁰, G. Punzi [ID](#)^{32,s}, W. Qian [ID](#)⁷,
 N. Qin [ID](#)⁴, S. Qu [ID](#)⁴, R. Quagliani [ID](#)⁴⁷, R.I. Rabadan Trejo [ID](#)⁵⁴, J.H. Rademacker [ID](#)⁵², M. Rama [ID](#)³²,
 M. Ramírez García [ID](#)⁷⁹, M. Ramos Pernas [ID](#)⁵⁴, M.S. Rangel [ID](#)³, F. Ratnikov [ID](#)⁴¹, G. Raven [ID](#)³⁶,
 M. Rebollo De Miguel [ID](#)⁴⁵, F. Redi [ID](#)⁴⁶, J. Reich [ID](#)⁵², F. Reiss [ID](#)⁶⁰, Z. Ren [ID](#)⁷, P.K. Resmi [ID](#)⁶¹,
 R. Ribatti [ID](#)^{32,r}, G. R. Ricart [ID](#)^{14,80}, D. Riccardi [ID](#)^{32,r}, S. Ricciardi [ID](#)⁵⁵, K. Richardson [ID](#)⁶²,
 M. Richardson-Slipper [ID](#)⁵⁶, K. Rinnert [ID](#)⁵⁸, P. Robbe [ID](#)¹³, G. Robertson [ID](#)⁵⁷, E. Rodrigues [ID](#)^{58,46},
 E. Rodriguez Fernandez [ID](#)⁴⁴, J.A. Rodriguez Lopez [ID](#)⁷², E. Rodriguez Rodriguez [ID](#)⁴⁴,
 A. Rogovskiy [ID](#)⁵⁵, D.L. Rolf [ID](#)⁴⁶, A. Rollings [ID](#)⁶¹, P. Roloff [ID](#)⁴⁶, V. Romanovskiy [ID](#)⁴¹,
 M. Romero Lamas [ID](#)⁴⁴, A. Romero Vidal [ID](#)⁴⁴, G. Romolini [ID](#)²³, F. Ronchetti [ID](#)⁴⁷, M. Rotondo [ID](#)²⁵, S.
 R. Roy [ID](#)¹⁹, M.S. Rudolph [ID](#)⁶⁶, T. Ruf [ID](#)⁴⁶, M. Ruiz Diaz [ID](#)¹⁹, R.A. Ruiz Fernandez [ID](#)⁴⁴,
 J. Ruiz Vidal [ID](#)^{78,z}, A. Ryzhikov [ID](#)⁴¹, J. Ryzka [ID](#)³⁷, J.J. Saborido Silva [ID](#)⁴⁴, R. Sadek [ID](#)¹⁴,
 N. Sagidova [ID](#)⁴¹, N. Sahoo [ID](#)⁵¹, B. Saitta [ID](#)^{29,j}, M. Salomoni [ID](#)^{28,o}, C. Sanchez Gras [ID](#)³⁵,
 I. Sanderswood [ID](#)⁴⁵, R. Santacesaria [ID](#)³³, C. Santamarina Rios [ID](#)⁴⁴, M. Santimaria [ID](#)²⁵,
 L. Santoro [ID](#)², E. Santovetti [ID](#)³⁴, A. Saputi [ID](#)^{23,46}, D. Saranin [ID](#)⁴¹, G. Sarpis [ID](#)⁵⁶, M. Sarpis [ID](#)⁷³,
 A. Sarti [ID](#)³³, C. Satriano [ID](#)^{33,t}, A. Satta [ID](#)³⁴, M. Saur [ID](#)⁶, D. Savrina [ID](#)⁴¹, H. Sazak [ID](#)¹¹,
 L.G. Scantlebury Smead [ID](#)⁶¹, A. Scarabotto [ID](#)¹⁵, S. Schael [ID](#)¹⁶, S. Scherl [ID](#)⁵⁸, A. M. Schertz [ID](#)⁷⁴,
 M. Schiller [ID](#)⁵⁷, H. Schindler [ID](#)⁴⁶, M. Schmelling [ID](#)¹⁸, B. Schmidt [ID](#)⁴⁶, S. Schmitt [ID](#)¹⁶, H. Schmitz⁷³,
 O. Schneider [ID](#)⁴⁷, A. Schopper [ID](#)⁴⁶, N. Schulte [ID](#)¹⁷, S. Schulte [ID](#)⁴⁷, M.H. Schune [ID](#)¹³,
 R. Schwemmer [ID](#)⁴⁶, G. Schwering [ID](#)¹⁶, B. Sciascia [ID](#)²⁵, A. Sciuccati [ID](#)⁴⁶, S. Sellam [ID](#)⁴⁴,
 A. Semennikov [ID](#)⁴¹, M. Senghi Soares [ID](#)³⁶, A. Sergi [ID](#)^{26,m}, N. Serra [ID](#)^{48,46}, L. Sestini [ID](#)³⁰,
 A. Seuthe [ID](#)¹⁷, Y. Shang [ID](#)⁶, D.M. Shangase [ID](#)⁷⁹, M. Shapkin [ID](#)⁴¹, R. S. Sharma [ID](#)⁶⁶,
 I. Shchemerov [ID](#)⁴¹, L. Shchutska [ID](#)⁴⁷, T. Shears [ID](#)⁵⁸, L. Shekhtman [ID](#)⁴¹, Z. Shen [ID](#)⁶, S. Sheng [ID](#)^{5,7},
 V. Shevchenko [ID](#)⁴¹, B. Shi [ID](#)⁷, E.B. Shields [ID](#)^{28,o}, Y. Shimizu [ID](#)¹³, E. Shmanin [ID](#)⁴¹, R. Shorkin [ID](#)⁴¹,
 J.D. Shupperd [ID](#)⁶⁶, R. Silva Coutinho [ID](#)⁶⁶, G. Simi [ID](#)³⁰, S. Simone [ID](#)^{21,h}, N. Skidmore [ID](#)⁶⁰,
 R. Skuza [ID](#)¹⁹, T. Skwarnicki [ID](#)⁶⁶, M.W. Slater [ID](#)⁵¹, J.C. Smallwood [ID](#)⁶¹, E. Smith [ID](#)⁶², K. Smith [ID](#)⁶⁵,
 M. Smith [ID](#)⁵⁹, A. Snoch [ID](#)³⁵, L. Soares Lavra [ID](#)⁵⁶, M.D. Sokoloff [ID](#)⁶³, F.J.P. Soler [ID](#)⁵⁷,
 A. Solomin [ID](#)^{41,52}, A. Solovev [ID](#)⁴¹, I. Solovyev [ID](#)⁴¹, R. Song [ID](#)¹, Y. Song [ID](#)⁴⁷, Y. Song [ID](#)⁴, Y. S.
 Song [ID](#)⁶, F.L. Souza De Almeida [ID](#)⁶⁶, B. Souza De Paula [ID](#)³, E. Spadaro Norella [ID](#)^{27,n},
 E. Spedicato [ID](#)²², J.G. Speer [ID](#)¹⁷, E. Spiridenkov⁴¹, P. Spradlin [ID](#)⁵⁷, V. Sriskaran [ID](#)⁴⁶, F. Stagni [ID](#)⁴⁶,

M. Stahl⁴⁶, S. Stahl⁴⁶, S. Stanislaus⁶¹, E.N. Stein⁴⁶, O. Steinkamp⁴⁸, O. Stenyakin⁴¹, H. Stevens¹⁷, D. Strelalina⁴¹, Y. Su⁷, F. Suljik⁶¹, J. Sun²⁹, L. Sun⁷¹, Y. Sun⁶⁴, P.N. Swallow⁵¹, F. Swystun⁵⁴, A. Szabelski³⁹, T. Szumlak³⁷, M. Szymanski⁴⁶, Y. Tan⁴, S. Taneja⁶⁰, M.D. Tat⁶¹, A. Terentev⁴⁸, F. Terzuoli^{32,v}, F. Teubert⁴⁶, E. Thomas⁴⁶, D.J.D. Thompson⁵¹, H. Tilquin⁵⁹, V. Tisserand¹¹, S. T'Jampens¹⁰, M. Tobin⁵, L. Tomassetti^{23,k}, G. Tonani^{27,n,46}, X. Tong⁶, D. Torres Machado², L. Toscano¹⁷, D.Y. Tou⁴, C. Trippel⁴², G. Tuci¹⁹, N. Tuning³⁵, L.H. Uecker¹⁹, A. Ukleja³⁷, D.J. Unverzagt¹⁹, E. Ursov⁴¹, A. Usachov³⁶, A. Ustyuzhanin⁴¹, U. Uwer¹⁹, V. Vagnoni²², A. Valassi⁴⁶, G. Valenti²², N. Valls Canudas⁴², H. Van Hecke⁶⁵, E. van Herwijnen⁵⁹, C.B. Van Hulse^{44,x}, R. Van Laak⁴⁷, M. van Veghel³⁵, R. Vazquez Gomez⁴³, P. Vazquez Regueiro⁴⁴, C. Vázquez Sierra⁴⁴, S. Vecchi²³, J.J. Velthuis⁵², M. Veltri^{24,w}, A. Venkateswaran⁴⁷, M. Vesterinen⁵⁴, M. Vieites Diaz⁴⁶, X. Vilasis-Cardona⁴², E. Vilella Figueras⁵⁸, A. Villa²², P. Vincent¹⁵, F.C. Volle¹³, D. vom Bruch¹², V. Vorobyev⁴¹, N. Voropaev⁴¹, K. Vos⁷⁶, G. Vouters¹⁰, C. Vrahas⁵⁶, J. Walsh³², E.J. Walton¹, G. Wan⁶, C. Wang¹⁹, G. Wang⁸, J. Wang⁶, J. Wang⁵, J. Wang⁴, J. Wang⁷¹, M. Wang²⁷, N. W. Wang⁷, R. Wang⁵², X. Wang⁶⁹, X. W. Wang⁵⁹, Y. Wang⁸, Z. Wang¹³, Z. Wang⁴, Z. Wang⁷, J.A. Ward^{54,1}, M. Waterlaet⁴⁶, N.K. Watson⁵¹, D. Websdale⁵⁹, Y. Wei⁶, B.D.C. Westhenry⁵², D.J. White⁶⁰, M. Whitehead⁵⁷, A.R. Wiederhold⁵⁴, D. Wiedner¹⁷, G. Wilkinson⁶¹, M.K. Wilkinson⁶³, M. Williams⁶², M.R.J. Williams⁵⁶, R. Williams⁵³, F.F. Wilson⁵⁵, W. Wislicki³⁹, M. Witek³⁸, L. Witola¹⁹, C.P. Wong⁶⁵, G. Wormser¹³, S.A. Wotton⁵³, H. Wu⁶⁶, J. Wu⁸, Y. Wu⁶, K. Wyllie⁴⁶, S. Xian⁶⁹, Z. Xiang⁵, Y. Xie⁸, A. Xu³², J. Xu⁷, L. Xu⁴, L. Xu⁴, M. Xu⁵⁴, Z. Xu¹¹, Z. Xu⁷, Z. Xu⁵, D. Yang⁴, S. Yang⁷, X. Yang⁶, Y. Yang^{26,m}, Z. Yang⁶, Z. Yang⁶⁴, V. Yeroshenko¹³, H. Yeung⁶⁰, H. Yin⁸, C. Y. Yu⁶, J. Yu⁶⁸, X. Yuan⁵, E. Zaffaroni⁴⁷, M. Zavertyaev¹⁸, M. Zdybal³⁸, M. Zeng⁴, C. Zhang⁶, D. Zhang⁸, J. Zhang⁷, L. Zhang⁴, S. Zhang⁶⁸, S. Zhang⁶, Y. Zhang⁶, Y. Z. Zhang⁴, Y. Zhao¹⁹, A. Zharkova⁴¹, A. Zhelezov¹⁹, X. Z. Zheng⁴, Y. Zheng⁷, T. Zhou⁶, X. Zhou⁸, Y. Zhou⁷, V. Zhovkovska⁵⁴, L. Z. Zhu⁷, X. Zhu⁴, X. Zhu⁸, V. Zhukov^{16,41}, J. Zhuo⁴⁵, Q. Zou^{5,7}, D. Zuliani³⁰, G. Zunica⁶⁰

¹ School of Physics and Astronomy, Monash University, Melbourne, Australia

² Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

³ Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

⁴ Center for High Energy Physics, Tsinghua University, Beijing, China

⁵ Institute Of High Energy Physics (IHEP), Beijing, China

⁶ School of Physics State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

⁷ University of Chinese Academy of Sciences, Beijing, China

⁸ Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China

⁹ Consejo Nacional de Rectores (CONARE), San Jose, Costa Rica

¹⁰ Université Savoie Mont Blanc, CNRS, IN2P3-LAPP, Annecy, France

¹¹ Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France

¹² Aix Marseille Univ, CNRS/IN2P3, CPPM, Marseille, France

¹³ Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France

¹⁴ Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

- ¹⁵ *LPNHE, Sorbonne Université, Paris Diderot Sorbonne Paris Cité, CNRS/IN2P3, Paris, France*
- ¹⁶ *I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany*
- ¹⁷ *Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany*
- ¹⁸ *Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany*
- ¹⁹ *Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*
- ²⁰ *School of Physics, University College Dublin, Dublin, Ireland*
- ²¹ *INFN Sezione di Bari, Bari, Italy*
- ²² *INFN Sezione di Bologna, Bologna, Italy*
- ²³ *INFN Sezione di Ferrara, Ferrara, Italy*
- ²⁴ *INFN Sezione di Firenze, Firenze, Italy*
- ²⁵ *INFN Laboratori Nazionali di Frascati, Frascati, Italy*
- ²⁶ *INFN Sezione di Genova, Genova, Italy*
- ²⁷ *INFN Sezione di Milano, Milano, Italy*
- ²⁸ *INFN Sezione di Milano-Bicocca, Milano, Italy*
- ²⁹ *INFN Sezione di Cagliari, Monserrato, Italy*
- ³⁰ *Università degli Studi di Padova, Università e INFN, Padova, Padova, Italy*
- ³¹ *INFN Sezione di Perugia, Perugia, Italy*
- ³² *INFN Sezione di Pisa, Pisa, Italy*
- ³³ *INFN Sezione di Roma La Sapienza, Roma, Italy*
- ³⁴ *INFN Sezione di Roma Tor Vergata, Roma, Italy*
- ³⁵ *Nikhef National Institute for Subatomic Physics, Amsterdam, Netherlands*
- ³⁶ *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, Netherlands*
- ³⁷ *AGH — University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ³⁸ *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ³⁹ *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ⁴⁰ *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ⁴¹ *Affiliated with an institute covered by a cooperation agreement with CERN*
- ⁴² *DS4DS, La Salle, Universitat Ramon Llull, Barcelona, Spain*
- ⁴³ *ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ⁴⁴ *Instituto Galego de Física de Altas Enerxías (IGFAE), Universidade de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴⁵ *Instituto de Física Corpuscular, Centro Mixto Universidad de Valencia - CSIC, Valencia, Spain*
- ⁴⁶ *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁴⁷ *Institute of Physics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁴⁸ *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁴⁹ *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁵⁰ *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁵¹ *University of Birmingham, Birmingham, U.K.*
- ⁵² *H.H. Wills Physics Laboratory, University of Bristol, Bristol, U.K.*
- ⁵³ *Cavendish Laboratory, University of Cambridge, Cambridge, U.K.*
- ⁵⁴ *Department of Physics, University of Warwick, Coventry, U.K.*
- ⁵⁵ *STFC Rutherford Appleton Laboratory, Didcot, U.K.*
- ⁵⁶ *School of Physics and Astronomy, University of Edinburgh, Edinburgh, U.K.*
- ⁵⁷ *School of Physics and Astronomy, University of Glasgow, Glasgow, U.K.*
- ⁵⁸ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, U.K.*
- ⁵⁹ *Imperial College London, London, U.K.*
- ⁶⁰ *Department of Physics and Astronomy, University of Manchester, Manchester, U.K.*
- ⁶¹ *Department of Physics, University of Oxford, Oxford, U.K.*
- ⁶² *Massachusetts Institute of Technology, Cambridge, MA, U.S.A.*
- ⁶³ *University of Cincinnati, Cincinnati, OH, U.S.A.*
- ⁶⁴ *University of Maryland, College Park, MD, U.S.A.*
- ⁶⁵ *Los Alamos National Laboratory (LANL), Los Alamos, NM, U.S.A.*

- ⁶⁶ *Syracuse University, Syracuse, NY, U.S.A.*
- ⁶⁷ *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ³*
- ⁶⁸ *School of Physics and Electronics, Hunan University, Changsha City, China, associated to ⁸*
- ⁶⁹ *Guangdong Provincial Key Laboratory of Nuclear Science, Guangdong-Hong Kong Joint Laboratory of Quantum Matter, Institute of Quantum Matter, South China Normal University, Guangzhou, China, associated to ⁴*
- ⁷⁰ *Lanzhou University, Lanzhou, China, associated to ⁵*
- ⁷¹ *School of Physics and Technology, Wuhan University, Wuhan, China, associated to ⁴*
- ⁷² *Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to ¹⁵*
- ⁷³ *Universität Bonn — Helmholtz-Institut für Strahlen und Kernphysik, Bonn, Germany, associated to ¹⁹*
- ⁷⁴ *Eotvos Lorand University, Budapest, Hungary, associated to ⁴⁶*
- ⁷⁵ *Van Swinderen Institute, University of Groningen, Groningen, Netherlands, associated to ³⁵*
- ⁷⁶ *Universiteit Maastricht, Maastricht, Netherlands, associated to ³⁵*
- ⁷⁷ *Tadeusz Kosciuszko Cracow University of Technology, Cracow, Poland, associated to ³⁸*
- ⁷⁸ *Department of Physics and Astronomy, Uppsala University, Uppsala, Sweden, associated to ⁵⁷*
- ⁷⁹ *University of Michigan, Ann Arbor, MI, U.S.A., associated to ⁶⁶*
- ⁸⁰ *Departement de Physique Nucleaire (SPhN), Gif-Sur-Yvette, France*

^a *Universidade de Brasília, Brasília, Brazil*

^b *Centro Federal de Educação Tecnológica Celso Suckow da Fonseca, Rio De Janeiro, Brazil*

^c *Hangzhou Institute for Advanced Study, UCAS, Hangzhou, China*

^d *School of Physics and Electronics, Henan University, Kaifeng, China*

^e *LIP6, Sorbonne Universite, Paris, France*

^f *Excellence Cluster ORIGINS, Munich, Germany*

^g *Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras*

^h *Università di Bari, Bari, Italy*

ⁱ *Università di Bologna, Bologna, Italy*

^j *Università di Cagliari, Cagliari, Italy*

^k *Università di Ferrara, Ferrara, Italy*

^l *Università di Firenze, Firenze, Italy*

^m *Università di Genova, Genova, Italy*

ⁿ *Università degli Studi di Milano, Milano, Italy*

^o *Università di Milano Bicocca, Milano, Italy*

^p *Università di Padova, Padova, Italy*

^q *Università di Perugia, Perugia, Italy*

^r *Scuola Normale Superiore, Pisa, Italy*

^s *Università di Pisa, Pisa, Italy*

^t *Università della Basilicata, Potenza, Italy*

^u *Università di Roma Tor Vergata, Roma, Italy*

^v *Università di Siena, Siena, Italy*

^w *Università di Urbino, Urbino, Italy*

^x *Universidad de Alcalá, Alcalá de Henares, Spain*

^y *Universidade da Coruña, Coruña, Spain*

^z *Department of Physics/Division of Particle Physics, Lund, Sweden*

[†] *Deceased*