

Differential gene expression in the *Longissimus dorsi* of Nguni and Bonsmara bulls finished on low and high energy diets

D.A. Linde^{1,2}, E. van Marle-Köster², M.M. Scholtz^{1,3}, M.G. Gonda⁴, J.L. Gonzalez-Hernandez^{5,6} & M. D. MacNeil^{1,3,7#}

¹ Agricultural Research Council, Animal Production Institute, Irene, South Africa

² Department of Animal and Wildlife Science, University of Pretoria, Pretoria, South Africa

³ Department of Animal, Wildlife and Grassland Sciences, University of the Free State, Bloemfontein, South Africa

⁴ Department of Animal Science, South Dakota State University, Brookings, SD, USA

⁵ Department of Agronomy, Horticulture, and Plant Science, South Dakota State University, Brookings, SD, USA

⁶ Research and Economic Division, South Dakota State University, Brookings, SD, USA

⁷ Delta G, Miles City, Montana, USA

(Submitted 24 September 2022; Accepted 9 January 2023; Published 11 April 2023)

Copyright resides with the authors in terms of the Creative Commons Attribution 4.0 South African Licence.

See: <http://creativecommons.org/licenses/by/4.0/za>

Condition of use: The user may copy, distribute, transmit and adapt the work, but must recognise the authors and the South African J. Anim. Sci.

Abstract

Objectives of this research were to examine differential gene expression profiles of Nguni and Bonsmara cattle fed diets differing in their energy density. The ultimate goal was to improve understanding of the mechanisms that underlie differences between these breeds and the potential interactions of the differences between breeds with the nutritive environment. The experiment was designed as a 2 × 2 factorial arrangement of breed and diet (12.5 MJ/kg DM vs. 10.9 MJ/kg DM). The initial feeding trial had 10 bull calves per treatment. However, financial constraints limited RNA sequencing to six animals per treatment and the RNA generated from one animal was of insufficient quality to be useful. Transcripts with false discovery rate *P*-values <0.05 and fold-changes >2.0 were considered significant. Bonsmara had a faster growth rate, heavier live and carcass weights, and better feed conversion compared to Nguni. However, lower levels of fat were observed in Nguni. Twenty different genes were differentially expressed, with three exhibiting interaction effects and all 20 having differences in transcript abundance between the breeds. A dietary effect was only observed for the one gene and that gene was also subject to an interaction effect with breed. Observed differences in gene expression between Bonsmara and Nguni by several genes affecting the structure or function of the mitochondria imply differences in energy metabolism between the breeds. Interaction effects on the abundance of some gene transcripts indicate the need to consider the diet when evaluating breed differences and conversely, consider breed when evaluating diets.

Keywords: hypothalamus–anterior pituitary axis, mitochondria, RNA sequence

Corresponding author: macneil.deltag@gmail.com

Introduction

The goal of a differential gene expression study is to determine which genes are expressed at different levels between the experimentally imposed conditions. These genes can offer biological insight into the processes affected by the condition(s) of interest. Identification of differentially expressed genes between specific conditions is a key in the understanding phenotypic variation (Costa-Silva *et al.*, 2017; McDermaid *et al.*, 2019). Examples of the successful use of this approach include providing biological insight into 1) parental genome effects on bovine foetal development between taurine and indicine breeds; and 2) differences in skeletal muscle differentiation between Wagyu and Piedmontese (Lehnert *et al.*, 2007). Higgins *et al.* (2019) used differential gene expression to investigate the hepatic transcriptome of Charolais and Holstein–Friesian steers divergent in their residual feed intake when fed concentrate and forage-based diets. Thus, the use of differential gene expression methodology was implemented to study differences between Nguni and Bonsmara cattle when fed diets that differed in their energy density.

Nguni cattle are a tropically adapted and admixed Sanga-type breed that are indigenous to southern Africa (Scholtz *et al.*, 2011a; Makina *et al.*, 2016). These cattle are also widely used in

URL: <http://www.sasas.co.za>

ISSN 0375-1589 (print), ISSN 2221-4062 (online)

Publisher: South African Society for Animal Science

<http://dx.doi.org/10.4314/sajas.v53i1.11>

communal and small holder farming systems on natural grazing where they contribute to the food security of resource-poor farmers in South Africa (Bester *et al.*, 2003; Mapiye *et al.*, 2009; Morokong, 2016). Females of this small-framed breed are often used in crossbreeding systems due to their low maintenance requirements, high fertility, and good mothering ability (Scholtz *et al.*, 2011b). However, because Nguni cattle have a propensity to deposit excess fat at less-than-optimal carcass weights when fed high-energy diets, they are not preferred for finishing under feedlot systems (Scholtz *et al.*, 2008; Strydom, 2008).

Bonsmara, while also considered to be a Sanga breed, were more recently developed using Milking Shorthorn, Hereford, and Afrikaner cattle to produce a locally-adapted, composite beef breed (Bonsma, 1980). Its current genetic composition is approximately 40% Afrikaner, 33% Shorthorn, 19% African zebu, and 5% Hereford (Makina *et al.*, 2016). In comparison to Nguni, Bonsmara cattle grow more rapidly and are heavier at all ages (Schoeman, 1996; Bergh *et al.*, 2010).

The majority of beef in South Africa is produced in feedlots using commercially-formulated, high energy diets, with preference given to medium- and large-framed and later maturing cattle (Scholtz *et al.*, 2008; Strydom, 2008). Due to the Nguni's early maturity and lower meat yield, it is not preferred as a feedlot animal (Scholtz *et al.*, 2008). At age-constant endpoints, du Plessis and Hoffman (2007) found Nguni steers were clearly fatter than Bonsmara-cross steers, even when finished on natural pastures. Nguni and Bonsmara cattle have been shown to produce similarly high quality meat (Strydom *et al.*, 2000; Muchenje *et al.*, 2008). One time-honoured strategy for feeding early maturing breeds to a heavier weight without depositing excess fat is to provide them a less energy-dense diet for a longer period of time (Owens *et al.*, 1995).

The objective of this research was to examine differences in the gene expression profiles in the longissimus thoracis et lumborum of Nguni and Bonsmara cattle fed diets of differing energy density. The ultimate goal of this research was to obtain an improved understanding of the molecular mechanisms that underlie the differences between these breeds and potential interactions with their nutritive environment. At this time, this is the only study of differential gene expression in livestock that are indigenous to South Africa other than a study of Boar and village goats under intensive and extensive management (Ncube *et al.*, 2022).

Materials & Methods

Ethical approval was received from both the Animal Ethical Committee of the Agricultural Research Council (ARC) and the AEC of the University of Pretoria (ec090-15).

Forty randomly-selected bull calves consisting of 20 Nguni and 20 Bonsmara from 10 to 14 months old were transported to the bull testing station of the ARC (Irene, South Africa). The bull-calves originated from two research farms (Vaalharts (27°57'19" S; 24°50'41" E) and Roodeplaat (25°34'11.27" S; 28°22'05.36" E). A number of bulls were used in both herds. However, parentage was not recorded and some of the calves may therefore be related.

At Irene, the calves were subjected to a 28-d adaptation period, followed by an 84-d feeding trial. The trial was designed as a 2 × 2 factorial arrangement of breed and feed level (high vs low energy diets). For each breed–diet treatment, there were 10 animals. The low energy and the high energy diets are shown in Table 1. The bulls were individually fed using Calan gates (American Calan, USA) and water was freely available.

Table 1 Composition of a low-energy and high-energy diet to be fed to Nguni and Bonsmara bulls prior to slaughter and the nutritional composition of the diets

Ingredients	Low energy diet (% DM)	High energy diet (% DM)
Hominy chop	13.0	6.0
Maize meal	17.0	49.5
Brewer's grain	20.0	12.0
Cottonseed oil cake meal	4.0	4.0
Molasses	9.0	8.0
Lucerne (alfalfa) hay	6.0	7.5
Wheat/maize straw	28.0	10.0
Mineral and Vitamin mix	3.0	3.0
Predicted Nutritional composition		
Metabolizable energy, MJ/kg DM	10.9	12.5
Crude protein, g/kg DM	14.7	14.3
Non-degradable protein, g/kg DM	50.6	49.0
Non-protein nitrogen, %	17.5	17.5
Acid detergent fibre, g/kg DM	17.6	11.4
Fat, g/kg DM	34.0	38.0
Calcium, g/kg DM	7.9	6.1
Phosphorus, g/kg DM	4.2	3.4
Sulphur, g/kg DM	1.9	1.9

DM: dry matter

The calves were weighed weekly and images of the longissimus muscle and rump were collected at the beginning and end of the feeding period with real-time ultrasound (RTU) (MyLabOne, Esaote, USA). Traits measured on the RTU images were longissimus muscle area (EMA), subcutaneous fat depths over the rib (RF) and rump (P8), and intramuscular fat content of the longissimus (IMF). After being fed 102 days, the animals were slaughtered at the local abattoir and the carcasses were weighed. The abattoir was 3 km from where the animals were fed and they were transported the morning of slaughter.

Data from the feeding trial were analysed using the PROC MIXED of SAS v 9.4 (SAS Institute, Inc., Cary, North Carolina, USA) as a 2 × 2 × 2 factorial of farm of origin, breed, and diet. All effects were tested for significance against the residual variance. Interaction effects that included farm of origin were generally not significant ($P > 0.05$) and were not considered further.

At slaughter, after each calf was stunned, hoisted and bled, a cut was made in its dorsal right side at the 13th rib. An 8 mm Miltex® biopsy punch (Integra LifeSciences, Princeton, New Jersey, USA) was used to sample the longissimus muscle. Forceps were used to place the muscle sample into a cryotube, which was then flash frozen in liquid nitrogen and the samples were stored in a -80 °C freezer until RNA extraction. Due to funding limitations, samples from 24 animals (six per treatment) were selected for RNA extraction. Three replicate tissue samples (120 mg each) from each animal were processed. TRIzol® Reagent (1 ml per sample) (Ambion, USA) was used to extract the RNA followed by homogenization. The sample was placed in a GenoGrinder 2010 (SPEX sample prep, USA) for 15–20 min at 1730 rpm. Chloroform was added to separate the DNA and the RNA. The homogenized and separated tissue was subsequently precipitated with isopropanol. In the final step, the precipitated pellet was washed with 1 ml 70% ethanol. Each sample was treated with the RNase-free DNase set (Qiagen, Hilden, Germany) to decrease the risk of genomic DNA contamination, and it was purified with the RNeasy mini kit according to the manufacturer's guidelines (Qiagen, Hilden, Germany). Extracted RNA from the samples was stored in a -80 °C freezer. The extracted RNA was quantified using a Qubit Fluorescent meter (Invitrogen, USA) according to manufacturer's protocols and with a 1% gel using ethidium bromide and Tris-borate-ethylenediaminetetraacetic acid buffer. Following quality control, the RNA from one Bonsmara calf that was fed the high energy diet was discarded. Samples from the remaining 23 animals had RNA with concentrations between 221 µg/ml and 570 µg/ml and were of sufficient quality for sequencing.

Sample preparation was done using the TruSeq stranded mRNA protocol (Illumina, USA) with 3–5 µg RNA for each sample. Sequencing was performed using a HiSeq 2500 (Illumina, USA) for generation of pair-ended 2 × 125 bp reads. Of the 69 sequences, 13 were removed due to poor quality

reads. Thus, there were 15, 3, and 5 animals with 3, 2, and 1 technical replicates, respectively. Technical replicate sequences were combined into a single file for each animal. Sequence reads were imported into CLC Genomics Workbench, version 22.0 (Qiagen, Aarhus, Denmark) for bioinformatics analysis. Reads were trimmed using Q = 20 as a quality threshold and adapters were removed prior to genome mapping. Reads were mapped to the ARS-UCD1.2 bovine genome assembly (Rosen *et al.*, 2020) using a length fraction of 0.8 and minimum similarity fraction of 0.8. Mapped read counts were normalized with the trimmed mean of M samples (TMM) method (Robinson and Oshlack, 2010). Pairwise comparisons between 1) breeds, 2) diets, and 3) diet-by-treatment interactions were completed with a generalized linear model assuming a negative binomial probability distribution of read counts (McCarthy *et al.*, 2012), with treatment as the independent variable. Transcripts with Benjamini-Hochberg False Discovery Rate (FDR) *P*-values <0.05 (Benjamini and Hochberg, 1995) and fold-changes >2.0 were considered statistically significant.

Results and Discussion

Average weights of the Nguni and Bonsmara bulls at the start of the trial were 193 ± 4.29 kg and 206.49 ± 4.53 kg, respectively. Live weight at slaughter, ADG, FCR, P8, and EMA differed markedly between the breeds (Table 2). Diet had an effect on IMF and carcass weight ($P < 0.05$). A breed by diet interaction ($P < 0.05$) also affected IMF. Farm of origin had no effect on the various traits ($P > 0.05$). Bonsmara had a faster growth rate, heavier live and carcass weights, and better feed conversion than Nguni cattle. However, lower levels of fat were observed in Nguni. Previous research has indicated that Nguni will be slower growing and weigh less at slaughter than Bonsmara, but the breeds will be similar in feed conversion and carcass fat content (Strydom *et al.*, 2001). Thus, the samples of the breeds used in the current study may be considered more or less consistent with their previous characterizations.

Table 2 Phenotypic characterization of Nguni and Bonsmara bulls fed diets differing in energy density

Traits	Bonsmara		Nguni	
	12.5 MJ/kg DM diet	10.9 MJ/kg DM diet	12.5 MJ/kg DM diet	10.9 MJ/kg DM diet
Live weight, kg	354.9 ± 5.86	330.6 ± 5.65	301.1 ± 5.86	306.6 ± 5.65
Average daily gain, kg/d	1.65 ± 0.04	1.50 ± 0.04	1.34 ± 0.04	1.30 ± 0.04
Feed conversion ratio	6.88 ± 0.21	7.32 ± 0.21	8.36 ± 0.21	8.30 ± 0.21
P8 fat depth, mm	5.3 ± 0.08	5.6 ± 0.07	5.3 ± 0.08	5.2 ± 0.07
Rib fat depth, mm	3.25 ± 0.13	3.29 ± 0.13	2.87 ± 0.13	2.89 ± 0.13
Longissimus area, cm ²	57.0 ± 1.07	57.5 ± 1.02	54.3 ± 1.07	54.7 ± 1.02
Intramuscular fat	3.27 ± 0.19	3.34 ± 0.17	3.40 ± 0.17	2.56 ± 0.17
Carcass weight, kg	187.70 ± 2.43	168.39 ± 2.34	159.11 ± 2.43	153.35 ± 2.34

Samples from the animals produced a median of 99.5 million RNA transcript reads in pairs. On average, 93.6% of the reads mapped to the ARS-UCD1.2 bovine genome assembly (Rosen *et al.*, 2020). Broken read pairs (those whose two reads didn't map to the same gene) were discarded, resulting in an average of 78.5% of reads subsequently being used. Of these counted reads, an average of 73.4% were mapped onto known genes (62.3% on exons and 11.3% on introns), while 26.6% mapped onto intergenic regions of the reference genome, suggesting the possibility of additional genes beyond those that are currently annotated.

Twenty genes were differentially expressed between Nguni and Bonsmara breeds, with 12 of these being more abundant in Nguni (Table 3). Differences in transcript abundance may be correlated with phenotypic differences in tissue growth and function. However, transcriptome differences should not be over-interpreted. Differences in transcript abundance do not always result in different protein abundance nor do they necessarily cause changes in phenotype. In this work, the transcripts were mapped to the genome and differences in gene expression were then quantified at the gene level. These gene transcripts corresponded to functions associated with ATP production and mitochondrial function, growth, appetite stimulation, and ribosome and cytoskeletal structure (Clark *et al.*, 2016).

Table 3 Differentially abundant transcripts between skeletal muscle tissue from Nguni and Bonsmara calves

Gene Abbreviation	Gene Name	Fold-Change ¹	FDR <i>P</i> -value
<i>RPL15</i> ²	Ribosomal Protein L15	59.85	5.5 x 10 ⁻⁹
<i>ENSBTAG00000031205</i>	Unidentified	38.97	3.7 x 10 ⁻⁷
<i>ENSBTAG00000019018</i>	Unidentified	86.66	1.3 x 10 ⁻⁵
<i>CHRND</i>	Cholinergic Receptor Nicotinic Delta Subunit	-4.64	7.5 x 10 ⁻⁴
<i>GHRH</i>	Growth Hormone Releasing Hormone	-156.86	7.9 x 10 ⁻⁴
<i>ENSBTAG00000019672</i>	Unidentified	-8.48	8.3 x 10 ⁻⁴
<i>MCHR1</i>	Melanin Concentrating Hormone Receptor 1	-3.42	1.1 x 10 ⁻³
<i>ENSBTAG00000055302</i>	Unidentified	5.01	2.4 x 10 ⁻³
<i>ERV PABL B-1</i> ²	Endogenous Retrovirus Group PABL B Member 1, Envelope	-170.24	7.8 x 10 ⁻³
<i>ZWINT</i> ²	ZW10 Interacting Kinetochore Protein	-5.85	7.8 x 10 ⁻³
<i>CYCS</i> ²	Cytochrome C	-185.49	7.8 x 10 ⁻³
<i>ENSBTAG00000049442</i>	Unidentified	-18.10	1.6 x 10 ⁻²
<i>PWWP3A</i>	PWWP Domain Containing 3A, DNA Repair Factor	-2.10	1.6 x 10 ⁻²
<i>RAB21_2</i>	RAB21, Member RAS Oncogene Family	24.66	1.9 x 10 ⁻²
<i>MRM2</i>	Mitochondrial rRNA Methyltransferase 2	-2.25	2.0 x 10 ⁻²
<i>C19H17orf98</i>	Uncharacterized Protein Chromosome 19	14.39	2.6 x 10 ⁻²
<i>DENND2A</i>	DENN Domain Containing 2A	-2.00	3.9 x 10 ⁻²
<i>SLC25A6</i> ²	Solute Carrier Family 25 Member 6	3.00	4.9 x 10 ⁻²
<i>UQCRC2</i> ²	Ubiquinol-Cytochrome C Reductase Core Protein 2	-7.61	4.9 x 10 ⁻²
<i>PECAM1</i>	Platelet and Endothelial Cell Adhesion Molecule 1	1.96	4.9 x 10 ⁻²

FDR: False discovery rate

¹ Positive values indicate greater abundance in Bonsmara and negative values indicate greater abundance in Nguni

² Identified by searching *Homo sapiens* and *Mus musculus* gene lists using Panther v17 (pantherdb.org)

Three gene transcripts were differentially abundant in the contrast that measured the breed by diet interaction. These transcripts illustrate that there may be instances when the joint consideration of both breed and diet are needed for the proper interpretation of treatment effects. The ubiquinol-cytochrome c reductase core protein 2 (*Uqcrc2*; fold-change = -19.1; FDR $P = 6.0 \times 10^{-7}$) is located in the mitochondrion, where it is part of the ubiquinol-cytochrome c reductase complex. This complex constitutes a part of the mitochondrial respiratory chain responsible for electron transport (Guo *et al.*, 2017). Previously, Kelly *et al.* (2011) failed to detect differences in the expression level of *Uqcrc2* in the longissimus dorsi muscle between groups of Limousin x Friesian heifers fed high and low forage diets. Expression of transcripts of *ENSBTAG00000031205* (fold-change = 31.5; FDR $P = 2.5 \times 10^{-5}$) that encode 40S ribosomal protein S4 were likewise affected by the breed by diet interaction. Wærp *et al.* (2019) previously observed transcripts of this gene to be differentially abundant in adipose tissue from dairy heifers at 12 months of age versus when they were seven months pregnant when the heifers were fed a high energy diet but not when a lower energy diet was fed. The *ENSBTAG00000049315* (fold-change = -178.5; FDR $P = 6.6 \times 10^{-3}$) was the final transcript affected by the breed by diet interaction. These interaction effects provide further evidence for the necessity of considering the diet when evaluating breed differences and conversely, the breed when evaluating different diets (da Costa *et al.*, 2013).

Only one uncharacterized gene transcript (*ENSBTAG00000031205*; fold-change = 19.8; FDR $P = 4.0 \times 10^{-3}$) was observed more frequently in animals that were fed the low energy diet. A plausible explanation for this was that the differences between the diets were not sufficient to lead to changes in gene expression. As discussed above, the number of transcripts of this gene was also affected by the breed by diet interaction, indicating the effect of diet on its level of expression to be breed-specific.

Four genes associated with mitochondria or the electron transport chain were differentially expressed between Nguni and Bonsmara. Previous reports have documented relationships of feed efficiency with mitochondrial function in cattle (Kolath *et al.*, 2006; Alam *et al.*, 2012). The cytochrome c (*Cycc*) gene encodes a protein that is directly involved in the electron transport system, which produces ATP (Dayhoff, 1972). Similarly, the ubiquinol-cytochrome c reductase core protein 2 (*Uqcrc2*) encodes a protein that is part of complex III of the electron transport chain (Guo *et al.*, 2017). Defects in electron transport also may be responsible for greater oxidized protein levels in less feed-efficient

phenotypes (Bottje *et al.*, 2006; Bottje and Carstens, 2009). The mitochondrial ribosomal RNA methyltransferase 2 (*Mrm2*) gene encodes a protein responsible for methylation of uridine in mitochondrial 21S rRNA (Pintard *et al.*, 2002); deletion of this gene in yeast leads to a defect in cell respiration (Garone *et al.*, 2017). The solute carrier family 25 member 6 (*Slc25a6*) gene encodes an ADP/ATP carrier that transports ADP/ATP across the mitochondrial membrane (Cl  men  on *et al.*, 2013). Three of these four gene transcripts were more highly abundant in Nguni skeletal muscle, the exception being *Slc25a6*. Higher abundance of gene transcripts involved in mitochondria and oxidative phosphorylation suggest that ATP (and energy) production might be higher in muscle of the less feed-efficient Nguni at time of slaughter. Higher ATP production might result from increased expression of oxidative phosphorylation genes, increased number of mitochondria, or both. Bottje and Kong (2013) found it logical that key genes associated with mitochondrial function, activities, or development would be differentially expressed between broilers that differed in feed efficiency.

Intriguingly, six genes associated with cytoskeleton structure or muscle fibres were differentially expressed between Nguni and Bonsmara skeletal muscle. Four of these gene transcripts (*Dennd2a*, *Pwwp3a*, *Zwint*, and *Chrmd*) were more highly abundant in Nguni. In chickens and turkeys, low feed efficiency has been associated with decreased abundance of transcripts associated with cytoskeleton structure and muscle fibres (Bottje and Kong, 2013; Pezeshkian *et al.*, 2022). The Nguni calves in this study had higher FCR (Table 2), suggesting that the greater expression of these genes in Nguni may contribute to their lower feed efficiency.

Two transcripts normally associated with the hypothalamus–anterior pituitary axis were more abundant in skeletal muscle tissue from Nguni calves. The growth hormone releasing hormone (*Ghrh*) gene encodes a peptide secreted from the hypothalamus that acts on the anterior pituitary gland to stimulate release of growth hormone (Grings *et al.*, 1988; Alba and Salvatori, 2004). Melanin concentrating hormone receptor-1 (*Mchr-1*) gene encodes a receptor that binds melanin-concentrating hormone, which regulates appetite (Ingvarstsen and Boisclair, 2001; Lord *et al.*, 2021). Increased abundance of these transcripts in Nguni skeletal muscle may suggest increased feed intake at this age compared to Bonsmara calves. However, the action of both genes in skeletal muscle tissue has not been characterized. Further, increased transcript abundance of *Ghrh* and *Mchr-1* may not lead to phenotypic differences in feed intake and growth rate.

Conclusions

Despite uncovering relatively few differentially expressed genes, fundamental differences in gene expression that underpin the metabolic processes of Bonsmara and Nguni were observed. In particular, several significant effects were noted for genes affecting the structure or function of the mitochondria. The presence of interaction effects on the abundance of some gene transcripts indicates the importance of considering the diet when evaluating breed differences and conversely, considering breed when evaluating diets.

Acknowledgements

The authors gratefully acknowledge Leon de Lange and the staff at the Bull Testing Centre of the Animal Production Campus of the ARC for the care and handling of the animals. This research was supported by Red Meat Research and Development South Africa and the National Research Foundation (UID 135438). The Northern Cape Department of Agriculture, Land Reform, and Rural Development provided animals for this study. The support of all entities is greatly appreciated.

Author' contributions

This manuscript represents work conducted by DAL as part of her MSc program of study. EvMK and MMS supervised DAL. DAL and MDM analysed the phenotypic data. MGG and JLGH completed the bioinformatics analysis of transcriptome data. DAL wrote the first draft of the manuscript, which was subsequently revised with contributions from all authors who have also read and approved the final version of the manuscript.

Conflicts of interest

The authors affirm that they have no conflicts of interest which would affect the research reported herein in any way.

References

- Alam, T., D. A. Keeny, T. Sweeney, F. Buckley, R. Prendville, M. McGee, and S. M. Waters. 2012. Expression of genes involved in energy homeostasis in the duodenum and liver of Holstein–Friesian and Jersey cows and their F1 hybrid. *Physiol. Genomics* 44:198–209. <https://doi.org/10.1152/physiolgenomics.00102.2011>
- Alba, M. & Salvatori, R., 2004. A mouse with targeted ablation of the growth hormone-releasing hormone gene: A new model of isolated growth hormone deficiency. *Endocrinology* 145(9), 4134-43. DOI: 10.1210/en.2004-0119

- Benjamini Y. & Hochberg, Y., 1995. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J R Stat Soc Series B.* 57(1), 289-300. <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>
- Bergh, L., Gerhard, R., Scholtz, M.M. & Mamabolo, M.J., 2010 Introduction to the information on beef and dual purpose breeds in South Africa. In: Scholtz, M.M. (editor), *Beef Breeding in South Africa*. Agricultural Research Council. Pretoria, South Africa.
- Bester, J., Matjuda, L.E., Rust, J.M. & Fourie, H.J., 2003. The Nguni: A Case Study. In: *Proc. Community-based Management of Animal Genetic Resources*. Mbabane, Swaziland, 7–11 May 2001. FAO. Rome, Italy. Pages 45-68. https://amatava.com/nguni_j_bester.htm
- Bonsma, J.C., 1980. Cross-breeding, breed creation and the genesis of the Bonsmara. In: *Livestock Production: A Global Approach*, J. C. Bonsma (ed.), Tafelberg Publishers Ltd., Cape Town South Africa. Pages 126–136.
- Bottje, W. G., and G. E. Carstens. 2009. Association of mitochondrial function and feed efficiency in poultry and livestock species. *J. Anim. Sci.* 87(E. Suppl.), E48–E63. DOI: 10.2527/jas.2008-1379
- Bottje, W. & Kong, B.-W., 2013. Feed efficiency: Mitochondrial function to global gene expression. *J. Anim. Sci.* 91, 1582–1593. DOI: 10.2527/jas2012-5787
- Bottje, W., N. Pumford, C. Ojano-Dirain, M. Iqbal, and K. Lassiter. 2006. Feed efficiency and mitochondrial function. *Poult. Sci.* 85, 8–14. DOI: 10.1093/ps/85.1.8
- Clark, K., Karsch-Mizrachi, I., Lipman, D.J., Ostell, J. & Sayers, E.W., 2016. GenBank. *Nucleic Acids Res.* 44(D1), D67-D72. DOI: 10.1093/nar/gkv1276
- Cléménçon, B., Babot, M. & Trézéguet, V., 2013. The mitochondrial ADP/ATP carrier (SLC25 family): Pathological implications of its dysfunction. *Mol. Aspects Med.* 34(2-3), 485-93. DOI: 10.1016/j.mam.2012.05.006
- Costa-Silva, J., Domingues, D. & Lopes, F.M., 2017. RNA-Seq differential expression analysis: An extended review and a software tool. *PLoS ONE* 12(12), e0190152. <https://doi.org/10.1371/journal.pone.0190152>
- Da Costa, A.S., Pires, V.M., Fontes, C.M. & Mestre Prates, J.A., 2013. Expression of genes controlling fat deposition in two genetically diverse beef cattle breeds fed high or low silage diets. *BMC Vet. Res.* 9, 118. doi: 10.1186/1746-6148-9-118
- Dayhoff, M.O. 1972. Cytochrome C group. *Atlas of Protein Sequence and Structure*, Volume 5. National Biomedical Research Foundation, Washington, DC, USA. pp. D7-D27.
- du Plessis, I. & Hoffman, L.C., 2007. Effect of slaughter age and breed on the carcass traits and meat quality of beef steers finished on natural pastures in the arid subtropics of South Africa. *S. Afr. J. Anim. Sci.* 37, 143-153. DOI: 10.4314/sajas.v37i3.4084
- Garone, C., D'Souza, A.R., Dallabona, C., Lodi, T., Rebelo-Guiomar, P., Rorbach, J., Donati, M.A., Procopio, E., Montomoli, M., Guerrini, R., Zeviani, M., Calvo, S.E., Mootha, V.K., DiMauro, S., Ferrero I. & Minczuk, M., 2017. Defective mitochondrial rRNA methyltransferase MRM2 causes MELAS-like clinical syndrome. *Hum. Mol. Genet.* 26(21), 4257-66. DOI: 10.1093/hmg/ddx314
- Grings, E.E., Scarborough, R., Schally, A.V. & Reeves, J.J., 1988. Response to a growth hormone-releasing hormone analog in heifers treated with recombinant growth hormone. *Domest. Anim. Endocrinol.* 5(1), 47-53. DOI: 10.1016/0739-7240(88)90025-2
- Guo, R., Zong, S., Wu, M., Gu, J. Yang, M., 2017. Architecture of human mitochondrial respiratory megacomplex I₂ III₂ IV₂. *Cell* 170(6), 1247-57. DOI: 10.1016/j.cell.2017.07.050
- Higgins, M.G., Kenny, D.A., Fitzsimons, C., Blackshields, G., Coyle, S., McKenna, C., McGee, M., Morris, D.W. & Waterset, S.M., 2019. The effect of breed and diet type on the global transcriptome of hepatic tissue in beef cattle divergent for feed efficiency. *BMC Genomics* 20, 525. <https://doi.org/10.1186/s12864-019-5906-8>
- Ingvarsen, K.L. & Boisclair, Y.R., 2001. Leptin and the regulation of food intake, energy homeostasis, and immunity with special focus on periparturient ruminants. *Domest. Anim. Endocrinol.* 21(4), 215-50. DOI: 10.1016/s0739-7240(02)00119-4
- Kelly, A.K., Waters, S.M., McGee, M., Fonseca, R.G., Carberry, C. & Kenny, D.A., 2011. mRNA expression of genes regulating oxidative phosphorylation in the muscle of beef cattle divergently ranked on residual feed intake. *Physiological Genomics* 43, 12–23. DOI: 10.1152/physiolgenomics.00213.2009.
- Kolath, W. H., M. S. Kerley, J. W. Golden, and D. H. Keister. 2006. The relationship between mitochondrial function and residual feed intake in steers. *J. Anim. Sci.* 84, 861–865. DOI: 10.2527/2006.844861x
- Lehnert, S.A., Reverter, A., Byrne, K.A. Wang, Y., Natrass, G.S., Hudson, N.J. & Greenwood, P.L., 2007. Gene expression studies of developing bovine longissimus muscle from two different beef cattle breeds. *BMC Dev. Biol.* 7, 95. <https://doi.org/10.1186/1471-213X-7-95>
- Liu, R., Tearle, R., Low, W.Y., Chen, T., Thomsen, D., Smith, T.P.L., Hiendleder, S. & Williams, J.L., 2021. Distinctive gene expression patterns and imprinting signatures revealed in reciprocal crosses between cattle sub-species. *BMC Genomics* 22(1), 410. doi: 10.1186/s12864-021-07667-2.
- Lord, M.N., Subramanian, K., Kanoski, S.E. & Noble, E.E., 2021. Melanin-concentrating hormone and food intake control: Sites of action, peptide interactions, and appetite. *Peptides* 137, 170476. DOI: 10.1016/j.peptides.2020.170476
- Makina, S.O., Whitacre, L.K., Decker, J.E. Taylor, J.F., MacNeil, M.D., Scholtz, M.M., van Marle-Köster, E., Muchadeyi, F.C., Makgahlela, M.L. & Maiwashe, A., 2016. Insight into the genetic composition of South African Sanga cattle using SNP data from cattle breeds worldwide. *Genet. Sel. Evol.* 48, 88 doi:10.1186/s12711-016-0266-1
- Mapiye, C., Chimonyo, M., Dzama, K., Raats, J.G. & Mapekula, M., 2009. Opportunities for improving Nguni cattle production in the smallholder farming systems of South Africa. *Livest. Sci.* 124, 196-204. <https://doi.org/10.1016/j.livsci.2009.01.013>

- McCarthy, D.J., Chen, Y. & Smyth, G.K., 2012. Differential expression analysis of multifactor RNA-seq experiments with respect to biological variation. *Nucleic Acids Res.* 10, 4288-4297. DOI: 10.1093/nar/gks042
- McDermaid, A., Monier, B., Zhao, J., Liu, B. & Ma, Q., 2019. Interpretation of differential gene expression results of RNA-seq data: Review and integration. *Brief Bioinform.* 20(6), 2044-2054. doi: 10.1093/bib/bby067
- Morokong, T., 2016. Sustainable options in communal beef cattle grazing systems in the Matatiele local municipality of the Eastern Cape, South Africa. MSc Thesis, Department of Agricultural Economics, Stellenbosch University.
- Muchenje, V., Dzama, K., Chimonyo, M., Raats, J.G. & Strydom, P.E., 2008. Meat quality of Nguni, Bonsmara, and Aberdeen Angus steers raised on natural pasture in the Eastern Cape, South Africa. *Meat Sci.* 79(1), 20-28. DOI: 10.1016/j.meatsci.2007.07.026
- Ncube, K.T., Dzomba, E.F., Rosen, B.D., Schroeder, S.G., Van Tassell, C.P. & Muchadeyi, F.C., 2022. Differential gene expression and identification of growth-related genes in the pituitary gland of South African goats. *Front. Genet.* 13 DOI=10.3389/fgene.2022.811193
- Owens, F.N., Gill, D.R., Secrist, D.S. & Coleman, S.W., 1995. Review of some aspects of growth and development of feedlot cattle. *J. Anim. Sci.* 73, 3152–3172. DOI: 10.2527/1995.73103152x
- Pezeshkian, Z., S.Z. Mirhoseini, S. Ghovvati, and E. Ebrahimie. 2022. Transcriptome analysis of breast muscle reveals pathways related to protein deposition in high feed efficiency of native turkeys. *Animals* 12(10): 1240. doi: 10.3390/ani12101240.
- Pintard, L., Bujnicki, J.M., Lapeyre, B. & Bonnerot, C., 2002. MRM2 encodes a novel yeast mitochondrial 21S rRNA methyltransferase. *EMBO J.* 21(5), 1139-47. DOI: 10.1093/emboj/21.5.1139
- Robinson, M.D. & Oshlack, A., 2010. A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol.* 11(3), R25. DOI: 10.1186/gb-2010-11-3-r25
- Rosen, B.D., Bickhart, D.M., Schnabel, R.D., Koren, S., Elsik, C.G., Tseng, E., Rowan, T.N., Low, W.Y., Zimin, A., Couldrey, C., Hall, R., Li, W., Rhie, A., Ghurye, J., McKay, S.D., Thibaud-Nissen, F., Hoffman, J., Murdoch, B.M., Snelling, W.M., McDanel, T.G., Hammond, J.A., Schwartz, J.C., Nandolo, W., Hagen, D.E., Dreischer, C., Schultheiss, S.J., Schroeder, S.G., Phillippy, A.M., Cole, J.B., Van Tassell, C.P., Liu, G., Smith, T.P.L. & Medrano, J.F., 2020. *De novo* assembly of the cattle reference genome with single-molecule sequencing. *GigaScience*, 9(3), g1aa021, <https://doi.org/10.1093/gigascience/g1aa021>
- Schoeman, S.J., 1996. Characterization of beef cattle breeds by virtue of their performances in the National Beef Cattle Performance and Progeny Testing Scheme. *S. Afr. J. Anim. Sci.* 26, 15-19.
- Scholtz, M.M., Gertenbach, W., & Hallowell, G., 2011a. History and origin of Nguni cattle. In: Scholtz M.M., Gertenbach W. & Hallowell G. (editors). *The Nguni breed of cattle: past, present and future*. The Nguni Cattle Breeders' Society, Pretoria, South Africa. p. 29–36.
- Scholtz, M.M., Bester, J., Mamabolo, J.M. & Ramsay, K.A., 2008. Results of the national cattle survey undertaken in South Africa, with emphasis on beef. *Appl. Anim. Husb. Rural Dev.* 1, 1-9.
- Scholtz, M.M., McManus, C., Okeyo, A.M. & Theunissen A., 2011b. Opportunities for beef production in developing countries of the southern hemisphere. *Livest. Sci.* 142,195-202. <https://doi.org/10.1016/j.livsci.2011.07.014>
- Strydom, P.E., 2008. Do indigenous Southern African cattle breeds have the right genetics for commercial production of quality meat? *Meat Sci.* 80, 86-93. DOI: 10.1016/j.meatsci.2008.04.017
- Strydom, P.E., Naude, R.T., Scholtz, M.M., & Van Wyk, J.B., 2000. Characterisation of indigenous African cattle breeds in relation to carcass characteristics. *Anim. Sci.* 70, 241-252. DOI: 10.1017/S1357729800054709
- Strydom, P.E., Naudé, R.T., Smith, M.F., Kotzé, A., Scholtz, M.M. & van Wyk, J.B., 2001. Relationships between production and product traits in subpopulations of Bonsmara and Nguni cattle. *S. Afr. J. Anim. Sci.* 31(3), 181-194. DOI: 10.4314/sajas.v31i3.3801
- Wærp, H.K.L., Waters, S.M., McCabe, M.S., Cormican, P. & Salte, R., 2019. Long-term effects of prior diets, dietary transition and pregnancy on adipose gene expression in dairy heifers. *PLoS ONE* 14(7): e0218723. <https://doi.org/10.1371/journal.pone.0218723>